1

EE5600 Assignment 1

Perabhattula Venkatesh AI20MTECH01004

Abstract—This document contains the solution to a Area of triangle problem.

Download all python codes from

https://github.com/venky-p/EE5600/Assignment 1

1 Problem

Problem Set: Vector2, Example II, Problem 5

1.1. Find the area of the triangle formed by the following points.

$$\mathbf{P} = \begin{pmatrix} a \\ c + a \end{pmatrix}, \mathbf{Q} = \begin{pmatrix} a \\ c \end{pmatrix}, \mathbf{R} = \begin{pmatrix} -a \\ c - a \end{pmatrix}$$
 (1.1.1)

2 Solution

We are going to solve this problem using vectors

$$\mathbf{P} - \mathbf{Q} = \begin{pmatrix} 0 \\ a \end{pmatrix} \tag{2.1.2}$$

$$\mathbf{P} - \mathbf{R} = \begin{pmatrix} 2a \\ 2a \end{pmatrix} \tag{2.1.3}$$

Area of given
$$\triangle le = \frac{1}{2} |\mathbf{P} - \mathbf{Q} \times \mathbf{P} - \mathbf{R}|$$
(2.1.4)

$$\mathbf{P} - \mathbf{Q} \times \mathbf{P} - \mathbf{R} = \begin{pmatrix} 0 & 0 & a \\ 0 & 0 & 0 \\ -a & 0 & 0 \end{pmatrix} \times \begin{pmatrix} 2a \\ 2a \\ 0 \end{pmatrix} \quad (2.1.5)$$

$$\implies \mathbf{P} - \mathbf{Q} \times \mathbf{P} - \mathbf{R} = \begin{pmatrix} 0 \\ 0 \\ -2a^2 \end{pmatrix} \qquad (2.1.6)$$

By substituting (2.1.6) in (2.1.4), we get

Area of given
$$\triangle le = \frac{1}{2} \begin{vmatrix} 0 \\ 0 \\ -2a^2 \end{vmatrix}$$
 (2.1.7)

$$\implies \frac{1}{2}\sqrt{0^2 + 0^2 + (-2a^2)^2} \qquad (2.1.8)$$

$$\implies \frac{1}{2}2a^2 \tag{2.1.9}$$

$$\implies a^2$$
 (2.1.10)

$$\therefore Area \ of \ given \ \triangle le = a^2 units^2$$
 (2.1.11)

3 Numerical Example

Let,

$$a = 2$$
 and $c = 4$ (3.1.12)

Solution: By substituting the given values in (1.1.1), We get

$$\mathbf{P} = \begin{pmatrix} 2 \\ 6 \end{pmatrix}, \mathbf{Q} = \begin{pmatrix} 2 \\ 4 \end{pmatrix}, \mathbf{R} = \begin{pmatrix} -2 \\ 2 \end{pmatrix}$$
 (3.1.13)




Fig. 3.1: Plot obtained from Python code Using (2.1.11),

Area of
$$\triangle le = a^2 = 2^2 = 4 \text{ units}^2$$
 (3.1.14)