


The First Experience of Transjugular Transcatheter Tricuspid Valve Replacement With LuX-Valve Plus System in Japan

Ryo Horita, MD; Daisuke Hachinohe, MD; Ryo Otake, MD;
Shah Sagar, MD; Hidemasa Shitan, MD; Kazuki Mizutani, MD, PhD



An 85-year-old woman with persistent atrial fibrillation (AF) was admitted for recurrent heart failure. Transthoracic echocardiography (TTE) demonstrated isolated massive tricuspid regurgitation (TR) due to annular dilatation with leaflet malcoaptation (**Figure A,B; Supplementary Movie 1**). Biventricular function was preserved (left ventricular ejection fraction: 62%, tricuspid annular plane systolic excursion: 20mm, fractional area change: 53%, right ventricular ejection fraction: 52%, tri-

cuspid annular plane systolic excursion/pulmonary artery systolic pressure ratio: 0.58).

Right heart catheterization revealed no significant pulmonary hypertension but an elevated right atrial (RA) v-wave consistent with massive TR. Given her advanced age, interstitial pneumonia and large coaptation gap, our heart team recommended transcatheter tricuspid valve replacement (TTVR). Preprocedural computed tomography suggested that the LuX-Valve Plus 30–45mm (Jenscare

Received January 31, 2025; accepted February 1, 2025; J-STAGE Advance Publication released online March 13, 2025 Time for primary review: 1 day

Department of Cardiology, Sapporo Heart Center, Sapporo Cardiovascular Clinic, Sapporo (R.H., D.H., R.O., S.S., H.S., K.M.), Japan; Baroda Heart Institute and Research Center, Vadodara (S.S.), India

Mailing address: Ryo Horita, MD, Department of Cardiology, Sapporo Heart Center, Sapporo Cardiovascular Clinic, 8-1, Kita 49, Higashi 16, Higashi-ku, Sapporo, Hokkaido 007-0849, Japan. email: h.ryo617@gmail.com

All rights are reserved to the Japanese Circulation Society. For permissions, please email: cr@j-circ.or.jp

ISSN-2434-0790



Scientific Co., Ningbo, China) would be anatomically suitable (**Figure C**).

Following institutional review board approval and informed consent, TTVR was performed via a transjugular approach. The LuX-Valve Plus was successfully implanted, achieving well-controlled TR with only a trivial paravalvular leak (PVL) (**Figure D,E; Supplementary Movie 2**). After implantation, the RA v-wave considerably improved (**Figure F**).

This case represents the first successful TTVR with the LuX-Valve Plus system in Japan. The device's flexible transjugular delivery system facilitates coaxial alignment within the tricuspid annulus, reducing the risk of PVL.¹ TTVR effectively controls TR and represents a potential option for high surgical risk patients. Preoperative assessment of right ventricular function and pulmonary artery pressure is associated with clinical outcomes; moreover, long-term durability remains a future concern. This case highlights the importance of patient selection through comprehensive evaluation by an experienced heart team.

Acknowledgment

We thank Dr. Su Xi for his full support and advice during the procedure.

Disclosures

The authors have nothing to disclose.

Reference

1. Wong L, Kam K, Lai L, Chow S, Lam Y, Wan S, et al. Step-by-step transcatheter tricuspid valve replacement using the LuX-Valve Plus system. *JACC Case Rep* 2024; **29**: 102699.

Supplementary Files

Supplementary Movie 1. Preprocedural transthoracic echocardiography showing massive tricuspid regurgitation.

Supplementary Movie 2. Postprocedural transthoracic echocardiography showing trivial paravalvular leak.

Please find supplementary file(s);
<https://doi.org/10.1253/circrep.CR-25-0020>