

(Continued from page 1)

"Most of the points are not new but are familiar to the American public because they were first advanced some years ago in statistical studies that admittedly are not supported by experimental evidence."

The PHS conclusions also said that "cigarette smoking is particularly associated" with the alleged increase and that persons who stopped smoking or never smoked reduce their risk of lung cancer.

These conclusions, Dr. Little's statement said, "place complete reliance on opinions published by those who are committed to the theory that tobacco has the principal causative role in lung cancer."

"The review recognizes that lung cancer afflicts non-smokers, and therefore can't all be blamed on smoking," he said. "It concedes the still-to-be evaluated role of air pollutants and occupational exposures. It admits differences in lung cancer incidence rates that cannot be explained by the smoking theory."

1. Finds that people described as the world's heaviest cigarette smokers have low lung cancer death rates compared with people who smoke less but have been long exposed to urban air pollution.

"3. Reveals that human lung tissues undergo changes, considered suspicious by some, that are found

A team of agricultural research scientists has reported on studies indicating that the two major organic insecticides used on growing tobacco plants do not get into cigarette smoke at levels considered harmful to man.

In discussing the work done under his direction, Dr. Kenneth R. Keller, director of the college's tobacco research program, was quoted by the *Associated Press* as saying: "Even with the best available chemical techniques, it has not been possible to detect even traces of endrin from the smoke of an entire carton of cigarettes."

The story quoted Dr. Keller as saying that a person would have to inhale, and retain, the smoke from a package of cigarettes a day for 250 years before he approached levels that might be considered harmful.

"Further research on chemical control has revealed that certain of the new experimental insecticides are heat degraded during flue-curing or smoking, resulting in an insecticide-free tobacco smoke," the AP said.

* "Insects in Residues on Tobacco." T. G. Harvey, W. E. Evans, F. B. Genthie and R. L. Rahn. *Journal of Agricultural and Food Chemistry*, Vol. 7, No. 10, October 1959.

among both young and old, non-smokers and smokers, while lung tissues 'may be perfectly normal in heavy smokers.'

"4. Confirms the long-established but little-publicized fact that lung cancer occurs more frequently in people who have a medical history of previous serious lung ailments, such as tuberculosis, pneumonia and influenza, indicating a relationship of possible significance."

(Note: Dr. Little's statement cited 23 scientific papers in support of these points. The full list and statement are available upon request to: The Editors, Tobacco and Health, 150 East 42nd Street, New York 17, N. Y.)

(Continued from page 1)

"Our failure during the past five years (as recorded in previous Annual Reports) to induce lung tumors in mice, rats and hamsters by exposure to strong concentrations of cigarette smoke is a striking negative result," he said.

"This has led to the careful examination of old breeding males, not subject to any smoking hazard, and six additional tumors have now been encountered in untreated animals," he said. "It is noteworthy that in each of these rats marked bronchiectasis has been an associated condition. The difficulty in the interpretation of this association is that, as in research institutes elsewhere, it is rare to make a post-mortem on any old rat without finding some more or less severe chronic bronchial lesions."

Sir Stanford Cade, head of the British cancer research effort, was quoted in news stories as saying that determining the exact relationship, if any, between smoking and lung cancer, was a "vexing problem" that required further research.

"If smoking alone causes lung cancer, we should expect to find many cases in Eskimos and Indians, who almost all smoke quite heavily, mostly cigarettes in recent decades. No bronchogenic carcinoma has been found, however, in Northern natives by our x-ray survey teams, now regularly combing the Mackenzie district and the Eastern and Western Arctic, or in the Charles Cammell Hospital and Mountain sanatorium. While lung cancer has been found in Southern Indians, it is not known to exist in Northern Indians who had not yet any occasion to inhale much air polluted by car exhausts and industrial fumes. . . ."

“Medical observations and problems in the Canadian Arctic.” Otto Schaefer, M.D., Edmonton, Alta. *Canadian Medical Association Journal*, Sept. 1, 1939.)