

THE THEORY OF HYPERGEOMETRIC STIRLING AND BELL NUMBERS

VENCISLAV POPOV

Department of Psychology, University of Zurich

ABSTRACT. We introduce a natural generalization of the Bell numbers: the m -Bell numbers $B_n^{(m)}$, characterized by invariance under m binomial transforms followed by an m -place left shift. Their exponential generating functions satisfy m -th order ordinary differential equations whose solutions are hypergeometric (hyper-Bessel) functions, specializing to the exponential function when $m = 1$ (classical Bell numbers) and to modified Bessel functions when $m = 2$ (yielding “Bessel–Bell” numbers). Mirroring the Bell–Stirling correspondence, we construct m -Stirling triangular arrays from the two-term recurrence $\llbracket \begin{smallmatrix} n+1 \\ k \end{smallmatrix} \rrbracket_m = m \lfloor k/m \rfloor \llbracket \begin{smallmatrix} n \\ k \end{smallmatrix} \rrbracket_m + \llbracket \begin{smallmatrix} n \\ k-1 \end{smallmatrix} \rrbracket_m$ and prove an elementary shift identity from which the central structure theorem follows: the row sums of the m -Stirling triangle reproduce $B_n^{(m)}$, and, more finely, the residue-class row sums are precisely the m primitive m -Bell sequences. The m -Stirling numbers come in dual pairs (with first-kind partners, generalized falling factorials, and Lah-type companions), serve as conversion operators between polynomial bases, admit Dobiński-like formulas, and count parity-constrained partitions in an urn model as well as restricted permutation insertion histories. Finally, we show that the m -Bell numbers govern the moments of the Conway–Maxwell–Poisson distribution with integer dispersion parameter $\nu = m$, exactly as the Bell numbers govern the moments of the Poisson distribution.

1. INTRODUCTION

Given a sequence of numbers $(a_n)_{n \geq 0} = (a_0, a_1, \dots)$, the binomial transform maps it to a new sequence $(b_n)_{n \geq 0} = (b_0, b_1, \dots)$ as follows:

$$b_n = \sum_{k=0}^n \binom{n}{k} a_k.$$

Repeated applications of the binomial transform on the resulting sequence can be represented with a single sum (see [20]):

$$b_n = \sum_{k_1=0}^n \binom{n}{k_1} \sum_{k_2=0}^{k_1} \binom{k_1}{k_2} \cdots \sum_{k_m=0}^{k_{m-1}} \binom{k_{m-1}}{k_m} a_{k_m} = \sum_{k=0}^n \binom{n}{k} m^{n-k} a_k,$$

where m is an integer that represents the number of times the binomial transform has been applied. Bernstein and Sloane [2] studied a number of what they call “eigen-sequences” of various such transformations—sequences (a_n) which, when transformed one or more times, shift by one or more places but are otherwise preserved. Such sequences show a “self-similarity” under an iterated transform, and understanding why this self-similarity occurs often reveals new properties or relations between different integer sequences and the combinatorial structures they enumerate.

Perhaps the most famous binomial-transform invariant sequence is that of the Bell numbers, entry A000110 in the Online Encyclopedia of Integer Sequences (OEIS): 1, 1, 2, 5, 15, 52, 203, 877, 4140, ... Applying the binomial transform to the Bell numbers

E-mail address: vencislav.popov@gmail.com.

2020 *Mathematics Subject Classification.* Primary 05A18, 11B73; Secondary 05A15, 33C10, 60E05.

Key words and phrases. Bell numbers, Stirling numbers, set partitions, binomial transform, Conway–Maxwell–Poisson distribution, modified Bessel functions, hypergeometric functions.

Computational results and other supporting information is available on GitHub and Wolfram Cloud.

produces an identical sequence, with the first element omitted and each subsequent element's index shifted once to the left:

$$(1) \quad B_{n+1} = \sum_{k=0}^n \binom{n}{k} B_k.$$

The Bell numbers count the total number of partitions of an n -element set, and they are part of a rich combinatorial structure that involves the Stirling numbers of the second kind (A008277), Touchard polynomials [22], the exponential function, and linear operators acting on it [8].

To simplify the notation, following Bernstein and Sloane [2], define the following operators on sequences:

$$\text{BINOM} \circ [a_0, a_1, a_2, \dots] = \left[\binom{0}{0} a_0, \binom{1}{0} a_0 + \binom{1}{1} a_1, \dots, \sum_{k=0}^n \binom{n}{k} a_k, \dots \right]$$

$$L \circ [a_0, a_1, a_2, \dots] = [a_1, a_2, \dots]$$

where BINOM is the binomial transform operator applied to sequence a , whereas L is the left-shift operator. The Bell numbers then form the unique sequence that satisfies the following equality with initial condition $a_0 = 1$:

$$\text{BINOM} \circ a = L \circ a.$$

Consider a new generalization of the Bell numbers defined by the following property:

$$(2) \quad B_{n+m}^{(m)} = \sum_{k=0}^n \binom{n}{k} m^{n-k} B_k^{(m)},$$

where the upper index (m) has no algebraic meaning and should be read merely as an “ m -Bell” number. Equivalently, these sequences satisfy the operator equation

$$(3) \quad \text{BINOM}^m \circ a = L^m \circ a.$$

These are sequences that shift to the left by m places after m applications of the binomial transform. Any such sequence is determined by its first m values $(a_0, \dots, a_{m-1})$, so the solution set of (3) is an m -dimensional lattice of integer sequences spanned by the m **primitive** solutions $B^{(m,r)}$, $r = 0, \dots, m-1$, whose initial conditions are $B_n^{(m,r)} = \delta_{n,r}$ for $0 \leq n < m$. We reserve the unadorned symbol $B^{(m)} = B^{(m,0)} + \dots + B^{(m,m-1)}$ for the **composite** sequence with initial conditions $(1, 1, \dots, 1)$. The case $m = 1$ recovers the Bell numbers, and $m = 2$ corresponds to sequences A007472, A351143 and A351028, which shift by 2 places left after 2 binomial transforms (the sequences for $m > 2$ are not currently present in the OEIS). Although the sequences for $m = 2$ are listed in the OEIS, little was previously known about their properties.

In this paper we show that the m -fold binomial-shift-invariance property that characterizes these sequences arises from a combinatorial structure that closely mirrors that of the classical Bell numbers. The m -Bell numbers have exponential generating functions (e.g.f.s) that are the solutions of ordinary differential equations of order m . The solutions to these equations are a class of hypergeometric (“hyper-Bessel”) functions, which reduce to the exponential function for $m = 1$ and to modified Bessel functions of the first and second kind for $m = 2$, motivating the name Bessel–Bell numbers for the 2-Bell case (Section 3; Theorem 2). These generalized Bell numbers admit Dobiński-like formulas (Theorem 3 and Theorem 30). Each m -Bell sequence arises from a Stirling-like triangular array generated by the two-term recurrence

$$\left[\begin{matrix} n+1 \\ k \end{matrix} \right]_m = m \lfloor k/m \rfloor \left[\begin{matrix} n \\ k \end{matrix} \right]_m + \left[\begin{matrix} n \\ k-1 \end{matrix} \right]_m,$$

motivating the name *m -Stirling numbers* (Sections 4 and 7). Our central structural result is an elementary shift identity for these triangles (Theorem 25), from which we deduce that the row sums of the m -Stirling triangle form the composite m -Bell sequence and—more finely—that the row sums along residue classes of $k \bmod m$ are precisely the m primitive m -Bell sequences (Corollary 26). For $m = 2$ this identifies A351143 and A351028 as the even- k and odd- k halves of a single triangle whose full row sums give A007472.

Like the classical Stirling numbers, the m -Stirling numbers come in dual pairs: there are companion triangles of the first kind, together with generalized falling factorials and Lah-type arrays, and the whole family acts as a system of conversion operators between polynomial bases (Section 6). Each m -Stirling array of the second kind also arises from the coefficients of polynomials generated by applying the exponential scaling operator to the hypergeometric e.g.f.s, once these are paired with appropriate *carrier functions* (Sections 4 and 7). On the combinatorial side, the m -Stirling and m -Bell numbers count parity-constrained partitions in an urn model with m -compartment urns (Section 5), while the first-kind numbers count permutations built by a parity-restricted insertion process (Section 6). Finally, just as the classical Bell numbers are the moments of the Poisson distribution with unit rate, the m -Bell numbers turn out to govern the moments of a well-known generalization of the Poisson distribution, the Conway–Maxwell–Poisson distribution [18], at integer dispersion parameter $\nu = m$ (Section 8). Appendix D collects the parallels between the classical framework and its generalization in a single reference table.

To clearly ground the analogy between the Bell–Stirling–Touchard framework and the novel results, we begin with a review of standard results and notation [5, 17].

2. BACKGROUND

The exponential generating function (e.g.f.) $\mathcal{A}(x)$ of a sequence $(a_n)_{n \geq 0}$ is a formal power series in x :

$$\mathcal{A}(x) = \sum_{n=0}^{\infty} a_n \frac{x^n}{n!}.$$

Proposition 1 (Functional equation for the e.g.f. of m -Bell numbers). *Let $\mathcal{A}(x)$ be the exponential generating function of a sequence $a = (a_n)$ satisfying (3). Then $\mathcal{A}(x)$ satisfies the m -th order ODE*

$$\left(\frac{d}{dx} \right)^m \mathcal{A}(x) - e^{mx} \mathcal{A}(x) = 0.$$

Proof. It is well known that both the binomial transform and the left-shift operator acting on a have simple effects on the e.g.f., $\mathcal{B}(x)$, of the resulting sequence b (e.g. [2]):

$$(4) \quad \begin{aligned} b = \text{BINOM} \circ a &\iff \mathcal{B}(x) = e^x \mathcal{A}(x) \\ b = \text{L} \circ a &\iff \mathcal{B}(x) = \mathcal{A}'(x) \end{aligned}$$

Applying BINOM m times corresponds to multiplying $\mathcal{A}(x)$ by e^{mx} . Applying L m times corresponds to taking the m -th derivative. Therefore, if $\text{BINOM}^m \circ a = \text{L}^m \circ a$, then

$$\mathcal{A}^{(m)}(x) = e^{mx} \mathcal{A}(x).$$

Rewriting this gives the result. □

For $m = 1$ the ODE $\mathcal{A}'(x) = e^x \mathcal{A}(x)$ with $\mathcal{A}(0) = 1$ integrates at once to the celebrated e.g.f. of the Bell numbers:

$$(5) \quad e^{e^x - 1} = \sum_{n=0}^{\infty} B_n \frac{x^n}{n!}.$$

Recall that the Bell numbers are the row sums of the triangular array formed by the Stirling numbers of the second kind:

$$B_n = \sum_{k=0}^n \left\{ \begin{matrix} n \\ k \end{matrix} \right\},$$

where the Stirling numbers of the second kind satisfy the two-term recurrence

$$\left\{ \begin{matrix} n+1 \\ k \end{matrix} \right\} = k \left\{ \begin{matrix} n \\ k \end{matrix} \right\} + \left\{ \begin{matrix} n \\ k-1 \end{matrix} \right\}, \quad n, k \geq 1.$$

A second classical recurrence, which will serve as the model for our main structural result (Theorem 25), expresses a downward shift of the whole triangle through binomial averaging [5, 12]:

$$(6) \quad \left\{ \begin{matrix} n+1 \\ k \end{matrix} \right\} = \sum_{j=0}^n \binom{n}{j} \left\{ \begin{matrix} j \\ k-1 \end{matrix} \right\}.$$

Summing (6) over k recovers the Bell recurrence (1); combinatorially, the binomial coefficient chooses which elements avoid the block containing the largest element.

The Stirling numbers of the second kind count the number of ways to partition n labeled objects into k unlabeled non-empty subsets and are the coefficients of the Touchard (also known as Bell or exponential) polynomials:

$$(7) \quad T_n(x) = \sum_{k=0}^n \left\{ \begin{matrix} n \\ k \end{matrix} \right\} x^k,$$

whose e.g.f. is a bi-variate generalization of the Bell numbers' e.g.f.:

$$(8) \quad \sum_{n=0}^{\infty} T_n(x) \frac{t^n}{n!} = e^{x(e^t-1)}.$$

Many useful identities involving Touchard polynomials and Stirling numbers can be shown via the action of the exponential scaling operator

$$(9) \quad e^{txD_x} f(x) = f(xe^t),$$

where D_x is the derivative operator with respect to x . Specifically, by applying this operator to the standard exponential function e^x we get precisely the e.g.f. for the Touchard polynomials [8]:

$$e^{-x} e^{txD_x} e^x = e^{x(e^t-1)}.$$

A final useful relation is the celebrated Dobiński formula [24] that lets us express $T_n(x)$ and the Bell numbers $B_n = T_n(1)$ as an infinite sum:

$$(10) \quad \begin{aligned} T_n(x) &= e^{-x} \sum_{k=0}^{\infty} \frac{x^k k^n}{k!} \\ B_n &= \frac{1}{e} \sum_{k=0}^{\infty} \frac{k^n}{k!} \end{aligned}$$

In the remainder of the paper we will see that the m -Bell sequences have properties analogous to all of those defined above, starting with the special case $m = 2$.

3. THE BESSEL-BELL NUMBERS ($m = 2$)

3.1. Deriving the e.g.f. Before solving the general case, let us focus on the $m = 2$ generalization of the Bell numbers, which satisfy the property that they shift by 2 places left after two binomial transforms:

$$(11) \quad B_{n+2}^{(2)} = \sum_{k=0}^n \binom{n}{k} 2^{n-k} B_k^{(2)}.$$

The following three sequences listed in the OEIS satisfy this property:

- A007472: 1, 1, 1, 3, 9, 29, 105, 431, 1969, ...
- A351143: 1, 0, 1, 2, 5, 16, 61, 258, 1177, ...
- A351028: 0, 1, 0, 1, 4, 13, 44, 173, 792, 4009, ...

The only difference between the three sequences is the pair of initial conditions (a_0, a_1) : $(1, 1)$, $(1, 0)$ and $(0, 1)$, respectively. In the terminology of Section 1, the primitive solutions for $m = 2$ are $A351143 = B^{(2,0)}$ and $A351028 = B^{(2,1)}$, whose element-wise sum produces the composite sequence $A007472 = B^{(2)}$. Owing to the form of their e.g.f.s, we will refer to the sequences satisfying (11) as Bessel-Bell numbers (not to be confused with the Bessel numbers of [4], which arise from a different structure).

Theorem 2. *The exponential generating functions of the 2-Bell sequences are solutions of the modified Bessel ODE composed with e^x : each is a linear combination*

$$\mathcal{A}(x) = p I_0(e^x) + q K_0(e^x)$$

of modified Bessel functions of the first (I_0) and second kind (K_0) of order 0, whose weights p, q are uniquely determined by the first two elements of the corresponding sequence. Specifically:

Sequence	Coefficient p	Coefficient q
A351143	$K_1(1) \approx 0.601907$	$I_1(1) \approx 0.565159$
A351028	$K_0(1) \approx 0.421024$	$-I_0(1) \approx -1.266066$
A007472	$K_0(1) + K_1(1) \approx 1.022932$	$I_1(1) - I_0(1) \approx -0.700907$

Proof. By Proposition 1, the e.g.f. $\mathcal{A}(x)$ satisfies the linear ODE

$$\mathcal{A}''(x) - e^{2x} \mathcal{A}(x) = 0.$$

Set $t = e^x$ (so $dt/dx = t$) and define $f(t) = \mathcal{A}(x)$. By the chain rule,

$$\mathcal{A}'(x) = t f'(t),$$

$$\mathcal{A}''(x) = t f'(t) + t^2 f''(t),$$

$$e^{2x} \mathcal{A}(x) = t^2 f(t).$$

Substituting in yields the standard form of the modified Bessel equation of order 0 (see 10.25.1) [10]

$$(12) \quad t^2 f'' + t f' - t^2 f = 0,$$

whose general solution is

$$f(t) = p I_0(t) + q K_0(t),$$

with I_ν and K_ν being the modified Bessel functions of the first and second kind, and p and q constants determined by the initial conditions. Therefore, we obtain the general solution for $\mathcal{A}(x)$, the e.g.f. of sequences A007472, A351143 and A351028, by substituting back $t = e^x$:

$$(13) \quad \mathcal{A}(x) = p I_0(e^x) + q K_0(e^x).$$

To determine p and q we use the known initial conditions. Sequence A007472 is the element-wise sum of A351143 and A351028, so it is sufficient to determine p and q for the latter two sequences only. For A351143 we have $a_0 = 1$ and $a_1 = 0$. Matching the first two Maclaurin coefficients of $\mathcal{A}(x)$ yields the following system of equations:

$$\mathcal{A}(0) = p I_0(1) + q K_0(1) = 1$$

$$\mathcal{A}'(x)|_{x=0} = p \frac{d}{dx} I_0(e^x)|_{x=0} + q \frac{d}{dx} K_0(e^x)|_{x=0} = 0,$$

where $f(x)|_{x=0}$ denotes evaluation at 0. The derivatives of I_0 and K_0 are thankfully straightforward (10.29.3) [10]:

$$(14) \quad \begin{aligned} I_0'(z) &= I_1(z) \\ K_0'(z) &= -K_1(z). \end{aligned}$$

By the chain rule we get

$$\frac{d}{dx} I_0(e^x) = e^x I_1(e^x)$$

$$\frac{d}{dx} K_0(e^x) = -e^x K_1(e^x).$$

Evaluating at $x = 0$ gives

$$\mathcal{A}(0) = p I_0(1) + q K_0(1) = 1$$

$$\mathcal{A}'(0) = p I_1(1) - q K_1(1) = 0.$$

After some standard algebraic torture we get the following expressions for p and q :

$$\begin{aligned} p &= \frac{1 - q K_0(1)}{I_0(1)} \\ q &= \frac{I_1(1)}{I_1(1) K_0(1) + I_0(1) K_1(1)} \end{aligned}$$

These expressions can be simplified further due to the following Bessel identity concerning the Wronskian of the modified Bessel functions (see 10.28.2):

$$(15) \quad I_\nu(z)K_{\nu+1}(z) + I_{\nu+1}(z)K_\nu(z) = 1/z,$$

which holds for any complex ν and z . In the special case when $\nu = 0$ and $z = 1$:

$$(16) \quad I_0(1)K_1(1) + I_1(1)K_0(1) = 1.$$

Therefore the constants for the e.g.f. of sequence A351143 are

$$q = I_1(1)$$

$$p = \frac{1 - I_1(1)K_0(1)}{I_0(1)} = \frac{I_0(1)K_1(1) + I_1(1)K_0(1) - I_1(1)K_0(1)}{I_0(1)} = K_1(1).$$

A very similar manipulation for the initial conditions $(0, 1)$ of A351028 gives $p = K_0(1)$ and $q = -I_0(1)$, and adding the two solutions gives the coefficients for A007472 stated in the table, which concludes the proof. $\square$

3.2. Why do the e.g.f.s of the 2-Bell numbers produce integer coefficients?

The e.g.f.s of the 2-Bell numbers are admittedly more complicated than the e.g.f. e^{e^x-1} of the Bell numbers (5) (although the form is quite similar, as we discuss at the end of this section). How is it that such a relatively complicated expression, involving four different special functions, produces integer coefficients? Let us explore the series expansion of the derived e.g.f.s. Consider first the case of sequence A351143:

$$\mathcal{A}_{351143}(x) = K_1(1)I_0(e^x) + I_1(1)K_0(e^x).$$

We have the following standard series for I_0 and K_0 (see 10.25.2 and 10.31.2) [10]:

$$(17) \quad I_0(z) = \sum_{k=0}^{\infty} \frac{(\frac{1}{2}z)^{2k}}{k!k!}$$

$$K_0(z) = -\log\left(\frac{z}{2}\right)I_0(z) + \sum_{k=0}^{\infty} \frac{\psi(k+1)(\frac{1}{2}z)^{2k}}{k!k!}$$

where ψ is the digamma function. Let us focus on the I_0 function. Substituting $z = e^x$ and then expanding the Taylor series for the exponential function we get

$$(18) \quad I_0(e^x) = \sum_{k=0}^{\infty} \frac{e^{2xk}}{2^{2k}k!k!} = \sum_{k=0}^{\infty} \frac{1}{2^{2k}k!k!} \sum_{n=0}^{\infty} \frac{x^n(2k)^n}{n!} = \sum_{n=0}^{\infty} \frac{x^n}{n!} \sum_{k=0}^{\infty} \frac{(2k)^n}{2^{2k}k!k!}.$$

(Interchanging the order of summation is justified by absolute convergence.) Let the inner sum be represented by $S(n) = \sum_{k=0}^{\infty} \frac{(2k)^n}{2^{2k}k!k!}$. This formula is reminiscent of the Dobiński formula for the standard Bell numbers (10), with an extra factorial in the denominator and extra powers of 2. Indeed, we can state an equivalent theorem:

Theorem 3 (Dobiński-like formula for the Bessel–Bell numbers). *Let $v_n = B_n^{(2,0)}$ and $u_n = B_n^{(2,1)}$ denote the primitive 2-Bell sequences A351143 and A351028. Then*

$$I_0(e^x) = \sum_{n=0}^{\infty} S(n) \frac{x^n}{n!}, \quad \text{where}$$

$$S(n) = \sum_{k=0}^{\infty} \frac{(2k)^n}{(2^k k!)^2} = v_n I_0(1) + u_n I_1(1).$$

Proof. Proceed by strong induction. First, establish the base cases:

$$S(0) = \sum_{k=0}^{\infty} \frac{1}{2^{2k}k!k!} = I_0(1)$$

$$S(1) = \sum_{k=0}^{\infty} \frac{2k}{2^{2k}k!k!} = \sum_{k=1}^{\infty} \frac{1}{2^{2k-1}(k-1)!k!} = \sum_{k=0}^{\infty} \frac{1}{2^{2k+1}k!(k+1)!} = I_1(1),$$

n	$S(n)$	v_n	u_n	$v_n + u_n$
0	$I_0(1)$	1	0	1
1	$I_1(1)$	0	1	1
2	$I_0(1)$	1	0	1
3	$2I_0(1) + I_1(1)$	2	1	3
4	$5I_0(1) + 4I_1(1)$	5	4	9
5	$16I_0(1) + 13I_1(1)$	16	13	29
6	$61I_0(1) + 44I_1(1)$	61	44	105

TABLE 1. Coefficients in the expansion of the Dobiński-like formula for the Bessel–Bell numbers (Theorem 3)

where in the case of $S(1)$ we used the series expansion of the modified Bessel function of general integer order $\nu \geq 0$ evaluated at 1 (10.25.2) [10]:

$$I_\nu(z) = \sum_{k=0}^{\infty} \frac{(z/2)^{2k+\nu}}{k! (k+\nu)!}.$$

These match the claim since $(v_0, u_0) = (1, 0)$ and $(v_1, u_1) = (0, 1)$. For the inductive step, suppose that $S(j) = v_j I_0(1) + u_j I_1(1)$ holds for every $j \leq n+1$. Then:

$$\begin{aligned}
 (19) \quad S(n+2) &= \sum_{k=0}^{\infty} \frac{(2k)^{n+2}}{2^{2k} k! k!} = \sum_{k=1}^{\infty} \frac{(2k)^n}{2^{2k-2} (k-1)! (k-1)!} = \sum_{k=0}^{\infty} \frac{(2k+2)^n}{2^{2k} k! k!} \\
 &= \sum_{k=0}^{\infty} \sum_{j=0}^n \binom{n}{j} 2^{n-j} \frac{(2k)^j}{2^{2k} k! k!} = \sum_{j=0}^n \binom{n}{j} 2^{n-j} \sum_{k=0}^{\infty} \frac{(2k)^j}{2^{2k} k! k!} \\
 &= \sum_{j=0}^n \binom{n}{j} 2^{n-j} S(j).
 \end{aligned}$$

First, notice that (19) has exactly the same form as the recurrence relation (11) that defines the 2-Bell numbers. Second, use the induction hypothesis and substitute $S(j)$:

$$\begin{aligned}
 S(n+2) &= \sum_{j=0}^n \binom{n}{j} 2^{n-j} (v_j I_0(1) + u_j I_1(1)) \\
 &= I_0(1) \left[\sum_{j=0}^n \binom{n}{j} 2^{n-j} v_j \right] + I_1(1) \left[\sum_{j=0}^n \binom{n}{j} 2^{n-j} u_j \right] \\
 &= v_{n+2} I_0(1) + u_{n+2} I_1(1),
 \end{aligned}$$

where the last step applies the defining recurrence (11) of the primitive sequences v and u . This completes the proof. Table 1 illustrates the first few cases. $\square$

The major difference between the Dobiński-like formula of Theorem 3 and the standard Dobiński formula (10) is that there are two different modified Bessel functions acting as **carriers** for the coefficients of the two primitive Bessel–Bell sequences, rather than a single exponential function. In the standard Dobiński formula one multiplies the infinite sum by $1/e$ to cancel the exponential carrier; here no single multiplicative factor can cancel two independent carriers at once. The concept of a carrier function will play a fundamental role in understanding how m -Bell and m -Stirling numbers are generated, and we formalize it in Section 4 (Definition 7).

What about the second Bessel function $K_0(e^x)$ appearing in the e.g.f.s? Its Maclaurin coefficients turn out to involve exactly the same integer sequences, with one sign change.

Theorem 4. *With v_n and u_n as in Theorem 3,*

$$K_0(e^x) = \sum_{n=0}^{\infty} \frac{x^n}{n!} (v_n K_0(1) - u_n K_1(1)).$$

Proof. Consider the two pairs of functions

$$F_0(x) = I_0(e^x), \quad F_1(x) = I_1(e^x), \quad G_0(x) = K_0(e^x), \quad G_1(x) = -K_1(e^x).$$

From the derivative formulas $I'_0 = I_1$, $I'_1(z) = I_0(z) - z^{-1}I_1(z)$, $K'_0 = -K_1$ and $K'_1(z) = -K_0(z) - z^{-1}K_1(z)$ (10.29.2) [10], the chain rule gives

$$\begin{aligned} F'_0 &= e^x F_1, & F'_1 &= e^x F_0 - F_1, \\ G'_0 &= e^x G_1, & G'_1 &= e^x G_0 - G_1. \end{aligned}$$

That is, (F_0, F_1) and (G_0, G_1) satisfy the *same* first-order linear system

$$(20) \quad X' = e^x Y, \quad Y' = e^x X - Y.$$

Define coefficient functions $a_n, b_n \in \mathbb{Z}[e^x]$ by the recursion

$$a_{n+1} = a'_n + e^x b_n, \quad b_{n+1} = e^x a_n + b'_n - b_n, \quad a_0 = 1, \quad b_0 = 0.$$

A one-line induction shows that *every* solution pair (X, Y) of (20) satisfies

$$(21) \quad X^{(n)} = a_n(x) X + b_n(x) Y \quad (n \geq 0),$$

since differentiating (21) and substituting (20) reproduces the recursion. We claim that $a_n(0) = v_n$ and $b_n(0) = u_n$. To see this, first note that any solution of (20) has first component satisfying the 2-Bell ODE: $X'' = (e^x Y)' = e^x Y + e^x Y' = X' + e^{2x} X - e^x Y = e^{2x} X$, with $X'(0) = Y(0)$. Comparing Maclaurin coefficients on both sides of $X'' = e^{2x} X$ shows that the sequence $(X^{(n)}(0))_n$ satisfies the 2-Bell recurrence (11). Now take the solution pair with initial data $(X(0), Y(0)) = (1, 0)$: then $X(0) = 1$, $X'(0) = 0$, so $(X^{(n)}(0))_n$ is precisely the primitive sequence A351143, i.e. $X^{(n)}(0) = v_n$; but (21) evaluated at 0 gives $X^{(n)}(0) = a_n(0)$. Hence $a_n(0) = v_n$, and the solution pair with initial data $(0, 1)$ gives $b_n(0) = u_n$ in the same way. Applying (21) to the pair (G_0, G_1) and evaluating at $x = 0$ yields

$$G_0^{(n)}(0) = a_n(0) K_0(1) + b_n(0) (-K_1(1)) = v_n K_0(1) - u_n K_1(1). \quad \square$$

Remark 5. *The same evaluation applied to the pair (F_0, F_1) re-derives Theorem 3, since $F_0^{(n)}(0) = S(n)$ by (18). The two proofs are complementary: the induction of Theorem 3 works directly with the series, while the present argument isolates the structural reason—I- and K-type functions solve the same differential system—why the same integer sequences appear in both expansions.*

Finally, by combining Theorem 3 and Theorem 4, we have

$$(22) \quad p I_0(e^x) + q K_0(e^x) = \sum_{n=0}^{\infty} \frac{x^n}{n!} \left[v_n (p I_0(1) + q K_0(1)) + u_n (p I_1(1) - q K_1(1)) \right].$$

Within this equation, an appropriate choice of p and q neutralizes the modified Bessel functions in the series thanks to the Wronskian identity (16). Specifically, $p = K_1(1)$, $q = I_1(1)$ makes the coefficient of v_n equal to 1 and the coefficient of u_n equal to 0, giving integer coefficients—the series A351143. Similarly for the other two choices of p and q established in Theorem 2: $(K_0(1), -I_0(1))$ isolates u_n , and the sum of the two choices produces $v_n + u_n$.

Thus the e.g.f. of a 2-Bell sequence is structurally similar to that of the 1-Bell (Bell) sequence: it is the composition of an exponential-like function with the exponential function, and the remaining constant factors are there just to ensure integer coefficients, exactly like the factor e^{-1} in $e^{-1}e^{e^x}$. With the integer-coefficient property now rigorously settled, let us shift to the combinatorial side: a triangular recurrence whose row sums reproduce precisely the Bessel–Bell sequences.

4. BESSEL–STIRLING NUMBERS OF THE SECOND KIND

The similarity of the Bessel–Bell numbers' e.g.f. to the standard Bell numbers' e.g.f. immediately invites the question of whether there is also a Bessel–Stirling equivalent of the triangular array of the Stirling numbers of the second kind, whose rows sum to the Bell numbers. There is, and the correct recurrence modifies the classical one in a single place.

	k=0	k=1	k=2	k=3	k=4	k=5	k=6	k=7	k=8	k=9	$\sum_{k=0}^n$
n=0	1										1
n=1	0	1									1
n=2	0	0	1								1
n=3	0	0	2	1							3
n=4	0	0	4	4	1						9
n=5	0	0	8	12	8	1					29
n=6	0	0	16	32	44	12	1				105
n=7	0	0	32	80	208	92	18	1			431
n=8	0	0	64	192	912	576	200	24	1		1969
n=9	0	0	128	448	3840	3216	1776	344	32	1	9785

TABLE 2. The Bessel–Stirling numbers of the second kind, $\llbracket n \rrbracket_k$ (OEIS A383235). The values in the upper triangle are all 0.

Definition 6 (Bessel–Stirling numbers of the second kind). *Let $\llbracket n \rrbracket_k$ be defined for integers $n, k \geq 0$ by the initial conditions $\llbracket 0 \rrbracket_k = \delta_{0,k}$, the convention $\llbracket n \rrbracket_k = 0$ for $k < 0$, and the two-term recurrence*

$$(23) \quad \llbracket n+1 \rrbracket_k = 2 \lfloor k/2 \rfloor \llbracket n \rrbracket_k + \llbracket n \rrbracket_{k-1}.$$

The notation follows the bracket style for binomial coefficients $\binom{n}{k}$ and Stirling numbers $\left\{ \begin{smallmatrix} n \\ k \end{smallmatrix} \right\}$ popularized by [12], with doubled delimiters and a subscript indicating the modulus (here 2); Section 7 treats general modulus m .

This recurrence is almost identical to that of the Stirling numbers of the second kind, except that the coefficient in front of $\llbracket n \rrbracket_k$ has parity—it is always even, with k rounded down to the nearest even integer. The numbers $\llbracket n \rrbracket_k$ form a triangular array whose first few rows are listed in Table 2; the triangle is available as OEIS entry A383235, which was submitted in the course of this work. An immediate consequence of the definition is that $\llbracket n \rrbracket_k = 0$ for $k > n$ (each application of (23) raises the maximal index k by one), that $\llbracket n \rrbracket_n = 1$, and that $\llbracket n \rrbracket_1 = \delta_{n,1}$, since the coefficient $2 \lfloor k/2 \rfloor$ vanishes for $k \in \{0, 1\}$.

The row sums of this array visibly reproduce the Bessel–Bell numbers A007472:

$$\sum_{k=0}^n \llbracket n \rrbracket_k = (1, 1, 1, 3, 9, 29, 105, 431, 1969, 9785, 52145, 296155, 1787385, \dots).$$

In this section we prove this and more via the operational calculus of *carrier systems*; Section 7 gives an independent, purely combinatorial proof valid for every m .

4.1. Carrier systems and Bessel–Touchard polynomials. Recall from (9) that Touchard polynomials arise by applying the exponential scaling operator e^{txD} to the carrier e^x , whose salient property is that the operator xD maps the module $\{P(x)e^x : P \text{ polynomial}\}$ to itself. Modified Bessel functions are not preserved individually by xD , but the *pair* (I_0, I_1) is, in the following precise sense. From the derivative relations (10.29.2) [10]:

$$(24) \quad \begin{aligned} (xD)I_0(x) &= xI_1(x) \\ (xD)I_1(x) &= xI_0(x) - I_1(x). \end{aligned}$$

We abstract exactly this structure:

Definition 7 (Carrier system). *Let $m \geq 1$. A carrier system of modulus m for the operator xD is a family of functions (or formal power series) $(f_0, f_1, \dots, f_{m-1})$ satisfying the cyclic first-order system*

$$(25) \quad (xD + r)f_r = x f_{(r+1) \bmod m}, \quad r = 0, 1, \dots, m-1.$$

The function f_0 is called the primary carrier of the system.

For $m = 1$ the single condition reads $(xD)f_0 = xf_0$, i.e. $f'_0 = f_0$, whose solution is the exponential carrier e^x of the classical theory. For $m = 2$, equations (24) say precisely that (I_0, I_1) is a carrier system: $(xD + 0)I_0 = xI_1$ and $(xD + 1)I_1 = xI_0$. The computation in the proof of Theorem 4 shows that $(K_0, -K_1)$ is a second, independent carrier system of modulus 2. Chaining the m conditions of (25) shows that a primary carrier always satisfies the eigenvalue-like equation

$$(26) \quad (xD)^m f_0 = x^m f_0,$$

which for $m = 2$ is the modified Bessel equation (12)—the same equation, we emphasize, that governs the e.g.f.s of the 2-Bell numbers under the substitution $t = e^x$.

The next theorem shows that iterating xD on a primary carrier manufactures the Bessel–Stirling triangle, in exactly the way that iterating xD on e^x manufactures the Stirling triangle. We state it for modulus 2; the statement and proof for general m are identical (Theorem 29).

Theorem 8 (Carrier expansion). *Let (f_0, f_1) be a carrier system of modulus 2 for xD . Then for all $n \geq 0$,*

$$(xD)^n f_0(x) = \sum_{k=0}^n \left[\begin{matrix} n \\ k \end{matrix} \right]_2 x^k f_{k \bmod 2}(x),$$

and consequently, by the scaling operator identity (9),

$$f_0(xe^t) = e^{txD} f_0(x) = \sum_{n=0}^{\infty} \frac{t^n}{n!} V_n(x), \quad V_n(x) = \sum_{k=0}^n \left[\begin{matrix} n \\ k \end{matrix} \right]_2 x^k f_{k \bmod 2}(x).$$

Proof. Induction on n . The case $n = 0$ is the initial condition $\left[\begin{matrix} 0 \\ k \end{matrix} \right]_2 = \delta_{0,k}$. For the step, apply xD to a single term $x^k f_r$ with $r = k \bmod 2$. Since xD is a derivation and $(xD)x^k = kx^k$,

$$\begin{aligned} (xD)(x^k f_r) &= kx^k f_r + x^k (xD)f_r = kx^k f_r + x^k (xf_{(r+1) \bmod 2} - rf_r) \\ &= (k - r)x^k f_r + x^{k+1} f_{(k+1) \bmod 2}. \end{aligned}$$

Because $r = k \bmod 2$, the factor $k - r$ equals $2 \lfloor k/2 \rfloor$. Summing over k with weights $\left[\begin{matrix} n \\ k \end{matrix} \right]_2$ and collecting the coefficient of $x^k f_{k \bmod 2}$ yields exactly the recurrence (23), i.e. the coefficient of $x^k f_{k \bmod 2}$ in $(xD)^{n+1} f_0$ is $2 \lfloor k/2 \rfloor \left[\begin{matrix} n \\ k \end{matrix} \right]_2 + \left[\begin{matrix} n \\ k-1 \end{matrix} \right]_2 = \left[\begin{matrix} n+1 \\ k \end{matrix} \right]_2$. $\square$

Taking $(f_0, f_1) = (I_0, I_1)$, the first few polynomials V_n are:

$$\begin{aligned} V_0(x) &= I_0(x) \\ V_1(x) &= x I_1(x) \\ V_2(x) &= x^2 I_0(x) \\ V_3(x) &= 2x^2 I_0(x) + x^3 I_1(x) \\ V_4(x) &= 4x^2 I_0(x) + 4x^3 I_1(x) + x^4 I_0(x) \\ V_5(x) &= 8x^2 I_0(x) + 12x^3 I_1(x) + 8x^4 I_0(x) + x^5 I_1(x) \end{aligned}$$

Notice two things. First, these polynomials are generated in very much the same way as the Touchard polynomials, but by applying the exponential scaling operator to $I_0(x)$ instead of to e^x . Second, each monomial x^k is accompanied by a modified Bessel carrier function: I_0 “carries” the even degrees of x and I_1 “carries” the odd degrees. The proper analogue of the Touchard polynomials should not have these carrier functions attached. Thankfully, the solution is the same one that produced integer coefficients for the e.g.f.s in Section 3.2: combine the two independent carrier systems, with weights chosen so that the general Wronskian identity (15) cancels all carriers.

Theorem 9 (Bessel–Touchard polynomials). *Define the bivariate generating function*

$$G(t, x) = x[K_0(x) + K_1(x)] I_0(xe^t) + x[I_1(x) - I_0(x)] K_0(xe^t).$$

Then, for $x > 0$,

$$G(t, x) = \sum_{n=0}^{\infty} \frac{t^n}{n!} \mathcal{B}_n(x), \quad \mathcal{B}_n(x) = \sum_{k=0}^n \left[\begin{matrix} n \\ k \end{matrix} \right]_2 x^k.$$

We call $\mathcal{B}_n(x)$ the *Bessel–Touchard* (or *Bessel–Bell*) polynomials.

Proof. Apply Theorem 8 to the two carrier systems (I_0, I_1) and $(K_0, -K_1)$:

$$I_0(xe^t) = \sum_{n=0}^{\infty} \frac{t^n}{n!} \sum_{k=0}^n \left[\begin{matrix} n \\ k \end{matrix} \right]_2 x^k I_{k \bmod 2}(x),$$

$$K_0(xe^t) = \sum_{n=0}^{\infty} \frac{t^n}{n!} \sum_{k=0}^n \left[\begin{matrix} n \\ k \end{matrix} \right]_2 x^k \varkappa_k(x), \quad \varkappa_k = \begin{cases} K_0 & k \text{ even} \\ -K_1 & k \text{ odd.} \end{cases}$$

Now form the combination $p(x)I_0(xe^t) + q(x)K_0(xe^t)$ with as yet undetermined weights. The coefficient of $\left[\begin{matrix} n \\ k \end{matrix} \right]_2 x^k$ in the t -expansion is

$$\begin{cases} p(x)I_0(x) + q(x)K_0(x) & k \text{ even,} \\ p(x)I_1(x) - q(x)K_1(x) & k \text{ odd.} \end{cases}$$

We want both expressions to equal 1 identically in x . Solving the linear system, the determinant is $-(I_0(x)K_1(x) + I_1(x)K_0(x)) = -1/x$ by the Wronskian identity (15), so the unique solution is

$$p(x) = x[K_0(x) + K_1(x)], \quad q(x) = x[I_1(x) - I_0(x)].$$

Indeed, one checks directly that

$$\begin{aligned} pI_0 + qK_0 &= x[K_1I_0 + I_1K_0] = 1, \\ pI_1 - qK_1 &= x[K_0I_1 + I_0K_1] = 1. \end{aligned}$$

With this choice every carrier collapses to 1 and the coefficient of $t^n/n!$ becomes $\sum_k \left[\begin{matrix} n \\ k \end{matrix} \right]_2 x^k$, as claimed. (All series converge absolutely for $x > 0$ and all real t , justifying the rearrangements.) $\square$

Corollary 10 (Row sums and parity row sums). *Setting $x = 1$ in Theorem 9 and comparing with Theorem 2:*

$$\mathcal{B}_n(1) = \sum_{k=0}^n \left[\begin{matrix} n \\ k \end{matrix} \right]_2 = B_n^{(2)} \quad (\text{A007472}),$$

since $G(t, 1) = [K_0(1) + K_1(1)]I_0(e^t) + [I_1(1) - I_0(1)]K_0(e^t)$ is precisely the e.g.f. of A007472. Moreover, choosing instead the weights $p(x) = xK_1(x)$, $q(x) = xI_1(x)$ (which map even-carrier coefficients to 1 and odd-carrier coefficients to 0) and $p(x) = xK_0(x)$, $q(x) = -xI_0(x)$ (the reverse) shows in the same way that

$$\sum_{\substack{k=0 \\ k \text{ even}}}^n \left[\begin{matrix} n \\ k \end{matrix} \right]_2 = B_n^{(2,0)} \quad (\text{A351143}), \quad \sum_{\substack{k=0 \\ k \text{ odd}}}^n \left[\begin{matrix} n \\ k \end{matrix} \right]_2 = B_n^{(2,1)} \quad (\text{A351028}).$$

Proof. For the first display, Theorem 2 identifies $G(t, 1)$ as the e.g.f. of A007472, while Theorem 9 expands the same function as $\sum_n \mathcal{B}_n(1)t^n/n!$. For the parity-restricted sums, the weights $p(x) = xK_1(x)$, $q(x) = xI_1(x)$ solve $pI_0 + qK_0 = 1$, $pI_1 - qK_1 = 0$ (again via the Wronskian), so the resulting bivariate function expands with only the even- k terms surviving; at $x = 1$ it equals $K_1(1)I_0(e^t) + I_1(1)K_0(e^t)$, the e.g.f. of A351143 by Theorem 2. The odd case is identical. $\square$

Corollary 10 reveals a satisfying structure: the two primitive 2-Bell sequences are interleaved inside a single triangle—A351143 collects the even-indexed columns and A351028 the odd-indexed ones, while their sum A007472 collects them all. In Section 7 (Corollary 26) we will prove by elementary means that this picture persists for every m , with the m primitive m -Bell sequences appearing as the residue-class row sums of the m -Stirling triangle.

The Bessel–Stirling numbers are a novel generalization of the Stirling numbers and join a broader family of two-term recurrences ([14, 15, 1, 19]). We return to the connection between this new array and previous generalizations of the Stirling numbers in Section 9. Note also that the Bessel–Stirling numbers are distinct from the r -Stirling numbers of Broder [3], despite the superficially similar subscript notation.

Due to the form of their recurrence, the Bessel–Stirling numbers admit many identities common to other such two-term-recurrence triangular arrays (see Appendices A and B). The most important one is an explicit formula (proof given in Appendix A):

Theorem 11 (Explicit formula for the Bessel–Stirling numbers of the second kind). *The Bessel–Stirling numbers can be calculated explicitly for $n \geq 2, k \geq 2$ via the following formula:*

$$\left[\begin{matrix} n \\ k \end{matrix} \right]_2 = \sum_{j=1}^{\lfloor k/2 \rfloor} (2j)^{n-k} (c_{k,j} + d_{k,j}(n-k+1)),$$

where d and c are determined as follows:

$$d_{k,r} = \frac{(-r)^{k-1}}{r! r! (\lfloor \frac{k}{2} \rfloor - r)! (\lfloor \frac{k-1}{2} \rfloor - r)!}$$

$$c_{k,r} = \begin{cases} -d_{k,r} \left(r(2H_{r-1} - H_{\lfloor \frac{k}{2} \rfloor - r} - H_{\lfloor \frac{k-1}{2} \rfloor - r}) - k + 3 \right) & \text{if } 1 \leq r < k/2 \\ \frac{r^{2r}}{r! r!} & \text{if } r = k/2 \\ 0 & \text{otherwise} \end{cases}$$

Alternatively, the following limit representation is valid for all $n \geq 0, k \geq 0$:

$$\left[\begin{matrix} n \\ k \end{matrix} \right]_2 = (-1)^{k-1} 2^{n-k} \sum_{j=0}^{\lfloor k/2 \rfloor} \lim_{t \rightarrow j} \left[t^{n-1} \frac{n-2-t \Psi_k(t)}{\Gamma(t+1)^2 \Gamma(\lfloor \frac{k}{2} \rfloor - t + 1) \Gamma(\lfloor \frac{k-1}{2} \rfloor - t + 1)} \right],$$

$$\Psi_k(t) = 2\psi(t) - \psi(\lfloor \frac{k}{2} \rfloor - t + 1) - \psi(\lfloor \frac{k-1}{2} \rfloor - t + 1),$$

where H_n is the n -th harmonic number, $\psi(x)$ is the digamma function, and $\Gamma(x)$ is the gamma function.

The proof proceeds by partial-fraction decomposition of the ordinary generating function of the k -th column,

$$(27) \quad A_k(x) = \sum_{n \geq 0} \left[\begin{matrix} n \\ k \end{matrix} \right]_2 x^n = \frac{x^k}{\prod_{j=1}^k (1 - 2 \lfloor j/2 \rfloor x)},$$

which itself follows directly from the recurrence (23) (see Appendix A). Two useful byproducts of (27) are worth recording here. First, explicit formulas for the low columns and the subdiagonal:

$$\begin{aligned} \left[\begin{matrix} n \\ 2 \end{matrix} \right]_2 &= 2^{n-2}, & n \geq 2 \quad (\text{rel. A000079}) \\ \left[\begin{matrix} n \\ 3 \end{matrix} \right]_2 &= (n-2)2^{n-3}, & n \geq 3 \quad (\text{rel. A001787}) \\ \left[\begin{matrix} n \\ 4 \end{matrix} \right]_2 &= 2^{2n-6} - (n-1)2^{n-4}, & n \geq 4 \quad (\text{rel. A100575}) \\ \left[\begin{matrix} n \\ 5 \end{matrix} \right]_2 &= n2^{n-5} + (n-6)2^{n-8}, & n \geq 5 \quad (\text{rel. A158681}) \\ \left[\begin{matrix} n \\ n-1 \end{matrix} \right]_2 &= \lfloor (n-1)^2/2 \rfloor, & n \geq 1 \quad (\text{rel. A007590}) \end{aligned}$$

(with the caveat that the sequences listed at the end of each line are offset or scaled). Second, expanding the product in (27) as a geometric series shows that $\left[\begin{matrix} n \\ k \end{matrix} \right]_2$ is the complete homogeneous symmetric polynomial h_{n-k} evaluated at the weights $2 \lfloor 1/2 \rfloor, 2 \lfloor 2/2 \rfloor, \dots, 2 \lfloor k/2 \rfloor$; we return to this symmetric-function viewpoint, and its dual for the first kind, in Section 6 (Proposition 18).

Now that we understand the analytic properties of the Bessel–Stirling and Bessel–Bell numbers and their basic recurrences, the natural next question is: what do these numbers count?

5. COMBINATORIAL INTERPRETATION OF THE BESSEL–STIRLING NUMBERS OF THE SECOND KIND AND THE BESSEL–BELL NUMBERS

The Bessel–Stirling numbers $\llbracket n \rrbracket_k_2$ enumerate the number of ways to place n labeled marbles into k unlabeled compartments following a parity-constrained urn model.

Theorem 12 (The two-compartment urn model). *The Bessel–Stirling number of the second kind $\llbracket n \rrbracket_2$ counts the number of ways to arrange n labeled marbles (placed one at a time in the order $1, 2, \dots, n$) in k total compartments distributed across urns with two unlabeled compartments each, subject to the following constraints:*

- You can place a marble in any compartment of any urn to begin
- You cannot place another marble in an urn’s used compartment until the other compartment of that urn also contains at least one marble
- You can freely place marbles in any compartment of urns where both compartments have been used
- You cannot open a new urn until both compartments of all previously used urns contain at least one marble each

Consequently, the Bessel–Bell numbers (A007472) count the total number of such parity-constrained placements of n labeled marbles, over all k .

Before diving into the proof, let us go through some examples to build intuition. Let $\frac{a}{b} \mid \frac{c}{d}$ represent 2 urns with vertical bars separating the urns and a horizontal bar separating the two compartments within an urn. The numbers that will replace the placeholder letters represent the marbles, labeled in the order in which they were placed (one at a time). For $\llbracket 3 \rrbracket_2$ we have two ways to place the marbles in two compartments of a single urn: $\frac{13}{2}$ or $\frac{1}{23}$. For $\llbracket 3 \rrbracket_3$ we have one way to place each marble in its own compartment (2 urns, 3 compartments used): $\frac{1}{2} \mid \frac{3}{}$. All valid partitions of up to 5 objects are presented in Table 3.

Proof. We proceed by induction on n . The base cases are straightforward:

- $\llbracket 0 \rrbracket_2 = 1$: There is exactly one way to arrange 0 marbles using 0 compartments.
- $\llbracket n \rrbracket_2 = 0$ for $n > 0$: It is impossible to arrange a positive number of marbles using 0 compartments.
- $\llbracket n \rrbracket_2 = 0$ for $k > n$: It is impossible to use more than n compartments when placing n marbles, since each used compartment contains at least one marble.
- $\llbracket n \rrbracket_2 = 1$ for $n > 0$: There is exactly one way to place n marbles into n different compartments (the constraints force the order in which compartments open).

For the inductive step, assume that for some $n \geq 1$, the number $\llbracket n \rrbracket_k_2$ correctly counts the arrangements of n labeled marbles into k compartments following our constraints. We need to show that $\llbracket n+1 \rrbracket_k_2 = 2\lfloor k/2 \rfloor \llbracket n \rrbracket_k_2 + \llbracket n \rrbracket_{k-1}_2$ counts the arrangements for $n+1$ marbles in k compartments. Consider the placement options for the $(n+1)$ -st marble:

- *Place in a new compartment:* The $(n+1)$ -st marble can be placed in a new (previously unused) compartment, making the total number of compartments used equal to k . This requires that the previous n marbles were arranged in $k-1$ compartments, and there are $\llbracket n \rrbracket_{k-1}_2$ such arrangements.
- *Place in a previously used compartment:* The $(n+1)$ -st marble can be placed in one of the compartments already containing marbles. Here, the key insight comes from the urn model’s constraints.

We need to count how many valid placement options exist among the k compartments already in use. This depends on the distribution of these compartments across urns:

- When k is even, say $k = 2j$, the k compartments must be distributed across exactly j fully-filled urns (both compartments used in each). This is because our constraints prohibit opening a new urn until all previous urns have at least one marble in each compartment. With j fully-filled urns, we have $2j = k$ valid placement options for the $(n+1)$ -st marble, since we can place it in any of the already-used compartments.
- When k is odd, say $k = 2j+1$, we must have j fully-filled urns (accounting for $2j$ compartments) plus one urn with exactly one compartment used. According to

$T(n,k)$	Count	Examples
$T(1,1)$	1	$\underline{1}$
$T(2,2)$	1	$\frac{1}{2}$
$T(3,2)$	2	$\frac{1}{23}, \frac{13}{2}$
$T(3,3)$	1	$\frac{1}{2} \frac{3}{2}$
$T(4,2)$	4	$\frac{14}{23}, \frac{134}{2}, \frac{1}{234}, \frac{13}{24}$
$T(4,3)$	4	$\frac{14}{2} \frac{3}{2}, \frac{1}{24} \frac{3}{2}, \frac{13}{2} \frac{4}{2}, \frac{1}{23} \frac{4}{2}$
$T(4,4)$	1	$\frac{1}{2} \frac{3}{4}$
$T(5,2)$	8	$\frac{145}{23}, \frac{14}{235}, \frac{1345}{2}, \frac{134}{25}, \frac{15}{234}, \frac{1}{2345}, \frac{135}{24}, \frac{13}{245}$
$T(5,3)$	12	$\frac{145}{2} \frac{3}{2}, \frac{14}{25} \frac{3}{2}, \frac{15}{24} \frac{3}{2}, \frac{1}{245} \frac{3}{2}, \frac{135}{2} \frac{4}{2}, \frac{13}{25} \frac{4}{2}, \frac{15}{23} \frac{4}{2}, \frac{1}{235} \frac{4}{2},$ $\frac{134}{2} \frac{5}{2}, \frac{1}{234} \frac{5}{2}, \frac{13}{24} \frac{5}{2}, \frac{14}{23} \frac{5}{2}$
$T(5,4)$	8	$\frac{13}{2} \frac{4}{5}, \frac{14}{2} \frac{3}{5}, \frac{15}{2} \frac{3}{4}, \frac{1}{23} \frac{4}{5}, \frac{1}{24} \frac{3}{5}, \frac{1}{25} \frac{3}{4}, \frac{1}{2} \frac{35}{4}, \frac{1}{2} \frac{3}{45}$
$T(5,5)$	1	$\frac{1}{2} \frac{3}{4} \frac{5}{2}$

TABLE 3. Valid Bessel–Stirling constrained partitions of n objects into k compartments of 2-compartment urns

the constraints, we cannot place another marble in this partially-filled urn’s used compartment until its other compartment also contains a marble. Therefore, we have only $2j = 2\lfloor k/2 \rfloor$ valid placement options in previously used compartments.

In either case, the number of valid placement options in previously used compartments is $2\lfloor k/2 \rfloor$. Since there are $\llbracket n \rrbracket_2$ ways to arrange the first n marbles in k compartments, we have $2\lfloor k/2 \rfloor \cdot \llbracket n \rrbracket_2$ ways to place the $(n+1)$ -st marble in a previously used compartment. Combining both cases, the total number of ways to arrange $n+1$ marbles in k compartments is

$$\llbracket n+1 \rrbracket_2 = 2\lfloor k/2 \rfloor \cdot \llbracket n \rrbracket_2 + \llbracket n \rrbracket_{k-1}.$$

This matches exactly the recurrence relation for the Bessel–Stirling numbers, completing the proof. (Two placement processes that produce the same final assignment of marbles to compartments coincide step by step, since the placement history can be recovered by removing marbles $n, n-1, \dots$ in reverse order; the count is therefore genuinely a count of distinct arrangements.) $\square$

This combinatorial interpretation generalizes verbatim to urns with m compartments, yielding the m -Stirling numbers and m -Bell sequences (Theorem 35). In the special case $m = 1$, each urn has only one compartment, so we are left with the standard interpretation of the Stirling numbers as counting partitions of labeled objects into unlabeled non-empty sets. The combinatorial model provides intuition for why the row sums shift two places to the left after applying the binomial transform twice: the constraint requiring both compartments of an urn to be filled before opening a new urn creates a “pairing effect”, which manifests as the shift-by-two property. This loose argument is made precise, in a purely algebraic way, by the shift identity of Theorem

25. Finally, an easy way to enumerate the combinatorial objects counted by the Bessel–Stirling numbers of the second kind is via succession rules and a generating tree, as introduced by West [23]. See Figure 1 and Appendix C.




FIGURE 1. First five levels of the generating tree for the Bessel–Stirling numbers of the second kind using the succession rule Ω of Appendix C.

6. BESSEL–STIRLING NUMBERS OF THE FIRST KIND AND BESSEL FALLING FACTORIALS

The regular Stirling numbers come in a dual pair. In addition to the Stirling numbers of the second kind, which we explored at length, there exists a partner triangular array that bears the name Stirling numbers of the first kind. The two arrays are orthogonal to each other, or inverses of one another [16]:

$$\sum_{j=k}^n s(n, j) S(j, k) = \sum_{j=k}^n (-1)^{n-j} \begin{bmatrix} n \\ j \end{bmatrix} \begin{Bmatrix} j \\ k \end{Bmatrix} = \delta_{n,k},$$

where $s(n, k) = (-1)^{n-k} \begin{bmatrix} n \\ k \end{bmatrix}$ denotes the *signed* Stirling numbers of the first kind, $\begin{bmatrix} n \\ k \end{bmatrix}$ the *unsigned* Stirling numbers of the first kind, and $\delta_{n,k}$ is the Kronecker delta. When represented as lower-triangular matrices, with S the matrix of Stirling numbers of the second kind and s the matrix of signed Stirling numbers of the first kind, we have $sS = Ss = I$. The recurrence relation for the unsigned Stirling numbers of the first kind has a factor of n rather than k :

$$\begin{bmatrix} n+1 \\ k \end{bmatrix} = n \begin{bmatrix} n \\ k \end{bmatrix} + \begin{bmatrix} n \\ k-1 \end{bmatrix}.$$

The Bessel–Stirling numbers also have a first-kind partner, defined by the analogous recurrence.

Definition 13 (Unsigned Bessel–Stirling numbers of the first kind). *Let $\|n\|_k$ be defined for $n, k \geq 0$ by $\|n\|_k = \delta_{0,k}$, the convention $\|n\|_k = 0$ for $k < 0$, and*

$$\|n+1\|_k = 2 \lfloor n/2 \rfloor \|n\|_k + \|n\|_{k-1}.$$

The first few rows of the unsigned Bessel–Stirling numbers of the first kind are shown in Table 4. This recurrence closely mirrors that of the standard Stirling numbers of the first kind, but with the factor n replaced by $2 \lfloor n/2 \rfloor$, reflecting the “even-step” structure inherent in the Bessel framework. Just as the ordinary Stirling numbers of the first kind arise in the context of a change of basis from ordinary powers to falling powers, these generalized coefficients appear as the change-of-basis matrix between ordinary powers and a novel class of falling factorials.

6.1. Bessel falling factorials. The signed Stirling numbers of the first kind are well known to be the coefficients in the expansion of the falling factorial:

$$x^n = \prod_{k=0}^{n-1} (x-k) = x(x-1)(x-2) \cdots (x-n+1) = \sum_{k=0}^n (-1)^{n-k} \begin{bmatrix} n \\ k \end{bmatrix} x^k.$$

In the Bessel case, we discover a strikingly analogous structure.

	k=0	k=1	k=2	k=3	k=4	k=5	k=6	k=7	k=8	k=9	$\sum_{k=0}^n$
n=0	1										1
n=1	0	1									1
n=2	0	0	1								1
n=3	0	0	2	1							3
n=4	0	0	4	4	1						9
n=5	0	0	16	20	8	1					45
n=6	0	0	64	96	52	12	1				225
n=7	0	0	384	640	408	124	18	1			1575
n=8	0	0	2304	4224	3088	1152	232	24	1		11025
n=9	0	0	18432	36096	28928	12304	3008	424	32	1	99225

TABLE 4. Unsigned Bessel–Stirling numbers of the first kind, $\left\|_k^n\right\|_2$.
Row sums give A000246.

	k=0	k=1	k=2	k=3	k=4	k=5	k=6	k=7	k=8	k=9	$\sum_{k=0}^n$
n=0	1										1
n=1	0	1									1
n=2	0	0	1								1
n=3	0	0	4	1							5
n=4	0	0	16	8	1						25
n=5	0	0	96	64	16	1					177
n=6	0	0	576	480	192	24	1				1273
n=7	0	0	4608	4416	2400	432	36	1			11893
n=8	0	0	36864	39936	28416	6720	864	48	1		112849
n=9	0	0	368640	436224	380928	109056	18816	1536	64	1	1315265

TABLE 5. Bessel–Lah numbers (Bessel–Stirling numbers of the third kind), $\left\|_k^n\right\|_2$.

Definition 14 (Bessel falling and rising factorials). *Define the Bessel falling and rising factorials (or even-step falling and rising factorials) as*

$$x^{\underline{n}} = \prod_{k=0}^{n-1} (x - 2 \lfloor k/2 \rfloor) = x \cdot x \cdot (x - 2)(x - 2)(x - 4)(x - 4) \cdots$$

$$x^{\overline{n}} = \prod_{k=0}^{n-1} (x + 2 \lfloor k/2 \rfloor) = x \cdot x \cdot (x + 2)(x + 2)(x + 4)(x + 4) \cdots$$

where each product has exactly n factors, so each factor is repeated twice except possibly the last: e.g. $x^{\underline{4}} = x^2(x - 2)^2$ but $x^{\underline{5}} = x^2(x - 2)^2(x - 4)$.

Theorem 15. *The Bessel–Stirling numbers of the first kind are the coefficients in the expansions of the Bessel falling and rising factorials:*

$$x^{\underline{n}} = \sum_{k=0}^n (-1)^{n-k} \left\|_k^n \right\|_2 x^k$$

$$x^{\overline{n}} = \sum_{k=0}^n \left\|_k^n \right\|_2 x^k.$$

Proof. Induction on n , rising version first. Multiplying $x^{\overline{n}} = \sum_k \left\|_k^n \right\|_2 x^k$ by the next factor $(x + 2 \lfloor n/2 \rfloor)$ sends the coefficient of x^k to $\left\|_{k-1}^n \right\|_2 + 2 \lfloor n/2 \rfloor \left\|_k^n \right\|_2$, which is exactly the recurrence of Definition 13. The falling version follows by substituting $x \mapsto -x$, since $x^{\underline{n}} = (-1)^n (-x)^{\overline{n}}$. $\square$

Here are the expansions of the first few Bessel rising factorials:

$$\begin{aligned} x^{\overline{1}} &= x = x \\ x^{\overline{2}} &= x^2 = x^2 \\ x^{\overline{3}} &= x^2(x+2) = 2x^2 + x^3 \\ x^{\overline{4}} &= x^2(x+2)^2 = 4x^2 + 4x^3 + x^4 \\ x^{\overline{5}} &= x^2(x+2)^2(x+4) = 16x^2 + 20x^3 + 8x^4 + x^5 \\ x^{\overline{6}} &= x^2(x+2)^2(x+4)^2 = 64x^2 + 96x^3 + 52x^4 + 12x^5 + x^6 \end{aligned}$$

Just as the regular falling powers form an alternative polynomial basis, which can be converted to and from the monomial basis using the Stirling numbers, the same is true for the Bessel falling powers. One recovers a monomial from them using the Bessel–Stirling numbers of the second kind:

Theorem 16.

$$x^n = \sum_{k=0}^n \left[\begin{matrix} n \\ k \end{matrix} \right]_2 x^{\underline{k}}.$$

Proof. From Definition 14, $x^{\overline{k+1}} = x^{\underline{k}}(x - 2 \lfloor k/2 \rfloor)$, so

$$(28) \quad x \cdot x^{\underline{k}} = x^{\overline{k+1}} + 2 \lfloor k/2 \rfloor x^{\underline{k}}.$$

Now induct on n : multiplying $x^n = \sum_k \left[\begin{matrix} n \\ k \end{matrix} \right]_2 x^{\underline{k}}$ by x and applying (28) term by term shows that the coefficient of $x^{\underline{k}}$ in x^{n+1} is $\left[\begin{matrix} n \\ k-1 \end{matrix} \right]_2 + 2 \lfloor k/2 \rfloor \left[\begin{matrix} n \\ k \end{matrix} \right]_2 = \left[\begin{matrix} n+1 \\ k \end{matrix} \right]_2$, by (23). Since $\{x^{\underline{k}}\}_{k \geq 0}$ has one polynomial of each degree, it is a basis and the coefficients are unique. $\square$

Comparing (28) with the carrier computation in the proof of Theorem 8 explains the coincidence of coefficients: multiplication by x acts on the falling-factorial basis exactly as xD acts on the carrier module $\{x^k f_{k \bmod 2}\}$ —both are “weighted shift” operators with weight sequence $2 \lfloor k/2 \rfloor$.

Proposition 17 (Orthogonality).

$$\sum_{j=k}^n (-1)^{n-j} \left[\begin{matrix} n \\ j \end{matrix} \right]_2 \left[\begin{matrix} j \\ k \end{matrix} \right]_2 = \delta_{n,k}, \quad \sum_{j=k}^n (-1)^{j-k} \left[\begin{matrix} n \\ j \end{matrix} \right]_2 \left[\begin{matrix} j \\ k \end{matrix} \right]_2 = \delta_{n,k}.$$

Proof. By Theorem 15, the matrix $((-1)^{n-k} \left[\begin{matrix} n \\ k \end{matrix} \right]_2)_{n,k}$ converts the monomial basis into the Bessel-falling basis, and by Theorem 16 the matrix $(\left[\begin{matrix} n \\ k \end{matrix} \right]_2)_{n,k}$ converts it back. The composition of the two changes of basis, in either order, is the identity. $\square$

Thus the two Bessel–Stirling triangles form a pair of lower-triangular inverse matrices (after attaching signs to one of them)—one converting monomials to generalized factorials, the other reversing the transformation, in complete parallel with the classical Stirling duality.

6.2. Symmetric function expressions and file placements. Both triangles are specializations of the classical bases of symmetric functions, evaluated on the “doubled” weight sequence.

Proposition 18. Let $w_j = 2 \lfloor j/2 \rfloor$, and let e_d and h_d denote the elementary and complete homogeneous symmetric polynomials of degree d . Then for $0 \leq k \leq n$:

$$\left[\begin{matrix} n \\ k \end{matrix} \right]_2 = e_{n-k}(w_0, w_1, \dots, w_{n-1}), \quad \left[\begin{matrix} n \\ k \end{matrix} \right]_2 = h_{n-k}(w_1, w_2, \dots, w_k).$$

Proof. The first identity reads off the coefficient of x^k in $x^{\overline{n}} = \prod_{j=0}^{n-1} (x + w_j)$ (Theorem 15). The second reads off the coefficient of x^n in the column generating function (27): expanding each geometric factor $(1 - w_j x)^{-1} = \sum_i w_j^i x^i$ and extracting $[x^{n-k}]$ gives precisely the sum of all monomials $w_{j_1} w_{j_2} \cdots w_{j_{n-k}}$ with $1 \leq j_1 \leq \cdots \leq j_{n-k} \leq k$, which is $h_{n-k}(w_1, \dots, w_k)$. $\square$

The classical Stirling numbers are the same specializations with weights $(0, 1, 2, \dots)$: $[n]_k = e_{n-k}(0, 1, \dots, n-1)$ and $\{n\}_k = h_{n-k}(1, 2, \dots, k)$ [5]. In rook-theoretic language [11], $e_{n-k}(w_0, \dots, w_{n-1})$ counts *file placements* of $n-k$ rooks on the “doubled staircase” board whose j -th column has height $w_j = 2 \lfloor j/2 \rfloor$ (rooks in distinct columns, one cell each, with no constraint along rows), giving the Bessel–Stirling numbers of the first kind a direct board-counting interpretation.

6.3. A combinatorial interpretation of the first kind: parity-restricted insertions. The unsigned Stirling numbers of the first kind $[n]_k$ count permutations of $[n] = \{1, \dots, n\}$ with exactly k cycles. The standard proof builds a permutation by inserting the elements $1, 2, \dots, n$ one at a time: element $i+1$ either opens a new cycle, or is inserted immediately after any of the i existing elements in its cycle. The Bessel–Stirling numbers of the first kind count the permutations produced when the insertion process is parity-restricted.

Theorem 19 (Restricted-insertion model). *Build permutations of $[n]$ by inserting elements $1, 2, \dots, n$ in order, where element $i+1$ either*

- *opens a new cycle as a fixed point, or*
- *is inserted immediately after one of the existing elements $1, \dots, i$ in cycle notation, except that when i is odd, insertion immediately after the largest element i is forbidden.*

Distinct insertion histories produce distinct permutations, and $\|n\|_k^2$ counts the resulting permutations with exactly k cycles. Equivalently, $\|n\|_k^2$ is the number of permutations π of $[n]$ with k cycles such that for every even $e \leq n$, the restriction of π to $\{1, \dots, e\}$ (obtained by deleting the elements larger than e from the cycle notation) does not map $e-1$ to e .

Proof. At the step inserting element $i+1$ there is 1 way to open a new cycle and $i - (i \bmod 2) = 2 \lfloor i/2 \rfloor$ permitted insertion positions, so the number of histories with k “new cycle” steps satisfies the recurrence of Definition 13 with the cycle count increasing exactly on new-cycle steps. Histories biject with outcomes because the history is recoverable from the final permutation: deleting the largest element (splicing it out of its cycle) undoes the last step, and iterating recovers every step. Finally, the insertion of $i+1$ immediately after i is precisely what makes the restriction of the final permutation to $\{1, \dots, i+1\}$ map i to $i+1$; with $e = i+1$ ranging over even values, the forbidden histories correspond exactly to the stated condition on π . $\square$

For general m the same model applies with “ i odd” replaced by the rule that insertion is only permitted after the $m \lfloor i/m \rfloor$ smallest elements (equivalently, insertion after the largest $i \bmod m$ elements is forbidden); the count of reachable permutations of $[n]$ with k cycles is then $\|n\|_k^m$ (Section 7).

6.4. Row sums of the first kind: permutations of odd order. Although the first-kind triangle was, prior to this work, also not present in the OEIS, its row sums yield a sequence with a rich combinatorial history: A000246, the number of permutations in the symmetric group S_n that have odd order, i.e. all of whose cycle lengths are odd: 1, 1, 1, 3, 9, 45, 225, 1575, 11025, 99225, $\dots$

Proposition 20.

$$\sum_{k=0}^n \|n\|_k^2 = \prod_{j=0}^{n-1} (2 \lfloor j/2 \rfloor + 1) = \#\{\pi \in S_n : \pi \text{ has odd order}\}.$$

Proof. Summing the recurrence of Definition 13 over k shows that the row sums s_n satisfy $s_{n+1} = (2 \lfloor n/2 \rfloor + 1)s_n$, $s_0 = 1$, which telescopes to the product. For the second equality, let a_n denote the number of permutations of $[n]$ all of whose cycles have odd length. By the exponential formula, $\sum_n a_n x^n / n! = \exp(\sum_{d \text{ odd}} x^d / d) = \exp(\frac{1}{2} \log \frac{1+x}{1-x}) = \sqrt{(1+x)/(1-x)} =: A(x)$. Differentiating, $(1-x^2)A'(x) = A(x)$, so extracting coefficients gives $a_{n+1} = a_n + n(n-1)a_{n-1}$. The product sequence s_n satisfies the same recurrence: $s_{n+1} - s_n = 2 \lfloor n/2 \rfloor s_n = 2 \lfloor n/2 \rfloor (2 \lfloor (n-1)/2 \rfloor + 1)s_{n-1} = n(n-1)s_{n-1}$, where the last equality holds for n even ($n \cdot (n-1)$) and n odd ($(n-1) \cdot n$) alike. Since $s_0 = a_0 = 1$ and $s_1 = a_1 = 1$, the sequences coincide. $\square$

We emphasize that while the row sums agree, the triangle $\llbracket n \rrbracket_2$ does *not* refine the odd-order permutations by their number of cycles: for $n = 5$ the odd-order permutations count $(24, 20, 1)$ by $(1, 3, 5)$ cycles, whereas row 5 of the triangle reads $(16, 20, 8, 1)$ for $k = (2, 3, 4, 5)$. Finding a statistic on odd-order permutations that is equidistributed with the k -statistic of Theorem 19—equivalently, a bijection between odd-order permutations and the parity-admissible permutations of Theorem 19—remains an open problem (Section 9).

6.5. Bessel–Lah numbers (the third kind). The classical Lah numbers $\lfloor n \rfloor_k = \binom{n-1}{k-1} \frac{n!}{k!}$ convert rising factorials into falling factorials, and equal the matrix product of the unsigned first-kind and second-kind Stirling triangles. The same construction goes through in the Bessel setting.

Definition 21 (Bessel–Lah numbers). *Define $\llbracket n \rrbracket_2$ as the connection coefficients between the Bessel rising and falling factorial bases:*

$$x^{\bar{n}} = \sum_{k=0}^n \llbracket n \rrbracket_k x^{\underline{k}}.$$

Theorem 22. *The Bessel–Lah numbers satisfy:*

- (i) $\llbracket n \rrbracket_2 = \sum_j \llbracket n \rrbracket_j \llbracket j \rrbracket_2$ (matrix product of the two Bessel–Stirling triangles);
- (ii) the two-term recurrence

$$\llbracket n+1 \rrbracket_k = (2 \lfloor n/2 \rfloor + 2 \lfloor k/2 \rfloor) \llbracket n \rrbracket_k + \llbracket n \rrbracket_{k-1},$$

$$\text{with } \llbracket 0 \rrbracket_k = \delta_{0,k}.$$

Proof. (i) Compose the expansions of Theorems 15 and 16: $x^{\bar{n}} = \sum_j \llbracket n \rrbracket_j x^j = \sum_j \llbracket n \rrbracket_j \sum_k \llbracket j \rrbracket_k x^{\underline{k}}$. (ii) Multiply the defining expansion by the next rising factor and use (28):

$$x^{\overline{n+1}} = (x + 2 \lfloor n/2 \rfloor) \sum_k \llbracket n \rrbracket_k x^{\underline{k}} = \sum_k \llbracket n \rrbracket_k \left(x^{\underline{k+1}} + 2 \lfloor k/2 \rfloor x^{\underline{k}} + 2 \lfloor n/2 \rfloor x^{\underline{k}} \right),$$

and collect the coefficient of $x^{\underline{k}}$, using uniqueness of coefficients in the basis $\{x^{\underline{k}}\}$. $\square$

The first few rows are shown in Table 5. Note how the recurrence interpolates the classical Lah recurrence $\lfloor n+1 \rfloor_k = (n+k) \lfloor n \rfloor_k + \lfloor n \rfloor_{k-1}$, doubling-and-flooring both parameters. The row sums $1, 1, 1, 5, 25, 177, 1273, 11893, 112849, 1315265, \dots$ (the Bessel analogue of A000262, the number of sets of lists) do not currently appear in the OEIS.

Figure 2 summarizes how all six triangles act as conversion operators among the five polynomial bases.

7. THE GENERAL CASE

Having explored the case $m = 2$ in depth, we now develop the theory for arbitrary $m \geq 1$. Remarkably, the central structural result—that the m -Stirling triangle’s row sums produce the m -Bell numbers, with the residue classes of $k \bmod m$ producing the primitive sequences—admits a completely elementary proof, independent of any special function theory. We give that proof first, and then develop the analytic counterparts (hyper-Bessel carrier functions, hypergeometric e.g.f.s, and a general Dobiński formula).

7.1. The m -Stirling triangle and the shift identity.

Definition 23 (m -Stirling numbers of the second kind). *For an integer $m \geq 1$, define $\llbracket n \rrbracket_m$ for $n, k \geq 0$ by $\llbracket 0 \rrbracket_m = \delta_{0,k}$, the convention $\llbracket n \rrbracket_m = 0$ for $k < 0$, and the recurrence*

$$(29) \quad \llbracket n+1 \rrbracket_k = m \lfloor k/m \rfloor \llbracket n \rrbracket_k + \llbracket n \rrbracket_{k-1}.$$

For $m = 1$ this is the classical Stirling triangle of the second kind; for $m = 2$, the Bessel–Stirling triangle of Section 4. Table 6 displays the triangle for $m = 3$.




FIGURE 2. Stirling numbers and Bessel–Stirling numbers as transformations between polynomial bases. The regular Stirling (and Lah) numbers convert between ordinary powers (middle), falling powers (bottom left), and rising powers (top left). The Bessel–Stirling numbers convert between ordinary powers, Bessel falling powers (bottom right), and Bessel rising powers (top right). An arrow from basis u to basis v labeled $c_{n,k}$ means $u_n = \sum_k c_{n,k} v_k$.

	k=0	k=1	k=2	k=3	k=4	k=5	k=6	k=7	k=8	k=9	$\sum_{k=0}^n$
n=0	1										1
n=1	0	1									1
n=2	0	0	1								1
n=3	0	0	0	1							1
n=4	0	0	0	3	1						4
n=5	0	0	0	9	6	1					16
n=6	0	0	0	27	27	9	1				64
n=7	0	0	0	81	108	54	15	1			259
n=8	0	0	0	243	405	270	144	21	1		1084
n=9	0	0	0	729	1458	1215	1134	270	27	1	4834

TABLE 6. The 3-Stirling numbers of the second kind, $\llbracket n \atop k \rrbracket_3$.

Lemma 24. $\llbracket n \atop k \rrbracket_m = 0$ for $k > n$, $\llbracket n \atop n \rrbracket_m = 1$, and the first $m+1$ rows form an identity block:

$$\llbracket n \atop k \rrbracket_m = \delta_{n,k} \quad \text{for } 0 \leq n \leq m.$$

Proof. The first two claims follow by induction as in the classical case. For the third, note that the coefficient $m \lfloor k/m \rfloor$ in (29) vanishes whenever $k < m$. Hence, for rows $n < m$, where only entries with $k \leq n < m$ can be nonzero, the recurrence degenerates to a pure shift $\llbracket n+1 \atop k \rrbracket_m = \llbracket n \atop k-1 \rrbracket_m$, which propagates the initial row $\delta_{0,k}$ diagonally. For row $n = m$, the only new entry is $\llbracket m \atop m \rrbracket_m = m \lfloor m/m \rfloor \llbracket m-1 \atop m \rrbracket_m + \llbracket m-1 \atop m-1 \rrbracket_m = 0 + 1 = 1$. $\square$

We now come to the central result of the paper, an m -fold analogue of the classical shift identity (6). It explains, at the level of individual triangle entries, why the row sums shift by m places after m binomial transforms.

Theorem 25 (Shift identity for m -Stirling numbers). *For all integers $n \geq 0$ and $k \geq 0$,*

$$(30) \quad \llbracket n+m \atop k \rrbracket_m = \sum_{j=0}^n \binom{n}{j} m^{n-j} \llbracket j \atop k-m \rrbracket_m.$$

For $m = 1$ this is the classical identity (6).

Proof. Induction on n , with k arbitrary. For $n = 0$ the claim reads $\llbracket m \atop k \rrbracket_m = \llbracket 0 \atop k-m \rrbracket_m = \delta_{m,k}$, which is Lemma 24. Assume (30) holds for some $n \geq 0$ and all k . Using Pascal's

rule $\binom{n+1}{j} = \binom{n}{j} + \binom{n}{j-1}$ and re-indexing the second sum,

$$\sum_{j=0}^{n+1} \binom{n+1}{j} m^{n+1-j} \left[\begin{matrix} j \\ k-m \end{matrix} \right]_m = m \sum_{j=0}^n \binom{n}{j} m^{n-j} \left[\begin{matrix} j \\ k-m \end{matrix} \right]_m + \sum_{j=0}^n \binom{n}{j} m^{n-j} \left[\begin{matrix} j+1 \\ k-m \end{matrix} \right]_m.$$

Expanding $\left[\begin{matrix} j+1 \\ k-m \end{matrix} \right]_m$ by the recurrence (29), and noting $\lfloor (k-m)/m \rfloor = \lfloor k/m \rfloor - 1$, gives

$$\left[\begin{matrix} j+1 \\ k-m \end{matrix} \right]_m = (m \lfloor k/m \rfloor - m) \left[\begin{matrix} j \\ k-m \end{matrix} \right]_m + \left[\begin{matrix} j \\ k-m-1 \end{matrix} \right]_m.$$

Substituting this into the previous display, the two terms proportional to $m \left[\begin{matrix} j \\ k-m \end{matrix} \right]_m$ cancel, leaving

$$\begin{aligned} \sum_{j=0}^{n+1} \binom{n+1}{j} m^{n+1-j} \left[\begin{matrix} j \\ k-m \end{matrix} \right]_m &= m \lfloor k/m \rfloor \sum_{j=0}^n \binom{n}{j} m^{n-j} \left[\begin{matrix} j \\ k-m \end{matrix} \right]_m \\ &\quad + \sum_{j=0}^n \binom{n}{j} m^{n-j} \left[\begin{matrix} j \\ k-1-m \end{matrix} \right]_m. \end{aligned}$$

By the induction hypothesis (applied at k and at $k-1$), the right side equals $m \lfloor k/m \rfloor \left[\begin{matrix} n+m \\ k \end{matrix} \right]_m + \left[\begin{matrix} n+m \\ k-1 \end{matrix} \right]_m$, which is $\left[\begin{matrix} n+m+1 \\ k \end{matrix} \right]_m$ by (29), completing the induction. (All manipulations remain valid at the boundaries thanks to the convention $\left[\begin{matrix} n \\ k \end{matrix} \right]_m = 0$ for $k < 0$.) $\square$

Corollary 26 (m -Bell numbers as row sums; primitive sequences as residue-class sums). *Fix $m \geq 1$ and let*

$$\rho_r(n) = \sum_{\substack{k \geq 0 \\ k \equiv r \pmod{m}}} \left[\begin{matrix} n \\ k \end{matrix} \right]_m \quad (0 \leq r < m), \quad \sigma(n) = \sum_{k=0}^n \left[\begin{matrix} n \\ k \end{matrix} \right]_m.$$

Then:

- (i) *each ρ_r satisfies the m -Bell recurrence (2) with initial conditions $\rho_r(n) = \delta_{n,r}$ for $0 \leq n < m$; hence $\rho_r = B^{(m,r)}$ is the r -th primitive m -Bell sequence;*
- (ii) *the full row sums satisfy $\sigma = \sum_{r=0}^{m-1} \rho_r = B^{(m)}$, the composite m -Bell sequence;*
- (iii) *every sequence satisfying (2) is the unique linear combination $\sum_r a_r B^{(m,r)}$ of the residue-class row sums determined by its initial values $a_0, \dots, a_{m-1}$.*

Proof. (i) Summing the shift identity (30) over all $k \equiv r \pmod{m}$, and noting that $k \mapsto k-m$ preserves the residue class, gives

$$\rho_r(n+m) = \sum_{j=0}^n \binom{n}{j} m^{n-j} \rho_r(j),$$

which is exactly (2). The initial values follow from Lemma 24: for $n < m$, row n contains the single entry 1 at $k = n$, contributing to ρ_r if and only if $n \equiv r$, i.e. $n = r$. (ii) Sum (i) over r . (iii) A solution of (2) is determined by its first m values, and the primitives form a basis for the space of initial conditions. $\square$

For $m = 2$, Corollary 26 recovers Corollary 10 by entirely elementary means. Table 7 lists the composite m -Bell numbers for $m \leq 4$ and the primitive sequences for $m = 3$.

7.2. The ODE in hypergeometric form and hyper-Bessel carriers. We next generalize the analytic theory of Sections 3 and 4. As in the case $m = 2$, set $t = e^x$ (so that $D_x = tD_t$) and $f(t) = \mathcal{A}(x)$, and recall the classical operator identity [5, 8]

$$(31) \quad (tD_t)^n = \sum_{k=0}^n \left\{ \begin{matrix} n \\ k \end{matrix} \right\} t^k D_t^k,$$

which is itself yet another manifestation of the Stirling numbers as change-of-basis coefficients (from the basis $(tD)^k$ to the basis $t^k D^k$ of the Weyl algebra).

Sequence	First terms
$B^{(1)}$ (Bell, A000110)	1, 1, 2, 5, 15, 52, 203, 877, 4140, 21147, 115975, 678570
$B^{(2)}$ (A007472)	1, 1, 1, 3, 9, 29, 105, 431, 1969, 9785, 52145, 296155
$B^{(3)}$	1, 1, 1, 1, 4, 16, 64, 259, 1084, 4834, 23635, 128377
$B^{(4)}$	1, 1, 1, 1, 1, 5, 25, 125, 625, 3129, 15745, 80265
$B^{(3,0)}$	1, 0, 0, 1, 3, 9, 28, 96, 387, 1864, 10242, 60534
$B^{(3,1)}$	0, 1, 0, 0, 1, 6, 27, 109, 426, 1728, 7858, 42084
$B^{(3,2)}$	0, 0, 1, 0, 0, 1, 9, 54, 271, 1242, 5535, 25759

TABLE 7. Composite m -Bell numbers for $m = 1, \dots, 4$, and the three primitive 3-Bell sequences (residue-class row sums of Table 6). The sequences for $m \geq 3$ are not currently in the OEIS.

Proposition 27. *The e.g.f. $\mathcal{A}(x)$ of a sequence satisfying (3) equals $f(e^x)$, where f satisfies the m -th order ODE*

$$(32) \quad (tD_t)^m f = t^m f, \quad \text{equivalently} \quad \sum_{k=0}^m \left\{ \begin{matrix} m \\ k \end{matrix} \right\} t^k f^{(k)}(t) = t^m f(t).$$

For $m = 1$ this is $tf' = tf$, solved by e^t ; for $m = 2$ it is the modified Bessel equation (12).

Proof. By Proposition 1, $D_x^m \mathcal{A} = e^{mx} \mathcal{A}$. Substituting $D_x = tD_t$ and $e^{mx} = t^m$ gives $(tD_t)^m f = t^m f$; expanding via (31) gives the second form. $\square$

Note that (32) is precisely the primary-carrier equation (26). The distinguished solution analytic at $t = 0$ is a generalized hypergeometric function, and the full carrier system consists of the *hyper-Bessel functions* of Delerue [9, 13]:

Proposition 28 (Hyper-Bessel carrier systems). *For $m \geq 1$ define*

$$C_0^{(m)}(x) = \sum_{k=0}^{\infty} \frac{(x/m)^{mk}}{(k!)^m} = {}_0F_{m-1} \left(; 1, \dots, 1; \left(\frac{x}{m} \right)^m \right),$$

$$C_r^{(m)}(x) = \sum_{k=0}^{\infty} \frac{(x/m)^{mk+m-r}}{(k!)^r ((k+1)!)^{m-r}} = \frac{(x/m)^{m-r}}{(1!)^{m-r}} {}_0F_{m-1} \left(; \underbrace{1, \dots, 1}_{r-1}, \underbrace{2, \dots, 2}_{m-r}; \left(\frac{x}{m} \right)^m \right)$$

for $1 \leq r \leq m-1$. Then $(C_0^{(m)}, \dots, C_{m-1}^{(m)})$ is a carrier system of modulus m in the sense of Definition 7:

$$(xD + r) C_r^{(m)} = x C_{(r+1) \bmod m}^{(m)}, \quad r = 0, \dots, m-1.$$

In particular $C_0^{(m)}$ satisfies $(xD)^m C_0^{(m)} = x^m C_0^{(m)}$. For $m = 1$, $C_0^{(1)} = e^x$; for $m = 2$, $(C_0^{(2)}, C_1^{(2)}) = (I_0, I_1)$.

Proof. Direct computation on the series. For $r = 0$: (xD) multiplies the k -th term of C_0 by mk , and

$$\frac{mk}{m^{mk}(k!)^m} x^{mk} = \frac{x \cdot x^{mk-1}}{m^{mk-1}(k-1)!(k!)^{m-1}},$$

which after the shift $k \mapsto k+1$ is x times the k -th term of C_1 . For $1 \leq r \leq m-1$: $(xD + r)$ multiplies the k -th term of C_r by $(mk + m - r) + r = m(k+1)$, and

$$\frac{m(k+1)}{m^{mk+m-r}(k!)^r ((k+1)!)^{m-r}} x^{mk+m-r} = \frac{x \cdot x^{mk+m-(r+1)}}{m^{mk+m-(r+1)}(k!)^{r+1} ((k+1)!)^{m-(r+1)}},$$

which is x times the k -th term of C_{r+1} when $r+1 \leq m-1$, and—since the case $r+1 = m$ reads $x x^{mk}/(m^{mk}(k!)^m)$ —is x times the k -th term of C_0 when $r = m-1$. The identification with ${}_0F_{m-1}$ follows by rewriting the factorials as Pochhammer symbols. The eigenvalue equation is (26). $\square$

In the notation of Delerue's hyper-Bessel functions $I_{\nu_1, \dots, \nu_{m-1}}^{(m-1)}$ [9, 13], we have $C_r^{(m)} = I_{(0, \dots, 0, 1, \dots, 1)}^{(m-1)}$ with $r-1$ zeros and $m-r$ ones. The carrier expansion theorem now holds verbatim:

Theorem 29 (General carrier expansion). *Let $(f_0, \dots, f_{m-1})$ be any carrier system of modulus m for xD . Then*

$$(xD)^n f_0(x) = \sum_{k=0}^n \left[\begin{matrix} n \\ k \end{matrix} \right]_m x^k f_{k \bmod m}(x), \quad f_0(xe^t) = \sum_{n=0}^{\infty} \frac{t^n}{n!} \sum_{k=0}^n \left[\begin{matrix} n \\ k \end{matrix} \right]_m x^k f_{k \bmod m}(x).$$

Proof. Identical to that of Theorem 8: since xD is a derivation, $(xD)(x^k f_r) = (k-r)x^k f_r + x^{k+1} f_{(r+1) \bmod m}$ for $r = k \bmod m$, and $k-r = m \lfloor k/m \rfloor$; the coefficients therefore evolve by (29). $\square$

7.3. A general Dobiński formula. Composing the primary hyper-Bessel carrier with the exponential produces a distinguished solution of the m -Bell ODE, whose Maclaurin coefficients generalize the Dobiński series—and the coefficients of the carriers in those coefficients are precisely the primitive m -Bell numbers.

Theorem 30 (Generalized Dobiński formula). *Fix $m \geq 1$ and write $C_r = C_r^{(m)}$. The function $\mathcal{A}(x) = C_0(e^x)$ satisfies $\mathcal{A}^{(m)} = e^{mx} \mathcal{A}$, and its Maclaurin coefficients*

$$S_m(n) := \sum_{k=0}^{\infty} \frac{(mk)^n}{(m^k k!)^m}$$

satisfy

$$S_m(n) = \sum_{r=0}^{m-1} B_n^{(m,r)} C_r(1),$$

where $B^{(m,r)}$ are the primitive m -Bell sequences. For $m=1$ this is Dobiński's formula $\sum_k k^n/k! = e B_n$; for $m=2$ it is Theorem 3.

Proof. Expanding the series of C_0 at e^x and swapping summation order (absolute convergence),

$$C_0(e^x) = \sum_{k=0}^{\infty} \frac{e^{mkx}}{m^{mk} (k!)^m} = \sum_{n=0}^{\infty} \frac{x^n}{n!} \sum_{k=0}^{\infty} \frac{(mk)^n}{m^{mk} (k!)^m} = \sum_{n=0}^{\infty} S_m(n) \frac{x^n}{n!}.$$

On the other hand, setting $x=1$ in Theorem 29 and using $e^{txD} f(x)|_{x=1} = f(e^t)$:

$$C_0(e^t) = \sum_{n=0}^{\infty} \frac{t^n}{n!} \sum_{k=0}^n \left[\begin{matrix} n \\ k \end{matrix} \right]_m C_{k \bmod m}(1) = \sum_{n=0}^{\infty} \frac{t^n}{n!} \sum_{r=0}^{m-1} \rho_r(n) C_r(1),$$

where $\rho_r(n)$ are the residue-class row sums. Comparing coefficients and applying Corollary 26(i) gives the result. That $\mathcal{A} = C_0(e^x)$ solves the ODE follows from Propositions 27 and 28. $\square$

Remark 31. The solution space of $\mathcal{A}^{(m)} = e^{mx} \mathcal{A}$ is m -dimensional and is spanned by the e.g.f.s of the m primitive sequences. For $m=2$ we exhibited this basis explicitly through I_0 and K_0 (Theorem 2), and the Wronskian identity let us cancel the carriers and certify integer coefficients (Theorem 9). For $m \geq 3$, the point $t=0$ of (32) is a regular singular point whose indicial polynomial has the single root 0 with multiplicity m ; hence $C_0(t)$ is the unique solution analytic at the origin (up to scaling), and the remaining $m-1$ Frobenius solutions involve powers of $\log t$ —these are hyper-Bessel functions of the second kind [13]. Deriving explicit weights, analogous to those of Theorem 2, that combine these singular solutions into the individual primitive e.g.f.s is left as an open problem; note that by Corollary 26 the integrality of all coefficients is already established without this analytic machinery.

Remark 32. The primitive/composite decomposition has a classical constant-coefficient shadow: replacing the operator equation $\text{BINOM}^m a = L^m a$ by $a = L^m a$ (i.e. dropping the transform) yields the ODE $\mathcal{A}^{(m)} = \mathcal{A}$, whose primitive solutions are the m -sections of the exponential function studied by Tauber [21]. The m -Bell theory developed here can thus be viewed as the “binomial-transform deformation” of the theory of exponential sections.

7.4. Duals, factorials, and combinatorics for general m . All the structures of Sections 5 and 6 generalize from $m = 2$ to arbitrary m , with the same proofs. We record the statements.

Definition 33. For $m \geq 1$ define:

- (i) the unsigned m -Stirling numbers of the first kind by $\left\| \begin{smallmatrix} 0 \\ k \end{smallmatrix} \right\|_m = \delta_{0,k}$ and

$$\left\| \begin{smallmatrix} n+1 \\ k \end{smallmatrix} \right\|_m = m \lfloor n/m \rfloor \left\| \begin{smallmatrix} n \\ k \end{smallmatrix} \right\|_m + \left\| \begin{smallmatrix} n \\ k-1 \end{smallmatrix} \right\|_m;$$

- (ii) the m -falling and m -rising factorials

$$x^{\underline{n}_m} = \prod_{k=0}^{n-1} (x - m \lfloor k/m \rfloor), \quad x^{\overline{n}_m} = \prod_{k=0}^{n-1} (x + m \lfloor k/m \rfloor),$$

in which each factor repeats m times (except possibly the last); when m is clear from context we drop the subscript, as we did for $m = 2$;

- (iii) the m -Lah numbers $\left\| \begin{smallmatrix} n \\ k \end{smallmatrix} \right\|_m$ by $x^{\overline{n}_m} = \sum_k \left\| \begin{smallmatrix} n \\ k \end{smallmatrix} \right\|_m x^{\underline{k}_m}$.

Theorem 34. For every $m \geq 1$ and all $0 \leq k \leq n$, with weights $w_j = m \lfloor j/m \rfloor$:

- (i) $x^{\overline{n}_m} = \sum_k \left\| \begin{smallmatrix} n \\ k \end{smallmatrix} \right\|_m x^k$, $x^{\underline{n}_m} = \sum_k (-1)^{n-k} \left\| \begin{smallmatrix} n \\ k \end{smallmatrix} \right\|_m x^k$, $x^n = \sum_k \left\| \begin{smallmatrix} n \\ k \end{smallmatrix} \right\|_m x^{\underline{k}_m}$;
- (ii) the signed first-kind and second-kind triangles are inverse matrices;
- (iii) $\left\| \begin{smallmatrix} n \\ k \end{smallmatrix} \right\|_m = e_{n-k}(w_0, \dots, w_{n-1})$ and $\left\| \begin{smallmatrix} n \\ k \end{smallmatrix} \right\|_m = h_{n-k}(w_1, \dots, w_k)$, and the column generating function of the second kind is $\sum_n \left\| \begin{smallmatrix} n \\ k \end{smallmatrix} \right\|_m x^n = x^k / \prod_{j=1}^k (1 - w_j x)$;
- (iv) $\left\| \begin{smallmatrix} n \\ k \end{smallmatrix} \right\|_m = \sum_j \left\| \begin{smallmatrix} n \\ j \end{smallmatrix} \right\|_m \left\| \begin{smallmatrix} j \\ k \end{smallmatrix} \right\|_m$, with recurrence

$$\left\| \begin{smallmatrix} n+1 \\ k \end{smallmatrix} \right\|_m = (m \lfloor n/m \rfloor + m \lfloor k/m \rfloor) \left\| \begin{smallmatrix} n \\ k \end{smallmatrix} \right\|_m + \left\| \begin{smallmatrix} n \\ k-1 \end{smallmatrix} \right\|_m;$$

- (v) the row sums of the first kind equal $\prod_{j=0}^{n-1} (m \lfloor j/m \rfloor + 1)$;
- (vi) $\left\| \begin{smallmatrix} n \\ k \end{smallmatrix} \right\|_m$ counts the permutations of $[n]$ with k cycles built by the restricted insertion process in which element $i+1$ either opens a new cycle or is inserted after one of the $m \lfloor i/m \rfloor$ smallest existing elements.

Proof. Identical, word for word, to the proofs of Theorems 15, 16 and 22, Propositions 17, 18 and 20 (first equality), and Theorem 19, with $2 \lfloor \cdot/2 \rfloor$ replaced by $m \lfloor \cdot/m \rfloor$ throughout. $\square$

Theorem 35 (The m -compartment urn model). $\left\| \begin{smallmatrix} n \\ k \end{smallmatrix} \right\|_m$ counts the arrangements of n labeled marbles, placed one at a time, into k compartments distributed across urns with m unlabeled compartments each, under the constraints of Theorem 12 (*mutatis mutandis*): within an urn, a compartment may receive a second marble only after every compartment of that urn is non-empty, and a new urn may be opened only when all previously used urns are completely non-empty. Consequently the composite m -Bell number $B_n^{(m)}$ counts all such arrangements of n marbles, and the primitive $B_n^{(m,r)}$ counts those in which the number of used compartments is $\equiv r \pmod{m}$.

Proof. As in Theorem 12: if $k = mj + r$ with $0 \leq r < m$ compartments are in use, then j urns are complete and one urn has r used compartments; the marbles in the incomplete urn's used compartments are frozen, so exactly $mj = m \lfloor k/m \rfloor$ old compartments can receive the next marble, while opening the next compartment (in the incomplete urn if $r > 0$, or in a fresh urn if $r = 0$) is the unique alternative. The final two claims follow from Corollary 26. $\square$

The succession rule of Appendix C also generalizes directly, encoding the recurrence as a generating tree:

Definition 36 (Succession rule for m -Stirling numbers of the second kind).

$$\Omega_m : \begin{cases} (0) \\ (mj + r) \rightarrow (mj + r)^{mj}, (mj + r + 1) \end{cases} \quad \text{for } 0 \leq r < m,$$

i.e. a node labeled (k) produces $m \lfloor k/m \rfloor$ children with its own label and one child labeled $(k+1)$. The number of nodes labeled (k) at level n equals $\llbracket \frac{n}{k} \rrbracket_m$, and the total number of nodes at level n equals $B_n^{(m)}$.

This representation further reinforces the structural unity of the m -Stirling family and provides an efficient way to generate and analyze these numbers through tree-based methods.

8. MOMENTS OF THE CONWAY–MAXWELL–POISSON DISTRIBUTION

Perhaps the most intriguing consequence of the present theory is a probabilistic one. The classical Bell–Stirling framework is inseparable from the Poisson distribution: if $X \sim \text{Poisson}(y)$, then the factorial moments are $\mathbb{E}[X^{\underline{n}}] = y^n$, the raw moments are the Touchard polynomials $\mathbb{E}[X^n] = T_n(y)$, and at $y = 1$ Dobinski's formula (10) states precisely that $\mathbb{E}[X^n] = B_n$. In this section we show that the entire package generalizes: the m -Bell and m -Stirling numbers stand in exactly the same relation to the *Conway–Maxwell–Poisson* (CMP) distribution with integer dispersion parameter $\nu = m$.

The CMP distribution [6, 18] is a two-parameter generalization of the Poisson distribution with probability mass function

$$\mathbb{P}(X = k) = \frac{\lambda^k}{(k!)^\nu Z(\lambda, \nu)}, \quad k = 0, 1, 2, \dots, \quad Z(\lambda, \nu) = \sum_{j=0}^{\infty} \frac{\lambda^j}{(j!)^\nu},$$

where $\lambda > 0$ and $\nu \geq 0$. The case $\nu = 1$ is the Poisson distribution; $\nu > 1$ models under-dispersion and $\nu < 1$ over-dispersion, which has made the distribution a popular tool in applied count modeling [18]. Our first observation is that at integer $\nu = m$ and with the parametrization $\lambda = (y/m)^m$, the normalizing constant is exactly the primary hyper-Bessel carrier of Proposition 28:

$$(33) \quad Z\left(\left(\frac{y}{m}\right)^m, m\right) = \sum_{j=0}^{\infty} \frac{(y/m)^{mj}}{(j!)^m} = C_0^{(m)}(y).$$

For $m = 2$ this is the familiar fact that $Z(\lambda, 2) = I_0(2\sqrt{\lambda})$ [18, 7].

It is well known (and immediate) that the CMP distribution has clean *powered* factorial moments:

Lemma 37. *If $X \sim \text{CMP}(\lambda, \nu)$ then $\mathbb{E}[(X^{\underline{j}})^\nu] = \lambda^j$ for all $j \geq 0$.*

$$\text{Proof.} \quad \sum_{k \geq j} \left(\frac{k!}{(k-j)!} \right)^\nu \frac{\lambda^k}{(k!)^\nu Z} = \frac{\lambda^j}{Z} \sum_{i \geq 0} \frac{\lambda^i}{(i!)^\nu} = \lambda^j. \quad \square$$

For $\nu = 1$, Lemma 37 is the Poisson factorial-moment identity $\mathbb{E}[X^{\underline{j}}] = \lambda^j$. The key discovery is that for $\nu = m$, the correct generalization of the falling factorial of the variable itself (rather than a power of it) is precisely the m -falling factorial of Definition 33, evaluated at mX —and its expectation picks up exactly one hyper-Bessel carrier ratio.

Theorem 38 (m -falling moments of the CMP distribution). *Let $m \geq 1$, $y > 0$, and $X \sim \text{CMP}((y/m)^m, m)$. Then for all $n \geq 0$, with $r = n \bmod m$,*

$$\mathbb{E}[(mX)^{\underline{n}_m}] = y^n \frac{C_r^{(m)}(y)}{C_0^{(m)}(y)}.$$

In particular, for $m = 1$: $\mathbb{E}[X^{\underline{n}}] = y^n$ (Poisson); and for $m = 2$:

$$\mathbb{E}[(2X)^{\underline{n}}] = \begin{cases} y^n & n \text{ even,} \\ y^n \frac{I_1(y)}{I_0(y)} & n \text{ odd.} \end{cases}$$

Proof. Write $n = qm + r$ with $0 \leq r < m$. In the product $(mX)^{\underline{n}_m} = \prod_{k=0}^{n-1} (mX - k)$, the factor $(mX - mi)$ appears m times for each $0 \leq i \leq q-1$ and the factor $(mX - mq)$ appears r times, so

$$(mX)^{\underline{n}_m} = m^n (X^{\underline{q}})^m (X - q)^r.$$

Taking expectations with $\lambda = (y/m)^m$ and $Z = C_0(y)$ (by (33)):

$$\mathbb{E}[(mX)^n] = \frac{m^n}{C_0(y)} \sum_{k \geq q} \left(\frac{k!}{(k-q)!} \right)^m (k-q)^r \frac{\lambda^k}{(k!)^m} = \frac{m^n \lambda^q}{C_0(y)} \sum_{i \geq 0} \frac{i^r \lambda^i}{(i!)^m}.$$

For $r = 0$ the remaining sum is $Z = C_0(y)$ and the result is $m^n \lambda^q = y^n$. For $1 \leq r \leq m-1$, use $i^r/(i!)^m = 1/(((i-1)!)^r (i!)^{m-r})$ for $i \geq 1$ and shift $i = k+1$:

$$\sum_{i \geq 1} \frac{i^r \lambda^i}{(i!)^m} = \sum_{k \geq 0} \frac{(y/m)^{m(k+1)}}{(k!)^r ((k+1)!)^{m-r}} = \left(\frac{y}{m} \right)^r C_r(y),$$

by the series in Proposition 28. Hence the expectation equals $m^n \lambda^q (y/m)^r C_r(y)/C_0(y) = y^{mq+r} C_r(y)/C_0(y) = y^n C_r(y)/C_0(y)$. $\square$

Theorem 39 (Raw moments of the CMP distribution). *With the assumptions of Theorem 38,*

$$m^n \mathbb{E}[X^n] = \frac{1}{C_0^{(m)}(y)} \sum_{k=0}^n \left[\begin{matrix} n \\ k \end{matrix} \right]_m y^k C_{k \bmod m}^{(m)}(y) = \frac{V_n(y)}{C_0^{(m)}(y)},$$

where V_n is the carrier-Touchard polynomial of Theorem 29 for the hyper-Bessel system. For $m = 1$ this is the classical $\mathbb{E}[X^n] = T_n(y)$, the Touchard polynomial evaluated at the rate.

Proof. Apply the basis conversion of Theorem 34(i) to the polynomial identity $(mX)^n = \sum_k \left[\begin{matrix} n \\ k \end{matrix} \right]_m (mX)^k$, take expectations, and insert Theorem 38. Alternatively and equivalently, $\mathbb{E}[(mX)^n] = \frac{1}{C_0(y)} \sum_k (mk)^n \lambda^k / (k!)^m = \frac{1}{C_0(y)} [(xD)^n C_0](y)$, and Theorem 29 expands the right-hand side. $\square$

As a check, the case $n = 1$, $m = 2$ gives the known mean of the CMP distribution with $\nu = 2$ [7]: $\mathbb{E}[X] = \frac{y}{2} I_1(y)/I_0(y) = \sqrt{\lambda} I_1(2\sqrt{\lambda})/I_0(2\sqrt{\lambda})$.

Corollary 40 (m -Bell numbers as CMP moments). *Let $X \sim \text{CMP}(m^{-m}, m)$, i.e. $y = 1$. Then*

$$m^n \mathbb{E}[X^n] = \sum_{r=0}^{m-1} B_n^{(m,r)} \frac{C_r^{(m)}(1)}{C_0^{(m)}(1)} = \frac{S_m(n)}{C_0^{(m)}(1)}.$$

For $m = 1$ ($X \sim \text{Poisson}(1)$) this is $\mathbb{E}[X^n] = B_n$; for $m = 2$ ($X \sim \text{CMP}(1/4, 2)$),

$$2^n \mathbb{E}[X^n] = B_n^{(2,0)} + B_n^{(2,1)} \frac{I_1(1)}{I_0(1)} = A351143(n) + A351028(n) \cdot \frac{I_1(1)}{I_0(1)}.$$

Proof. Set $y = 1$ in Theorem 39 and group the terms by $k \bmod m$, applying Corollary 26; the second expression restates Theorem 30. $\square$

Thus the scaled moments of the $\text{CMP}(m^{-m}, m)$ distribution are $\mathbb{Z}$ -linear combinations of the fixed “Bessel quotients” $C_r(1)/C_0(1)$, and the integer coefficients are exactly the primitive m -Bell numbers. In this precise sense, the Conway–Maxwell–Poisson distribution is the natural probabilistic home of the m -Bell numbers, just as the Poisson distribution is for the Bell numbers. It seems fitting that a distribution introduced for queuing systems with state-dependent service rates [6], and revived for fitting under- and over-dispersed count data [18], turns out to encode a natural generalization of the partition-counting apparatus of enumerative combinatorics.

9. FINAL REMARKS

We conclude with remarks situating the m -Stirling framework within the literature, and a list of open problems. Appendix D presents a reference table cataloguing the parallels between the classical theory, the Bessel case $m = 2$, and general m .

9.1. Relation to other generalizations of Stirling numbers. Triangular arrays defined by two-term recurrences of the form $T(n+1, k) = f(n, k)T(n, k) + T(n, k-1)$ have been studied at great generality—as “Galton arrays” by Neuwirth [15], through bivariate generating function PDEs by Barbero et al. [1], and via explicit-formula machinery by Spivey [19] and Mansour, Mulay and Shattuck [14]. The m -Stirling triangles fit these frameworks formally but sit outside their sharpest results: the coefficient $m \lfloor k/m \rfloor$ is not of the affine form $\alpha + \beta n + \gamma k$ covered by the complete classification in [1], and the general explicit formula of [14] requires the column weights b_k to be pairwise distinct, whereas ours occur with multiplicity m —this is exactly why the partial-fraction analysis in Appendix A involves higher-order poles and produces the harmonic-number corrections in Theorem 11. The m -Stirling numbers are also distinct from Broder’s r -Stirling numbers [3] (which restrict where the first r elements sit) and from the Bessel numbers of Cheon et al. [4] (connection coefficients of Bessel polynomials); the appearance of Bessel functions in our $m = 2$ case is structural (the e.g.f. ODE), not definitional.

9.2. The sequence A000246 and a missing bijection. Proposition 20 shows that the row sums of the Bessel–Stirling triangle of the first kind count the permutations of odd order, while Theorem 19 exhibits the triangle itself as counting parity-admissible permutations by cycles. Since the two 9-element sets for $n = 4$ already differ, the equinumerosity is genuinely non-trivial at the level of the statistic. A bijection between odd-order permutations and parity-admissible permutations—ideally one translating a natural statistic on odd-order permutations into the cycle count of the admissible ones—would give a fully combinatorial account of Table 4.

9.3. Open problems.

Problem 41. Find a direct bijective proof of the shift identity (30) in the urn model of Theorem 35: the right-hand side suggests distinguishing, in a placement of $n + m$ marbles, the j marbles that avoid the “last” urn, with the factor m^{n-j} recording an m -fold choice for each of the remaining marbles. Such a proof would make the “pairing effect” heuristic of Section 5 precise.

Problem 42. Determine explicit closed forms, analogous to Theorem 2, for the e.g.f.s of the individual primitive m -Bell sequences when $m \geq 3$, in terms of hyper-Bessel functions of the second kind (Remark 31).

Problem 43. The Bell numbers satisfy the Touchard congruence $B_{p+n} \equiv B_n + B_{n+1} \pmod{p}$. The m -Stirling triangles show visible m -adic structure (for instance $\lfloor \lfloor n/m \rfloor \rfloor_m = m^{n-m}$ for $n \geq m$, by the column generating function); a systematic study of congruence properties of the m -Bell sequences is open.

Problem 44. The Hankel determinants of the Bell numbers are the superfactorials $\prod_{i=0}^n i!$. The Hankel determinants of A007472 begin $1, 0, -4, -48, 0, 442368, \dots$: they vanish infinitely often and alternate in sign irregularly, so the sequence is not a classical moment sequence, yet Corollary 40 exhibits it as an integer part of a moment sequence. Explaining this structure (e.g. via continued fractions or orthogonal polynomials for the CMP weight) is open.

Problem 45. Develop q -analogues: both the urn model (Theorem 35) and the file-placement interpretation (Proposition 18) suggest natural inversion statistics, and the Garsia–Remmel q -rook framework applies to the doubled staircase boards.

Problem 46. Determine the asymptotics of $B_n^{(m)}$ for fixed $m \geq 2$, generalizing the Moser–Wyman asymptotics of the Bell numbers; the Dobiński-type series of Theorem 30 is the natural starting point for a saddle-point analysis.

REFERENCES

- [1] J. Fernando Barbero G., Jesús Salas, and Eduardo J. S. Villaseñor. “Bivariate generating functions for a class of linear recurrences: General structure”. In: *Journal of Combinatorial Theory, Series A* 125 (July 2014), pp. 146–165. ISSN: 0097-3165. DOI: 10.1016/j.jcta.2014.02.007. URL: <https://www.sciencedirect.com/science/article/pii/S0097316514000272> (visited on 04/07/2025).

- [2] M. Bernstein and N.J.A. Sloane. “Some canonical sequences of integers”. In: *Linear Algebra and its Applications* 226-228 (Sept. 1995), pp. 57–72. DOI: 10.1016/0024-3795(94)00245-9. URL: <https://linkinghub.elsevier.com/retrieve/pii/0024379594002459>.
- [3] Andrei Z. Broder. “The r-Stirling numbers”. In: *Discrete Mathematics* 49.3 (1984), pp. 241–259. DOI: 10.1016/0012-365X(84)90161-4.
- [4] Gi-Sang Cheon, Ji-Hwan Jung, and Louis W. Shapiro. “Generalized Bessel Numbers and Some Combinatorial Settings”. In: *Discrete Mathematics* 313.20 (Oct. 2013), pp. 2127–2138. ISSN: 0012365X. DOI: 10.1016/j.disc.2013.05.001. URL: <https://linkinghub.elsevier.com/retrieve/pii/S0012365X13002094> (visited on 05/03/2025).
- [5] Louis Comtet. *Advanced Combinatorics*. DOI: 10.1007/978-94-010-2196-8. Dordrecht: Springer Netherlands, 1974. URL: <http://link.springer.com/10.1007/978-94-010-2196-8>.
- [6] Richard W. Conway and William L. Maxwell. “A queuing model with state dependent service rates”. In: *Journal of Industrial Engineering* 12 (1962), pp. 132–136.
- [7] Fraser Daly and Robert E. Gaunt. “The Conway–Maxwell–Poisson distribution: distributional theory and approximation”. In: *ALEA, Latin American Journal of Probability and Mathematical Statistics* 13 (2016), pp. 635–658. DOI: 10.30757/ALEA.v13-25.
- [8] G. Dattoli et al. *Touchard like Polynomials and Generalized Stirling Numbers*. Oct. 28, 2010. DOI: 10.48550/arXiv.1010.5934. arXiv: 1010.5934 [math]. URL: <http://arxiv.org/abs/1010.5934> (visited on 04/13/2025). Pre-published.
- [9] Paul Delerue. “Sur le calcul symbolique à n variables et les fonctions hyperbesséliennes (II)”. fr. In: *Annales de la Société Scientifique de Bruxelles, Série I* 67 (1953), pp. 229–274.
- [10] *NIST Digital Library of Mathematical Functions*. <https://dlmf.nist.gov/>, Release 1.2.4 of 2025-03-15. F. W. J. Olver, A. B. Olde Daalhuis, D. W. Lozier, B. I. Schneider, R. F. Boisvert, C. W. Clark, B. R. Miller, B. V. Saunders, H. S. Cohl, and M. A. McClain, eds. URL: <https://dlmf.nist.gov/>.
- [11] Jay R. Goldman, J. T. Joichi, and Dennis E. White. “Rook theory. I. Rook equivalence of Ferrers boards”. In: *Proceedings of the American Mathematical Society* 52 (1975), pp. 485–492. DOI: 10.1090/S0002-9939-1975-0429578-4.
- [12] Ronald L. Graham, Donald E. Knuth, and Oren Patashnik. *Concrete Mathematics: A Foundation for Computer Science*. en. 2nd. Reading, MA: Addison-Wesley, 1994.
- [13] Virginia Kiryakova. *Generalized Fractional Calculus and Applications*. Pitman Research Notes in Mathematics 301. Harlow: Longman Scientific & Technical, 1994.
- [14] Toufik Mansour, Shashikant Mulay, and Mark Shattuck. “A general two-term recurrence and its solution”. en. In: *European Journal of Combinatorics* 33.1 (Jan. 2012), pp. 20–26. ISSN: 01956698. DOI: 10.1016/j.ejc.2011.07.008. URL: <https://linkinghub.elsevier.com/retrieve/pii/S019566981100134X> (visited on 04/01/2025).
- [15] Erich Neuirth. “Recursively defined combinatorial functions: extending Galton’s board”. In: *Discrete Mathematics* 239.1 (Aug. 2001), pp. 33–51. ISSN: 0012-365X. DOI: 10.1016/S0012-365X(00)00373-3. URL: <https://www.sciencedirect.com/science/article/pii/S0012365X00003733> (visited on 04/06/2025).
- [16] John Riordan. “Inverse Relations and Combinatorial Identities”. In: *The American Mathematical Monthly* 71.5 (1964). Publisher: Mathematical Association of America, pp. 485–498. ISSN: 0002-9890. DOI: 10.2307/2312584. URL: <https://www.jstor.org/stable/2312584> (visited on 05/10/2025).
- [17] J. Sándor and B. Crstici. “Stirling, bell, bernoulli, euler and eulerian numbers”. In: ed. by J. Sándor and B. Crstici. DOI: 10.1007/1-4020-2547-5_5. Dordrecht: Springer Netherlands, 2004, pp. 459–618. URL: https://doi.org/10.1007/1-4020-2547-5_5.
- [18] Galit Shmueli et al. “A Useful Distribution for Fitting Discrete Data: Revival of the Conway–Maxwell–Poisson Distribution”. In: *Journal of the Royal Statistical*

- Society Series C: Applied Statistics* 54.1 (Jan. 1, 2005), pp. 127–142. ISSN: 0035-9254. DOI: 10.1111/j.1467-9876.2005.00474.x. URL: <https://doi.org/10.1111/j.1467-9876.2005.00474.x> (visited on 04/13/2025).
- [19] Michael Z. Spivey. “On Solutions to a General Combinatorial Recurrence”. In: *J. Integer Seq* 14.9 (2011). URL: <http://ftp.gwdg.de/pub/EMIS/journals/JIS/VOL14/Spivey/spivey31.pdf> (visited on 04/06/2025).
- [20] Michael Z. Spivey and Laura L. Steil. “The K-Binomial Transforms and the Hankel Transform”. In: *J. Integer Seq* 9.1 (2006), p. 19. URL: <https://cs.uwaterloo.ca/journals/JIS/VOL9/Spivey/spivey7.pdf> (visited on 05/04/2025).
- [21] S. Tauber. “On Generalizations of the Exponential Function”. In: *The American Mathematical Monthly* 67.8 (1960), pp. 763–767. ISSN: 0002-9890. DOI: 10.2307/2308653. JSTOR: 2308653. URL: <https://www.jstor.org/stable/2308653> (visited on 04/03/2025).
- [22] Eric W. Weisstein. *Bell Polynomial*. Publisher: Wolfram Research, Inc. URL: <https://mathworld.wolfram.com/BellPolynomial.html>.
- [23] Julian West. “Generating trees and the Catalan and Schröder numbers”. en. In: *Discrete Mathematics* 146.1-3 (Nov. 1995), pp. 247–262. ISSN: 0012365X. DOI: 10.1016/0012-365X(94)00067-1. URL: <https://linkinghub.elsevier.com/retrieve/pii/0012365X94000671> (visited on 05/04/2025).
- [24] Herbert S. Wilf. *Generatingfunctionology: Third Edition*. CRC Press, Dec. 20, 2005. 254 pp. ISBN: 978-1-4398-6439-5. Google Books: bVFuBwAAQBAJ.

APPENDIX A: EXPLICIT FORMULA FOR THE BESSEL-STIRLING NUMBERS (PROOF OF THEOREM 11)

Let A_k be the ordinary generating function for the k -th column over x^n :

$$A_k(x) = \sum_{n \geq 0} \left[\begin{matrix} n \\ k \end{matrix} \right]_2 x^n.$$

We first multiply each side of the recurrence (23) by x^n and sum over n :

$$\begin{aligned} \sum_{n \geq 1} \left[\begin{matrix} n \\ k \end{matrix} \right]_2 x^n &= 2 \lfloor k/2 \rfloor \sum_{n \geq 1} \left[\begin{matrix} n-1 \\ k \end{matrix} \right]_2 x^n + \sum_{n \geq 1} \left[\begin{matrix} n-1 \\ k-1 \end{matrix} \right]_2 x^n \\ A_k - \left[\begin{matrix} 0 \\ k \end{matrix} \right]_2 &= 2 \lfloor k/2 \rfloor x A_k + x A_{k-1}, \quad k \geq 1. \end{aligned}$$

Since $\left[\begin{matrix} 0 \\ k \end{matrix} \right]_2 = 0$ when $k \neq 0$, this simplifies to the following recurrence for the column generating function:

$$A_k = \frac{x}{(1 - 2 \lfloor k/2 \rfloor x)} A_{k-1}.$$

We know from the initial conditions that $A_0(x) = 1$, hence:

$$\begin{aligned} A_0 &= 1 \\ A_1 &= x \\ A_2 &= \frac{x^2}{1 - 2x} \\ A_3 &= \frac{x^3}{(1 - 2x)^2} \\ A_4 &= \frac{x^4}{(1 - 2x)^2(1 - 4x)} \\ A_5 &= \frac{x^5}{(1 - 2x)^2(1 - 4x)^2}, \end{aligned}$$

and so on. More generally, the ordinary generating function for the k -th column is

$$(34) \quad A_k(x) = \frac{x^k}{\prod_{j=1}^k (1 - 2 \lfloor j/2 \rfloor x)},$$

which is Equation (27) of the main text. In order to get an explicit formula for $\left[\begin{matrix} n \\ k \end{matrix} \right]_2$, use the technique on p. 19 of [24]:

$$\left[\begin{matrix} n \\ k \end{matrix} \right]_2 = [x^n] A_k(x),$$

where $[x^n]$ is the coefficient extraction operator. Start with a partial fraction decomposition of $A_k(x)$. First, denote

$$A_k(x) = x^k P_k(x),$$

where

$$P_k(x) = \prod_{1 \leq j \leq k} \frac{1}{1 - 2 \lfloor j/2 \rfloor x}.$$

Hence

$$(35) \quad \left[\begin{matrix} n \\ k \end{matrix} \right]_2 = [x^n] x^k P_k(x) = [x^{n-k}] P_k(x).$$

We can express $P_k(x)$ as a sum of partial fractions with unknown coefficients (note the slight difference in ranges for odd and even k):

$$(36) \quad \begin{aligned} P_{2m}(x) &= \sum_{1 \leq j \leq m} \frac{c_{2m,j}}{1 - 2jx} + \sum_{1 \leq j \leq m-1} \frac{d_{2m,j}}{(1 - 2jx)^2} \\ P_{2m+1}(x) &= \sum_{1 \leq j \leq m} \frac{c_{2m+1,j}}{1 - 2jx} + \sum_{1 \leq j \leq m} \frac{d_{2m+1,j}}{(1 - 2jx)^2} \end{aligned}$$

It is easy to calculate the d coefficients with the standard Heaviside cover-up method:

$$\begin{aligned}
 d_{k,r} &= (1 - 2rx)^2 P_k(x) \Big|_{x=\frac{1}{2r}} \\
 &= (1 - 2rx)^2 \prod_{1 \leq j \leq k} \frac{1}{1 - 2 \lfloor j/2 \rfloor x} \Big|_{x=\frac{1}{2r}} \\
 &= \prod_{1 \leq j < 2r} \frac{1}{1 - \lfloor j/2 \rfloor / r} \prod_{2r < j \leq k} \frac{1}{1 - \lfloor j/2 \rfloor / r} \\
 &= \prod_{1 \leq j < 2r} \frac{r}{r - \lfloor j/2 \rfloor} \prod_{2r < j \leq k} \frac{r}{r - \lfloor j/2 \rfloor},
 \end{aligned}$$

which simplifies to

$$(37) \quad d_{k,r} = \frac{(-r)^{k-1}}{r! r! \left(\lfloor \frac{k}{2} \rfloor - r \right)! \left(\lfloor \frac{k-1}{2} \rfloor - r \right)!}.$$

$d_{k,r}$ varies only slightly for even and odd k :

$$\begin{aligned}
 d_{2m,r} &= \frac{-r^{2m-1}}{r! r! (m-r)! (m-1-r)!} \\
 d_{2m+1,r} &= \frac{r^{2m}}{r! r! (m-r)! (m-r)!}
 \end{aligned}$$

For the $c_{k,j}$ coefficients, we have two main cases to consider. When $k = 2m$ is even and $j = m$, $P_k(x)$ has a simple pole at $x = \frac{1}{2m}$, so we get $c_{2m,m}$ easily by the same method:

$$(38) \quad c_{2m,m} = (1 - 2mx) P_{2m}(x) \Big|_{x=\frac{1}{2m}} = \frac{m^{2m}}{m! m!}$$

For the remaining c coefficients we have a pole of $P_k(x)$ of multiplicity 2 at $x = \frac{1}{2j}$, so we need to use the derivative method. Specifically:

$$-2r c_{k,r} = \frac{d}{dx} (1 - 2rx)^2 P_k(x) \Big|_{x=\frac{1}{2r}}$$

Let

$$\begin{aligned}
 Q_r &= (1 - 2rx)^2 P_k(x) = \prod_j \frac{1}{(1 - 2jx)^{e_j}}, \\
 e_j &= \begin{cases} 2, & \text{if } 1 \leq 2j < k \wedge j \neq r \\ 1, & \text{if } 2j = k \wedge j \neq r \\ 0, & \text{otherwise} \end{cases}
 \end{aligned}$$

Then $c_{k,r} = -\frac{1}{2r} Q'_r \left(\frac{1}{2r} \right)$, and we can use the logarithmic derivative to simplify the calculation:

$$\begin{aligned}
 \frac{Q'_r}{Q_r} &= \frac{d}{dx} \log(Q_r) = \frac{d}{dx} \left[- \sum_j e_j \log(1 - 2jx) \right] = \sum_j 2e_j \frac{j}{1 - 2jx} \\
 c_{k,r} &= -\frac{1}{2r} Q'_r \left(\frac{1}{2r} \right) = -\frac{1}{2r} Q_r \left(\frac{1}{2r} \right) 2 \sum_j e_j \frac{j}{1 - j/r} \\
 &= -\frac{1}{2r} 2d_{k,r} \sum_j e_j \frac{rj}{r-j} = -d_{k,r} \sum_j e_j \frac{j}{r-j}
 \end{aligned}$$

Let $s_k = \sum_j e_j \frac{j}{r-j}$ and simplify. First, consider the case $k = 2m + 1$:

$$\begin{aligned}
s_{2m+1} &= 2 \sum_{\substack{1 \leq j \leq m \\ j \neq r}} \frac{j}{r-j} \\
&= 2 \left[\frac{1}{r-1} + \frac{2}{r-2} + \cdots + \frac{r-1}{1} - \frac{r+1}{1} - \frac{r+2}{2} - \cdots - \frac{m}{m-r} \right] \\
&= 2 \sum_{1 \leq j \leq r-1} \frac{r-j}{j} - 2 \sum_{1 \leq j \leq m-r} \frac{j+r}{j} \\
&= 2r \left(\sum_{1 \leq j \leq r-1} \frac{1}{j} \right) - 2(r-1) - 2r \left(\sum_{1 \leq j \leq m-r} \frac{1}{j} \right) - 2(m-r) \\
&= 2r(H_{r-1} - H_{m-r}) - 2m + 2,
\end{aligned}$$

where H_n is the n -th harmonic number. We proceed similarly for $k = 2m$, with some care about $e_m = 1$:

$$\begin{aligned}
s_{2m} &= \sum_j e_j \frac{j}{r-j} = \frac{m}{r-m} + 2 \sum_{\substack{1 \leq j \leq m-1 \\ j \neq r}} \frac{j}{r-j} \\
&= \frac{m}{r-m} + 2 \sum_{1 \leq j \leq r-1} \frac{j}{r-j} - 2 \sum_{r+1 \leq j \leq m-1} \frac{j}{j-r} \\
&= \frac{m}{r-m} + 2 \sum_{1 \leq j \leq r-1} \frac{r-j}{j} - 2 \sum_{1 \leq j \leq m-r-1} \frac{j+r}{j} \\
&= \frac{m}{r-m} + 2rH_{r-1} - 2(r-1) - 2rH_{m-r-1} - 2(m-r-1) \\
&= 2rH_{r-1} - 2rH_{m-r-1} - 2(m-2) - \frac{m}{m-r} \\
&= 2rH_{r-1} - 2rH_{m-r-1} - 2(m-2) - \frac{r}{m-r} - 1 \\
&= r(2H_{r-1} - H_{m-r-1} - H_{m-r}) - 2m + 3
\end{aligned}$$

Generally, when $r \neq m$:

$$(39) \quad s_k = r(2H_{r-1} - H_{\lfloor k/2 \rfloor - r} - H_{\lfloor (k-1)/2 \rfloor - r}) - k + 3.$$

Combining everything we have:

$$\begin{aligned}
d_{k,r} &= \frac{(-r)^{k-1}}{r! r! (\lfloor k/2 \rfloor - r)! (\lfloor (k-1)/2 \rfloor - r)!} \\
s_{k,r} &= r(2H_{r-1} - H_{\lfloor k/2 \rfloor - r} - H_{\lfloor (k-1)/2 \rfloor - r}) - k + 3 \\
(40) \quad c_{k,r} &= \begin{cases} -d_{k,r} s_{k,r} & \text{if } 1 \leq r < k/2 \\ \frac{r^{2r}}{r! r!} & \text{if } r = k/2 \\ 0 & \text{otherwise} \end{cases}
\end{aligned}$$

Next, let us find an explicit formula for $\llbracket n \rrbracket_2$. From (35) and (36):

$$\begin{aligned}
\llbracket n \rrbracket_2 &= [x^{n-k}] P_k(x) \\
&= [x^{n-k}] \left[\sum_{1 \leq j \leq \lfloor k/2 \rfloor} \frac{c_{k,j}}{1-2jx} + \sum_{1 \leq j \leq \lfloor (k-1)/2 \rfloor} \frac{d_{k,j}}{(1-2jx)^2} \right] \\
&= \sum_{1 \leq j \leq \lfloor k/2 \rfloor} c_{k,j} [x^{n-k}] \frac{1}{1-2jx} + \sum_{1 \leq j \leq \lfloor (k-1)/2 \rfloor} d_{k,j} [x^{n-k}] \frac{1}{(1-2jx)^2}
\end{aligned}$$

We have the standard geometric series expansions:

$$\begin{aligned} [x^{n-k}] \frac{1}{1-2jx} &= [x^{n-k}] \sum_{i \geq 0} (2j)^i x^i = (2j)^{n-k} \\ [x^{n-k}] \frac{1}{(1-2jx)^2} &= [x^{n-k}] \sum_{i \geq 0} (i+1)(2j)^i x^i = (n-k+1)(2j)^{n-k}. \end{aligned}$$

This results in the following final set of formulas, valid for $n \geq 2, k \geq 2$:

$$\begin{aligned} \left[\begin{matrix} n \\ k \end{matrix} \right]_2 &= \sum_{1 \leq j \leq \lfloor \frac{k}{2} \rfloor} (2j)^{n-k} (c_{k,j} + d_{k,j}(n-k+1)) \\ (41) \quad d_{k,r} &= \frac{(-r)^{k-1}}{r! r! (\lfloor \frac{k}{2} \rfloor - r)! (\lfloor \frac{k-1}{2} \rfloor - r)!} \\ c_{k,r} &= \begin{cases} -d_{k,r} (r(2H_{r-1} - H_{\lfloor \frac{k}{2} \rfloor - r} - H_{\lfloor \frac{k-1}{2} \rfloor - r}) - k + 3) & \text{if } 1 \leq r < k/2 \\ \frac{r^{2r}}{r! r!} & \text{if } r = k/2 \\ 0 & \text{otherwise} \end{cases} \end{aligned}$$

This piecewise construction is necessary because the harmonic numbers have singularities at negative integer indices, which occur when $r \geq k/2$ or $r \leq 0$. This problem can be avoided by replacing the harmonic numbers via the identity $H_n = \psi(n+1) + \gamma$, where γ is the Euler–Mascheroni constant and $\psi(x)$ is the digamma function, and by replacing the factorials with the gamma function. Since we have digamma functions in the numerator and gamma functions in the denominator of the resulting expression, the limit of their ratio as the argument approaches non-positive integers is well defined. Specifically, using the residues of the digamma and gamma functions around the negative integer poles it can be shown that

$$\lim_{x \rightarrow -n} \frac{\psi(x)}{\Gamma(x)} = (-1)^{n+1} n!, \quad n \in \mathbb{N}^+.$$

After incorporating the coefficient definitions into the expression for $\left[\begin{matrix} n \\ k \end{matrix} \right]_2$ (41) and performing simplifications using properties of these special functions, we arrive at the limit representation stated in Theorem 11:

$$\begin{aligned} \left[\begin{matrix} n \\ k \end{matrix} \right]_2 &= (-1)^{k-1} 2^{n-k} \sum_{j=0}^{\lfloor k/2 \rfloor} \lim_{t \rightarrow j} \left[t^{n-1} \frac{n-2-t \Psi_k(t)}{\Gamma(t+1)^2 \Gamma(\lfloor \frac{k}{2} \rfloor - t + 1) \Gamma(\lfloor \frac{k-1}{2} \rfloor - t + 1)} \right], \\ \Psi_k(t) &= 2\psi(t) - \psi(\lfloor \frac{k}{2} \rfloor - t + 1) - \psi(\lfloor \frac{k-1}{2} \rfloor - t + 1), \end{aligned}$$

This formula provides a single, unified expression for $\left[\begin{matrix} n \\ k \end{matrix} \right]_2$ that is valid for all integers $n, k \geq 0$. The inclusion of the $j = 0$ term in the sum is essential, as the limit process correctly evaluates its contribution, ensuring validity for boundary cases like $\left[\begin{matrix} 0 \\ 0 \end{matrix} \right]_2 = \left[\begin{matrix} 1 \\ 1 \end{matrix} \right]_2 = 1$, and ensuring $\left[\begin{matrix} n \\ k \end{matrix} \right]_2 = 0$ for $k > n$. $\square$

APPENDIX B: BESSEL-STIRLING NUMBER IDENTITIES

In all cases below $b = 2 \lfloor k/2 \rfloor$ is used to simplify the notation.

Lemma 47 (Distant hockey-stick-like identity).

$$\left[\begin{matrix} n+j \\ k \end{matrix} \right]_2 = b^j \left[\begin{matrix} n \\ k \end{matrix} \right]_2 + \sum_{i=0}^{j-1} b^{j-1-i} \left[\begin{matrix} n+i \\ k-1 \end{matrix} \right]_2$$

Proof. The base case for $j = 1$ is the basic recurrence (23):

$$\left[\begin{matrix} n+1 \\ k \end{matrix} \right]_2 = b \left[\begin{matrix} n \\ k \end{matrix} \right]_2 + \left[\begin{matrix} n \\ k-1 \end{matrix} \right]_2.$$

Assume the identity holds for j (induction hypothesis). Then from the basic recurrence we have

$$\left[\begin{matrix} n+j+1 \\ k \end{matrix} \right]_2 = b \left[\begin{matrix} n+j \\ k \end{matrix} \right]_2 + \left[\begin{matrix} n+j \\ k-1 \end{matrix} \right]_2.$$

From the induction hypothesis:

$$\begin{aligned} \left[\begin{matrix} n+j+1 \\ k \end{matrix} \right]_2 &= b \left(b^j \left[\begin{matrix} n \\ k \end{matrix} \right]_2 + \sum_{i=0}^{j-1} b^{j-1-i} \left[\begin{matrix} n+i \\ k-1 \end{matrix} \right]_2 \right) + \left[\begin{matrix} n+j \\ k-1 \end{matrix} \right]_2 \\ &= b^{j+1} \left[\begin{matrix} n \\ k \end{matrix} \right]_2 + \sum_{i=0}^j b^{j-i} \left[\begin{matrix} n+i \\ k-1 \end{matrix} \right]_2, \end{aligned}$$

which completes the proof. $\square$

Proposition 48 (Hockey-stick-like identity).

$$\left[\begin{matrix} n \\ k \end{matrix} \right]_2 = \sum_{j=k}^n b^{n-j} \left[\begin{matrix} j-1 \\ k-1 \end{matrix} \right]_2$$

Proof. Set $n = 0$ in Lemma 47:

$$\left[\begin{matrix} j \\ k \end{matrix} \right]_2 = b^j \left[\begin{matrix} 0 \\ k \end{matrix} \right]_2 + \sum_{i=0}^{j-1} b^{j-1-i} \left[\begin{matrix} i \\ k-1 \end{matrix} \right]_2.$$

Due to the initial conditions $\left[\begin{smallmatrix} 0 \\ 0 \end{smallmatrix} \right]_2 = 1$, $\left[\begin{smallmatrix} n \\ 0 \end{smallmatrix} \right]_2 = \left[\begin{smallmatrix} 0 \\ k \end{smallmatrix} \right]_2 = 0$ ($n, k \geq 1$), the first term on the right-hand side vanishes for $k \geq 1$. Then we can simplify, rewriting the indices:

$$\left[\begin{matrix} n \\ k \end{matrix} \right]_2 = \sum_{j=0}^{n-1} b^{n-j-1} \left[\begin{matrix} j \\ k-1 \end{matrix} \right]_2,$$

and since $\left[\begin{smallmatrix} n \\ k \end{smallmatrix} \right]_2 = 0$ when $k > n$, we can start j from k and shift it by 1 to match the proposition's form:

$$\left[\begin{matrix} n \\ k \end{matrix} \right]_2 = \sum_{j=k}^n b^{n-j} \left[\begin{matrix} j-1 \\ k-1 \end{matrix} \right]_2. \quad \square$$

Both identities, with $b = m \lfloor k/m \rfloor$, hold verbatim for the general m -Stirling numbers of the second kind, by the same proofs.

APPENDIX C: SUCCESSION RULES AND A GENERATING TREE FOR BESSEL–STIRLING NUMBERS

The Bessel–Stirling numbers of the second kind can also be elegantly described using the formalism of succession rules and generating trees, as introduced by West [23]. This representation provides yet another perspective on the structure of these numbers and their recurrence relation.

Definition 49 (Succession rule for Bessel–Stirling numbers of the second kind). *The Bessel–Stirling triangle of the second kind can be generated using the following succession rule:*

$$\Omega : \begin{cases} (0) \\ (2k) \rightarrow (2k)^{2k}, (2k+1) \\ (2k+1) \rightarrow (2k+1)^{2k}, (2k+2) \end{cases}$$

where the notation $(a) \rightarrow (b)^c, (d)$ means that a node labeled (a) produces c children labeled (b) and one child labeled (d) .

Starting from the root node labeled (0) , we apply this rule iteratively to build a generating tree. The labels at level n of this tree encode the Bessel–Stirling numbers $\llbracket n \rrbracket_2$: the number of nodes with label (k) at level n corresponds precisely to $\llbracket n \rrbracket_2$. For example, the first few levels of the generating tree produce the following multisets of labels:

Level 0 : (0)
 Level 1 : (1)
 Level 2 : (2)
 Level 3 : $(2)^2(3)$
 Level 4 : $(2)^4(3)^4(4)$
 Level 5 : $(2)^8(3)^{12}(4)^8(5)$

The total number of labels at level n is precisely the n -th Bessel–Bell number (sequence A007472). This succession rule directly encodes the recurrence relation for the Bessel–Stirling numbers. For even values $k = 2j$, a node labeled $(2j)$ produces $2j$ children with the same label plus one child with label $(2j+1)$, corresponding to the term $2j \cdot \llbracket n \rrbracket_{2j}$ in the recurrence relation. Similarly, for odd values $k = 2j+1$, a node produces $2j$ children with the same label, corresponding to the term $2j \cdot \llbracket n \rrbracket_{2j+1}$, plus one child with label $(2j+2)$.

This succession rule formalism provides a direct connection to the combinatorial interpretation: the label (k) represents using k compartments, and the number of children with the same label reflects the number of valid placements in already-used compartments, while the single child with a new label represents opening a new compartment. The generalization Ω_m for arbitrary m is given in Definition 36.

Property	Classical ($m = 1$)	Bessel case ($m = 2$)	General m
Defining shift property	$B_{n+1} = \sum_k \binom{n}{k} B_k$	$B_{n+2}^{(2)} = \sum_k \binom{n}{k} 2^{n-k} B_k^{(2)}$	$B_{n+m}^{(m)} = \sum_k \binom{n}{k} m^{n-k} B_k^{(m)}$
Operator equation	$\text{BINOM } a = L a$	$\text{BINOM}^2 a = L^2 a$	$\text{BINOM}^m a = L^m a$
e.g.f. ODE	$\mathcal{A}' = e^x \mathcal{A}$	$\mathcal{A}'' = e^{2x} \mathcal{A}$	$\mathcal{A}^{(m)} = e^{mx} \mathcal{A}$
ODE at $t = e^x$	$t f' = t f$	$t^2 f'' + t f' = t^2 f$	$(tD)^m f = t^m f$
Primary carrier	e^t	$I_0(t)$	$C_0^{(m)}(t) = {}_0F_{m-1}(\cdot; 1, \dots, 1; (t/m)^m)$
Carrier system	$\{e^x\}$	$\{I_0, I_1\}$	$\{C_0^{(m)}, \dots, C_{m-1}^{(m)}\}$ (hyper-Bessel)
e.g.f.	$e^{-1} e^{e^x}$	$p I_0(e^x) + q K_0(e^x)$	span of m primitive solutions
Triangle recurrence	$\left\{ \begin{smallmatrix} n+1 \\ k \end{smallmatrix} \right\} = k \left\{ \begin{smallmatrix} n \\ k \end{smallmatrix} \right\} + \left\{ \begin{smallmatrix} n \\ k-1 \end{smallmatrix} \right\}$	$\llbracket \begin{smallmatrix} n+1 \\ k \end{smallmatrix} \rrbracket_2 = 2 \lfloor \frac{k}{2} \rfloor \llbracket \begin{smallmatrix} n \\ k \end{smallmatrix} \rrbracket_2 + \llbracket \begin{smallmatrix} n \\ k-1 \end{smallmatrix} \rrbracket_2$	$\llbracket \begin{smallmatrix} n+1 \\ k \end{smallmatrix} \rrbracket_m = m \lfloor \frac{k}{m} \rfloor \llbracket \begin{smallmatrix} n \\ k \end{smallmatrix} \rrbracket_m + \llbracket \begin{smallmatrix} n \\ k-1 \end{smallmatrix} \rrbracket_m$
Shift identity	$\left\{ \begin{smallmatrix} n+1 \\ k \end{smallmatrix} \right\} = \sum_j \binom{n}{j} \left\{ \begin{smallmatrix} j \\ k-1 \end{smallmatrix} \right\}$	$\llbracket \begin{smallmatrix} n+2 \\ k \end{smallmatrix} \rrbracket_2 = \sum_j \binom{n}{j} 2^{n-j} \llbracket \begin{smallmatrix} j \\ k-2 \end{smallmatrix} \rrbracket_2$	$\llbracket \begin{smallmatrix} n+m \\ k \end{smallmatrix} \rrbracket_m = \sum_j \binom{n}{j} m^{n-j} \llbracket \begin{smallmatrix} j \\ k-m \end{smallmatrix} \rrbracket_m$
Row sums	B_n (A000110)	A007472; parity sums A351143, A351028	$B^{(m)}$; residue sums $= B^{(m,r)}$
Dobiński formula	$B_n = e^{-1} \sum_k \frac{k^n}{k!}$	$\sum_k \frac{(2k)^n}{(2^k k!)^2} = v_n I_0(1) + u_n I_1(1)$	$\sum_k \frac{(mk)^n}{(m^k k!)^m} = \sum_r B_n^{(m,r)} C_r^{(m)}(1)$
Polynomials	$T_n(x) = \sum_k \left\{ \begin{smallmatrix} n \\ k \end{smallmatrix} \right\} x^k$	$\mathcal{B}_n(x) = \sum_k \llbracket \begin{smallmatrix} n \\ k \end{smallmatrix} \rrbracket_2 x^k$	$\mathcal{B}_n^{(m)}(x) = \sum_k \llbracket \begin{smallmatrix} n \\ k \end{smallmatrix} \rrbracket_m x^k$
Operational rep.	$e^{-x} e^{txD} e^x = e^{x(e^t-1)}$	$G(t, x)$ of Thm. 9	carrier expansion (Thm. 29)
Falling factorial	$x^{\underline{n}} = \prod (x - k)$	$x^{\underline{n}} = \prod (x - 2 \lfloor k/2 \rfloor)$	$x^{\underline{n}}_m = \prod (x - m \lfloor k/m \rfloor)$
First kind recurrence	$n \begin{bmatrix} n \\ k \end{bmatrix} + \begin{bmatrix} n \\ k-1 \end{bmatrix}$	$2 \lfloor \frac{n}{2} \rfloor \llbracket \begin{smallmatrix} n \\ k \end{smallmatrix} \rrbracket_2 + \llbracket \begin{smallmatrix} n \\ k-1 \end{smallmatrix} \rrbracket_2$	$m \lfloor \frac{n}{m} \rfloor \llbracket \begin{smallmatrix} n \\ k \end{smallmatrix} \rrbracket_m + \llbracket \begin{smallmatrix} n \\ k-1 \end{smallmatrix} \rrbracket_m$
First kind row sums	$n!$ (all permutations)	A000246 (odd-order permutations)	$\prod_{j < n} (m \lfloor j/m \rfloor + 1)$
Lah recurrence	$(n+k) \begin{bmatrix} n \\ k \end{bmatrix} + \begin{bmatrix} n \\ k-1 \end{bmatrix}$	$2(\lfloor \frac{n}{2} \rfloor + \lfloor \frac{k}{2} \rfloor) \llbracket \begin{smallmatrix} n \\ k \end{smallmatrix} \rrbracket_2 + \llbracket \begin{smallmatrix} n \\ k-1 \end{smallmatrix} \rrbracket_2$	$m(\lfloor \frac{n}{m} \rfloor + \lfloor \frac{k}{m} \rfloor) \llbracket \begin{smallmatrix} n \\ k \end{smallmatrix} \rrbracket_m + \llbracket \begin{smallmatrix} n \\ k-1 \end{smallmatrix} \rrbracket_m$
Symmetric functions	$h_{n-k}(1..k), e_{n-k}(0..n-1)$	h, e at weights $2 \lfloor j/2 \rfloor$	h, e at weights $m \lfloor j/m \rfloor$
Combinatorial model	set partitions	2-compartment urns	m -compartment urns
Distribution	$\text{Poisson}(y)$	$\text{CMP}((y/2)^2, 2)$	$\text{CMP}((y/m)^m, m)$
Factorial moments	$\mathbb{E}[X^{\underline{n}}] = y^n$	$\mathbb{E}[(2X)^{\underline{n}}] = y^n \frac{I_{n \bmod 2}(y)}{I_0(y)}$	$\mathbb{E}[(mX)^{\underline{n}}_m] = y^n \frac{C_{n \bmod m}^{(m)}(y)}{C_0^{(m)}(y)}$
Raw moments	$\mathbb{E}[X^n] = T_n(y)$	$2^n \mathbb{E}[X^n] = V_n(y)/I_0(y)$	$m^n \mathbb{E}[X^n] = V_n(y)/C_0^{(m)}(y)$