



SVD and PCA in HD

December 5, 2023

Motivation - Gene Expression



Transcription factors



PCA - reduction of dimensionality of "omics" data

$X_{n \times p}$ - data matrix (e.g. gene expressions), $n = k \times 100$,
 $p \approx 20000$ - number of genes

PCA - reduction of dimensionality of "omics" data

$X_{n \times p}$ - data matrix (e.g. gene expressions), $n = k \times 100$,
 $p \approx 20000$ - number of genes

PCA - reduction of dimensionality of "omics" data

$X_{n \times p}$ - data matrix (e.g. gene expressions), $n = k \times 100$,
 $p \approx 20000$ - number of genes

Assumptions : $X = M + E$, where M is of a low rank and E is a random noise

PCA - reduction of dimensionality of "omics" data

$X_{n \times p}$ - data matrix (e.g. gene expressions), $n = k \times 100$,

$p \approx 20000$ - number of genes

Assumptions : $X = M + E$, where M is of a low rank and E is a random noise

We usually assume that $e_{ij} \sim N(0, \sigma)$

PCA - reduction of dimensionality of "omics" data

$X_{n \times p}$ - data matrix (e.g. gene expressions), $n = k \times 100$,

$p \approx 20000$ - number of genes

Assumptions : $X = M + E$, where M is of a low rank and E is a random noise

We usually assume that $e_{ij} \sim N(0, \sigma)$

Mathematical goal - recovering M , separation of the signal from noise

PCA - reduction of dimensionality of "omics" data

$X_{n \times p}$ - data matrix (e.g. gene expressions), $n = k \times 100$,
 $p \approx 20000$ - number of genes

Assumptions : $X = M + E$, where M is of a low rank and E is a random noise

We usually assume that $e_{ij} \sim N(0, \sigma)$

Mathematical goal - recovering M , separation of the signal from noise

Practical goal - data compression, several basis vectors
[Principal Components] may contain most of the information
and be applied for prediction (of the patient's response to the therapy)

Principal Components Analysis (2)

Method - Singular Value Decomposition:

$$X = U_{n \times I} D_{I \times I} V_{I \times p}^T ,$$

$U^T U = I_{I \times I}$, $V^T V = I_{I \times I}$, $I = \min\{n, p\}$, and D is the diagonal matrix

Principal Components Analysis (2)

Method - Singular Value Decomposition:

$$X = U_{n \times I} D_{I \times I} V_{I \times p}^T ,$$

$U^T U = I_{I \times I}$, $V^T V = I_{I \times I}$, $I = \min\{n, p\}$, and D is the diagonal matrix

Eckhart-Young theorem: SVD truncated to k largest singular values (i.e. setting $D_{k+1,k+1} = \dots = D_{I,I} = 0$) is the best rank k approximation to X in the Frobenius norm, i.e.

$$\hat{M}_k = U_{n \times k} D_{k \times k} V_{k \times p}^T = \arg \min_{M: \text{rank}(M)=k} \sum_{ij} (X_{ij} - M_{ij})^2$$

Principal Components Analysis (2)

Method - Singular Value Decomposition:

$$X = U_{n \times I} D_{I \times I} V_{I \times p}^T ,$$

$U^T U = I_{I \times I}$, $V^T V = I_{I \times I}$, $I = \min\{n, p\}$, and D is the diagonal matrix

Eckhart-Young theorem: SVD truncated to k largest singular values (i.e. setting $D_{k+1,k+1} = \dots = D_{I,I} = 0$) is the best rank k approximation to X in the Frobenius norm, i.e.

$$\hat{M}_k = U_{n \times k} D_{k \times k} V_{k \times p}^T = \arg \min_{M: \text{rank}(M)=k} \sum_{ij} (X_{ij} - M_{ij})^2$$

\hat{M}_k is the maximum likelihood estimator of M under the restriction $\text{rank}(M) = k$

Principal Components Analysis (2)

Method - Singular Value Decomposition:

$$X = U_{n \times l} D_{l \times l} V_{l \times p}^T ,$$

$U^T U = I_{l \times l}$, $V^T V = I_{l \times l}$, $l = \min\{n, p\}$, and D is the diagonal matrix

Eckhart-Young theorem: SVD truncated to k largest singular values (i.e. setting $D_{k+1,k+1} = \dots = D_{l,l} = 0$) is the best rank k approximation to X in the Frobenius norm, i.e.

$$\hat{M}_k = U_{n \times k} D_{k \times k} V_{k \times p}^T = \arg \min_{M: \text{rank}(M)=k} \sum_{ij} (X_{ij} - M_{ij})^2$$

\hat{M}_k is the maximum likelihood estimator of M under the restriction $\text{rank}(M) = k$

Statistical Goal - determining rank k of matrix M

PESEL (PEnalized SEmi-integrated Likelihood)

Sobczyk, Bogdan, Josse, Journal of Computational Graphical Statistics, 2017

Bayesian Information Criterion (BIC) (1)

$A_1 \in A_2 \in A_3 \dots$ - nested sequence of statistical models

Bayesian Information Criterion (BIC) (1)

$A_1 \in A_2 \in A_3 \dots$ - nested sequence of statistical models

Bayesian Information Criterion (BIC) (1)

$A_1 \in A_2 \in A_3 \dots$ - nested sequence of statistical models

In our example A_k - $\text{rank}(M) \leq k$

Bayesian Information Criterion (BIC) (1)

$A_1 \in A_2 \in A_3 \dots$ - nested sequence of statistical models

In our example A_k - $\text{rank}(M) \leq k$

θ - vector of parameters of A_k :

elements of $U_k \in S_{k,n}$, $V_k \in S_{k,p}$, D_k , i σ

$S_{k,n}$ - Stiefel manifold of orthonormal matrices of dimension $n \times k$

Bayesian Information Criterion (BIC) (1)

$A_1 \in A_2 \in A_3 \dots$ - nested sequence of statistical models

In our example A_k - $\text{rank}(M) \leq k$

θ - vector of parameters of A_k :

elements of $U_k \in S_{k,n}$, $V_k \in S_{k,p}$, D_k , i σ

$S_{k,n}$ - Stiefel manifold of orthonormal matrices of dimension $n \times k$

$l(X, \theta)$ - likelihood function (density of the distribution describing the data)

Bayesian Information Criterion (BIC) (2)

In general situation BIC suggests selecting the model maximizing

$$\max_{\theta \in A_k} \log l(X, \theta) - 1/2 \dim(A_k) \log N$$

where N is the number of independent observations.

Bayesian Information Criterion (BIC) (2)

In general situation BIC suggests selecting the model maximizing

$$\max_{\theta \in A_k} \log l(X, \theta) - 1/2 \dim(A_k) \log N$$

where N is the number of independent observations.

BIC is justified (consistent) where $\dim(A_k) = \text{const}$ when $N \rightarrow \infty$

Bayesian Information Criterion (BIC) (2)

In general situation BIC suggests selecting the model maximizing

$$\max_{\theta \in A_k} \log l(X, \theta) - 1/2 \dim(A_k) \log N$$

where N is the number of independent observations.

BIC is justified (consistent) where $\dim(A_k) = \text{const}$ when $N \rightarrow \infty$

In our case $N = np$, so $\dim(A_k)$ increases with n and p

Bayesian Information Criterion (BIC) (2)

In general situation BIC suggests selecting the model maximizing

$$\max_{\theta \in A_k} \log l(X, \theta) - 1/2 \dim(A_k) \log N$$

where N is the number of independent observations.

BIC is justified (consistent) where $\dim(A_k) = \text{const}$ when $N \rightarrow \infty$

In our case $N = np$, so $\dim(A_k)$ increases with n and p

Idea - reduction of the number of parameters by integrating them out with respect to some prior distribution

PESEL for large p

Assume that $M = TW^T$, where

$T = [t_{i,l}]_{n \times k}$ is the matrix of "hidden factors",

$W = [w_{i,l}]_{p \times k}$ is the matrix of coefficients

PESEL for large p

Assume that $M = TW^T$, where

$T = [t_{i,l}]_{n \times k}$ is the matrix of "hidden factors",

$W = [w_{i,l}]_{p \times k}$ is the matrix of coefficients

prior distribution -

$$w_{j.} \sim N(0, I_k) \text{ ,}$$

which implies, that $x_{.1}, \dots, x_{.p}$ are iid random vectors from the distribution

$$x_{.j} \sim N(0; TT^T + \sigma^2 I_n) \text{ .}$$

PESEL for large p

Assume that $M = TW^T$, where

$T = [t_{i,l}]_{n \times k}$ is the matrix of "hidden factors",

$W = [w_{i,l}]_{p \times k}$ is the matrix of coefficients

prior distribution -

$$w_{j.} \sim N(0, I_k) \text{ ,}$$

which implies, that $x_{.1}, \dots, x_{.p}$ are iid random vectors from the distribution

$$x_{.j} \sim N(0; TT^T + \sigma^2 I_n) \text{ .}$$

Now we have p independent vectors and the number of parameters does not depend on p - we can apply BIC if only $p \gg n$

PESEL for large p

Assume that $M = TW^T$, where

$T = [t_{i,l}]_{n \times k}$ is the matrix of "hidden factors",

$W = [w_{i,l}]_{p \times k}$ is the matrix of coefficients

prior distribution -

$$w_{j\cdot} \sim N(0, I_k) \text{ ,}$$

which implies, that $x_{\cdot 1}, \dots, x_{\cdot p}$ are iid random vectors from the distribution

$$x_{\cdot j} \sim N(0; TT^T + \sigma^2 I_n) \text{ .}$$

Now we have p independent vectors and the number of parameters does not depend on p - we can apply BIC if only $p \gg n$

The related criterion is called PESEL (Penalized SEmi-integrated Likelihood).

Consistency of PESEL

Let $\hat{k}_0(p, n)$ be the PESEL estimator of the rank of M .

Consistency of PESEL

Let $\hat{k}_0(p, n)$ be the PESEL estimator of the rank of M .

Let $\frac{1}{p}MM' \rightarrow L$ with $\text{rank}(L) = k_0$.

Consistency of PESEL


Let $\hat{k}_0(p, n)$ be the PESEL estimator of the rank of M .

Let $\frac{1}{p}MM' \rightarrow L$ with $\text{rank}(L) = k_0$.

Then, for n fixed and under mild regularity conditions it holds

$$P(\exists p_0 \forall p > p_0 \quad \hat{k}_0(p, n) = k_0) = 1.$$

Errors from the log-normal distribution



Multiple Latent Component Clustering, Sobczyk, Wilczyński,
Bogdan, Graczyk, Josse, Panloup, Seegers, 2020

Multiple Latent Component Clustering, Sobczyk, Wilczyński, Bogdan, Graczyk, Josse, Panloup, Seegers, 2020

Awards for young scientists P. Sobczyk (Vienna workshop on simulation, 2015) i S. Wilczyński (International Conference on Biometrics and Bio-Pharmaceutical Statistics, Wiedeń 2017)

Multiple Latent Component Clustering, Sobczyk, Wilczyński, Bogdan, Graczyk, Josse, Panloup, Seegers, 2020

Awards for young scientists P. Sobczyk (Vienna workshop on simulation, 2015) i S. Wilczyński (International Conference on Biometrics and Bio-Pharmaceutical Statistics, Wiedeń 2017)

Goal: Identification of groups of co-regulated variables (genetic pathways) and selection of appropriate Principal Components.

Varclust

Multiple Latent Component Clustering, Sobczyk, Wilczyński, Bogdan, Graczyk, Josse, Panloup, Seegers, 2020

Awards for young scientists P. Sobczyk (Vienna workshop on simulation, 2015) i S. Wilczyński (International Conference on Biometrics and Bio-Pharmaceutical Statistics, Wiedeń 2017)

Goal: Identification of groups of co-regulated variables (genetic pathways) and selection of appropriate Principal Components.

Mathematics: clustering of variables into groups, such that each of them is spanned by just few of "hidden" variables.

Varclust

Multiple Latent Component Clustering, Sobczyk, Wilczyński, Bogdan, Graczyk, Josse, Panloup, Seegers, 2020

Awards for young scientists P. Sobczyk (Vienna workshop on simulation, 2015) i S. Wilczyński (International Conference on Biometrics and Bio-Pharmaceutical Statistics, Wiedeń 2017)

Goal: Identification of groups of co-regulated variables (genetic pathways) and selection of appropriate Principal Components.

Mathematics: clustering of variables into groups, such that each of them is spanned by just few of "hidden" variables.

Package *varclust* by P. Sobczyk and S. Wilczyński- Algorithm K-centroids around PCs. Estimation of the number of clusters and their dimensions by modifications of BIC.

Methodology

K-centroids algorithm

Methodology

K-centroids algorithm

Centers - PCs, distance - BIC

Methodology

K-centroids algorithm

Centers - PCs, distance - BIC

Estimation of clusters dimensions by PESEL

Methodology

K-centroids algorithm

Centers - PCs, distance - BIC

Estimation of clusters dimensions by PESEL

Repeat for different K and estimate K by mBIC

Overview of the algorithm

Algorithm: Multiple Latent Clustering Components

Input: n - number of individuals, p - number of variables,
 $X_{n \times p} = (x_1, \dots, x_p)$ - data set, d - maximal subspace
dimension, N - number of runs of the algorithm

Scale X to have columns with mean 0 and unit variance

for $i \in \{1, \dots, N\}$ **do**

Find the model using K-means and store its value of mBIC

end for

Choose the model with the highest value of mBIC and return
the model (segmentation, mBIC, factors) as the result. =0

K-means step

1. Initialize clusters' centres

K-means step

1. Initialize clusters' centres
2. Until convergence or maximal number of iterations is reached repeat:

K-means step

1. Initialize clusters' centres
2. Until convergence or maximal number of iterations is reached repeat:
 - ▶ For every variable x_j and every cluster factors $F_{j'}$ fit a linear regression model without intercept $lm(x_j \sim F_{j'})$ and store BIC value as $BIC_{jj'}$
 - ▶ Assign variable x_j to the cluster M_q where

$$q = \arg \max_{j' \in \{1, \dots, K\}} BIC_{jj'}$$

K-means step

1. Initialize clusters' centres
2. Until convergence or maximal number of iterations is reached repeat:
 - ▶ For every variable x_j and every cluster factors $F_{j'}$ fit a linear regression model without intercept $lm(x_j \sim F_{j'})$ and store BIC value as $BIC_{jj'}$
 - ▶ Assign variable x_j to the cluster M_q where

$$q = \arg \max_{j' \in \{1, \dots, K\}} BIC_{jj'}$$

- ▶ For every cluster M_i use PESEL to estimate its dimensionality k_i with an upper bound of d . Use PCA to compute the first k_i principal components and store them in F_i

Convergence (1)

Both the partition (BIC) and the center's selection (PESEL) steps lead to increase of Laplace approximations to the model posterior probability.

Convergence (1)


Both the partition (BIC) and the center's selection (PESEL) steps lead to increase of Laplace approximations to the model posterior probability.

For sufficiently large n and p each step of the algorithm leads to increase of mBIC.


Convergence (2)

$$p = 750p = 3000$$

Value of mBIC after given number of iterations



Value of mBIC after given number of iterations



Informative prior distribution and mBIC

- ▶ The problem with BIC (non-informative prior)
- ▶ Prior distribution taking into account the number of clusters and maximal dimension of the subspace

$$P(M) = \frac{1}{K^p} \frac{1}{d^K}$$
$$mBIC = \sum_{i=1}^K \ln \left(\hat{P}(X_i | M_i) \right) - p \ln(K) - K \ln(d)$$

Informative prior distribution and mBIC


- ▶ The problem with BIC (non-informative prior)
- ▶ Prior distribution taking into account the number of clusters and maximal dimension of the subspace

$$P(M) = \frac{1}{K^p} \frac{1}{d^K}$$
$$mBIC = \sum_{i=1}^K \ln \left(\hat{P}(X_i | M_i) \right) - p \ln(K) - K \ln(d)$$

Application

mBIC can be used to compare different models and to choose the number of clusters in the data.

Figure: Estimation of the number of clusters. Simulation parameters:
 $n = 100$, $p = 600$, $d = 3$, $SNR = 1$ mode : *not shared*.



Compared methods

1. **Sparse Subspace Clustering (SSC)**
2. Low Rank Subspace Clustering (LRSC)
3. **MLCC with random initialization (MLCC)**
4. **MLCC with initialization by the result of SSC**
(MLCC_{aSSC})
5. MLCC with initialization by sparse PCA (MLCC_{sPCA})
6. ClustOfVar (COV)

Data generation - shared factors

Input: n, SNR, K, p, d

Number of factors $m \leftarrow K \frac{d}{2}$

Factors $F = (f_1, \dots, f_m), f_i \sim N(0, I_n)$

Draw subspaces' dimension d_1, \dots, d_K uniformly from $\{1, \dots, d\}$

for $i = 1, \dots, K$ **do**

$F_i \leftarrow$ sample of size d_i from columns of F

Draw matrix of coefficients C_i from

$U(0.1, 1) \cdot \text{sgn}(U(-1, 1))$

Variables in the i -th subspace are $X_i \leftarrow F_i C_i$

end for

Scale matrix $X = (X_1, \dots, X_K)$ (columns with unit variance)

return $X + Z$ where $Z \sim N(0, \frac{1}{SNR} I_n) = 0$

Data generation - independent subspaces

Remark

To generate data without shared factors we draw independently i -th subspaces basis F_i as sample of size d_i from standard multivariate normal distribution

Measures of effectiveness

Compare two partitions $A = (A_1, \dots, A_n)$, $B = (B_1, \dots, B_m)$

- ▶ Adjusted Rand Index (ARI)
- ▶ Integration
- ▶ Acontamination
- ▶ $\text{ARI} \in [-1, 1]$, Integration, Acontamination $\in [0, 1]$.

Measures of effectiveness

Compare two partitions $A = (A_1, \dots, A_n)$, $B = (B_1, \dots, B_m)$


- ▶ Adjusted Rand Index (ARI)
- ▶ Integration
- ▶ Acontamination
- ▶ $\text{ARI} \in [-1, 1]$, Integration, Acontamination $\in [0, 1]$.

Remark


The bigger the indices, the better the clustering.

Mode

Values of ARI, Integration and Acontamination




Values of ARI, Integration and Acontamination




Number of variables

Values of ARI, Integration and Acontamination




Values of ARI, Integration and Acontamination




Signal to noise ratio

Values of ARI, Integration and Acontamination





Values of ARI, Integration and Acontamination




Execution time

Figure: Comparison of the execution time of the methods with respect to p and K . Simulation parameters: $n = 100$, $d = 3$, $SNR = 1$ mode : shared.

With respect to the number of variables With respect to the number of clusters



Particle matter measurements clustered using varclust



Sparse PCA (1)

maximize $a' \Sigma a$, subject to $\|a\| = 1$ and

1. $\|a\|_0 \leq k$
2. $\|a\|_1 \leq t$ (ScoTLASS)

Sparse PCA in R (*sparsepca*)

$$\operatorname{argmin}_B \frac{1}{2} \|X - XBA'\| + \phi(B) ,$$

where $AA' = I$ and $\phi(B)$ is the sparsity inducing penalty (LASSO or elastic net)