



**University of  
Zurich**<sup>UZH</sup>

**Zurich Open Repository and  
Archive**

University of Zurich  
University Library  
Strickhofstrasse 39  
CH-8057 Zurich  
[www.zora.uzh.ch](http://www.zora.uzh.ch)

---

Year: 2016

---

## **Knowing the past to forecast the future: a case study on a relictual, endemic species of the SW Alps, *Berardia subacaulis***

Guerrina, Maria ; Conti, Elena ; Minuto, Luigi ; Casazza, Gabriele

**Abstract:** Future climate change may lead to a substantial loss of biodiversity, particularly affecting mountain regions, including the Alps. Range-size reduction in high mountain plant species is predicted to be more pronounced for endemic species. Investigating the broad temporal spectrum of range shifts is important for the conservation of biodiversity, since learning how species responded to climate change in the past provides useful insights on how they might react to warming trends in the present and future. Using species distribution models and an ensemble forecasting approach, we explored how the distribution of *Berardia subacaulis*, a monospecific genus endemic of the south-west Alps, may be affected by past and future projected climate change. During the last interglacial, the habitat suitability of *Berardia* was lower than present and a progressive increase was observed from the last glacial maximum until now. In the future, *Berardia* appears to lose more than 80 % of its range, becoming endangered by 2050. Our results suggest that *Berardia* probably survived past warmer periods in situ, expanding its distributional range during cooler periods. The severe future range contraction predicted for *Berardia* reflects similar results for other endemic species. As *Berardia* represents an interesting model species to evaluate the effects of climate warming on range size and shifts, demographic and precise range monitoring may be undertaken on this species.

DOI: <https://doi.org/10.1007/s10113-015-0816-z>

Posted at the Zurich Open Repository and Archive, University of Zurich

ZORA URL: <https://doi.org/10.5167/uzh-111309>

Journal Article


Published Version

Originally published at:

Guerrina, Maria; Conti, Elena; Minuto, Luigi; Casazza, Gabriele (2016). Knowing the past to forecast the future: a case study on a relictual, endemic species of the SW Alps, *Berardia subacaulis*. *Regional Environmental Change*, 16(4):1035-1045.

DOI: <https://doi.org/10.1007/s10113-015-0816-z>

# Knowing the past to forecast the future: a case study on a relictual, endemic species of the SW Alps, *Berardia subacaulis*

Maria Guerrina<sup>1</sup>  · Elena Conti<sup>2</sup> · Luigi Minuto<sup>1</sup> · Gabriele Casazza<sup>1</sup>

Received: 22 December 2014 / Accepted: 10 May 2015  
© Springer-Verlag Berlin Heidelberg 2015

**Abstract** Future climate change may lead to a substantial loss of biodiversity, particularly affecting mountain regions, including the Alps. Range-size reduction in high mountain plant species is predicted to be more pronounced for endemic species. Investigating the broad temporal spectrum of range shifts is important for the conservation of biodiversity, since learning how species responded to climate change in the past provides useful insights on how they might react to warming trends in the present and future. Using species distribution models and an ensemble forecasting approach, we explored how the distribution of *Berardia subacaulis*, a monospecific genus endemic of the south-west Alps, may be affected by past and future projected climate change. During the last interglacial, the habitat suitability of *Berardia* was lower than present and a progressive increase was observed from the last glacial

maximum until now. In the future, *Berardia* appears to lose more than 80 % of its range, becoming endangered by 2050. Our results suggest that *Berardia* probably survived past warmer periods in situ, expanding its distributional range during cooler periods. The severe future range contraction predicted for *Berardia* reflects similar results for other endemic species. As *Berardia* represents an interesting model species to evaluate the effects of climate warming on range size and shifts, demographic and precise range monitoring may be undertaken on this species.

**Keywords** Palaeoendemic plant species · Ecological niche modelling · Glaciations · Global warming · Past climate change

Editor: Wolfgang Cramer.

**Electronic supplementary material** The online version of this article (doi:10.1007/s10113-015-0816-z) contains supplementary material, which is available to authorized users.

✉ Maria Guerrina  
maria.guerrina@edu.unige.it

Elena Conti  
econti@systbot.uzh.ch

Luigi Minuto  
minuto@dipteris.unige.it

Gabriele Casazza  
gabriele.casazza@unige.it

<sup>1</sup> DISTAV, University of Genoa, C.so Europa 26,  
I-16132 Genoa, Italy

<sup>2</sup> Institute of Systematic Botany and Botanic Garden,  
University of Zurich, Zollikerstrasse 107, CH-8008 Zurich,  
Switzerland

## Introduction

In the last decades, global climate has undergone dramatic changes, which are expected to continue into the twenty-first century (Solomon et al. 2007), inducing a substantial loss in biodiversity (McCann 2000). Predictions play a crucial role in alerting scientists and decision-makers to the potential future effects of global warming, with the possibility to develop proactive strategies aimed at mitigating these risks (Pereira et al. 2010; Bellard et al. 2012). Although the application of species distribution models (SDMs—Guisan and Zimmermann 2000; Guisan and Thuiller 2005) to climate change analyses has some limitations, major improvements have been achieved, allowing the development of ever more accurate models (Elith et al. 2006; Guisan and Thuiller 2005; Wisz and Guisan 2009). Therefore, SDMs currently represent one of the main tools for assessing the impacts of predicted

climate change on species distribution (Guisan and Thuiller 2005).

According to SDMs, future climate change may lead to substantial losses of biodiversity (Thuiller et al. 2005b; Thomas et al. 2004), affecting especially mountain regions, including the Alps, Pyrenees, French Cevennes, Carpathians, ranges in central Spain and the Balkan peninsula (Thuiller et al. 2005b). The effects of ongoing global warming are already visible, for instance, in the significant upward shift of plant distributional ranges documented in the western European mountains for the last century (Lenoir et al. 2008).

Mountains often constitute centres of endemism for biodiversity, harbouring endangered species and ecosystems. It has long been assumed that the loss of endemic species will have limited impacts on ecosystem functioning in the short term and at local scales, due to their frequently low abundance in communities (Grime 1998). However, Mouillot et al. (2013) observed that species with restricted geographic distribution tend to support vulnerable functions and to increase the level of functional diversity within the communities, which in turn sustains local ecosystem processes. Additionally, species with restricted distribution, which usually occupy specialized habitats, are by definition more susceptible to changes in their habitats (Thuiller et al. 2005a). These species are also likely to be more dispersal-limited and less genetically variable and thus less able to rapidly adapt to climate change than species with broader distribution (Hu and Jiang 2011).


The SW Alps have been identified as the region with the highest diversity of plant species, specifically of endemics, in the entire Alpine mountain chain (Pawlowski 1970). This highest diversity of plants reflects the richness, topography and history of the SW Alps (Médail and Verlaque 1997), and it derives also from its situation as a biogeographical crossroad between the Mediterranean and Alpine regions (Médail and Verlaque 1997). In the SW Alps, the effects of glaciations were less severe, and this area served as a refugium where some species escaped extinction caused by Pleistocene glaciations (Diadema et al. 2005; Minuto et al. 2006; Casazza et al. 2008). Here, numerous endemic species survived (e.g. *Acanthoprasium frutescens*, *Acis nicaeensis*, *Berardia subacaulis*, *Moehringia lebrunii*, *Phyteuma cordatum*, *Potentilla saxifraga*, *Saxifraga florulenta*—Casazza et al. 2005).

Pleistocene glaciations have strongly determined the distribution of narrow-ranging plants (Tribsch 2004). Climatic fluctuations have led to range restrictions, forcing species into refugia (Hewitt 1996; Tribsch and Schönswetter 2003), where climate conditions remained favourable during unfavourable periods (Araujo et al. 2008). Given this scenario of past range fluctuations in response to climatic cycles, investigating the broad temporal spectrum

of range shifts can provide useful information for the conservation of endemic species (Sillero and Carretero 2012).

Plants endemic to the Alps provide excellent models for such analyses, since they have experienced repeated cycles of Quaternary oscillations, often surviving in refugia when climate was adverse. *Berardia subacaulis* Vill. belongs to a monospecific genus endemic to the SW Alps (Fig. 1), grows exclusively on high-altitude calcareous screes and is viewed as an element of the palaeoflora, which was formed by a mosaic of tropical and subtropical plants mixed with temperate plants (Ozenda 2009). Contrary to the majority of palaeoendemic species that became extinct during Quaternary glaciations (Ozenda 2009), *B. subacaulis* probably survived in its peripheral refugium, located in the southern part of the SW Alps (Susanna and Garcia-Jacas 2009). The relictual nature of *B. subacaulis*, first proposed on the basis of its morphological features intermediate between Cardueae and Mutisieae tribes, of the monospecificity of the genus and of its position on an isolated long branch of the phylogenetic tree of the Cardueae (Susanna et al. 2006; Susanna and Garcia-Jacas 2009), was recently confirmed by molecular phylogenetic analyses that dated its origin in the early Oligocene about 33 Ma (95 % confidence interval 28.95–37.92 Ma) (Barres et al. 2013).

The present study is aimed at using *B. subacaulis* as a model to investigate how past climate change and future global warming are predicted to affect the survival of a palaeoendemic species restricted to the SW Alps. To achieve this goal, we used SDMs to infer the suitable habitats under a range of climate change scenarios for different time slices: last interglacial, last glacial maximum, current, 2050 and 2070. More specifically we asked three main questions: (1) How did past climate changes affect the distributional range of *B. subacaulis*? (2) Will projected climate change alter suitable habitat availability



**Fig. 1** Distributional range of *Berardia subacaulis* (continuous black line)

for *B. subacaulis*? (3) To what extent will *B. subacaulis* be threatened by future climate change? More broadly, this study contributes to ongoing debates on how species respond to climatic cycles, and, more specifically, whether ancient and narrowly distributed mountain species will go extinct or persist.

## Materials and methods

### Study species

*Berardia subacaulis* is a perennial herb growing on calcareous screes and stony slopes at high altitudes (between 1700 and 2700 m), and it is currently restricted to the SW Alps. Despite the species originated from a tropical and subtropical flora, it has some typical adaptations of plants to high mountains: acaulescence, dense wooly indument and stout woody rootstock (Garcia-Jacas et al. 2002). The flowers are protandrous, favouring the cross-fertilization, but self-fertilization is allowed (personal observations). The flowers are pollinated by a wide array of insects, but the visits are scarce (personal observations). The population size varies considerably from small population, with fewer than ten individuals, to very large population, with thousand of plants. Currently, *B. subacaulis* is not listed in the IUCN Red List at global scale. In Italy, at regional scale, *B. subacaulis* is listed as “critically endangered” (CR) (Conti et al. 1997). In France, the species has been included in Annex I of 92/43/EEC Directive (Council of European Communities 1992). Several populations are in conserved areas, such as SCIs in the Nature 2000 network or Natural Parks.

### Species occurrence data

We collected all occurrence data from the database “SILENE” of the Conservatoire Botanique National Méditerranéen de Porquerolles (CBNMED) and “FLORE” of the Conservatoire Botanique National Alpin de Gap (CBNA) and from our own field surveys (additional data are given in Online Resource 1). To perform detailed distribution models, we used only the data collected since 1995 with a GPS instrument from field surveys, which ensure enhanced geographic accuracy, for a final data set of 1184 records.

### Environmental layers at different time slices

Climatic variables related to temperature and precipitation play a significant role, directly and indirectly, in plant physiology (Körner 2003). Nineteen bioclimatic variables for five time slices were downloaded from the WorldClim

website (Hijmans et al. 2005; available at <http://www.worldclim.org>): last interglacial (LIG, ~140–120 kyr BP), current, 2050 and 2070 at 30-s spatial resolution; last glacial maximum (LGM, ~21.000 kyr BP) at 2.5-min spatial resolution.

For SDM analyses of current climatic layers, we performed a pairwise Pearson correlation among bioclimatic predictors aimed at reducing multicollinearity and minimizing model overfitting, retaining only predictors that were not highly correlated ( $r \leq 0.80$ ; see recommendation of Elith et al. 2006). Six bioclimatic variables were retained for the further analyses: BIO3, isothermality (monthly/annual temperature range); BIO8, mean temperature of wettest quarter; BIO10, mean temperature of warmest quarter; BIO15, precipitation seasonality; BIO17, precipitation of the coldest quarter; and BIO18, precipitation of the warmest quarter (additional information is given in Online Resource 2).

Because *B. subacaulis* grows only on specific calcareous substrates, for all the time slices we added a layer reporting the presence/absence of suitable substrate, based on the global lithological map data set, GLiM (Hartmann and Moosdorf 2012; in Online Resource 3).

### Climate change scenarios

For SDM predictions in the LGM, we used data climate layers from both the Community Climate System Model (CCSM) and the Model for Interdisciplinary Research on Climate (MIROC).

For SDM predictions in the future, we chose three general circulation models (GCM): one provided by the Max Planck Institute (MPI-ESM-LR; Online Resource 4), one by the UK Met Office (HadGEM2-ES; Online Resource 4) and one developed by the Institut Pierre Simon Laplace (IPSL-CM5A-LR; Online Resource 4). For each GCM and each time slice, we applied two representative concentration pathways (RCP2.6 the low mitigation scenario and RCP8.5 the high emission scenario—Van Vuuren et al. 2011; Online Resource 4) adopted by the Intergovernmental Panel on Climate Change for its fifth Assessment Report (AR5).

### Species distribution modelling

Species occurrence data and the seven selected variables (Online Resource 2) were used to construct distribution models using the BIOMOD2 package (Thuiller et al. 2009) implemented in R (R Development Core Team 2008). We applied six SDM techniques: multivariate adaptive regression splines (MARS—Friedman 1991), generalized linear models (GLM—McCullagh and Nelder 1989), classification tree analysis (CTA—Breiman et al. 1984),

flexible discriminant analysis (FDA—Hastie et al. 1994), random forest (RF—Breiman 2001) and Maxent (Phillips et al. 2006). These techniques belong to three different categories of models (i.e. regression methods—MARS and GLM; classification methods—CTA and FDA; and machine learning algorithms—RF and Maxent; Barbet-Massin et al. 2012).

For each SDM technique, the number of pseudo-absences was selected according to the recommendation of Barbet-Massin et al. (2012) (Online Resource 5). The predictive performance of the models was evaluated for each pseudo-absence run by repeating a split-sample cross-validation ten times, using a random subset (70 %) of the initial data set each time to calibrate the models, while the remaining 30 % were used to evaluate the models. The models predicted suitability values between 0 and 1000 at each site and were downscaled at 0–1 range. Two different measures for the evaluation of the models were calculated: the area under the curve (AUC) of a receiver operating characteristic (ROC) plot and the true skill statistic (TSS) (Thuiller et al. 2009). To transform the inferred continuous probability values to binary presence–absence form, we used the TSS. For the final ensemble projections, we considered the average among all the models for the same time slices.

### Species extinction risks

The following thresholds were used to assign the species to different threat categories for future SDM projections, according to the IUCN Red List criterion A3 (IUCN 2012): projected range loss of 100 %, extinct (EX); projected range loss of >80 %, critically endangered (CR); projected range loss of >50 %, endangered (EN); and projected range loss of >30 %, vulnerable (VU). Although this Red Listing approach is simplistic because it considers only the role of climatic variables on species survival, it provides a synthetic overview of biotic threats driven specifically by climate change (Thuiller et al. 2005b).

### Spatial index for potential impacts of climate change

Following Hu and Jiang (2011), we used two approaches to assess the impact of climate change on potential habitat range. First, we estimated the percentage of predicted future range change ( $C$ ) by cell using  $C = 100(RG - RL)/PR$ , where  $RG$  is the range gain by cell,  $RL$  is the range loss by cell and  $PR$  is the predicted present range by cell. A negative  $C$  value indicates a loss in overall range, whereas a positive value indicates an increase in overall range size. The future range turnover ( $T$ ) by cell of the climate envelope range was estimated using  $T = 100(RL + RG)/(PR + RG)$ . A  $T$  value of 0 indicates no shift in range, whereas a value of 100 indicates a complete range shift

when compared with the predicted range under current conditions. The full-dispersal scenario assumes that a species could reach all geographical areas that are predicted to be suitable in future climate projections (i.e.  $RG \geq 0$ ). On the contrary, the no-dispersal scenario assumes no dispersion from current to future suitable sites (i.e.  $RG = 0$ ), implying that future sites that might become suitable for a species remain unoccupied. The no-dispersal scenario was calculated as the percentage of range lost (percentage of  $RL$ ), while the full-dispersal scenario was calculated as percentage of range changed ( $C$ ), as the result of the difference between  $RG$  and  $RL$  related to the present range ( $PR$ ). The full-dispersal scenario is likely to be unrealistic because it assumes that a species can colonize all locations without physiological, environmental or geographical limitations, while the no-dispersal scenario is likely to be more appropriate for poor dispersers as endemic species frequently are.

Second, we extracted the probability of occupancy for known localities of *B. subacaulis* for the past and future time slices considered. We then characterized the trends in the projected probability of occurrences using raster packages for R (R Development Core Team 2008).


### Results

Under current climate conditions, AUC and TSS, interpreted using the classification of the BIOMOD manual (Thuiller et al. 2008), indicated excellent model performance for all modelling techniques, making them suitable for deriving past and future projections (additional information are given in Online Resources 6 and 7).

The potential range of *B. subacaulis* changed substantially among the five investigated time slices. During the LIG, the model identified habitat suitability in the area of SW Alps (Fig. 2a). During the LGM, the model highlighted an overall increase in habitat suitability in the study area, with higher values of habitat suitability mainly found in the southern part of SW Alps (Fig. 2b). Under climatic current conditions, the highest suitability values were detected in the entire area of the SW Alps (Fig. 2c). For the future projections, a progressive decrease in suitability values was observed in the SW Alps (Fig. 2d, e). According to these results, the probability of occurrence for known localities of *B. subacaulis* increased from the past to the present time, varying from 0.5 by LIG to 0.7 by LGM and 0.99 at present, and is predicted to decrease to 0.5 by 2050 and 0.4 by 2070 (Fig. 3).

The potential range of *B. subacaulis* was discernibly impacted by projected climate change. Across the GCMs and the two RCP scenarios, it was clear that the strength of the impacts increased with time. In the RCP2.6, the average





**Fig. 2** Predicted distribution of *Berardia subacaulis* for five time slices: LIG (**a** ~ 120 years BP), LGM (**b** ~ 21 years BP), current (**c**), 2050 (**d**) and 2070 (**e**). LGM model is obtained with an ensemble forecasting approach across two palaeoclimate models (MIROC and CCSM). Future models are obtained with an ensemble forecasting approach across three general circulation models (MPI-ESM,


HadGEM2-ES and IPSL-CM5A-LR) and two representative concentration pathways (RCP2.6 and RCP8.5). The values of habitat suitability are represented in a grey scale. The current coastal line is added to all the projections. The black colour represents the highest value of habitat suitability

percentage of range loss was 75.74 and 75.76 %, while in the RCP8.5, it was 90.79 and 97.55 %, for the years 2050 and 2070, respectively (Fig. 4a, b). The average percentage of range gain was 8.69 and 8.35 % in RCP2.6, while in the RCP8.5, it was 17.58 and 14.81 %, for the years 2050 and 2070, respectively. The high range loss and the low range gain values resulted in high range change values (Fig. 4c). This predicted a strong turnover (Fig. 4d) in range, and small range increases were projected for all future time slices. Under the full migration assumption, *B. subacaulis* becomes endangered (>50 % range loss) by 2050 and critically endangered (>80 % range loss) starting from 2070.

## Discussion

In this study, we present an analysis of the potential effects of past and future climate change on the distributional range of *B. subacaulis*. Our results should be considered with caution

due to some limitations of the approach we adopted. First, climate is only one of several determinants of species distribution (Nogués-Bravo 2009; Thuiller et al. 2009; Chuine 2010; Swab et al. 2012). For example, a restricted distribution may result from dispersal limitation, competition with established vegetation, effects of past, large-scale events and human impacts (Normand et al. 2011; Nogués-Bravo 2009, Svenning and Skov 2004). Nevertheless, the predictive power of climatic distribution models is high for species with small geographic ranges and limited environmental tolerance (Brotons et al. 2004; Elith et al. 2006; Hernandez et al. 2006; Rebelo and Jones 2010), such as *B. subacaulis*. Moreover, models of species with a restricted geographical distribution usually obtain higher values of model evaluation compare to species with wider distribution, because the former species, occupying a smaller part of the environmental space, have a stronger contrast between presence and pseudo-absences in that space (Jiménez-Valverde et al. 2008; Stokland et al. 2011). According to this statement, we found high values of




**Fig. 3** Predicted probability of occurrence for known localities of *Berardia subacaulis* at different time slices. The probability of occurrence is extracted on the base of predicted distributions from an ensemble forecast approach. The *black line* represents the median, *black circle* represents the mean, *edge box* corresponds to the first and third quartiles (the 25th and 75th percentiles), *whiskers* are  $1.5 \times \text{IQR}$  (where IQR is the inter-quartile range), and *circles* are outliers

both AUC and TSS. Second, coarse-scale models might underestimate the presence of microrefugia (Randin et al. 2009; Ashcroft 2010; Franklin et al. 2013; Patsiou et al. 2014). However, we produced our models at roughly  $1 \text{ km}^2$  of resolution to avoid the problem of hidden local refugia, as suggested by Maiorano et al. (2011) and Franklin et al. (2013). Consequently, we can be reasonably confident that our models for the distribution of *B. subacaulis* are as accurate as currently possible.

### Past and future models

According to our results, *B. subacaulis* experienced cycles of severe reduction of habitat suitability, followed by expansion, during the Pleistocene climatic oscillations (Fig. 2). In particular, during the LGM ( $\sim 21,000$  years BP, Fig. 2b), SDMs exhibited a suitable area wider than during the LIG ( $\sim 120,000$  years BP, Fig. 2a). The low habitat suitability inferred for the warmer interglacial period and the higher values inferred for the colder glacial period (Figs. 2, 3) may explain the survival

**Fig. 4** Projected impacts of climate change on the distribution of *Berardia subacaulis*. Percentage of range loss (a), range gain (b), range change (c) and range turnover (d) as predicted using three general circulation models (MPI-ESM, HadGEM2-ES and IPSL-CM5A-LR) and two representative concentration pathways (RCP2.6 and RCP8.5), for two time slices (2050 and 2070). The *black line* represents the median, *black circle* represents the mean, *edge box* corresponds to the first and third quartiles (the 25th and 75th percentiles), *whiskers* are  $1.5 \times \text{IQR}$  (where IQR is the inter-quartile range), and *circles* are outliers



of this species, suggesting that *B. subacaulis* is a cold-loving plant. Contrarily, on the basis of fossil evidence and molecular analyses, range contraction during the glacial period is recorded for other species belonging to the same flora, such as *Ramonda myconi* (Dubreuil et al. 2008), *Frangula alnus* (Hampe et al. 2003), *Tetraclinis articulata* (Médail and Diadema 2009) and *Zelkova sicula* (Di Pasquale et al. 1992). The pattern observed in *B. subacaulis* seems to be more similar to the one recorded for low temperature adapted species, such as *Juniperus thurifera* (Terrab et al. 2008), belonging to the flora differentiated at the end of the Miocene epoch, when the global cooling occurred. This observation is congruent with the morphological adaptations to high mountains conditions detected in *B. subacaulis*. In addition, the pattern observed in *B. subacaulis* fits with the ‘displacement refugia model’ (Kropf et al. 2003) predicting that European high-elevation plants faced range fragmentation mostly during the warmer, short interglacial periods. A survival in refugia during interglacial periods, such as *B. subacaulis*, has been inferred also for other plant (*Dryas octopetala* and *Betula nana*—Stewart et al. 2009; *Potentilla matsumurae*—Ikeda et al. 2006) and animal species (*Spermophilus citellus*—Musil 1985; *Stenobothrus cotticus*—Berger et al. 2010) on the basis of multidisciplinary analyses. According to Theurillat and Guisan (2001), cold-loving Tertiary relictual species and cold-resistant species probably survived in situ uninterruptedly since the late Tertiary. This hypothesis seems to be likely for *B. subacaulis*, which probably survived the interglacial periods in refugia situated at high altitude in the SW Alps, as shown in the LIG projection (Fig. 2a). As observed by Patsiou et al. (2014) in *Saxifraga florulenta*, *B. subacaulis* could have contracted in microrefugia during adverse periods. During glacial maxima, *B. subacaulis* probably expanded its putative distributional range within SW Alps due to more favourable climatic conditions and to the less severe ice effect in this part of the Alps (Médail and Diadema 2009). In the future, phylogeographical studies will be useful to clarify the history of the species and to confirm the results of the SDMs, since for *B. subacaulis*, as for most of the herbaceous taxa, there are no fossil records. In general, species such as *B. subacaulis* that prefer cooler conditions are of most immediate concern as they face increased threat with further rises in global temperatures (Skov and Svenning 2004). Phylogeographical studies on *B. subacaulis* will elucidate its genetic structure and diversity, two elements that are thought to be important factors for persistence of populations against changing environments (Barret and Kohn 1991). Moreover, genetic studies can improve conservation activities by identifying the populations that have very low genetic diversity. They can also provide information

regarding the extent of gene flow among fragmented populations (Frankham et al. 2010).

*Berardia subacaulis* currently seems to be at its maximum range expansion, while a severe reduction in habitat suitability is projected for the next few decades (Figs. 2, 3, 4). Similarly, severe reductions have been predicted for both plant and animal species; in particular, 22 % of the European plant species are expected to lose more than 80 % of their range by 2080 (Thuiller et al. 2005b). Greater habitat loss is projected for higher- than lower-elevation species; depending on future climate scenarios, 36–55 % of alpine species will lose more than 80 % of their suitable habitat by 2070–2100, as compared to 31–51 % of subalpine species and 19–46 % of montane species (Engler et al. 2011). Range-size reductions in high mountain plant species are predicted to be more pronounced for endemic (72–76 %) than non-endemic (39–48 %) species (Dullinger et al. 2012). For example, the alpine species *Lilium pomponium*, endemic to the Maritime and Ligurian Alps, is expected to lose more than 80 % of their native range by 2050 (Casazza et al. 2014). Similar studies on vertebrates in the Mediterranean Basin show that endemic species will be severely threatened by global warming (Maiorano et al. 2011, 2013). Moreover, the projected future distribution of plant species may be strongly affected, and even hampered, by the dispersal capability of individual species, which influences their potential to reach new, suitable areas (Engler et al. 2009; Normand et al. 2011).

### Conservation implications

*Berardia subacaulis* appears to lose more than 80 % of its range by 2070 (Fig. 4a) under the full-spread assumption (the possibility to reach all new potential habitat range). The range gain results very low and relatively stable through future time slices (Fig. 4b). Consequently, the increase in range turnover over the next few decades (Fig. 4d) is mainly due to the increase in habitat loss. Under the full-spread assumption, the species will become endangered by 2050. Our results corroborate those of previous, similar studies in the Alps. Using a range of climate change scenarios, Thuiller et al. (2005b) estimated that up to 48.5 % of mountain plant species in the European Alps are at risk of extinction. In general, global warming may result in a significant upward shift in species optimum elevation (Lenoir et al. 2008). However, marginal chains in the Alps have relatively low summit heights, making their alpine plant populations particularly vulnerable to mountaintop extinctions (Dullinger et al. 2012). Fitting this scenario, an altitudinal shift towards higher altitudinal areas does not appear to be a viable option for *B.*



*subacaulis*, for it has already reached the top of the mountains in the SW Alps (Theurillat and Guisan 2001). Additionally, the increase in competition due to upward shifts in species optimum elevations (Camenisch 2002; Burga et al. 2003; Lenoir et al. 2008; Engler et al. 2009, 2011) may strongly alter species interactions (Suttle et al. 2007; Engler et al. 2009), likely affecting species survival. This phenomenon, however, did not prevent *B. subacaulis* from surviving adverse climatic conditions since its estimated origin in the Oligocene (Barres et al. 2013), for it persisted in extreme habitats characterized by very limited, interspecific competition. Due to the severity and imminence of forecasted, climatic changes, *B. subacaulis* appears to be a useful species to study the direct and indirect effects of global warming on endemic species distribution and the ability of species to adapt to these changes.

## Conclusions

As suggested by Schwartz (2012), SDMs are more useful to identify conservation opportunities in newly available habitats under changing climate than they are for asserting where current habitat will no longer exist under future climate conditions. Our results suggest that distinguishing the relative roles of present and past climates in determining species' distributions is not only of theoretical interest, but a key to understand species' responses to climate change, as already observed by Araújo and Pearson (2005). The paucity of local cold-climate refugia at high altitudes, combined with low mobility, exposes alpine endemics to extreme risks of extinction under climate warming (Dullinger et al. 2012). This seems to be the case of *B. subacaulis*: its preference for cool temperature conditions, the low likelihood of upward shift and its low-dispersal capability disproportionately increase extinction risk. Moreover, since *B. subacaulis* is on a long isolated branch on the phylogenetic tree (Susanna et al. 2006; Susanna and García-Jacas 2009; Barres et al. 2013), its loss would represent a large loss of evolutionary history (Faith 2008). Due to its ancient origin and the high percentage of range loss forecasted, *B. subacaulis* seems to be an especially useful model species to study how cold-loving plants survived past warm periods and evaluate the effects of climate warming on their survival.

**Acknowledgments** We are grateful to the Conservatoire Botanique National Alpin de Gap and to the Conservatoire Botanique National Méditerranéen de Porquerolles for allowing us to use the occurrence points from the SILENE and FLORE database. We are grateful to the two anonymous reviewers for their helpful comments. This work was funded by Parco Naturale Regionale delle Alpi Marittime and by Interreg Alcotra programme 2007–2013, project n. 192 BIODIVAM.

## References

- Araújo MB, Pearson RG (2005) Equilibrium of species' distribution with climate. *Ecography* 28:693–695. doi:[10.1111/j.2005.0906-7590.04253.x](https://doi.org/10.1111/j.2005.0906-7590.04253.x)
- Araújo MB, Nogués-Bravo J, Diniz-Filho AF, Haywood AM, Valdes PJ, Rahbek C (2008) Quaternary climate changes explain diversity among reptiles and amphibians. *Ecography* 31:8–15. doi:[10.1111/j.2007.0906-7590.05318.x](https://doi.org/10.1111/j.2007.0906-7590.05318.x)
- Ashcroft MB (2010) Identifying refugia from climate change. *J Biogeogr* 37:1407–1413. doi:[10.1111/j.1365-2699.2010.02300.x](https://doi.org/10.1111/j.1365-2699.2010.02300.x)
- Barbet-Massin M, Jiguet F, Albert CH, Thuiller W (2012) Selecting pseudo-absences for species distribution models: how, where and how many? *Methods Ecol Evol* 3:327–338. doi:[10.1111/j.2041-210X.2011.00172.x](https://doi.org/10.1111/j.2041-210X.2011.00172.x)
- Barres L, Sanmartín I, Anderson CL, Susanna A, Buerki S, Galbany-Casals M, Vilatersana R (2013) Reconstructing the evolution and biogeographic history of tribe Cardueae (Compositae). *Am J Bot* 5:867–882. doi:[10.3732/ajb.1200058](https://doi.org/10.3732/ajb.1200058)
- Barret SCH, Kohn JR (1991) Genetic and evolutionary consequences of small population size in plants: implications for conservation. In: Falk DA, Holsinger KE (eds) *Genetics and conservation of rare plants*. Oxford University Press, New York, pp 3–30
- Bellard C, Bertelsmeier C, Leadley P, Thuiller W, Courchamp F (2012) Impacts of climate change on the future of biodiversity. *Ecol Lett* 15:365–377. doi:[10.1111/j.1461-0248.2011.01736.x](https://doi.org/10.1111/j.1461-0248.2011.01736.x)
- Berger D, Chobanov DP, Mayer F (2010) Interglacial refugia and range shifts of the alpine grasshopper *Stenobothrus coticus* (Orthoptera: Acrididae: Gomphocerinae). *Org Divers Evol* 10:123–133. doi:[10.1007/s13127-010-0004-4](https://doi.org/10.1007/s13127-010-0004-4)
- Breiman L (2001) Statistical modeling: the two cultures. *Stat Sci* 16:199–215. doi:[10.1214/ss/1009213726](https://doi.org/10.1214/ss/1009213726)
- Breiman L, Friedman JH, Olshen RA, Stone CJ (1984) *Classification and regression trees*. Chapman and Hall, New York
- Brotans L, Thuiller W, Araújo MB, Hirzel AH (2004) Presence-absence versus presence-only modelling methods for predicting bird habitat suitability. *Ecography* 27:437–448. doi:[10.1111/j.0906-7590.2004.03764.x](https://doi.org/10.1111/j.0906-7590.2004.03764.x)
- Burga CA, Haeberli W, Krummenacher B, Walther GR (2003) Abiotische und biotische Dynamik in Gebirgsräumen—Status quo und Zukunftsperspektiven. In: Jeanneret F, Wastl-Walter D, Wiesmann U, Schwyn M (eds) *Welt der Alpen—Gebirge der Welt. Jahrbuch der Geographischen Gesellschaft Bern* 61. Haupt, Bern, pp 25–37
- Camenisch M (2002) Veränderungen der Gipfflora im Bereich des Schweizerischen Nationalparks: ein Vergleich über die letzten 80 Jahre. *Jahresber Nat forsch Ges Graubünden* 111:27–37
- Casazza G, Barberis G, Minuto L (2005) Ecological characteristics and rarity of endemic plants of the Italian Maritime Alps. *Biol Conserv* 123:361–371. doi:[10.1016/j.biocon.2004.12.005](https://doi.org/10.1016/j.biocon.2004.12.005)
- Casazza G, Zappa E, Mariotti MG, Médail F, Minuto L (2008) Ecological and historical factors affecting distribution pattern and richness of endemic plant species: the case of Maritime and Ligurian Alps hotspot. *Divers Distrib* 14:47–58. doi:[10.1111/j.1472-4642.2007.00412.x](https://doi.org/10.1111/j.1472-4642.2007.00412.x)
- Casazza G, Giordani P, Benesperi R, Foggi B, Viciani D, Filigheddu R, Farris E, Bagella S, Pisanu S, Mariotti MG (2014) Climate change hastens the conservation urgency of range-restricted plant species in the central-northern Mediterranean region. *Biol Conserv* 179:129–138. doi:[10.1016/j.biocon.2014.09.015](https://doi.org/10.1016/j.biocon.2014.09.015)
- Chuine I (2010) Why does phenology drive species distribution? *Philos Trans R Soc B* 365:3149–3160. doi:[10.1098/rstb.2010.0142](https://doi.org/10.1098/rstb.2010.0142)

- Conti F, Manzi A, Pedrotti F (1997) Liste rosse regionali delle piante d'Italia. WWF Italia. Società Botanica Italiana, Università di Camerino, Camerino
- Council of European Communities (1992) Council directive 92/43/EEC of 21 May 1992 on the conservation of natural habitats and of wild fauna and flora. Off J Eur Communities 35:7–50
- Di Pasquale G, Garfi G, Quézel P (1992) Sur la presence d'un *Zelkova* nouveau en Sicile sud-orientale (Ulmaceae). Biocosme Méditerranéen 8–9:401–409
- Diadema K, Bretagnolle F, Affre L, Yuan YM, Médail F (2005) Geographic structure of molecular variation of *Gentiana ligustica* (Gentianaceae) in the Maritime and Ligurian regional hotspot, inferred from ITS sequences. Taxon 54:887–894. doi:10.2307/25065475
- Dubreuil M, Riba M, Mayol M (2008) Genetic structure and diversity in *Ramonda myconi* (Gesneriaceae): effects of historical climate change on a pre-glacial relict species. Am J Bot 95:577–587. doi:10.3732/ajb.2007320
- Dullinger S, Gatttringer A, Thuiller W, Moser D, Zimmermann NE, Guisan A, Willner W, Plutzer C, Leitner M, Mang T, Caccianiga M, Dirnböck T, Ertl S, Fischer A, Lenoir J, Svenning JC, Psomas A, Schmatz DR, Silc U, Vittoz P, Hülber K (2012) Extinction debt of high-mountain plants under twenty-first-century climate change. Nat Clim Change 2:619–622. doi:10.1038/nclimate1514
- Elith J, Graham CH, Anderson RP, Dudik M, Ferrier S, Guisan A, Hijmans RJ, Huettmann F, Leathwick JR, Lehmann A, Li J, Lohmann LG, Loiselle BA, Manion G, Moritz C, Nakamura M, Nakazawa Y, Overton JMcC, Peterson AT, Phillips SJ, Richardson K, Scachetti-Pereira R, Schapire RE, Soberón J, Williams S, Wisz MS, Zimmermann NE (2006) Novel methods improve prediction of species' distributions from occurrence data. Ecography 29:129–151. doi:10.1111/j.2006.0906-7590.04596.x
- Engler R, Randin CF, Vittoz P, Czaka T, Beniston M, Zimmermann NE, Guisan A (2009) Predicting future distributions of mountain plants under climate change: does dispersal capacity matter? Ecography 32:34–45. doi:10.1111/j.1600-0587.2009.05789.x
- Engler R, Randin CF, Thuiller W, Dullinger S, Zimmermann NE, Araújo MB, Pearman PB, Le Lay G, Piedallu C, Albert CH, Choler P, Coldea G, De Lamo X, Dirnböck T, Gégout JC, Gómez-García D, Grytnes JA, Heegaard E, Høistad F, Nogués-Bravo D, Normand S, Puşcas M, Sebastià MT, Stanisci A, Theurillat JP, Trivedi MR, Vittoz P, Guisan A (2011) 21st century climate change threatens mountain flora unequally across Europe. Glob Change Biol 17:2330–2341. doi:10.1111/j.1365-2486.2010.02393.x
- Faith D (2008) Threatened species and the potential loss of phylogenetic diversity: conservation scenarios based on estimated extinction probabilities and phylogenetic risk analysis. Biol Conserv 22:1461–1470. doi:10.1111/j.1523-1739.2008.01068.x
- Frankham R, Ballou JD, Briscoe DA (2010) Introduction to conservation genetics, 2nd edn. Cambridge University Press, Cambridge
- Franklin J, Davis FW, Ikegami M, Syphard AD, Flint LE, Flint AL, Hannah L (2013) Modeling plant species distributions under future climates: how fine scale do climate projections need to be? Glob Change Biol 19:473–483. doi:10.1111/gcb.12051
- Friedman J (1991) Multivariate adaptive regression splines. Ann Stat 19:1–141
- García-Jacas N, Garnatje T, Susanna A, Vilatersana R (2002) Tribal and subtribal delimitation and phylogeny of the Cardueae (Asteraceae): a combined nuclear and chloroplast DNA analysis. Mol Phylogenet Evol 22:51–64. doi:10.1006/mpev.2001.1038
- Grime GP (1998) Benefits of plant diversity to ecosystems: immediate, filter and founder effects. J Ecol 86:902–910. doi:10.1046/j.1365-2745.1998.00306.x
- Guisan A, Thuiller W (2005) Predicting species distribution: offering more than simple habitat models. Ecol Lett 8:993–1009. doi:10.1111/j.1461-0248.2005.00792.x
- Guisan A, Zimmermann NE (2000) Predictive habitat distribution models in ecology. Ecol Model 135:147–186. doi:10.1016/S0304-3800(00)00354-9
- Hampe A, Arroyo J, Jordano P, Petit RJ (2003) Rangewide phylogeography of a bird-dispersed Eurasian shrub: contrasting Mediterranean and temperate glacial refugia. Mol Ecol 12:3415–3426. doi:10.1046/j.1365-294X.2003.02006.x
- Hartmann J, Moosdorf N (2012) The new global lithological map database GLiM: a representation of rock properties at the earth surface. Geochem Geophys Geosyst 13:1525–2027. doi:10.1029/2012GC004370
- Hastie T, Tibshirani R, Buja A (1994) Flexible discriminant analysis by optimal scoring. J Am Stat Assoc 89:1255–1270. doi:10.1080/01621459.1994.10476866
- Hernandez PA, Graham CH, Master LL, Albert DL (2006) The effect of sample size and species characteristics on performance of different species distribution modeling methods. Ecography 29:773–785. doi:10.1111/j.0906-7590.2006.04700.x
- Hewitt CM (1996) Some genetic consequences of ice ages, and their role in divergence and speciation. Biol J Linn Soc 58:247–276. doi:10.1111/j.1095-8312.1996.tb01434.x
- Hijmans RJ, Cameron SE, Parra JL, Jones PG, Jarvis A (2005) Very high resolution interpolated climate surfaces for global land areas. Int J Climatol 25:1965–1978. doi:10.1002/joc.1276
- Hu J, Jiang Z (2011) Climate change hastens the conservation urgency of an endangered ungulate. Plos One 6(8):e22873. doi:10.1371/journal.pone.0022873
- Ikeda H, Senni K, Fujii N, Setoguchi H (2006) Refugia of *Potentilla matsumurae* (Rosaceae) located at high mountains in the Japanese archipelago. Mol Ecol 15:3731–3740. doi:10.1111/j.1365-294X.2006.03054.x
- IUCN (2012) Guidelines for application of IUCN Red List criteria at regional and national levels: version 4.0. IUCN, Gland
- Jiménez-Valverde A, Lobo JM, Hortal J (2008) Not good as they seem: the importance of concepts in species distribution modelling. Divers Distrib 14:885–890. doi:10.1111/j.1472-4642.2008.00496.x
- Körner C (2003) Alpine plant life: functional plant ecology of high mountain ecosystems. Springer, Berlin
- Kropf M, Kadereit JW, Comes HP (2003) Differential cycles of range contraction and expansion in European high mountain plants during the Late Quaternary: insights from *Pritzelago alpina* (L.) O. Kuntze (Brassicaceae). Mol Ecol 12:931–949. doi:10.1046/j.1365-294X.2003.01781.x
- Lenoir J, Gégout JC, Marquet PA, de Ruffray P, Brisse H (2008) A significant upward shift in plant species optimum elevation during the 20th century. Science 320:1768–1771. doi:10.1126/science.1156831
- Maiorano L, Falcucci A, Zimmermann NE, Psomas A, Pottier J, Baisero D, Rondinini C, Guisan A, Boitani L (2011) The future of terrestrial mammals in the Mediterranean basin under climate change. Philos Trans R Soc B 366:2681–2692. doi:10.1098/rstb.2011.0121
- Maiorano L, Amori G, Capula M, Falcucci A, Masi M, Montemaggiore A, Pottier J, Psomas A, Rondinini C, Russo D, Zimmermann NE, Boitani L, Guisan A (2013) Threats from climate change to terrestrial vertebrate hotspots in Europe. PLoS One 8:e74989. doi:10.1371/journal.pone.0074989
- McCann KS (2000) The diversity–stability debate. Nature 405:228–233. doi:10.1038/35012234
- McCullagh P, Nelder JA (1989) Generalized linear models. Chapman and Hall, London

- Médail F, Diadema K (2009) Glacial refugia influence plant diversity patterns in the Mediterranean Basin. *J Biogeogr* 36:1333–1345. doi:[10.1111/j.1365-2699.2008.02051.x](https://doi.org/10.1111/j.1365-2699.2008.02051.x)
- Médail F, Verlaque R (1997) Ecological characteristics and rarity of endemic plants from southeast France and Corsica: implications for biodiversity conservation. *Biol Conserv* 80:269–281. doi:[10.1016/S0006-3207\(96\)00055-9](https://doi.org/10.1016/S0006-3207(96)00055-9)
- Minuto L, Grassi F, Casazza G (2006) Ecogeographic and genetic evaluation of endemic species in the Maritime Alps: the case of *Moehringia lebrunii* and *M. sedoides* (Caryophyllaceae). *Pl Biosyst* 140:146–155. doi:[10.1080/11263500600756348](https://doi.org/10.1080/11263500600756348)
- Mouillot D, Bellwood DR, Baraloto C, Chave J, Galzin R, Harmelin-Vivien M, Kulbicki M, Lavergne S, Lavorel S, Mouquet N, Paine CET, Renaud J, Thuiller W (2013) Rare species support vulnerable functions in high-diversity ecosystems. *Plos One* 11(5):e1001569. doi:[10.1371/journal.pbio.1001569](https://doi.org/10.1371/journal.pbio.1001569)
- Musil R (1985) Paleobiography of terrestrial communities in Europe during the last glacial. *Acta Musei Natl Pragae XLI B* 1–2:25–56
- Nogués-Bravo D (2009) Predicting the past distribution of species climatic niches. *Global Ecol Biogeogr* 18:521–531. doi:[10.1111/j.1466-8238.2009.00476.x](https://doi.org/10.1111/j.1466-8238.2009.00476.x)
- Normand S, Ricklefs RE, Skov F, Bladt J, Tackenberg O, Svenning JC (2011) Postglacial migration supplements climate in determining plant species ranges in Europe. *Proc R Soc* 278:3644–3653. doi:[10.1098/rspb.2010.2769](https://doi.org/10.1098/rspb.2010.2769)
- Ozenda P (2009) On the genesis of the plant population in the Alps: new or critical aspects. *C R Biol* 332:1092–1103. doi:[10.1016/j.crvi.2009.09.018](https://doi.org/10.1016/j.crvi.2009.09.018)
- Patsiou T, Conti E, Zimmermann NE, Theodoridis S, Randin C (2014) Topo-climatic microrefugia explain the persistence of a rare endemic plant in the Alps during the last 21 millennia. *Glob Change Biol* 20:2286–2300. doi:[10.1111/gcb.12515](https://doi.org/10.1111/gcb.12515)
- Pawlowski B (1970) Remarques sur l'endémisme dans les Alpes et les Carpates. *Vegetatio* 21:181–243
- Pereira HM, Leadley PW, Proença V, Alkemade R et al (2010) Scenarios for global biodiversity in 21st century. *Science* 330:1469–1501. doi:[10.1126/science.1196624](https://doi.org/10.1126/science.1196624)
- Phillips SJ, Anderson RP, Schapire RE (2006) Maximum entropy modeling of species geographic distributions. *Ecol Model* 190:231–259. doi:[10.1016/j.ecolmodel.2005.03.026](https://doi.org/10.1016/j.ecolmodel.2005.03.026)
- R Development Core Team (2008) R: A language and environment for statistical computing. R Foundation for Statistical Computing, Vienna
- Randin CF, Engler R, Normand S, Zappa M, Zimmermann NE, Pearman PB, Vittoz P, Thuiller W, Guisan A (2009) Climate change and plant distribution: local models predict high-elevation persistence. *Glob Change Biol* 15:1557–1569. doi:[10.1111/j.1365-2486.2008.01766.x](https://doi.org/10.1111/j.1365-2486.2008.01766.x)
- Rebello H, Jones G (2010) Ground validation of presence-only modelling with rare species: a case study on barbastelles *Barbastella barbastellus* (Chiroptera: Vespertilionidae). *J Appl Ecol* 47:410–420. doi:[10.1111/j.1365-2664.2009.01765.x](https://doi.org/10.1111/j.1365-2664.2009.01765.x)
- Schwartz MW (2012) Using niche models with climate projections to inform conservation management decisions. *Biol Conserv* 155:149–156. doi:[10.1016/j.biocon.2012.06.011](https://doi.org/10.1016/j.biocon.2012.06.011)
- Sillero N, Carretero MA (2012) Modelling the past and future distribution of contracting species. The Iberian lizard *Podarcis carbonelli* (Squamata: Lacertidae) as a case study. *Zool Anz* 252:289–298. doi:[10.1016/j.jcz.2012.08.004](https://doi.org/10.1016/j.jcz.2012.08.004)
- Skov F, Svenning JC (2004) Potential impact of climatic change on the distribution of forest herbs in Europe. *Ecography* 27:366–380. doi:[10.1111/j.0906-7590.2004.03823.x](https://doi.org/10.1111/j.0906-7590.2004.03823.x)
- Solomon S, Qin D, Manning M, Alley RB, Bernsten T, Bindoff NL, Chen Z, Chidthaisong A, Gregory JM, Hegerl GC, Heimann M, Hewitson B, Hoskins BJ, Joos F, Jouzel J, Kattsov V, Lohmann U, Matsuno T, Molina M, Nicholls N, Overpeck J, Raga G, Ramaswamy V, Ren J, Rusticucci M, Somerville R, Stocker TF, Whetton P, Wood RA, Wratt D (2007) Technical Summary. In: Solomon S, Qin D, Manning M, Chen Z, Marquis M, Averyt KB, Tignor M, Miller HL (eds) *Climate change 2007: the physical science basis. Contribution of working group I to the fourth assessment report of the intergovernmental panel on climate change*. Cambridge University Press, Cambridge
- Stewart JR, Lister AM, Barnes I, Dalén L (2009) Refugia revisited: individualistic responses of species in space and time. *Proc R Soc B* 277:661–671. doi:[10.1098/rspb.2009.1272](https://doi.org/10.1098/rspb.2009.1272)
- Stokland JK, Halvorsen R, Stoa B (2011) Species distribution modelling—effect of design and sample size of pseudo-absence observations. *Ecol Model* 222:1800–1809. doi:[10.1016/j.ecolmodel.2011.02.025](https://doi.org/10.1016/j.ecolmodel.2011.02.025)
- Susanna A, Garcia-Jacas N (2009) The tribe Cardueae. In: Funk VA, Susanna A, Stuessy T, Bayer R (eds) *Systematics, evolution, and biogeography of the compositae*. IAPT, Bratislava, pp 293–313
- Susanna A, Garcia-Jacas N, Hidalgo O, Vilatersana R, Garnatje T (2006) The Cardueae (Compositae) revisited: insights from ITS, trnL-trnF, and matK nuclear and chloroplast DNA analysis. *Ann Mo Bot Gard* 93:150–171. doi:[10.3417/0026-6493\(2006\)93\[150:TCCRIF\]2.0.CO;2](https://doi.org/10.3417/0026-6493(2006)93[150:TCCRIF]2.0.CO;2)
- Suttle KB, Thomsen MA, Power ME (2007) Species interactions reverse grassland responses to climate change. *Science* 315:640–642. doi:[10.1126/science.1136401](https://doi.org/10.1126/science.1136401)
- Svenning JC, Skov F (2004) Limited filling of the potential range in European tree species. *Ecol Lett* 7:565–573. doi:[10.1111/j.1461-0248.2004.00614.x](https://doi.org/10.1111/j.1461-0248.2004.00614.x)
- Swab RM, Regan HM, Keith DA, Regan TJ, Ooi MKJ (2012) Niche models tell half the story: spatial context and life history traits influence species responses to global change. *J Biogeogr* 39:1266–1277. doi:[10.1111/j.1365-2699.2012.02690.x](https://doi.org/10.1111/j.1365-2699.2012.02690.x)
- Terrab A, Schönswetter P, Talavera S, Vela E, Stuessy TF (2008) Range-wide phylogeography of *Juniperus thurifera* L., a presumptive keystone species of western Mediterranean vegetation during cold stages of the Pleistocene. *Mol Phylogenet Evol* 48:94–102. doi:[10.1016/j.ympev.2008.03.018](https://doi.org/10.1016/j.ympev.2008.03.018)
- Theurillat JP, Guisan A (2001) Potential impact of climate change on vegetation in the European Alps—a review. *Clim Change* 50:77–109. doi:[10.1023/A:1010632015572](https://doi.org/10.1023/A:1010632015572)
- Thomas CD, Cameron A, Green R, Bakkenes M et al (2004) Extinction risk from climate change. *Nature* 427:145–148. doi:[10.1038/nature02121](https://doi.org/10.1038/nature02121)
- Thuiller W, Lavorel S, Araújo MB (2005a) Niche properties and geographical extent as predictors of species sensitivity to climate change. *Glob Ecol Biogeogr* 14:347–357. doi:[10.1111/j.1466-822X.2005.00162.x](https://doi.org/10.1111/j.1466-822X.2005.00162.x)
- Thuiller W, Lavorel S, Araújo MB, Sykes MT, Prentice IC (2005b) Climate change threats to plant diversity in Europe. *PNAS* 102:8245–8250. doi:[10.1073/pnas.0409902102](https://doi.org/10.1073/pnas.0409902102)
- Thuiller W, Lafourcade B, Araújo MB (2008) Modelling manual for BIOMOD. Laboratoire d'Ecologie Alpine, Université Joseph Fourier, Grenoble
- Thuiller W, Lafourcade B, Engler R, Araújo MB (2009) BIOMOD: a platform for ensemble forecasting of species distribution. *Ecography* 32:369–373. doi:[10.1111/j.1600-0587.2008.05742.x](https://doi.org/10.1111/j.1600-0587.2008.05742.x)
- Tribsch A (2004) Areas of endemism of vascular plants in the Eastern Alps in relation to Pleistocene glaciation. *J Biogeogr* 31:747–760. doi:[10.1111/j.1365-2699.2004.01065.x](https://doi.org/10.1111/j.1365-2699.2004.01065.x)
- Tribsch A, Schönswetter P (2003) Patterns of endemism and comparative phylogeography confirm paleo-environmental evidence for Pleistocene refugia in the Eastern Alps. *Taxon* 52:477–497. doi:[10.2307/3647447](https://doi.org/10.2307/3647447)
- Van Vuuren DP, Edmonds J, Kainuma M, Riahi K, Thomson A, Hibbard K, Hurtt GC, Kram T, Krey V, Lamarque JF, Masui T, Meinshausen M, Nakicenovic N, Smith SJ, Rose SK (2011) The

- representative concentration pathways: an overview. *Clim Change* 109:5–31. doi:[10.1007/s10584-011-0148-z](https://doi.org/10.1007/s10584-011-0148-z)
- Wisz MS, Guisan A (2009) Do pseudo-absence selection strategies influence species distribution models and their predictions? *An information-theoretic approach based on simulated data. BMC Ecol* 9:8. doi:[10.1186/1472-6785-9-8](https://doi.org/10.1186/1472-6785-9-8)