



**University of
Zurich**^{UZH}

**Zurich Open Repository and
Archive**

University of Zurich
University Library
Strickhofstrasse 39
CH-8057 Zurich
www.zora.uzh.ch

Year: 2016

Microbes and asthma: Opportunities for intervention

Smits, Hermelijn H ; Hiemstra, Pieter S ; Prazeres da Costa, Clarissa ; Ege, Markus ; Edwards, Michael ; Garn, Holger ; Howarth, Peter H ; Jartti, Tuomas ; de Jong, Esther C ; Maizels, Rick M ; Marsland, Ben J ; McSorley, Henry J ; Müller, Anne ; Pfefferle, Petra I ; Savelkoul, Huub ; Schwarze, Jürgen ; Unger, Wendy W J ; von Mutius, Erika ; Yazdanbakhsh, Maria ; Taube, Christian

Abstract: The worldwide incidence and prevalence of asthma continues to increase. Asthma is now understood as an umbrella term for different phenotypes or endotypes, which arise through different pathophysiologic pathways. Understanding the many factors contributing to development of the disease is important for the identification of novel therapeutic targets for the treatment of certain asthma phenotypes. The hygiene hypothesis has been formulated to explain the increasing prevalence of allergic disease, including asthma. This hypothesis postulates that decreased exposure at a young age to certain infectious agents as a result of improved hygiene, increased antibiotic use and vaccination, and changes in lifestyle and dietary habits is associated with changes in the immune system, which predispose subjects to allergy. Many microbes, during their coevolution with human subjects, developed mechanisms to manipulate the human immune system and to increase their chances of survival. Improving models of asthma, as well as choosing adequate end points in clinical trials, will lead to a more complete understanding of the underlying mechanisms, thus providing an opportunity to devise primary and secondary interventions at the same time as identifying new molecular targets for treatment. This article reports the discussion and conclusion of a workshop under the auspices of the Netherlands Lung Foundation to extend our understanding of how modulation of the immune system by bacterial, parasitic, and viral infections might affect the development of asthma and to map out future lines of investigation.

DOI: <https://doi.org/10.1016/j.jaci.2016.01.004>

Posted at the Zurich Open Repository and Archive, University of Zurich

ZORA URL: <https://doi.org/10.5167/uzh-124568>

Journal Article

Published Version

Originally published at:

Smits, Hermelijn H; Hiemstra, Pieter S; Prazeres da Costa, Clarissa; Ege, Markus; Edwards, Michael; Garn, Holger; Howarth, Peter H; Jartti, Tuomas; de Jong, Esther C; Maizels, Rick M; Marsland, Ben J; McSorley, Henry J; Müller, Anne; Pfefferle, Petra I; Savelkoul, Huub; Schwarze, Jürgen; Unger, Wendy W J; von Mutius, Erika; Yazdanbakhsh, Maria; Taube, Christian (2016). Microbes and asthma: Opportunities for intervention. *Journal of Allergy and Clinical Immunology*, 137(3):690-697.

DOI: <https://doi.org/10.1016/j.jaci.2016.01.004>

Microbes and asthma: Opportunities for intervention



Hermelijn H. Smits, PhD,^a Pieter S. Hiemstra, PhD,^b Clarissa Prazeres da Costa, MD, PhD,^c Markus Ege, MPH,^d Michael Edwards, PhD,^e Holger Garn, PhD,^f Peter H. Howarth, MD,^g Tuomas Jartti, MD,^h Esther C. de Jong, PhD,ⁱ Rick M. Maizels, PhD,^j Ben J. Marsland, PhD,^k Henry J. McSorley, PhD,^j Anne Müller, PhD,^l Petra I. Pfefferle, PhD,^m Huub Savelkoul, PhD,ⁿ Jürgen Schwarze, MD,^o Wendy W. J. Unger, PhD,^p Erika von Mutius, MD, MSc,^d Maria Yazdanbakhsh, PhD,^a and Christian Taube, MD, PhD^b
 Leiden, Amsterdam, Wageningen, and Rotterdam, The Netherlands, Munich and Marburg, Germany, London, Edinburgh, Southampton, and Glasgow, United Kingdom, Turku, Finland, and Epalinges and Zurich, Switzerland

The worldwide incidence and prevalence of asthma continues to increase. Asthma is now understood as an umbrella term for different phenotypes or endotypes, which arise through different pathophysiologic pathways. Understanding the many factors contributing to development of the disease is important for the identification of novel therapeutic targets for the treatment of certain asthma phenotypes. The hygiene hypothesis has been formulated to explain the increasing prevalence of allergic disease, including asthma. This hypothesis postulates that decreased exposure at a young age to certain infectious agents as a result of improved hygiene, increased antibiotic use and vaccination, and changes in lifestyle and dietary habits is associated with changes in the immune system, which predispose subjects to allergy. Many microbes, during their coevolution with human subjects, developed mechanisms to manipulate the human

immune system and to increase their chances of survival. Improving models of asthma, as well as choosing adequate end points in clinical trials, will lead to a more complete understanding of the underlying mechanisms, thus providing an opportunity to devise primary and secondary interventions at the same time as identifying new molecular targets for treatment. This article reports the discussion and conclusion of a workshop under the auspices of the Netherlands Lung Foundation to extend our understanding of how modulation of the immune system by bacterial, parasitic, and viral infections might affect the development of asthma and to map out future lines of investigation. (*J Allergy Clin Immunol* 2016;137:690-7.)

Key words: Hygiene hypothesis, asthma, sensitization, microbes, microbiome, helminths, viruses, immune regulation

From the Departments of ^aParasitology and ^bPulmonology, Leiden University Medical Center; ^cthe Institute of Medical Microbiology, Immunology and Hygiene, Technische Universität München; ^dDr von Hauner Children's Hospital, Ludwig-Maximilians-Universität München; ^ethe Department of Respiratory Medicine & Wright-Fleming Institute of Infection and Immunity, Imperial College London; ^fthe Institute for Laboratory Medicine and Pathobiochemistry, Philipps University of Marburg; ^gthe Academic Unit of Clinical and Experimental Sciences, University of Southampton Faculty of Medicine, University Hospital Southampton; ^hthe Department of Pediatrics, Turku University Hospital; ⁱthe Department of Cell Biology and Histology, Academic Medical Center, University of Amsterdam; ^jthe Institute of Infection, Immunity and Inflammation, University of Glasgow; ^kthe Faculty of Biology and Medicine, University of Lausanne, Service de Pneumologie, CHUV, Epalinges; ^lthe Institute of Molecular Cancer Research, University of Zurich; ^mComprehensive Biomaterial Bank Marburg (CBBMR), Philipps University Marburg; ⁿthe Cell Biology and Immunology Group, Wageningen University; ^othe MRC-Centre for Inflammation Research, Queens Medical Research Institute, University of Edinburgh; and ^pthe Department of Paediatrics, Erasmus Medical Center, Rotterdam.

Disclosure of potential conflict of interest: H. H. Smits has received grants and travel support from the Dutch Lung Foundation. P. S. Hiemstra has received travel support from Lung Foundation Netherlands and grants from Boehringer Ingelheim and Galapagos. C. Prazeres da Costa has received travel support from Longfonds, Netherlands. M. Ege has received grants from the German Federal Ministry of Research (BMBF), the German Research Foundation (DFG), the European Commission, the European Research Council, and Friesland Campina and has received royalties from Protectimmun GmbH. M. Edwards is a grant reviewer for NMRC Singapore; has consultant arrangements with Chiesi; is employed by GlaxoSmithKline; has received grants from GlaxoSmithKline, Chiesi, Merck, and Pfizer; has received payment for lectures from Pfizer; and has received travel support from the European Academy of Allergy and Clinical Immunology, the European Respiratory Society, and the American Academy of Allergy, Asthma & Immunology. H. Garn has received a grant from the German Research Foundation (DFG; SFB/TR22) and has

received travel support from Lung Foundation Netherlands. T. Jartti has received a grant from the Academy of Finland. E. C. de Jong is employed by AMC, University of Amsterdam. R. M. Maizels is employed by the University of Edinburgh and has received grants from Wellcome Trust and the Rainin Foundation. B. Marsland has received travel support from the Netherlands Lung Foundation. H. J. McSorley has received a grant from Asthma UK. J. Schwarze has received travel support from the Netherlands Lung Foundation; has consultant arrangements with MEDA, GlaxoSmithKline, and Bausch & Lomb; is employed by the University of Edinburgh; has received grants from the Medical Research Council and the Wellcome Trust; has received payment for lectures from Abbvie and Thermo Fisher; and has received support for allergy study days for clinicians from Thermo Fisher Scientific, MEDA Pharmaceuticals, Nutricia, Mead Johnson Nutrition, GlaxoSmithKline, Abbott, IMed-Emerade, Allergy Therapeutics, and Stallergenes. E. von Mutius has received grants from European Research Council, the German Research Foundation, and Friesland-Campina; is on the Editorial Board for the *New England Journal of Medicine*; has served as an Associate Editor for the *Journal of Allergy and Clinical Immunology*; has provided expert testimony for the European Research Council and European Commission, UK Research Excellence Framework, AUKCAR, Messerli Research Institute, and University of Tampere; and has received payment for lectures from Novartis, Mundipharma, DOC Congress SRL, Oekosoziales Forum Oberösterreich. The rest of the authors declare that they have no relevant conflicts of interest.

Received for publication October 18, 2015; revised January 6, 2016; accepted for publication January 19, 2016.

Corresponding author: Hermelijn H. Smits, PhD, Department of Parasitology, Leiden University Medical Center, 2333 ZA Leiden, The Netherlands. E-mail: h.h.smits@lumc.nl

The CrossMark symbol notifies online readers when updates have been made to the article such as errata or minor corrections

0091-6749/\$36.00

© 2016 American Academy of Allergy, Asthma & Immunology

<http://dx.doi.org/10.1016/j.jaci.2016.01.004>

Abbreviations used

DC: Dendritic cell
RSV: Respiratory syncytial virus
TLR: Toll-like receptor
Treg: Regulatory T

In recent decades, there has been a marked increase in the incidence of many noncommunicable diseases, including asthma, which is now estimated to affect 300 million persons worldwide.¹ Patients with asthma experience a variable degree of airflow obstruction, breathlessness, and bronchial hyperresponsiveness associated with chronic airway inflammation and excessive mucus production. Various specific and unspecific triggers have been identified that can lead to an increase in inflammation, obstruction, and symptoms. Traditionally, asthma, especially allergic asthma, has been considered an inflammatory disease associated with T_H2 cells, production of IgE antibodies, accumulation of eosinophils in the lungs, and goblet cell hyperplasia. It is now recognized that asthma is a complex syndrome in which many different phenotypes exist, including early-onset allergic asthma, late-onset eosinophilic asthma, and exercise-induced, obesity-related, and noneosinophilic asthma.²

Recently, the definition of asthma has shifted further with the introduction of endotypes, which distinguish asthma variants by their underlying molecular mechanisms. Probably the best described endotype is the type 2–induced form of the disease.³ Other endotypes are less well defined and include patients without type 2–induced airway inflammation (probably driven by T_H1 or T_H17 cells) and allergic bronchopulmonary mycosis as an asthma endotype.⁴

Most asthmatic patients have a mild form of the disease, which can be managed with inhaled corticosteroids and long-acting β -agonists. However, patients with more severe disease and particularly those with a non-T_H2 endotype might not respond well to currently available therapies. Particularly in asthmatic patients, personalized medicine might open novel approaches to accommodate the heterogeneity of the disease. Better understanding of mechanisms and endotypes will provide opportunities for both prevention and causal treatment.

In the last years, interactions of microbes, including worm parasites, with their host have been identified: exposure to microorganisms not only triggers but also effectively suppresses immune responses, and beneficial effects of microorganisms are increasingly recognized and mechanistically understood. Strategies are emerging to potentially implement these effects in novel interventions to prevent or treat allergic diseases, such as allergic asthma (Fig 1). A better understanding of the disease in its many guises at a basic level is needed to endorse such strategies and improve and refine interventions. In this context a group of clinicians and basic scientists with wide-ranging fields of expertise convened in Amersfoort, The Netherlands, under the auspices of the Netherlands Lung Foundation for a workshop to assess our current understanding of the disease and identify challenges and opportunities for the prevention and treatment of asthma, with microbial intervention as the guiding theme for the workshop.

HYGIENE HYPOTHESIS AND “OLD FRIENDS” HYPOTHESIS

The so-called hygiene hypothesis is frequently invoked to help explain the increasing prevalence of asthma. The hypothesis has

its origins in observations published in 1989 by Strachan,⁵ who noted that decreasing family size was associated with hay fever in developed countries and suggested that this might be related to a lower degree of sibling-related childhood infections and microbial exposure. In extension of the hygiene hypothesis, Rook⁶ has postulated the “old friends” hypothesis, in which many infectious agents and microbes in their coevolution with human subjects have developed mechanisms to modulate and evade the host immune system (Fig 1). Immunomodulatory microorganisms have been described to activate various cells of the regulatory network, such as regulatory T (Treg) cells and regulatory B cells, and to modulate or even reprogram certain antigen-presenting cells, leading to tolerogenic dendritic cells (DCs), alternatively activated macrophages, or both. A more detailed understanding of how these infectious agents accomplish this can provide indicators for primary prevention strategies and might help to identify new molecular targets for novel treatments. This is especially relevant because in patients with various noncommunicable inflammatory diseases, such as asthma, these regulatory networks seemed to be underrepresented and poorly developed.

Rural exposure and “archaic” microbiome

As discussed above, microbes (“old friends”) form a central part of the (extended) hygiene hypothesis.^{6,7} Interestingly, this has not so much to do with personal hygiene (as often interpreted from the hygiene hypothesis) because a recent study showed that personal or home cleanliness was not associated with a risk of asthma or allergy.⁸ The “old friends” mostly represent a group of microbes with which the human race has coevolved and that in the past 50 years were rapidly lost because of changes in lifestyle, living conditions, or occupations. Prime candidates are microbes associated with rural living, such as farming, and various members of an archaic microbiome responsible for a richer composition of our personal microbial hemisphere, which includes compartments such as the gut, lung, and skin. From this perspective, helminths are regarded as a natural and ancient (evolutionary conserved) partner of the microbial community, which is still the case in many parts of the world but no longer in westernized countries. The likelihood that archaic microbes play an essential role in protection against asthma and allergic diseases is framed by several landmark studies.

Several studies noted that living on farms offers a protective effect against atopy, hay fever, and asthma, especially in children.⁹ Further analysis suggested a link with increased exposure to a variety of bacteria and fungi related to farming and protection from asthma.¹⁰ Interestingly, several gene-environment interactions were found for early farm exposure. A number of single nucleotide polymorphisms in children living in rural Europe were linked to farming, such as in the genes transcribing CD14 or Toll-like receptors (TLRs).¹¹ Remarkably, a recent farm study also reports on associations with the asthma risk alleles on chromosome 17q21, suggesting that the same genotype both constitutes a genetic risk to asthma and, at the same time, is susceptible to environmental influences.¹² This would imply options for future preventive strategies.

In addition, in particular, farm exposure during pregnancy seems to influence gene expression patterns by means of DNA methylation in specific asthma- and allergy-related genes, further contributing to its protective effect.¹³ In contrast, a higher




FIG 1. Schematic overview of the hygiene/"old friends" hypothesis. Several microbial signals, such as environmental exposure (rural/farm environment) and an archaic microbiota (containing more diverse and abundant bacteria and typical helminth parasites), provide strong signals to develop regulatory responses (tolerance), whereas an urbanized environment and/or a westernized microbiota does not, tipping the immune balance more toward inflammation. Tolerance leads to good health, whereas in its absence, an inflammatory profile leads to airway diseases, such as respiratory allergies and asthma. Composition and diversity of the microbiota are affected by diet. Viral infections, sinusitis, and air pollution trigger the immune system to move toward inflammation, whereas a diverse microbiota might reduce these inflammatory signals.

prevalence of asthma and a higher morbidity can be found in children living in inner cities. Various environmental risk factors have been recognized, including high levels of indoor allergen, pollutant, and endotoxin exposure,¹⁴ although part of this effect might be explained by differences in ethnicity and demographics.¹⁵

Also, the composition of the gut microbiota influences the development of allergic diseases, such as asthma. Indeed, intestinal gene expression and ultimately immune system development is different in "germ-free" mice, which lack a gut microbiome, compared with conventional mice.¹⁶ In addition, germ-free mice exhibit increased susceptibility to allergen-induced airway disease, and this has been linked to different mechanisms, such as increased IgE production and basophil numbers at mucosal sites, as well as induction of natural killer T cells.¹⁷⁻¹⁹ Also, in human subjects the composition of gut microbiota has been linked to airway disease because stool samples from babies that later have allergies and asthma have a different composition and contain less lactobacilli, Bacteroidetes and bifidobacteria.²⁰

Finally, early and chronic parasitic worm infections of or linked to the gastrointestinal tract protected against autoimmune diseases, such as multiple sclerosis²¹ and inflammatory bowel disease.²² This effect has also been observed for respiratory allergies and, in the case of hookworm infections, also for asthma.²³ Nevertheless, not all parasitic worm infections are protective: in particular, early and chronic infections tend to be protective, whereas in some studies low-burden and sporadic infections were associated with enhanced allergic reactions.²⁴ Interestingly, some studies have suggested a possible interaction between gut parasites and the microbiota because they both inhabit the same organ. For example, helminth infection in human subjects is linked to an increased diversity of the microbiota.^{25,26} Furthermore, murine *Heligmosomoides polygyrus* promotes the colonization of *Lactobacillus* species. Interestingly, helminth-modified microbiota mediated protection against experimental

allergic asthma.^{27,28} Knowledge on the interaction between worm parasites and the microbiota is still in its infancy, and identification of the factors that are crucially involved is the subject of future research.

Respiratory viruses

Unlike parasitic and bacterial infections, viral respiratory tract infections are not associated with protection against atopy or asthma. In contrast, bronchiolitis induced by respiratory syncytial virus (RSV) or rhinovirus has been consistently associated with increased risk of later asthma in numerous studies.^{29,30} For example, in a prospective cohort study of children hospitalized for RSV-induced bronchiolitis in the first year of life, Sigurs et al³¹ found that viral bronchiolitis resulting in hospitalization was significantly associated with asthma (defined as 3 episodes of bronchial obstruction) at 3 years of age, which persisted into early adulthood.³² The Childhood Origins of Asthma study³³ found that in children selected for a high risk of asthma, rhinovirus-induced wheeze was the strongest predictor of wheezing at the age of 3 years,³⁴ and this effect was carried over to 6 years of age.³⁵ Importantly, allergic sensitization (as indicated by measurement of allergen-specific IgE) preceded the onset of recurrent viral wheeze. These findings, also supported by others,^{36,37} also suggest that rhinovirus-induced wheezing can be used as a marker for children at risk of asthma.

Interestingly, some recent studies have suggested a putative interplay between viruses and pathogenic bacteria in the noses and upper respiratory tracts of young children based on positive associations between the bacteria *Haemophilus influenzae*, *Streptococcus pneumoniae*, *Staphylococcus aureus*, and *Moraxella catarrhalis* and/or rhinoviruses and RSV.^{38,39} Importantly, the presence of these pathogenic bacteria was associated with increased respiratory symptoms and asthma exacerbations.⁴⁰ However, the question remains whether the colonization of those pathogenic bacteria precedes viral respiratory tract infections and

asthma symptoms or whether this is the consequence of persistent viral infection in the upper airways. New preventive strategies could be designed based on disrupting those reinforcing interactions between viruses and pathogenic bacteria.

OPPORTUNITIES FOR INTERVENTION

Several new therapies in the pipelines of the pharmaceutical industry will enter the market in the coming years. These therapies cover a number of different molecular targets, thereby enabling the tailoring of therapy according to a particular endotype. However, many of these new therapies focus on controlling T_H2 induction and thus are unlikely to cover all patient needs.⁴¹ Therefore prevention of asthma might be a more efficient and sustainable approach to reducing disease burden in the long term.

Primary prevention

As discussed above, exposure to microorganisms might afford a potential protective effect, and a number of mechanisms have been proposed to explain the effect. An important question is whether these mechanisms can be harnessed to develop interventions to reduce the burden of asthma. One of the most promising approaches seems to be primary prevention. However, timing is crucial: after birth, the window of opportunity for primary prevention rapidly closes as the immune system matures. It is important to identify subjects who are at risk of asthma at an early stage to target such interventions effectively. Some studies suggest that preventive strategies should already be started before birth to increase their efficacy. Prenatal priming might induce or reset long-term epigenetic checkpoints, allowing the immature immune system to respond sufficiently to microbial stimuli and quickly develop a strong and sustained regulatory network.

Farm bacteria. A possible source of the bacterial diversity encountered in farm living is stable dust, a rich source of a highly diverse bacterial ecosystem. From this large pool of farm-related microorganisms, 2 species in particular, *Acinetobacter lwoffii* F78 and *Lactococcus lactis* G121, have been tested and potentially inhibited allergic reactions in mice.⁴² Interestingly, a recent study shows that farm dust reduced the production of innate type 2 cytokines by epithelial cells, which was attributed to the ubiquitin-modifying enzyme A20 in lung epithelium.⁴³ Not only neonate but also maternal exposure to microorganisms can reduce the risk of offspring having allergic diseases, such as asthma. Epigenetic changes after farm exposure might be responsible for increasing the number and function of cord blood Treg cells.⁴⁴ This might then lead to lower T_H2 cytokine secretion and lymphocyte proliferation on innate exposure. In the specific case of exposure to *A lwoffii* F78, a murine model suggested that maternal bacterial exposures were directly related to functional maternal TLR signaling, resulting in asthma protection in the progeny.^{45,46} The mechanism was IFN- γ dependent, possibly through protection from loss of IFN- γ promoter-associated histone 4 acetylation.⁴⁶

Bacteria in the lungs. Also, the lungs support a complex microbiota originating from inhaled microbes and flora from the digestive system.⁴⁷ Low microbial diversity in the lung has been found in patients with diseases such as asthma, with an increased proportion of Proteobacteria, such as *Haemophilus*, *Neisseria*, or *Streptococcus* species. Absence of a lung microbiota in germ-free

mice is associated with increased T_H2 responses and increased allergic airway disease, suggesting a strong inhibitory effect of the lung microbiota on T_H2 development.⁴⁸ Furthermore, new studies have linked the composition of the lung microbiome to therapy responses to corticosteroids by uncovering a difference in microbiota composition in steroid-responsive and steroid-resistant patients.⁴⁹ New therapies targeting the lung microbiome would be an interesting approach in the prevention or early treatment of asthma. However, in this stage more information is needed on the ideal composition of a “healthy” lung microbiome or alternatively which bacteria species should be avoided or removed before new therapeutic strategies can be designed.

Gut bacterial biodiversity and bacterial intestinal infections. Recently, published work on a murine model suggests that the gut microbiota metabolizes dietary fibers, resulting in increased circulating short-chain fatty acids.⁵⁰ Importantly, the authors demonstrated that short-chain fatty acids, through ligation of G protein-coupled receptor 41, ultimately induced seeding of the lungs with DCs with an impaired ability to promote T_H2 cell effector function (but high phagocytic capacity). Therefore these results suggest a mechanism whereby diet, especially fiber content, in association with the intestinal microbiota could have a direct influence on the development of asthma.

In addition to gut microbial biodiversity and diet, individual gut bacteria, such as *Helicobacter pylori*, have received considerable attention recently. *H pylori* infection is better known for its pathogenic properties, being linked to conditions such as peptic ulcer and gastric cancer.⁵¹ However, infection by *H pylori*, especially in early childhood, might confer benefits because protective effects of *H pylori* infection against the development of asthma and allergies have been described.^{52,53} Also, when mice were infected during the neonatal period, they were subsequently protected against the development of allergic airway disease.⁵⁴ The mechanisms by which this protection is enforced involve induction of Treg cells and reprogramming of DCs toward a tolerogenic phenotype.⁵⁵ Various persistence determinants of *H pylori*, such as the γ -glutamyl transpeptidase, vacuolating cytotoxin, and urease, have been shown to be critically important to these protective effects and are being evaluated for further therapeutic applications.^{56,57}

Evidence from clinical trials for the effectiveness of primary prevention by using probiotic strains is patchy at present. For example, trials are heterogeneous with respect to the timing of the intervention, bacterial strain, and use of monotherapy or combination therapy.⁵⁸ End points are also somewhat varied, with few studies focusing on asthma itself. Findings can vary according to the population included, as evidenced by the different results with *Lactobacillus GG* reported with the same treatment protocol and end points. In these studies an effect was seen for mothers who had at least 1 first-degree relative (or partner) with atopic eczema, allergic rhinitis, or asthma⁵⁹ but not when at least 1 family member (mother, father, or child) had atopic disease.⁶⁰ Taken together, the body of evidence to date suggests that a combined antenatal and postnatal approach is the most promising, with monotherapy with lactobacilli seeming to be the most promising agent, although there is plenty of room for improvement in the quality of the clinical trials. In particular, there is a need for validated surrogate end points for the development of asthma to provide a readout more quickly than would be possible with asthma itself. Finally, the ethics and safety concerns of prenatal exposure need to be taken into account.

Worm parasites. It is not surprising that organisms that coevolved with human subjects have developed mechanisms for modulating human inflammatory responses to promote their own survival. In general, worm parasites seem to be able to manipulate both innate immunity (by affecting TLR-induced responses and triggering the inflammasome)⁶¹ and adaptive immunity.⁶² In the case of adaptive immunity, unlike bacteria, helminths strongly induce T_H2 responses and increased IgE levels.⁶³ Interestingly, however, this T_H2 response is not associated with an increased predisposition to asthma. In fact, the IgE detected is primarily cross-reactive to carbohydrate epitopes present in parasites and not to the protein component of allergens and does not lead to mast cell degranulation. Currently, it is hypothesized that cross-reactive IgE might help prevent atopic sensitization and the development of allergic diseases in children with helminth infections. Knowledge on the processes that lead to IgE antibodies against cross-reactive carbohydrates might help implement these strategies in children at risk and prevent the development of high-affinity IgE molecules against proteins of allergens and thereby clinical symptoms.⁶⁴

Helminths are also master inducers of immunoregulatory processes.⁶⁵ Detailed studies in schistosome-infected mice or human subjects have shown increased numbers of regulatory B and Treg cells with an enhanced regulatory capacity, which, at least in mice, were crucial for protection against allergic airway disease.⁶⁶⁻⁶⁹ Also, infections with *H. polygyrus* or exposure to its excretory/secretory antigens (collected from cultures of live adults) could prevent experimentally induced airway allergy,^{70,71} and this has been linked to suppression of IL-33 release.⁷² Ultimately, the goal of future work in this line of research is to identify the immunomodulators within the *H. polygyrus* excretory/secretory fraction and translate these molecules into novel preventive therapies.

Interestingly, prenatal parasite exposure of the mother might also confer protection against allergic diseases, as illustrated by the higher incidence of eczema among Ugandan children born to schistosome-infected Ugandan mothers who had been treated with the deworming agent praziquantel during pregnancy.⁷³ Furthermore, mouse studies of maternal schistosome infection have pointed in the same direction and showed a crucial role for the maternal cytokine milieu within the placenta, ultimately determining whether allergic responses are promoted or suppressed in offspring.^{74,75}

Secondary and tertiary prevention

Given the effects of certain microbes in the context of primary interventions, the possibilities for secondary and tertiary prevention are less well investigated. Secondary prevention aims to detect and treat disease that has not yet become symptomatic, whereas tertiary prevention is directed at those who already have symptomatic disease to prevent further deterioration. These approaches include anti-inflammatory treatment, treatment with antiviral agents, and administration of nonviable lysates from the bacterial causative agents of acute respiratory tract illnesses. Recent trials suggest that systemic corticosteroid treatment is beneficial in children with first-time wheezing susceptible to rhinovirus infection, especially in those with high viral loads at presentation.^{76,77} The marked reduction of relapses and asthma during long-term follow-up supports the role for high rhinovirus load as an important marker of those children with early

pulmonary inflammation who might benefit from early intervention with anti-inflammatory treatment. The importance of airway inflammatory or allergen exposure control is also supported by a trial with the anti-IgE omalizumab, which nearly abolished autumn peaks in exacerbations typically caused by viral infections.⁷⁸

Prevention of repeated RSV-associated wheezing, which has also been associated with asthma, is another potential area for intervention. Monthly injections of palivizumab, a humanized anti-RSV mAb, has been shown to effectively reduce recurrent wheeze during the first year of life.⁷⁹ Because children with recurrent infant wheeze are at a high risk of asthma development, it could be speculated that the intervention also reduces the risk of asthma. Although this trial is undoubtedly valuable for understanding the mechanisms by which viral infections can lead to asthma, such treatments for RSV are expensive. Furthermore, in the case of rhinovirus, the variability in the virus (>100 serotypes) is a barrier to developing effective mAbs against infection.

Finally, evidence from mouse models suggests that bacterial lysates can reduce airway inflammation and lead to selective recruitment of Treg cells in the tracheal compartment,⁸⁰ whereas evidence from clinical studies suggests that treatment increases secretory IgA levels at mucosal surfaces. Analysis of clinical trials in this setting provide at best only weak support for a beneficial effect of the intervention. The most promising agent seems to be the bacterial lysate (OM-85 BV), which has been shown to prevent wheezing attacks provoked by acute respiratory tract illnesses in children.⁸¹ Similarly, studies have been conducted to treat patients with allergic rhinitis with *Trichuris suis* eggs⁸² or asthmatic patients with hookworm larvae⁸³ but did not show major clinical improvement of disease symptoms. It is unclear whether the lack of effect can be explained by the type of worm, dosing, timing, duration, or disease state.^{24,84} In general, the studies conducted to date often are inadequately designed or lack statistical power.

Treatment of asthma

Increasing knowledge of the immunomodulatory effects of microbial-host interactions might also offer a chance to develop novel therapeutic treatments for asthmatic patients. Indeed, treatment of the disease is mainly based on a one-size-fits-all approaches, mostly relying on inhaled corticosteroids as anti-inflammatory treatment. One example of transferring the understanding of the pathophysiologic mechanism into therapeutic approaches has been the development and use of mAbs for the treatment of severe asthma. Indeed, several novel opportunities are now available to specifically target type 2–driven inflammation.⁴¹ The increasing knowledge of the effect of microbial-host interactions on allergy and asthma might also result in novel therapeutic approaches. Whether microbial-based interventions in patients are useful first needs to be assessed in therapeutic mouse models. Indeed, one promising study showed that microbial interventions also work as a treatment for experimental allergic airway inflammation.⁵⁰ Treatment using the short-chain fatty acid propionate modulates the intestinal microbiome and effectively treats allergic airway disease in adult mice, suggesting that microbial interventions might not only be used for preventive approaches.⁵⁰ However, thus far, only few data are available from human patients. Bacteria-derived lysates have been used in infancy to prevent the development of atopy,⁸⁵ but only few studies have

investigated these compounds in patients with already established disease. Compounds targeting certain TLRs have been used in patients with asthma, with somewhat conflicting results. Administration of a TLR9 agonist resulted in improved asthma control during steroid reduction in patients receiving moderate- or high-dose inhaled steroids,⁸⁶ whereas a different TLR9 agonist showed no additional benefit in patients with insufficiently controlled moderate-to-severe allergic asthma.⁸⁷ Whether microbial compounds will be a useful addition to the current therapeutic options needs to be determined in further studies.

CONCLUSIONS

For the foreseeable future, asthma will remain a disease difficult to prevent or cure. Microbial infections and certain microbial metabolites and secretions seem to point at protection against asthma and offer numerous interesting opportunities. Primary or secondary prevention seems the most promising approach to reduce the overall burden of disease. Identification of populations at risk is of utmost importance to ensure that primary prevention is delivered where it is most needed. Indeed, there is some evidence to support the effectiveness of interventions in pregnant women, although ethical and safety issues would need to be carefully addressed, and the risk/balance equation would need to be accurately assessed. Beyond the primary prevention setting, there are opportunities for secondary or tertiary intervention, but earlier interventions are more likely to be successful. Finally, in the treatment setting the goal of new asthma treatments should be disease modification. In this sense a focus on underlying immune mechanisms would seem the approach most likely to deliver promising results, but the effects of alterations of the structural compartment should not be ignored. A shift to animal models that better represent the complex conditions and phenotypes of human asthma would not only enhance our understanding of the disease but also help identify better therapeutic candidates. Given that asthma develops over a long period of time, robust clinical trials can be challenging but are needed to assess the effectiveness of novel therapeutic approaches. In addition, it is necessary to elucidate whether these novel approaches only offer prevention of airway disease or whether they also offer a therapeutic benefit in patients with already established disease.

We thank the Lung Foundation Netherlands for hosting and organizing this 2-day workshop in Amersfoort, The Netherlands.

REFERENCES

1. Boulet LP, FitzGerald JM, Reddel HK. The revised 2014 GINA strategy report: opportunities for change. *Curr Opin Pulm Med* 2015;21:1-7.
2. Wenzel SE. Complex phenotypes in asthma: current definitions. *Pulm Pharmacol Ther* 2013;26:710-5.
3. Agache I, Sugita K, Morita H, Akdis M, Akdis CA. The complex type 2 endotype in allergy and asthma: from laboratory to bedside. *Curr Allergy Asthma Rep* 2015;15:29.
4. Lotvall J, Akdis CA, Bacharier LB, Björner L, Casale TB, Custovic A, et al. Asthma endotypes: a new approach to classification of disease entities within the asthma syndrome. *J Allergy Clin Immunol* 2011;127:355-60.
5. Strachan DP. Hay fever, hygiene, and household size. *BMJ* 1989;299:1259-60.
6. Rook GA. 99th Dahlem conference on infection, inflammation and chronic inflammatory disorders: Darwinian medicine and the 'hygiene' or 'old friends' hypothesis. *Clin Exp Immunol* 2010;160:70-9.
7. Brown EM, Arrieta MC, Finlay BB. A fresh look at the hygiene hypothesis: how intestinal microbial exposure drives immune effector responses in atopic disease. *Semin Immunol* 2013;25:378-87.
8. Weber J, Illi S, Nowak D, Schierl R, Holst O, von Mutius E, et al. Asthma and the hygiene hypothesis. Does cleanliness matter? *Am J Respir Crit Care Med* 2015;191:522-9.
9. von Mutius E, Vercelli D. Farm living: effects on childhood asthma and allergy. *Nat Rev Immunol* 2010;10:861-8.
10. Ege MJ, Mayer M, Normand AC, Genuneit J, Cookson WO, Braun-Fahrlander C, et al. Exposure to environmental microorganisms and childhood asthma. *N Engl J Med* 2011;364:701-9.
11. Lluís A, Schaub B. Lesson from the farm environment. *Curr Opin Allergy Clin Immunol* 2012;12:158-63.
12. Loss GJ, Depner M, Hose AJ, Genuneit J, Karvonen AM, Hyvärinen A, et al. The early development of wheeze: environmental determinants and genetic susceptibility at 17q21. *Am J Respir Crit Care Med* 2015 [Epub ahead of print].
13. Michel S, Busato F, Genuneit J, Pekkanen J, Dalphin JC, Riedler J, et al. Farm exposure and time trends in early childhood may influence DNA methylation in genes related to asthma and allergy. *Allergy* 2013;68:355-64.
14. Gergen PJ, Teach SJ, Togias A, Busse WW. Reducing exacerbations in the inner city: lessons from the Inner-City Asthma Consortium (ICAC). *J Allergy Clin Immunol Pract* 2016;4:22-6.
15. Keet CA, McCormack MC, Pollack CE, Peng RD, McGowan E, Matsui EC. Neighborhood poverty, urban residence, race/ethnicity, and asthma: Rethinking the inner-city asthma epidemic. *J Allergy Clin Immunol* 2015;135:655-62.
16. Gaboriau-Routhiau V, Rakotobe S, Lecuyer E, Mulder I, Lan A, Bridonneau C, et al. The key role of segmented filamentous bacteria in the coordinated maturation of gut helper T cell responses. *Immunity* 2009;31:677-89.
17. Cahenzli J, Koller Y, Wyss M, Geuking MB, McCoy KD. Intestinal microbial diversity during early-life colonization shapes long-term IgE levels. *Cell Host Microbe* 2013;14:559-70.
18. Hill DA, Artis D. The influence of commensal bacteria-derived signals on basophil-associated allergic inflammation. *Gut Microbes* 2013;4:76-83.
19. Olszak T, An D, Zeissig S, Vera MP, Richter J, Franke A, et al. Microbial exposure during early life has persistent effects on natural killer T cell function. *Science* 2012;336:489-93.
20. Huang YJ, Boushey HA. The microbiome in asthma. *J Allergy Clin Immunol* 2015;135:25-30.
21. Correale J, Farez MF. The impact of parasite infections on the course of multiple sclerosis. *J Neuroimmunol* 2011;233:6-11.
22. Koloski NA, Bret L, Radford-Smith G. Hygiene hypothesis in inflammatory bowel disease: a critical review of the literature. *World J Gastroenterol* 2008;14:165-73.
23. Leonardi-Bee J, Pritchard D, Britton J. Asthma and current intestinal parasite infection: systematic review and meta-analysis. *Am J Respir Crit Care Med* 2006;174:514-23.
24. Wammes LJ, Mpairwe H, Elliott AM, Yazdanbakhsh M. Helminth therapy or elimination: epidemiological, immunological, and clinical considerations. *Lancet Infect Dis* 2014;14:1150-62.
25. Lee SC, Tang MS, Lim YA, Choy SH, Kurtz ZD, Cox LM, et al. Helminth colonization is associated with increased diversity of the gut microbiota. *PLoS Negl Trop Dis* 2014;8:e2880.
26. Cantacessi C, Giacomini P, Croese J, Zakrzewski M, Sotillo J, McCann L, et al. Impact of experimental hookworm infection on the human gut microbiota. *J Infect Dis* 2014;210:1431-4.
27. Reynolds LA, Smith KA, Filbey KJ, Hargrett-Young JP, Redpath SA, et al. Commensal-pathogen interactions in the intestinal tract: lactobacilli promote infection with, and are promoted by, helminth parasites. *Gut Microbes* 2014;5:522-32.
28. Zaiss MM, Rapin A, Lebon L, Dubey LK, Mosconi I, Sarter K, et al. The intestinal microbiota contributes to the ability of helminths to modulate allergic inflammation. *Immunity* 2015;43:998-1010.
29. Feldman AS, Hartert TV, Gebretsadik T, Carroll KN, Minton PA, Woodward KB, et al. Respiratory severity score separates upper versus lower respiratory tract infections and predicts measures of disease severity. *Pediatr Allergy Immunol Pulmonol* 2015;28:117-20.
30. Jartti T, Korppi M. Rhinovirus-induced bronchiolitis and asthma development. *Pediatr Allergy Immunol* 2011;22:350-5.
31. Sigurs N, Bjarnason R, Sigurbergsson F, Kjellman B, Björkstén B. Asthma and immunoglobulin E antibodies after respiratory syncytial virus bronchiolitis: a prospective cohort study with matched controls. *Pediatrics* 1995;95:500-5.
32. Sigurs N, Aljassim F, Kjellman B, Robinson PD, Sigurbergsson F, Bjarnason R, et al. Asthma and allergy patterns over 18 years after severe RSV bronchiolitis in the first year of life. *Thorax* 2010;65:1045-52.
33. Lemanske RF Jr. The childhood origins of asthma (COAST) study. *Pediatr Allergy Immunol* 2002;13(suppl 15):38-43.

34. Lemanske RF Jr, Jackson DJ, Gangnon RE, Evans MD, Li Z, Shult PA, et al. Rhinovirus illnesses during infancy predict subsequent childhood wheezing. *J Allergy Clin Immunol* 2005;116:571-7.
35. Jackson DJ, Gangnon RE, Evans MD, Roberg KA, Anderson EL, Pappas TE, et al. Wheezing rhinovirus illnesses in early life predict asthma development in high-risk children. *Am J Respir Crit Care Med* 2008;178:667-72.
36. Jartti T, Kuusipalo H, Vuorinen T, Soderlund-Venermo M, Allander T, Waris M, et al. Allergic sensitization is associated with rhinovirus-, but not other virus-, induced wheezing in children. *Pediatr Allergy Immunol* 2010;21:1008-14.
37. Soto-Quiros M, Avila L, Platts-Mills TA, Hunt JF, Erdman DD, Carper H, et al. High titers of IgE antibody to dust mite allergen and risk for wheezing among asthmatic children infected with rhinovirus. *J Allergy Clin Immunol* 2012;129:1499-505.
38. van den Bergh MR, Biesbroek G, Rossen JW, de Steenhuijsen Piters WA, Bosch AA, van Gils EJ, et al. Associations between pathogens in the upper respiratory tract of young children: interplay between viruses and bacteria. *PLoS One* 2012;7:e47711.
39. Skevaki CL, Tsialta P, Trochoutsou AI, Logotheti I, Makrinioti H, Taka S, et al. Associations between viral and bacterial potential pathogens in the nasopharynx of children with and without respiratory symptoms. *Pediatr Infect Dis J* 2015;34:1296-301.
40. Kloefer KM, Lee WM, Pappas TE, Kang TJ, Vrtis RF, Evans MD, et al. Detection of pathogenic bacteria during rhinovirus infection is associated with increased respiratory symptoms and asthma exacerbations. *J Allergy Clin Immunol* 2014;133:1301-7.
41. van Buul AR, Taube C. Treatment of severe asthma: entering the era of targeted therapy. *Expert Opin Biol Ther* 2015;15:1713-25.
42. Debarry J, Garn H, Hanuszkiewicz A, Dickgreber N, Blumer N, von Mutius E, et al. *Acinetobacter lwoffii* and *Lactococcus lactis* strains isolated from farm cowsheds possess strong allergy-protective properties. *J Allergy Clin Immunol* 2007;119:1514-21.
43. Schuijs MJ, Willart MA, Vergote K, Gras D, Deswarte K, Ege MJ, et al. Farm dust and endotoxin protect against allergy through A20 induction in lung epithelial cells. *Science* 2015;349:1106-10.
44. Schaub B, Liu J, Hoppler S, Schleich I, Huehn J, Olek S, et al. Maternal farm exposure modulates neonatal immune mechanisms through regulatory T cells. *J Allergy Clin Immunol* 2009;123:774-82.
45. Brand S, Teich R, Dicke T, Harb H, Yildirim AO, Tost J, et al. Epigenetic regulation in murine offspring as a novel mechanism for transmaternal asthma protection induced by microbes. *J Allergy Clin Immunol* 2011;128:618-25.
46. Conrad ML, Ferstl R, Teich R, Brand S, Blumer N, Yildirim AO, et al. Maternal TLR signaling is required for prenatal asthma protection by the nonpathogenic microbe *Acinetobacter lwoffii* F78. *J Exp Med* 2009;206:2869-77.
47. Marsland BJ, Yadava K, Nicod LP. The airway microbiome and disease. *Chest* 2013;144:632-7.
48. Gollwitzer ES, Saglani S, Trompette A, Yadava K, Sherburn R, McCoy KD, et al. Lung microbiota promotes tolerance to allergens in neonates via PD-L1. *Nat Med* 2014;20:642-7.
49. Goleva E, Jackson LP, Harris JK, Robertson CE, Sutherland ER, Hall CF, et al. The effects of airway microbiome on corticosteroid responsiveness in asthma. *Am J Respir Crit Care Med* 2013;188:1193-201.
50. Trompette A, Gollwitzer ES, Yadava K, Sichelstiel AK, Sprenger N, Ngom-Bru C, et al. Gut microbiota metabolism of dietary fiber influences allergic airway disease and hematopoiesis. *Nat Med* 2014;20:159-66.
51. Malfertheiner P, Megraud F, O'Morain CA, Atherton J, Axon AT, Bazzoli F, et al. Management of *Helicobacter pylori* infection—the Maastricht IV/Florence Consensus Report. *Gut* 2012;61:646-64.
52. Amberbir A, Medhin G, Erku W, Alem A, Simms R, Robinson K, et al. Effects of *Helicobacter pylori*, geohelminth infection and selected commensal bacteria on the risk of allergic disease and sensitization in 3-year-old Ethiopian children. *Clin Exp Allergy* 2011;41:1422-30.
53. Chen Y, Blaser MJ. *Helicobacter pylori* colonization is inversely associated with childhood asthma. *J Infect Dis* 2008;198:553-60.
54. Arnold IC, Dehzad N, Reuter S, Martin H, Becher B, Taube C, et al. *Helicobacter pylori* infection prevents allergic asthma in mouse models through the induction of regulatory T cells. *J Clin Invest* 2011;121:3088-93.
55. Oertli M, Sundquist M, Hitzler I, Engler DB, Arnold IC, Reuter S, et al. DC-derived IL-18 drives Treg differentiation, murine *Helicobacter pylori*-specific immune tolerance, and asthma protection. *J Clin Invest* 2012;122:1082-96.
56. Oertli M, Noben M, Engler DB, Semper RP, Reuter S, Maxeiner J, et al. *Helicobacter pylori* gamma-glutamyl transpeptidase and vacuolating cytotoxin promote gastric persistence and immune tolerance. *Proc Natl Acad Sci U S A* 2013;110:3047-52.
57. Koch KN, Hartung ML, Urban S, Kyburz A, Bahlmann AS, Lind J, et al. *Helicobacter urease*-induced activation of the TLR2/NLRP3/IL-18 axis protects against asthma. *J Clin Invest* 2015;125:3297-302.
58. Pfefferle PI, Prescott SL, Kopp M. Microbial influence on tolerance and opportunities for intervention with prebiotics/probiotics and bacterial lysates. *J Allergy Clin Immunol* 2013;131:1453-63.
59. Kalliomaki M, Salminen S, Arvilommi H, Kero P, Koskinen P, Isolauri E. Probiotics in primary prevention of atopic disease: a randomised placebo-controlled trial. *Lancet* 2001;357:1076-9.
60. Kopp MV, Hennemuth I, Heinzmann A, Urbanek R. Randomized, double-blind, placebo-controlled trial of probiotics for primary prevention: no clinical effects of *Lactobacillus GG* supplementation. *Pediatrics* 2008;121:e850-6.
61. Ritter M, Gross O, Kays S, Ruland J, Nimmerjahn F, Saijo S, et al. *Schistosoma mansoni* triggers Dectin-2, which activates the Nlrp3 inflammasome and alters adaptive immune responses. *Proc Natl Acad Sci U S A* 2010;107:20459-64.
62. Maizels RM, Yazdanbakhsh M. T-cell regulation in helminth parasite infections: implications for inflammatory diseases. *Chem Immunol Allergy* 2008;94:112-23.
63. Yazdanbakhsh M, Kremsner PG, van Ree R. Allergy, parasites, and the hygiene hypothesis. *Science* 2002;296:490-4.
64. Hamid F, Amoah AS, van RR, Yazdanbakhsh M. Helminth-induced IgE and protection against allergic disorders. *Curr Top Microbiol Immunol* 2015;388:91-108.
65. McSorley HJ, Maizels RM. Helminth infections and host immune regulation. *Clin Microbiol Rev* 2012;25:585-608.
66. Amu S, Saunders SP, Kronenberg M, Mangan NE, Atzberger A, Fallon PG. Regulatory B cells prevent and reverse allergic airway inflammation via FoxP3-positive T regulatory cells in a murine model. *J Allergy Clin Immunol* 2010;125:1114-24.
67. van der Vlugt LE, Labuda LA, Ozir-Fazalalikhani A, Lievers E, Gloudemans AK, Liu KY, et al. Schistosomes induce regulatory features in human and mouse CD1dhi B cells: inhibition of inflammation by IL-10 and enhanced regulatory T cells. *PLoS One* 2012;7:e30883.
68. Layland LE, Straubinger K, Ritter M, Loffredo-Verde E, Garn H, Sparwasser T, et al. *Schistosoma mansoni*-mediated suppression of allergic airway inflammation requires patency and Foxp3+ Treg cells. *PLoS Negl Trop Dis* 2013;7:e2379.
69. van der Vlugt LE, Zinsou JF, Ozir-Fazalalikhani A, Kremsner PG, Yazdanbakhsh M, Adegnik AA, et al. Interleukin 10 (IL-10)-producing CD1dhi regulatory B cells from *Schistosoma haematobium*-infected individuals induce IL-10-positive T cells and suppress effector T-cell cytokines. *J Infect Dis* 2014;210:1207-16.
70. Wilson MS, Taylor MD, Balic A, Finney CA, Lamb JR, Maizels RM. Suppression of allergic airway inflammation by helminth-induced regulatory T cells. *J Exp Med* 2005;202:1199-212.
71. McSorley HJ, O'Gorman MT, Blair N, Sutherland TE, Filbey KJ, Maizels RM. Suppression of type 2 immunity and allergic airway inflammation by secreted products of the helminth *Heligmosomoides polygyrus*. *Eur J Immunol* 2012;42:2667-82.
72. McSorley HJ, Blair NF, Smith KA, McKenzie AN, Maizels RM. Blockade of IL-33 release and suppression of type 2 innate lymphoid cell responses by helminth secreted products in airway allergy. *Mucosal Immunol* 2014;7:1068-78.
73. Mpairwe H, Ndibazza J, Webb EL, Nampijja M, Muhangi L, Apule B, et al. Maternal hookworm modifies risk factors for childhood eczema: results from a birth cohort in Uganda. *Pediatr Allergy Immunol* 2014;25:481-8.
74. Straubinger K, Paul S, Prazeres da Costa CU, Ritter M, Buch T, Busch DH, et al. Maternal immune response to helminth infection during pregnancy determines offspring susceptibility to allergic airway inflammation. *J Allergy Clin Immunol* 2014;134:1271-9.
75. Smits HH, Akdis CA. In utero priming by worms protects against respiratory allergies. *J Allergy Clin Immunol* 2014;134:1280-1.
76. Lukkarinen M, Lukkarinen H, Lehtinen P, Vuorinen T, Ruuskanen O, Jartti T. Prednisolone reduces recurrent wheezing after first rhinovirus wheeze: a 7-year follow-up. *Pediatr Allergy Immunol* 2013;24:237-43.
77. Jartti T, Nieminen R, Vuorinen T, Lehtinen P, Vahlberg T, Gern J, et al. Short- and long-term efficacy of prednisolone for first acute rhinovirus-induced wheezing episode. *J Allergy Clin Immunol* 2015;135:691-8.
78. Busse WW, Morgan WJ, Gergen PJ, Mitchell HE, Gern JE, Liu AH, et al. Randomized trial of omalizumab (anti-IgE) for asthma in inner-city children. *N Engl J Med* 2011;364:1005-15.
79. Blanken MO, Rovers MM, Molenaar JM, Winkler-Seinstra PL, Meijer A, Kimpfen JL, et al. Respiratory syncytial virus and recurrent wheeze in healthy pre-term infants. *N Engl J Med* 2013;368:1791-9.

80. Navarro S, Cossalter G, Chiavaroli C, Kanda A, Fleury S, Lazzari A, et al. The oral administration of bacterial extracts prevents asthma via the recruitment of regulatory T cells to the airways. *Mucosal Immunol* 2011;4:53-65.
81. Schaad UB. OM-85 BV, an immunostimulant in pediatric recurrent respiratory tract infections: a systematic review. *World J Pediatr* 2010;6:5-12.
82. Bourke CD, Mutapi F, Nausch N, Photiou DM, Poulsen LK, Kristensen B, et al. *Trichuris suis* ova therapy for allergic rhinitis does not affect allergen-specific cytokine responses despite a parasite-specific cytokine response. *Clin Exp Allergy* 2012;42:1582-95.
83. Feary JR, Venn AJ, Mortimer K, Brown AP, Hooi D, Falcone FH, et al. Experimental hookworm infection: a randomized placebo-controlled trial in asthma. *Clin Exp Allergy* 2010;40:299-306.
84. Fleming JO, Weinstock JV. Clinical trials of helminth therapy in autoimmune diseases: rationale and findings. *Parasite Immunol* 2015;37:277-92.
85. Lau S, Gerhold K, Zimmermann K, Ockeloen CW, Rossberg S, Wagner P, et al. Oral application of bacterial lysate in infancy decreases the risk of atopic dermatitis in children with 1 atopic parent in a randomized, placebo-controlled trial. *J Allergy Clin Immunol* 2012;129:1040-7.
86. Beeh KM, Kannies F, Wagner F, Schilder C, Naudts I, Hammann-Haenni A, et al. The novel TLR-9 agonist QbG10 shows clinical efficacy in persistent allergic asthma. *J Allergy Clin Immunol* 2013;131:866-74.
87. Casale TB, Cole J, Beck E, Vogelmeier CF, Willers J, Lassen C, et al. CYT003, a TLR9 agonist, in persistent allergic asthma—a randomized placebo-controlled Phase 2b study. *Allergy* 2015;70:1160-8.