

# Astro2020 Science White Paper

## Cosmological Probes of Dark Matter Interactions: The Next Decade

**Thematic area:** Cosmology and Fundamental Physics.

**Principal Author:**

Vera Gluscevic

Institution: University of Florida

Email: [v.gluscevic@ufl.edu](mailto:v.gluscevic@ufl.edu)

Phone: 1-352-392-8754

**Co-Authors:**

Yacine Ali-Haïmoud<sup>1</sup>, Kimberly K. Boddy<sup>2</sup>, Céline Boehm<sup>3,4</sup>, Jens Chluba<sup>5</sup>, Francis-Yan Cyr-Racine<sup>6,7</sup>, Cora Dvorkin<sup>7</sup>, Vera Gluscevic<sup>8,9</sup>, Daniel Grin<sup>10</sup>, Julien Lesgourgues<sup>11</sup>, Julian B. Muñoz<sup>7</sup>, Ethan O. Nadler<sup>12</sup>, Vivian Poulin<sup>13</sup>, Sarah Shandera<sup>14</sup>, Katelin Schutz<sup>15</sup>, Tracy R. Slatyer<sup>16</sup>

<sup>1</sup> Department of Physics, New York University

<sup>2</sup> Department of Physics and Astronomy, Johns Hopkins University

<sup>3</sup> School of Physics, The University of Sydney, Australia

<sup>4</sup> LAPTH, France; Perimeter institute, Canada

<sup>5</sup> Jodrell Bank Center for Astrophysics, UK

<sup>6</sup> Department of Physics and Astronomy, University of New Mexico

<sup>7</sup> Department of Physics, Harvard University

<sup>8</sup> Department of Physics, University of Florida

<sup>9</sup> Department of Physics, Princeton University

<sup>10</sup> Department of Physics and Astronomy, Haverford College

<sup>11</sup> RWTH Aachen University, Germany

<sup>12</sup> KIPAC and Department of Physics, Stanford University

<sup>13</sup> France

<sup>14</sup> Department of Physics, Penn State University

<sup>15</sup> Center for Theoretical Physics, Massachusetts Institute of Technology

<sup>16</sup> Department of Physics, University of California Berkeley

**Endorsers:**

**Abstract**

Cosmological observations offer unique and robust avenues for probing the fundamental nature of dark matter particles—they broadly test a range of compelling theoretical scenarios, often surpassing or complementing the reach of terrestrial and other experiments. We discuss observational and theoretical advancements that will play a pivotal role in realizing a strong program of cosmological searches for the identity of dark matter in the coming decade. Specifically, we focus on measurements of the cosmic-microwave-background anisotropy and spectral distortions, and tracers of structure (such as the Lyman- $\alpha$  forest, galaxies, and cosmological 21-cm signal).

# 1 Key Question: What is Dark Matter?

Observations of the Universe, from our Galactic neighborhood, to the cosmological horizon, consistently testify that  $\sim 85\%$  of matter behaves as a cold non-collisional fluid that sources gravitational potentials and underpins structure on virtually all observable scales. **Over the past decades, it was confidently established that the main constituents of the dark matter (DM) component *cannot* be any known baryonic particles.** The existence of DM thus implies new physics whose investigation centrally drives research at the intersection of modern astrophysics, cosmology, and particle physics.

Following its discovery, a versatile range of laboratory experiments was built worldwide to directly detect and/or produce some of the best-motivated particle candidates that could account for cosmological DM: WIMPs, WIMP-like particles, axions, etc; so far, this effort is without success. Astrophysical and cosmological observations provide the only evidence for the existence of DM and source a large portion of what is known about its properties: its stability on cosmological time scales, its apparent non-collisional nature, and its central role in the formation and growth of structure. Recently, those observations have also emerged as a powerful probe of DM microphysics, complementary in reach to laboratory experiments.

We focus on cosmological searches, whose key goal is to detect the effects of DM interactions on the structure and thermal history of the Universe, and use them to pin down particle identity of DM. **Cosmology is a versatile tool that can test broad classes of theoretical scenarios:** DM interactions with known particles (baryons, photons, and neutrinos), interactions within the DM sector itself, and DM annihilations and decays.<sup>1</sup> **The next decade of observations will see a tremendous leap in sensitivity to each of them,** in some cases, by many orders of magnitude in DM mass and interaction cross sections. Multiple observables—cosmic microwave background (CMB) anisotropy and spectral distortions, tracers of large-scale structure, and objects in our Galactic neighborhood—can all probe the same underlying microphysics throughout cosmic history, on a broad range of physical scales (Fig. 1). **Establishing DM signal discovery and robustly testing its fundamental nature will rely on confirmation and consistency checks between different data sets.** This will necessitate progress in understanding synergies between observables, and development of frameworks that enable their joint analyses to probe the same underlying DM microphysics. In addition, the DM search program with next-generation cos-




Figure 1: Approximate cosmological epochs and physical scales corresponding to different observables (spectral distortions are omitted for compactness of presentation; their origin lies to the left of the plot).

<sup>1</sup>We note that other science white papers focus on scenarios in which DM consists of ultra-light axion-like particles, warm DM, primordial black holes, etc.

mological data will rely on accurate theoretical treatment of the formation and evolution of structure in non-standard cosmologies, and modeling of baryonic and other non-linear effects that present systematic uncertainties for DM parameter inference [1–3]. We summarize the future promise of cosmological measurements in search for the identity of DM, and conclude with a list of advancements that will be pivotal to its success in the coming decade.

## 2 Theory and Observations

### 2.1 Scattering with Baryons

In the standard model of cosmology DM is non-collisional, but elastic scattering between DM and visible particles is commonplace in some of the best-motivated DM models, including the WIMPs and WIMP-like particles. For this reason, terrestrial DM searches with direct detection experiments are looking for scattering of Galactic-halo DM on nuclei in underground targets. However, the same scattering processes can occur in a cosmological setting and lead to an exchange of heat and momentum between DM and baryons (i. e. the visible particles), absent in the standard (non-collisional) cosmology [4–13]. This can affect both the thermal history and the evolution of cosmological perturbations, captured by various observables.

**CMB spectral distortions.** DM can drain heat from the primordial plasma, by scattering with either protons, electrons, or photons in the early Universe [14]. Heat exchange occurring later than two months after the Big Bang cannot fully thermalize and thus leads to distortions of the CMB frequency spectrum away from a perfect blackbody [15, 16]. The CMB frequency spectrum is most sensitive to light particles that thermally decouple as early as  $z \sim 2 \times 10^6$ . The existing COBE FIRAS measurements provide upper limits on DM-baryon and DM-photon interactions for DM masses lower than  $\sim 0.1$  MeV. These limits are similarly stringent but independent of those derived from CMB anisotropy from *Planck*. Future measurements that can detect a fractional distortion of order  $\sim 10^{-8}$  would be sensitive to interacting DM particles as massive as  $\sim 1$  GeV [14]. Spectral distortions in conjunction with power suppression in CMB anisotropy in future data could yield robust evidence for DM physics taking place in the very early Universe.

**CMB anisotropy.** DM-baryon scattering prior to recombination ( $z \sim 1100$ ) induces a drag force and momentum transfer between the DM and baryon fluids. This results in power suppression in density fluctuations, which is more prominent for progressively smaller scales and stronger interactions. As perturbations grow, the lack of small-scale structure translates into an underabundance of luminous DM tracers throughout cosmic history (Fig. 1).

*Planck* measurements of the CMB temperature anisotropy currently provide the most pristine cosmological bound on the DM-proton scattering cross section [4–9]. They already sensitively probe interactions of DM particles with masses outside the detection limits of most existing direct detection experiments ( $\gtrsim 1$  keV) [5], taking place as far back in time as when the Universe was a thousand years old. However, the first high-signal-to-noise measurements of the CMB polarization and lensing on small ( $\sim$ arcminute) angular scales—to be delivered by the next-generation CMB experiments—can enable a leap in sensitivity to DM scattering cross sections by up to *several orders of magnitude*, as compared to *Planck* [17–20]. Interestingly, measurement of DM-baryon scattering with CMB anisotropy is insensitive to uncertainties on cosmological parameters, and the relevant signals are distinct from those

of other new physics targeted by future CMB experiments (e.g. the neutrino mass). However, in spite of the robustness of these measurements, the CMB has a limited grasp of small scales, where the effects of the interactions are more prominent; small scales are accessible to other observables, but are more prone to systematic modeling and measurement uncertainties.

**Lyman- $\alpha$  forest.** For example, SDSS Lyman- $\alpha$  forest measurements trace the matter power spectrum on comoving scale of about 1 Mpc; their analysis set the most stringent to-date cosmological bound on DM-baryon interactions, surpassing the CMB [7, 21]. This bound implies that a baryon within a Milky-Way-like galaxy does *not* scatter with DM over the age of the galaxy. Future spectroscopic surveys could improve upon these limits by reconstructing even smaller scales with similar measurements. However, inference of fundamental physics from these data relies on accurate modeling of the Lyman- $\alpha$  forest and non-linear evolution of small scales [22]. Consistency checks and future joint analyses with high-precision CMB measurements and other probes could alleviate such uncertainties.

**21-cm cosmology.** Over the coming decade, measurements of the hyperfine 21-cm line transition in atomic hydrogen during cosmic Dark Ages and the Epoch of Reionization will start to probe redshifts  $10 \lesssim z \lesssim 200$ , far beyond galaxy surveys. The strength of the 21-cm signal is proportional to the difference between the temperature of the gas (i. e. the baryons) and that of the CMB, and acts as a calorimeter that can measure the thermal history at these redshifts. If DM-baryon scattering occurs *after* recombination is complete, it can alter temperature of baryons at that time. The sky-averaged signal and its fluctuations are both sensitive to this effect [23–27]. Such a late-time scattering scenario arises, for example, in a simple and well-motivated model whereby DM particles carry a small electric charge and exhibit Coulomb-like interactions (“millicharged” DM [28, 29]). This scenario is challenging to test in direct detection experiments [30]; cosmological probes like the 21-cm signal may thus be the optimal detection channel. However, 21-cm cosmology is both rapidly-advancing and exceptionally challenging, since it necessitates removal of an overwhelming Galactic foreground signal. As was demonstrated with the recent analyses of the EDGES result [31], cross-checks with other probes will likely be essential for interpretation of future anomalies in the 21-cm signal as new physics [32–34].

**Galaxies.** Galaxies trace structure on even smaller scales, and thus hold a bold promise as a probe of DM physics [1]. Given the physical scales involved, studies of the abundances of dwarf and satellite galaxies, in particular, could unlock orders of magnitude improvement in sensitivity to DM interactions, compared to the high-redshift and large-scale probes discussed earlier. However, robust inference of fundamental physics from studies of galaxies will crucially depend on improvements in modeling and numerical simulation of their formation and evolution within cosmologies with interacting DM, and understanding of baryonic effects—a challenge that will need much attention in the coming decade [1].<sup>2</sup>

## 2.2 Scattering with Neutrinos

Neutrinos reside in one of the least well understood parts of the Standard Model of particle physics; indeed, non-vanishing neutrino masses require new physics to explain. In addition, terrestrial neutrino experiments and probes of astrophysical neutrinos have seen a number of

---

<sup>2</sup>Observables such as galaxy clustering and weak lensing are also relevant for this discussion and could potentially test the same physics; however, they are a prime focus of other science white papers.

anomalies that may indicate new physics [35–40]. The idea that DM may communicate with visible matter through interactions with neutrinos is an intriguing possibility, but laboratory tests of this scenario are currently unfeasible. Meanwhile, the high abundance of cosmological neutrinos allows for observational tests of a possible link between neutrinos DM.

If neutrinos efficiently scatter with DM seconds after the Big Bang, the DM-neutrino fluid undergoes acoustic oscillations and diffusion damping, and is smoother than ordinary free-streaming neutrinos, with a smaller sound speed. CMB anisotropy captures this physics through a suppression of power and a shift of the acoustic peaks towards smaller angular scales, all used to probe DM-neutrino scattering with current CMB data [41–44]. The small-scale power in matter fluctuations is also suppressed in this scenario, and can, in principle, probe even weaker interactions than the CMB; the best current bounds on DM-neutrino scattering cross section come from Lyman- $\alpha$  forest observations [45]. As for DM-baryon scattering, future CMB, Lyman- $\alpha$ , and galaxy surveys will increase sensitivity to DM-neutrino scattering by mapping out perturbations and structure on even smaller scales.

## 2.3 Annihilation and Decay

If DM interacts with visible particles, it may also annihilate or decay, further altering the thermal history of the Universe. If annihilation or decay products include electromagnetically interacting particles, these particles can generically heat and/or ionize the baryonic gas. Cosmological searches have distinct advantages over classic indirect searches for DM annihilation and decay, as they do not suffer from astrophysical backgrounds and large uncertainties in the distribution of DM in the target systems. Furthermore, they can probe processes with no detectable signals in terrestrial-scale experiments and in the local Universe; for example, the decays with lifetimes comparable to the age of the Universe, and decays into invisible channels, such as neutrinos or new dark particles [46, 47].

Increasing the ionization fraction near the time of recombination can affect the CMB anisotropy [48–50]. *Planck* measurements of CMB temperature and polarization anisotropy on degree angular scales provide some of the strongest and most robust bounds on annihilations and decays of sub-GeV DM, complementing indirect searches that probe heavier DM candidates [51, 52]. Measurements from the next-generation ground-based CMB experiments can improve sensitivity to DM annihilation cross section and lifetime by a factor of several. Furthermore, as discussed in context of DM-baryon interactions, future observations of the cosmological 21-cm signal from atomic hydrogen can sensitively track gas temperature during cosmic Dark Ages and Reionization. As such, they can capture any new energy injection from DM decays and annihilations in the post-recombination Universe, likely surpassing the sensitivity of the CMB anisotropy to the same processes [47, 53–56].

## 2.4 Interactions with Dark Radiation

Models in which the dark sector is complex and contains not only dark matter, but also dark radiation (DR) that interacts with DM particles, are motivated in several ways: they generically arise in theories proposed to explain the hierarchy problem of the Standard Model of particles [57, 58]; they could explain the anomalously low large-scale amplitude of matter fluctuations seen in some weak-lensing surveys [59–62]; and they present a specific case of self-interacting DM, proposed as a possible solution for putative anomalies in DM structure

on sub-galactic scales [63–66]. Cosmology offers a unique way to probe DM-DR interactions, especially if DM resides in a secluded sector that only weakly couples to known particles.

Similar to how photon pressure prohibits the growth of baryon fluctuations until recombination, DM interacting with DR in the early Universe experiences suppressed growth of structure, as compared to a scenario with no DR. The resulting suppression of power is captured in the CMB anisotropy and all tracers of large-scale structure (similar to the case of DM-baryon interactions): galaxy clustering, galaxy weak lensing, the Lyman- $\alpha$  forest, and cosmological 21-cm signal [67–69]. Notably, cosmological signatures of DR are distinct from signatures of relativistic particles that do not couple to DM, such as the standard free-streaming neutrinos [70–72]. High-precision measurements of small-angular-scale CMB polarization anisotropy with the next-generation ground-based experiments could robustly detect or rule out DM-DR interactions taking place in the early Universe, even in the case where only a small fraction (less than 5%) of the overall DM density couples to DR. As in the previously discussed theoretical scenarios, galaxies, the Lyman- $\alpha$  forest, and 21-cm observations could probe even smaller scales and earlier times.

### 3 Recommendations

The next decade of cosmological observations will open avenues to broadly probe the fundamental nature of DM particles in context of compelling theoretical scenarios, complementing the reach of terrestrial experiments. Many individual observations (e. g. the next-generation CMB measurements, galaxy surveys, 21-cm line intensity mapping) should reach sufficient raw sensitivity to potentially uncover evidence of DM interactions that are hidden from the present-day observations. However, a linchpin to their robustness as probes of fundamental physics will be in their synergies; cross-validation and consistency across the observables is necessary to establish a discovery of new DM physics in cosmological data.

The following advancements will all play central roles in cosmological searches for the identity of DM in the coming decade:

- **Observations:**

- Next-generation measurements of high-multipole CMB anisotropy.
- Measurements of thermal history (global 21-cm signal, CMB spectral distortions).
- Surveys of large-scale structure (21-cm tomography, Lyman- $\alpha$  forest, galaxy clustering and weak lensing).
- Near-field cosmology (local probes of small-scale structure).

- **Theory and analysis:**

- High-accuracy theoretical prediction of cosmological DM signals.
- Simulation and modeling of non-linearities in novel DM cosmologies.
- Understanding of baryonic systematic uncertainties in DM-related inference.
- Frameworks for joint analyses of cosmological probes.

## References

- [1] A. Drlica-Wagner, Y.-Y. Mao, and et al., “Probing the Fundamental Nature of Dark Matter with the Large Synoptic Survey Telescope,” *arXiv e-prints*, p. arXiv:1902.01055, Feb 2019.
- [2] R. Murgia, V. Iršič, and M. Viel, “Novel constraints on noncold, nonthermal dark matter from Lyman- $\alpha$  forest data,” *Phys. Rev. D*, vol. 98, p. 083540, Oct 2018.
- [3] Y. Ali-Haïmoud, “Boltzmann-Fokker-Planck formalism for dark matter-baryon scattering,” *Phys. Rev. D*, vol. 99, p. 023523, Jan. 2019.
- [4] K. K. Boddy and V. Gluscevic, “First Cosmological Constraint on the Effective Theory of Dark Matter-Proton Interactions,” *Phys. Rev.*, vol. D98, no. 8, p. 083510, 2018.
- [5] V. Gluscevic and K. K. Boddy, “Constraints on Scattering of keV–TeV Dark Matter with Protons in the Early Universe,” *Phys. Rev. Lett.*, vol. 121, no. 8, p. 081301, 2018.
- [6] K. K. Boddy, V. Gluscevic, V. Poulin, E. D. Kovetz, M. Kamionkowski, and R. Barkana, “A Critical Assessment of CMB Limits on Dark Matter-Baryon Scattering: New Treatment of the Relative Bulk Velocity,” 2018.
- [7] W. L. Xu, C. Dvorkin, and A. Chael, “Probing sub-GeV Dark Matter-Baryon Scattering with Cosmological Observables,” *Phys. Rev.*, vol. D97, no. 10, p. 103530, 2018.
- [8] T. R. Slatyer and C.-L. Wu, “Early-Universe constraints on dark matter-baryon scattering and their implications for a global 21 cm signal,” *Phys. Rev.*, vol. D98, no. 2, p. 023013, 2018.
- [9] C. Dvorkin, K. Blum, and M. Kamionkowski, “Constraining Dark Matter-Baryon Scattering with Linear Cosmology,” *Phys. Rev.*, vol. D89, no. 2, p. 023519, 2014.
- [10] C. Boehm, P. Fayet, and R. Schaeffer, “Constraining dark matter candidates from structure formation,” *Physics Letters B*, vol. 518, pp. 8–14, Oct 2001.
- [11] C. Boehm and R. Schaeffer, “Constraints on Dark Matter interactions from structure formation: damping lengths,” *Astron. Astrophys.*, vol. 438, pp. 419–442, Aug 2005.
- [12] C. Boehm and P. Fayet, “Scalar dark matter candidates,” *Nuclear Physics B*, vol. 683, pp. 219–263, Apr 2004.
- [13] K. Sigurdson, M. Doran, A. Kurylov, R. R. Caldwell, and M. Kamionkowski, “Dark-matter electric and magnetic dipole moments,” vol. 70, p. 083501, Oct. 2004. [Erratum: *Phys. Rev. D* 73, 089903 (2006)].
- [14] Y. Ali-Haïmoud, J. Chluba, and M. Kamionkowski, “Constraints on Dark Matter Interactions with Standard Model Particles from Cosmic Microwave Background Spectral Distortions,” *Physical Review Letters*, vol. 115, p. 071304, Aug. 2015.



- [15] W. Hu and J. Silk, “Thermalization and spectral distortions of the cosmic background radiation,” *Phys. Rev. D*, vol. 48, pp. 485–502, July 1993.
- [16] J. Chluba and R. A. Sunyaev, “The evolution of CMB spectral distortions in the early Universe,” *Mon. Not. R. Astron. Soc.*, vol. 419, pp. 1294–1314, Jan. 2012.
- [17] Z. Li, V. Gluscevic, K. K. Boddy, and M. S. Madhavacheril, “Disentangling dark physics with cosmic microwave background experiments,” *Phys. Rev. D*, vol. 98, p. 123524, Dec 2018.
- [18] The Simons Observatory collaboration, “The Simons Observatory: science goals and forecasts,” *Journal of Cosmology and Astro-Particle Physics*, vol. 2019, p. 056, Feb 2019.
- [19] K. N. Abazajian *et al.*, “CMB-S4 Science Book, First Edition,” 2016.
- [20] S. Hanany and et al., “PICO: Probe of Inflation and Cosmic Origins,” *arXiv e-prints*, p. arXiv:1902.10541, Feb 2019.
- [21] C. Dvorkin, K. Blum, and M. Zaldarriaga, “Perturbed Recombination from Dark Matter Annihilation,” *Phys. Rev.*, vol. D87, no. 10, p. 103522, 2013.
- [22] V. Iršič, M. Viel, M. G. Haehnelt, J. S. Bolton, S. Cristiani, G. D. Becker, V. D’Odorico, G. Cupani, T.-S. Kim, T. A. M. Berg, S. López, S. Ellison, L. Christensen, K. D. Denney, and G. Worseck, “New constraints on the free-streaming of warm dark matter from intermediate and small scale Lyman- $\alpha$  forest data,” *Phys. Rev. D*, vol. 96, p. 023522, Jul 2017.
- [23] J. B. Muñoz, E. D. Kovetz, and Y. Ali-Haïmoud, “Heating of baryons due to scattering with dark matter during the dark ages,” *Phys. Rev. D*, vol. 92, p. 083528, Oct. 2015.
- [24] R. Barkana, “Possible interaction between baryons and dark-matter particles revealed by the first stars,” *Nature (London)*, vol. 555, pp. 71–74, Mar. 2018.
- [25] A. Fialkov, R. Barkana, and A. Cohen, “Constraining Baryon-Dark-Matter Scattering with the Cosmic Dawn 21-cm Signal,” *Physical Review Letters*, vol. 121, p. 011101, July 2018.
- [26] J. B. Muñoz, C. Dvorkin, and A. Loeb, “21-cm Fluctuations from Charged Dark Matter,” *Physical Review Letters*, vol. 121, p. 121301, Sept. 2018.
- [27] J. B. Muñoz and A. Loeb, “Insights on Dark Matter from Hydrogen during Cosmic Dawn,” *arXiv e-prints*, p. arXiv:1802.10094, Feb 2018.
- [28] M. Battaglieri and et al., “US Cosmic Visions: New Ideas in Dark Matter 2017: Community Report,” *arXiv e-prints*, p. arXiv:1707.04591, Jul 2017.
- [29] C. Dvorkin, T. Lin, and K. Schutz, “Making dark matter out of light: freeze-in from plasma effects,” *arXiv e-prints*, p. arXiv:1902.08623, Feb 2019.



- [30] S. D. McDermott, H.-B. Yu, and K. M. Zurek, “Turning off the lights: How dark is dark matter?,” *Phys. Rev. D*, vol. 83, p. 063509, Mar. 2011.
- [31] J. D. Bowman, A. E. E. Rogers, R. A. Monsalve, T. J. Mozdzen, and N. Mahesh, “An absorption profile centred at 78 megahertz in the sky-averaged spectrum,” *Nature (London)*, vol. 555, pp. 67–70, Mar. 2018.
- [32] R. Barkana, “Possible interaction between baryons and dark-matter particles revealed by the first stars,” vol. 555, no. 7694, pp. 71–74, 2018.
- [33] E. D. Kovetz, V. Poulin, V. Gluscevic, K. K. Boddy, R. Barkana, and M. Kamionkowski, “Tighter Limits on Dark Matter Explanations of the Anomalous EDGES 21cm Signal,” 2018.
- [34] A. Berlin, D. Hooper, G. Krnjaic, and S. D. McDermott, “Severely Constraining Dark Matter Interpretations of the 21-cm Anomaly,” 2018.
- [35] C. Athanassopoulos *et al.*, “Evidence for anti-muon-neutrino  $\rightarrow$  anti-electron-neutrino oscillations from the LSND experiment at LAMPF,” *Phys. Rev. Lett.*, vol. 77, pp. 3082–3085, 1996.
- [36] A. Aguilar-Arevalo *et al.*, “Evidence for neutrino oscillations from the observation of anti-neutrino(electron) appearance in a anti-neutrino(muon) beam,” *Phys. Rev.*, vol. D64, p. 112007, 2001.
- [37] A. A. Aguilar-Arevalo *et al.*, “A Search for electron neutrino appearance at the  $\Delta m^2 \sim 1\text{eV}^2$  scale,” *Phys. Rev. Lett.*, vol. 98, p. 231801, 2007.
- [38] MiniBooNE Collaboration, “Significant Excess of ElectronLike Events in the Mini-BooNE Short-Baseline Neutrino Experiment,” *arXiv e-prints*, p. arXiv:1805.12028, May 2018.
- [39] P. W. Gorham *et al.*, “Characteristics of Four Upward-pointing Cosmic-ray-like Events Observed with ANITA,” *Phys. Rev. Lett.*, vol. 117, no. 7, p. 071101, 2016.
- [40] P. W. Gorham *et al.*, “Observation of an Unusual Upward-going Cosmic-ray-like Event in the Third Flight of ANITA,” *Phys. Rev. Lett.*, vol. 121, no. 16, p. 161102, 2018.
- [41] G. Mangano, A. Melchiorri, P. Serra, A. Cooray, and M. Kamionkowski, “Cosmological bounds on dark matter-neutrino interactions,” *Phys. Rev.*, vol. D74, p. 043517, 2006.
- [42] M. Escudero, O. Mena, A. C. Vincent, R. J. Wilkinson, and C. Boehm, “Exploring dark matter microphysics with galaxy surveys,” *JCAP*, vol. 1509, no. 09, p. 034, 2015.
- [43] E. Di Valentino, C. Boehm, E. Hivon, and F. R. Bouchet, “Reducing the  $H_0$  and  $\sigma_8$  tensions with Dark Matter-neutrino interactions,” *Phys. Rev.*, vol. D97, no. 4, p. 043513, 2018.
- [44] J. A. D. Diacoumis and Y. Y. Wong, “Prior dependence of cosmological constraints on dark matter-radiation interactions,” 2018.

- [45] R. J. Wilkinson, C. Boehm, and J. Lesgourgues, “Constraining Dark Matter-Neutrino Interactions using the CMB and Large-Scale Structure,” *JCAP*, vol. 1405, p. 011, 2014.
- [46] V. Poulin, P. D. Serpico, and J. Lesgourgues, “A fresh look at linear cosmological constraints on a decaying dark matter component,” *JCAP*, vol. 1608, no. 08, p. 036, 2016.
- [47] V. Poulin, J. Lesgourgues, and P. D. Serpico, “Cosmological constraints on exotic injection of electromagnetic energy,” *JCAP*, vol. 1703, no. 03, p. 043, 2017.
- [48] J. A. Adams, S. Sarkar, and D. W. Sciama, “CMB anisotropy in the decaying neutrino cosmology,” *Mon. Not. Roy. Astron. Soc.*, vol. 301, pp. 210–214, 1998.
- [49] X.-L. Chen and M. Kamionkowski, “Particle decays during the cosmic dark ages,” *Phys. Rev.*, vol. D70, p. 43502, 2004.
- [50] N. Padmanabhan and D. P. Finkbeiner, “Detecting Dark Matter Annihilation with CMB Polarization: Signatures and Experimental Prospects,” *Phys. Rev.*, vol. D72, p. 23508, 2005.
- [51] N. Aghanim *et al.*, “Planck 2018 results. VI. Cosmological parameters,” 2018.
- [52] T. R. Slatyer and C.-L. Wu, “General Constraints on Dark Matter Decay from the Cosmic Microwave Background,” *Phys. Rev.*, vol. D95, no. 2, p. 023010, 2017.
- [53] S. R. Furlanetto, S. P. Oh, and E. Pierpaoli, “The Effects of Dark Matter Decay and Annihilation on the High-Redshift 21 cm Background,” *Phys. Rev.*, vol. D74, p. 103502, 2006.
- [54] M. Valdes, A. Ferrara, M. Mapelli, and E. Ripamonti, “Constraining DM through 21 cm observations,” *Mon. Not. Roy. Astron. Soc.*, vol. 377, pp. 245–252, 2007.
- [55] C. Evoli, A. Mesinger, and A. Ferrara, “Unveiling the nature of dark matter with high redshift 21 cm line experiments,” *JCAP*, vol. 1411, no. 11, p. 024, 2014.
- [56] L. Lopez-Honorez, O. Mena, A. Molin<sup>’</sup>e, S. Palomares-Ruiz, and A. C. Vincent, “The 21 cm signal and the interplay between dark matter annihilations and astrophysical processes,” *JCAP*, vol. 1608, no. 08, p. 004, 2016.
- [57] N. Arkani-Hamed, T. Cohen, R. T. D’Agnolo, A. Hook, H. D. Kim, and D. Pinner, “Solving the Hierarchy Problem at Reheating with a Large Number of Degrees of Freedom,” *Phys. Rev. Lett.*, vol. 117, no. 25, p. 251801, 2016.
- [58] Z. Chacko, D. Curtin, M. Geller, and Y. Tsai, “Cosmological Signatures of a Mirror Twin Higgs,” *JHEP*, vol. 09, p. 163, 2018.
- [59] J. Lesgourgues, G. Marques-Tavares, and M. Schmaltz, “Evidence for dark matter interactions in cosmological precision data?,” *JCAP*, vol. 1602, no. 02, p. 037, 2016.
- [60] Z. Chacko, Y. Cui, S. Hong, T. Okui, and Y. Tsai, “Partially Acoustic Dark Matter, Interacting Dark Radiation, and Large Scale Structure,” *JHEP*, vol. 12, p. 108, 2016.

- [61] M. A. Buen-Abad, M. Schmaltz, J. Lesgourgues, and T. Brinckmann, “Interacting Dark Sector and Precision Cosmology,” *JCAP*, vol. 1801, no. 01, p. 008, 2018.
- [62] R. Krall, F.-Y. Cyr-Racine, and C. Dvorkin, “Wandering in the Lyman-alpha Forest: A Study of Dark Matter-Dark Radiation Interactions,” *JCAP*, vol. 1709, no. 09, p. 003, 2017.
- [63] S. Tulin, H.-B. Yu, and K. M. Zurek, “Resonant Dark Forces and Small Scale Structure,” *Phys. Rev. Lett.*, vol. 110, p. 111301, 2013.
- [64] S. Tulin, H.-B. Yu, and K. M. Zurek, “Beyond Collisionless Dark Matter: Particle Physics Dynamics for Dark Matter Halo Structure,” *Phys. Rev. D*, vol. 87, p. 115007, 2013.
- [65] M. Kaplinghat, S. Tulin, and H.-B. Yu, “Dark Matter Halos as Particle Colliders: Unified Solution to Small-Scale Structure Puzzles from Dwarfs to Clusters,” *Phys. Rev. Lett.*, vol. 116, no. 4, p. 041302, 2016.
- [66] J. S. Bullock and M. Boylan-Kolchin, “Small-Scale Challenges to the  $\Lambda$ CDM Paradigm,” *Ann. Rev. Astron. Astrophys.*, vol. 55, pp. 343–387, 2017.
- [67] C. Boehm, A. Riazuelo, S. H. Hansen, and R. Schaeffer, “Interacting dark matter disguised as warm dark matter,” *Phys. Rev. D*, vol. 66, p. 083505, 2002.
- [68] F.-Y. Cyr-Racine, R. de Putter, A. Raccanelli, and K. Sigurdson, “Constraints on Large-Scale Dark Acoustic Oscillations from Cosmology,” *Phys. Rev.*, vol. D89, no. 6, p. 063517, 2014.
- [69] F.-Y. Cyr-Racine, K. Sigurdson, J. Zavala, T. Bringmann, M. Vogelsberger, and C. Pfrommer, “ETHOS – an effective theory of structure formation: From dark particle physics to the matter distribution of the Universe,” *Phys. Rev.*, vol. D93, p. 123527, June 2016.
- [70] S. Bashinsky and U. Seljak, “Neutrino perturbations in CMB anisotropy and matter clustering,” *Phys. Rev. D*, vol. 69, p. 083002, 2004.
- [71] B. Follin, L. Knox, M. Millea, and Z. Pan, “First Detection of the Acoustic Oscillation Phase Shift Expected from the Cosmic Neutrino Background,” *Phys. Rev. Lett.*, vol. 115, no. 9, p. 091301, 2015.
- [72] D. Baumann, D. Green, J. Meyers, and B. Wallisch, “Phases of New Physics in the CMB,” *JCAP*, vol. 1601, p. 007, 2016.