

# Mathematics Exam Solutions

## 1 Elementary algebra

**Problem 1.1.** Simplify

$$\frac{x^{32}}{x^9 \cdot x^2} \cdot \frac{x^7}{x^2} = \frac{x^{39}}{x^{13}} = x^{26}$$

**Problem 1.2.** Solve for  $x$ :

$$8^2 \cdot 4^x \cdot 2^x = 8^4$$

$$2^6 \cdot 2^{2x} \cdot 2^x = 2^{12}$$

$$2^{3x} = 2^6$$

$$3 \cdot x = 6$$

$$x = 2$$

**Problem 1.3.** Calculate the missing value. If  $\frac{x}{y}$  is 3, then  $x^{-4}y^4 = \dots$

$$x = 3 \cdot y$$

$$(3 \cdot y)^{-4}y^4 = 3^{-4} \cdot y^{-4} \cdot y^4 = \frac{1}{3^4} = \frac{1}{81} \approx 0.012$$

**Problem 1.4.** Calculate

$$\frac{\sqrt{4^{15}}}{\sqrt{16^7}} = \sqrt{\frac{4^{15}}{4^{14}}} = \sqrt{4} = 2$$

**Problem 1.5.** True or False ( $x$  and  $y$  and  $z$  are real numbers):

(a)  $x + (y + z) = (y + x) + z$     TRUE

(b)  $y(x + z) = xy + zy$     TRUE

(c)  $x^{y+z} = x^z + x^y$     FALSE

(d)  $\frac{x^z}{x^y} = x^{y-z}$     FALSE

**Problem 1.6.** Find the solution set for the inequality below:

$$\ln(x) \geq e$$

$$e^{\ln(x)} \geq e^e$$

$$x \geq e^e$$

## 2 Functions of one variable

**Problem 2.1 (Based on SYD 2.5.6).** The relationship between temperatures measured in Celsius and Fahrenheit is linear.  $0^{\circ}\text{C}$  is equivalent to  $32^{\circ}\text{F}$  and  $100^{\circ}\text{C}$  is the same as  $212^{\circ}\text{F}$ . Which temperature is measured by the same number on both scales?

Equations:

$$(a) \quad 100 = 212 \cdot a + b$$

$$(b) \quad 0 = 32 \cdot a + b$$

Thus  $32 \cdot a = -b$  and:

$$100 = 212 \cdot a - 32 \cdot a = 180 \cdot a$$

Thus  $b = -32 \cdot \frac{100}{180}$  and:

$$y = \frac{100}{180} \cdot x - 32 \cdot \frac{100}{180}$$

If  $y = x$ :

$$x - \frac{100}{180} \cdot x = -32 \cdot \frac{100}{180}$$

$$\frac{180}{100} \cdot x - x = -32$$

$$x = -40$$

**Problem 2.2.** Take the following function  $f(x) = 3x - 12$ . Find  $y$  if  $f(y) = 0$ .

$$f(y) = 3y - 12 = 0$$

$$3y = 12$$

$$y = 4$$

**Problem 2.3.** Find all values of  $x$  that satisfy:

$$9^{x^2-6x+2} = 81$$

$$9^{x^2-6x+2} = 9^2$$

$$x^2 - 6x + 2 = 2$$

$$x^2 - 6x = 0$$

$$x \cdot (x - 6) = 0$$

So  $x_1 = 0$ ,  $x_2 = 6$ .

**Problem 2.5.** Calculate the following value

$$\log_{\pi} \left( \frac{1}{\pi^5} \right)$$

$$\log_{\pi} (\pi^{-5}) = -5$$

### 3 Calculus

**Problem 3.1.** Calculate the following sum

$$\sum_{i=0}^{\infty} \left( \frac{1}{5^i} + 0.3^i \right) = \sum_{i=0}^{\infty} \left( \frac{1}{5^i} \right) + \sum_{i=0}^{\infty} 0.3^i$$

Applying the formula for infinite geometric series:

$$\sum_{i=0}^{\infty} \left( \frac{1}{5^i} \right) = \frac{5}{4}$$

$$\sum_{i=0}^{\infty} \left( \frac{3}{10} \right)^i = \frac{10}{7}$$

$$\sum_{i=0}^{\infty} \left( \frac{1}{5^i} + 0.3^i \right) = \frac{5}{4} + \frac{10}{7} \approx 2.67857$$

**Problem 3.2.** Find the following limit

$$\begin{aligned} \lim_{x \rightarrow 5} \frac{x^2 - 25}{x - 5} \\ \lim_{x \rightarrow 5} \frac{(x - 5)(x + 5)}{x - 5} \\ \lim_{x \rightarrow 5} (x + 5) = 10 \end{aligned}$$

**Problem 3.3.** Find the slope of the function  $f(x) = x^3 - 4$  at  $(-2, -12)$ .

$$f'(x) = 3x^2$$

$$f'(-2) = 12 = m$$

$$y = m \cdot x$$

$$y - y_1 = m \cdot (x - x_1)$$

$$y + 12 = 12 \cdot (x + 2)$$

$$y = 12x + 12$$

**Problem 3.4.** Find the derivative of the following function:

$$f(x) = \frac{x^5 + 3}{x^2 - 1}$$

$$f'(x) = \frac{5x^4 \cdot (x^2 - 1) - (x^5 + 3) \cdot 2x}{(x^2 - 1)^2}$$

**Problem 3.5.** Find the second derivative of the following function:

$$f(x) = x^9 + 3$$

$$f'(x) = 9x^8$$

$$f''(x) = 72x^7$$

**Problem 3.6.** Is the function  $f(x) = \frac{1}{x}$  continuous at 0? Why?

It is not because division by zero is undefined therefore as  $x$  approaches 0 from the right,  $y$  approaches infinity and as  $x$  approaches 0 from the left,  $y$  approaches negative infinity.

**Problem 3.7.** Consider the following function. Find all of its local minima, local maxima or inflection points.

$$\begin{aligned}f(x) &= 4x^3 - 12x \\f'(x) &= 12x^2 - 12 = 0 \\x^2 &= 1 \\x &= \pm 1\end{aligned}$$

Possible local minima/maxima at  $x = \pm 1$ .

$$f''(x) = 24x$$

For  $x = 1$ ,  $f''(x) = 24$  so it's a local minimum.

For  $x = -1$ ,  $f''(x) = -24$  so it's a local maximum.

The function also has an inflection point at  $x = 0$ .

**Problem 3.8.** Let  $f(x, y) = x^3 - y^2$ . Calculate  $f(2, 3)$

$$f(2, 3) = 8 - 9 = -1$$

**Problem 3.9.** Consider the following function:  $f(x, y) = \ln(x - 3y)$ . For what combinations of  $x$  and  $y$  is this function defined?

$$\begin{aligned}x - 3y &> 0 \\x &> 3y\end{aligned}$$

**Problem 3.10.** Find the following partial derivative:

$$\frac{\partial}{\partial x} \left( x^5 y^7 + \frac{x^2}{y^3} \right) = \left( 5x^4 y^7 + 2x \cdot \frac{1}{y^3} \right)$$

**Problem 3.12.** Solve the following constrained optimization problem using Lagrange's method:  $\max x^2 y^2$   
s.t.  $2x + y = 9$

$$\begin{aligned}L(x, y, \lambda) &= f(x, y) - \lambda g(x, y) = x^2 y^2 - \lambda(2x + y - 9) \\ \frac{\partial L}{\partial x} &= 2xy^2 - 2\lambda = 0 \\ \frac{\partial L}{\partial y} &= 2yx^2 - \lambda = 0 \\ \frac{\partial L}{\partial \lambda} &= 2x + y - 9 = 0 \\ xy^2 &= \lambda \\ 2yx^2 &= \lambda \\ y &= 2x \\ x &= \frac{9}{4} \\ y &= \frac{9}{2}\end{aligned}$$

## 4 Linear algebra

**Problem 4.1.** Take the following matrices:

$$A = \begin{bmatrix} 2 & 5 \\ 2 & 1 \\ 7 & 6 \end{bmatrix}$$

$$B = \begin{bmatrix} 1 & 0 & 1 \\ 9 & 1 & 5 \end{bmatrix}$$

What is  $B \cdot A$ ?

$$B \cdot A = \begin{bmatrix} 9 & 11 \\ 55 & 76 \end{bmatrix}$$

**Problem 4.2.** Take the following matrices:

$$A = \begin{bmatrix} 5 & 3 \\ 0 & 1 \\ 1 & 2 \end{bmatrix}$$

$$B = \begin{bmatrix} 8 & 4 & 0 \\ 2 & 1 & 2 \end{bmatrix}$$

What is  $A \cdot B$ ?

$$A \cdot B = \begin{bmatrix} 46 & 23 & 6 \\ 2 & 1 & 2 \\ 12 & 6 & 4 \end{bmatrix}$$

**Problem 4.3.** What is the transpose of the following matrix?

$$A = \begin{bmatrix} e & 93 & 4.7 \\ 2 & 6.1 & 4.22 \\ 4 & \pi & 0 \end{bmatrix}$$

$$A^T = \begin{bmatrix} e & 2 & 4 \\ 93 & 6.1 & \pi \\ 4.7 & 4.22 & 0 \end{bmatrix}$$

**Problem 4.4.** Calculate the determinant of

$$A = \begin{bmatrix} 2 & 6 \\ 2 & 8 \end{bmatrix}$$

$$\det(A) = 16 - 12 = 4$$

## 5 Probability theory

**Problem 5.1.** You run an experiment where you toss a dice two times. Each time you get either 1, 2, 3, 4, 5 or 6. What is the sample space of your experiment?

Sample space:  $6 \cdot 6 = 36$ .

**Problem 5.2.** Assume that in a certain country 0.1% of the population uses a certain drug. You have a way to test drug use, which will give you a positive result in 98% of the cases where the individual is indeed a drug user and a negative result in 99.7% of the cases where the individual doesn't use the drug. What is the probability that someone with a positive drug test is indeed a drug user?

Let's take the following events:

- $A$ : positive drug test
- $A^C$ : negative drug test
- $B$ : drug user,  $P(B) = 0.001$
- $B^C$ : not a drug user,  $P(B^C) = 0.999$

$$P(B|A) = \frac{P(A|B) \cdot P(B)}{P(A|B) \cdot P(B) + P(A|B^C) \cdot P(B^C)} = \frac{0.98 \cdot 0.001}{0.98 \cdot 0.001 + 0.003 \cdot 0.999} = 0.2464 \rightarrow 24.64\%$$

**Problem 5.3.** You run an experiment in which you toss a dice 20 times and record how many times you ended up with a 1, 2, 3, 4, 5 or 6. Your random variable is the number of times you ended up with a 5. What is expected value of this random variable?

$X$  = number of times we ended up with a 5

$$E(X) = 20 \cdot \frac{1}{6} \approx 3.33$$