

# Seminar 2024-2025

## Stochastic Loewner Evolutions

Jean Vereecke

University of Rennes

January 7, 2025

Last modified: January 16, 2025 (11:19:29)

# Introduction and motivation

Goal: model the scaling limits of interfaces for models of 2D statistical mechanics

# Introduction and motivation

Goal: model the scaling limits of interfaces for models of 2D statistical mechanics




# Introduction and motivation

Goal: model the scaling limits of interfaces for models of 2D statistical mechanics




# Introduction and motivation

Goal: model the scaling limits of interfaces for models of 2D statistical mechanics



# Introduction and motivation

Goal: model the scaling limits of interfaces for models of 2D statistical mechanics



# Introduction and motivation

Expectations:

- ▶ Scale invariance:  $(K_t)_{t \geq 0} \sim (\lambda K_{\lambda^{-2}t})_{t \geq 0}$
- ▶ Domain Markov property

# Introduction and motivation

Expectations:

- ▶ Scale invariance:  $(K_t)_{t \geq 0} \sim (\lambda K_{\lambda^{-2}t})_{t \geq 0}$
- ▶ Domain Markov property



# Introduction and motivation


Expectations:

- ▶ Scale invariance:  $(K_t)_{t \geq 0} \sim (\lambda K_{\lambda^{-2}t})_{t \geq 0}$
- ▶ Domain Markov property

# Introduction and motivation

Expectations:

- ▶ Scale invariance:  $(K_t)_{t \geq 0} \sim (\lambda K_{\lambda^{-2}t})_{t \geq 0}$
- ▶ Domain Markov property



`../dossier/rust8.0.png`

# Introduction and motivation

Create links between probability, geometry and analysis

# Table of contents

# Conformal transformations

## Definition (Conformal transformation)

Let  $U$  and  $V$  be open domains of  $\mathbb{C}$  and let  $f : U \rightarrow V$  be a map. We say that  $f$  is a *conformal transformation* if and only if  $f$  is holomorphic and bijective.

## Examples

- ▶  $f : \mathbb{D} \ni w \mapsto \lambda \frac{w+z}{1+\bar{z}w} \in \mathbb{D}$  where  $\lambda \in \mathbb{U}$  and  $z \in \mathbb{D}$ .
- ▶  $g : \mathbb{D} \ni z \mapsto i \frac{1+z}{1-z} \in \mathbb{H}$ . Its inverse is  $f : \mathbb{H} \ni z \mapsto \frac{z-i}{z+i} \in \mathbb{D}$ .
- ▶  $f(z) = \frac{az+b}{cz+d}$  where  $ad - bc = 1$ .

# Conformal transformations

## Definition (Conformal transformation)

Let  $U$  and  $V$  be open domains of  $\mathbb{C}$  and let  $f : U \rightarrow V$  be a map. We say that  $f$  is a *conformal transformation* if and only if  $f$  is holomorphic and bijective.

## Examples

- ▶  $f : \mathbb{D} \ni w \mapsto \lambda \frac{w+z}{1+\bar{z}w} \in \mathbb{D}$  where  $\lambda \in \mathbb{U}$  and  $z \in \mathbb{D}$ .
- ▶  $g : \mathbb{D} \ni z \mapsto i \frac{1+z}{1-z} \in \mathbb{H}$ . Its inverse is  $f : \mathbb{H} \ni z \mapsto \frac{z-i}{z+i} \in \mathbb{D}$ .
- ▶  $f(z) = \frac{az+b}{cz+d}$  where  $ad - bc = 1$ .

# Conformal transformations

## Definition (Conformal transformation)

Let  $U$  and  $V$  be open domains of  $\mathbb{C}$  and let  $f : U \rightarrow V$  be a map. We say that  $f$  is a *conformal transformation* if and only if  $f$  is holomorphic and bijective.

## Examples

- ▶  $f : \mathbb{D} \ni w \mapsto \lambda \frac{w+z}{1+\bar{z}w} \in \mathbb{D}$  where  $\lambda \in \mathbb{U}$  and  $z \in \mathbb{D}$ .
- ▶  $g : \mathbb{D} \ni z \mapsto i \frac{1+z}{1-z} \in \mathbb{H}$ . Its inverse is  $f : \mathbb{H} \ni z \mapsto \frac{z-i}{z+i} \in \mathbb{D}$ .
- ▶  $f(z) = \frac{az+b}{cz+d}$  where  $ad - bc = 1$ .

# Conformal transformations

## Definition (Conformal transformation)

Let  $U$  and  $V$  be open domains of  $\mathbb{C}$  and let  $f : U \rightarrow V$  be a map. We say that  $f$  is a *conformal transformation* if and only if  $f$  is holomorphic and bijective.

## Examples

- ▶  $f : \mathbb{D} \ni w \mapsto \lambda \frac{w+z}{1+\bar{z}w} \in \mathbb{D}$  where  $\lambda \in \mathbb{U}$  and  $z \in \mathbb{D}$ .
- ▶  $g : \mathbb{D} \ni z \mapsto i \frac{1+z}{1-z} \in \mathbb{H}$ . Its inverse is  $f : \mathbb{H} \ni z \mapsto \frac{z-i}{z+i} \in \mathbb{D}$ .
- ▶  $f(z) = \frac{az+b}{cz+d}$  where  $ad - bc = 1$ .



# Riemann mapping theorem

## Theorem (Riemann mapping theorem)

*Suppose that  $U$  is a simply connected domain with  $U \neq \mathbb{C}$  and let  $z \in U$ . Then there exists a unique conformal transformation  $f : \mathbb{D} \rightarrow U$  with  $f(0) = z$  and  $f'(0) > 0$ .*

## Example

►  $f : \mathbb{D} \ni w \mapsto \frac{w+z}{1+\bar{z}w} \in \mathbb{D}$  where  $z \in \mathbb{D}$

# Riemann mapping theorem

## Theorem (Riemann mapping theorem)

*Suppose that  $U$  is a simply connected domain with  $U \neq \mathbb{C}$  and let  $z \in U$ . Then there exists a unique conformal transformation  $f : \mathbb{D} \rightarrow U$  with  $f(0) = z$  and  $f'(0) > 0$ .*

## Example

►  $f : \mathbb{D} \ni w \mapsto \frac{w+z}{1+\bar{z}w} \in \mathbb{D}$  where  $z \in \mathbb{D}$

# Table of contents

# Loewner equation

For  $t \geq 0$ , let  $H_t = \mathbb{H} \setminus [0, 2\sqrt{t}i]$  and  $g_t : H_t \ni z \rightarrow \sqrt{z^2 + 4t} \in \mathbb{H}$  is a conformal transformation  $H_t \rightarrow \mathbb{H}$ .

# Loewner equation

For  $t \geq 0$ , let  $H_t = \mathbb{H} \setminus [0, 2\sqrt{t}i]$  and  $g_t : H_t \ni z \rightarrow \sqrt{z^2 + 4t} \in \mathbb{H}$  is a conformal transformation  $H_t \rightarrow \mathbb{H}$ .

Two observations:

- ▶  $|g_t(z) - z| = \left| \sqrt{z^2 + 4t} - z \right| \xrightarrow{z \rightarrow \infty} 0;$
- ▶ For each  $z \in \mathbb{H}$  fixed,

$$\partial_t g_t(z) = \frac{1}{2\sqrt{z^2 + 4t}} \cdot 4 = \frac{2}{g_t(z)}.$$

# Loewner equation

For  $t \geq 0$ , let  $H_t = \mathbb{H} \setminus [0, 2\sqrt{t}i]$  and  $g_t : H_t \ni z \rightarrow \sqrt{z^2 + 4t} \in \mathbb{H}$  is a conformal transformation  $H_t \rightarrow \mathbb{H}$ .

Two observations:

- ▶  $|g_t(z) - z| = \left| \sqrt{z^2 + 4t} - z \right| \xrightarrow{z \rightarrow \infty} 0;$
- ▶ For each  $z \in \mathbb{H}$  fixed,

$$\partial_t g_t(z) = \frac{1}{2\sqrt{z^2 + 4t}} \cdot 4 = \frac{2}{g_t(z)}.$$

# Loewner equation

For  $t \geq 0$ , let  $H_t = \mathbb{H} \setminus [0, 2\sqrt{t}i]$  and  $g_t : H_t \ni z \rightarrow \sqrt{z^2 + 4t} \in \mathbb{H}$  is a conformal transformation  $H_t \rightarrow \mathbb{H}$ .

Two observations:

- ▶  $|g_t(z) - z| = \left| \sqrt{z^2 + 4t} - z \right| \xrightarrow{z \rightarrow \infty} 0;$
- ▶ For each  $z \in \mathbb{H}$  fixed,

$$\partial_t g_t(z) = \frac{1}{2\sqrt{z^2 + 4t}} \cdot 4 = \frac{2}{g_t(z)}.$$

# Loewner equation

For  $t \geq 0$ , let  $H_t = \mathbb{H} \setminus [0, 2\sqrt{t}i]$  and  $g_t : H_t \ni z \rightarrow \sqrt{z^2 + 4t} \in \mathbb{H}$  is a conformal transformation  $H_t \rightarrow \mathbb{H}$ .

Two observations:

- $|g_t(z) - z| = \left| \sqrt{z^2 + 4t} - z \right| \xrightarrow{z \rightarrow \infty} 0;$
- For each  $z \in \mathbb{H}$  fixed,

$$\partial_t g_t(z) = \frac{1}{2\sqrt{z^2 + 4t}} \cdot 4 = \frac{2}{g_t(z)}.$$

This is a special case of Loewner's theorem.



# Loewner equation

## Definition

We say that  $A \subset \mathbb{H}$  is a *compact  $\mathbb{H}$ -hull* iff  $A$  and  $\mathbb{H} \setminus A$  are simply connected.

## Proposition

*For all compact  $\mathbb{H}$ -hull  $A$ , there exists a unique conformal transformation  $g_A : \mathbb{H} \setminus A \rightarrow \mathbb{H}$  with*

$$|g_A(z) - z| \xrightarrow{|z| \rightarrow \infty} 0.$$

*It also verifies*

$$g_A(z) = z + \frac{\text{hcap}(A)}{z} + O\left(\frac{\text{hcap}(A) \text{rad}(A)}{|z|^2}\right).$$

# Loewner equation

## Definition

We say that  $A \subset \mathbb{H}$  is a *compact  $\mathbb{H}$ -hull* iff  $A$  and  $\mathbb{H} \setminus A$  are simply connected.

## Proposition

For all compact  $\mathbb{H}$ -hull  $A$ , there exists a unique conformal transformation  $g_A : \mathbb{H} \setminus A \rightarrow \mathbb{H}$  with

$$|g_A(z) - z| \xrightarrow{|z| \rightarrow \infty} 0.$$

It also verifies

$$g_A(z) = z + \frac{\text{hcap}(A)}{z} + O\left(\frac{\text{hcap}(A) \text{rad}(A)}{|z|^2}\right).$$

## Definition

Let  $(A_t)_{t \geq 0}$  be a family of compact  $\mathbb{H}$ -hulls, we say that  $A$  is:

- ▶ *non-decreasing* if and only if  $0 \leq s \leq t < \infty$ , implies  $A_s \subset A_t$ ;
- ▶ *locally growing* if and only if for every  $T, \varepsilon > 0$ , there exists  $\delta > 0$  such that  $0 \leq s \leq t \leq s + \delta \leq T$  implies that  $\text{diam}(g_s(A_t \setminus A_s)) \leq \varepsilon$ ;
- ▶ *parametrized by half-plane capacity* if and only if for all  $t \geq 0$ ,  $\text{hcap}(A_t) = 2t$ .

Let  $\mathcal{A}$  be the set of all families of compact  $\mathbb{H}$ -hulls verifying all three conditions.

## Definition

Let  $(A_t)_{t \geq 0}$  be a family of compact  $\mathbb{H}$ -hulls, we say that  $A$  is:

- ▶ *non-decreasing* if and only if  $0 \leq s \leq t < \infty$ , implies  $A_s \subset A_t$ ;
- ▶ *locally growing* if and only if for every  $T, \varepsilon > 0$ , there exists  $\delta > 0$  such that  $0 \leq s \leq t \leq s + \delta \leq T$  implies that  $\text{diam}(g_s(A_t \setminus A_s)) \leq \varepsilon$ ;
- ▶ *parametrized by half-plane capacity* if and only if for all  $t \geq 0$ ,  $\text{hcap}(A_t) = 2t$ .

Let  $\mathcal{A}$  be the set of all families of compact  $\mathbb{H}$ -hulls verifying all three conditions.

## Definition

Let  $(A_t)_{t \geq 0}$  be a family of compact  $\mathbb{H}$ -hulls, we say that  $A$  is:

- ▶ *non-decreasing* if and only if  $0 \leq s \leq t < \infty$ , implies  $A_s \subset A_t$ ;
- ▶ *locally growing* if and only if for every  $T, \varepsilon > 0$ , there exists  $\delta > 0$  such that  $0 \leq s \leq t \leq s + \delta \leq T$  implies that  $\text{diam}(g_s(A_t \setminus A_s)) \leq \varepsilon$ ;
- ▶ *parametrized by half-plane capacity* if and only if for all  $t \geq 0$ ,  $\text{hcap}(A_t) = 2t$ .

Let  $\mathcal{A}$  be the set of all families of compact  $\mathbb{H}$ -hulls verifying all three conditions.

## Definition

Let  $(A_t)_{t \geq 0}$  be a family of compact  $\mathbb{H}$ -hulls, we say that  $A$  is:

- ▶ *non-decreasing* if and only if  $0 \leq s \leq t < \infty$ , implies  $A_s \subset A_t$ ;
- ▶ *locally growing* if and only if for every  $T, \varepsilon > 0$ , there exists  $\delta > 0$  such that  $0 \leq s \leq t \leq s + \delta \leq T$  implies that  $\text{diam}(g_s(A_t \setminus A_s)) \leq \varepsilon$ ;
- ▶ *parametrized by half-plane capacity* if and only if for all  $t \geq 0$ ,  $\text{hcap}(A_t) = 2t$ .

Let  $\mathcal{A}$  be the set of all families of compact  $\mathbb{H}$ -hulls verifying all three conditions.

## Definition

Let  $(A_t)_{t \geq 0}$  be a family of compact  $\mathbb{H}$ -hulls, we say that  $A$  is:

- ▶ *non-decreasing* if and only if  $0 \leq s \leq t < \infty$ , implies  $A_s \subset A_t$ ;
- ▶ *locally growing* if and only if for every  $T, \varepsilon > 0$ , there exists  $\delta > 0$  such that  $0 \leq s \leq t \leq s + \delta \leq T$  implies that  $\text{diam}(g_s(A_t \setminus A_s)) \leq \varepsilon$ ;
- ▶ *parametrized by half-plane capacity* if and only if for all  $t \geq 0$ ,  $\text{hcap}(A_t) = 2t$ .

Let  $\mathcal{A}$  be the set of all families of compact  $\mathbb{H}$ -hulls verifying all three conditions.

## Theorem (Loewner equation)

*Suppose that  $(A_t)_t$  is in  $\mathcal{A}$  with  $A_0 = \emptyset$ . For each  $t \geq 0$ , let  $g_t = g_{A_t}$ . There exists  $U : [0, \infty) \rightarrow \mathbb{R}$  continuous such that*

$$\partial_t g_t(z) = \frac{2}{g_t(z) - U_t}, \quad g_0(z) = z.$$

*The function  $U$  is called the “Loewner driving function” for  $A$ .*



## Definition

Suppose that  $(A_t)_t$  is a random family in  $\mathcal{A}$  encoded with the Loewner driving function  $U$ . We say that  $(A_t)_t$  satisfies the *conformal Markov property* if and only if for each  $t \geq 0$ , we denote  $\mathcal{F}_t = \sigma(U_s : s \leq t)$  and

1. (Markov property)

$$\mathcal{L}((g_t(A_{t+s}) - U_t)_{s \geq 0} | \mathcal{F}_t) = \mathcal{L}((A_s)_{s \geq 0})$$

$$(g_t(A_{t+s}) - U_t)_{s \geq 0} \perp\!\!\!\perp (\mathcal{F}_s)_{s \geq 0};$$

2. (scale invariance)

$$\forall r > 0, (rA_{t/r^2})_{t \geq 0} \sim (A_t)_{t \geq 0}.$$

## Definition

Suppose that  $(A_t)_t$  is a random family in  $\mathcal{A}$  encoded with the Loewner driving function  $U$ . We say that  $(A_t)_t$  satisfies the *conformal Markov property* if and only if for each  $t \geq 0$ , we denote  $\mathcal{F}_t = \sigma(U_s : s \leq t)$  and

1. (Markov property)

$$\mathcal{L}((g_t(A_{t+s}) - U_t)_{s \geq 0} | \mathcal{F}_t) = \mathcal{L}((A_s)_{s \geq 0})$$

$$(g_t(A_{t+s}) - U_t)_{s \geq 0} \perp\!\!\!\perp (\mathcal{F}_s)_{s \geq 0};$$

2. (scale invariance)

$$\forall r > 0, (rA_{t/r^2})_{t \geq 0} \sim (A_t)_{t \geq 0}.$$

## Definition

Suppose that  $(A_t)_t$  is a random family in  $\mathcal{A}$  encoded with the Loewner driving function  $U$ . We say that  $(A_t)_t$  satisfies the *conformal Markov property* if and only if for each  $t \geq 0$ , we denote  $\mathcal{F}_t = \sigma(U_s : s \leq t)$  and

1. (Markov property)

$$\mathcal{L}((g_t(A_{t+s}) - U_t)_{s \geq 0} | \mathcal{F}_t) = \mathcal{L}((A_s)_{s \geq 0})$$

$$(g_t(A_{t+s}) - U_t)_{s \geq 0} \perp\!\!\!\perp (\mathcal{F}_s)_{s \geq 0};$$

2. (scale invariance)

$$\forall r > 0, (rA_{t/r^2})_{t \geq 0} \sim (A_t)_{t \geq 0}.$$

## Theorem (Schramm)

*If  $(A_t)$  satisfies the conformal Markov property, then there exists  $\kappa \geq 0$  such that  $U_t = \sqrt{\kappa}B_t$  where  $B$  is a Brownian motion i.e.*

$$\partial_t g_t(z) = \frac{2}{g_t(z) - \sqrt{\kappa}B_t}, \quad g_0(z) = z.$$

# Table of contents

../dossier/SLE0.0.png

../dossier/SLE1.0.png

../dossier/SLE2.0.png



../dossier/SLE3.0.png

../dossier/SLE4.0.png

../dossier/SLE5.0.png

../dossier/SLE6.0.png

../dossier/SLE7.0.png

../dossier/SLE8.0.png

../dossier/SLE9.0.png

../dossier/SLE10.0.png



# Conclusion

Thank you for your attention ! Do you have any question ?

- [KN04] Wouter Kager and Bernard Nienhuis. “A Guide to Stochastic Löwner Evolution and Its Applications”. In: *Journal of Statistical Physics* 115.5/6 (June 2004), pp. 1149–1229. ISSN: 0022-4715. DOI: 10.1023/b:joss.0000028058.87266.be. URL: <http://dx.doi.org/10.1023/B:J0SS.0000028058.87266.be>.
- [Ken09] Tom Kennedy. “Numerical Computations for the Schramm-Loewner Evolution”. In: *Journal of Statistical Physics* 137.5–6 (Nov. 2009), pp. 839–856. ISSN: 1572-9613. DOI: 10.1007/s10955-009-9866-2. URL: <http://dx.doi.org/10.1007/s10955-009-9866-2>.
- [Bef15] Vincent Beffara. *Schramm–Loewner Evolution*. Draft notes as of January 20, 2015. 2015. URL: <https://www.newton.ac.uk/files/seminar/20150113133014301-284588.pdf>.
- [Mil19] Jason Miller. *Notes on Schramm-Loewner Evolution (SLE)*. Lecture notes, University of Cambridge, Lent Term 2019. 2019. URL: [https://www.statslab.cam.ac.uk/~jpm205/teaching/lent2019/sle\\_notes.pdf](https://www.statslab.cam.ac.uk/~jpm205/teaching/lent2019/sle_notes.pdf).
- [FLM22] James Foster, Terry Lyons, and Vlad Margarint. “An Asymptotic Radius of Convergence for the Loewner Equation and Simulation of  $SLE_{\kappa}$  Traces via Splitting”. In: *Journal of Statistical Physics* 189.2 (Sept. 2022). ISSN: 1572-9613. DOI: 10.1007/s10955-022-02979-3. URL: <http://dx.doi.org/10.1007/s10955-022-02979-3>.
- [Fos24] James M. Foster. *sle.py: Simulation of Schramm-Loewner Evolution (SLE)*. <https://github.com/james-m-foster/sle-simulation/blob/master/sle.py>. Accessed: 2024-12-28. 2024.