



Modelling monotonic effects of ordinal predictors in Bayesian regression models

Paul-Christian Bürkner^{1*}  and Emmanuel Charpentier²

¹Department of Computer Science, Aalto University, Finland

²Assistance publique – Hôpitaux de Paris, France

Ordinal predictors are commonly used in regression models. They are often incorrectly treated as either nominal or metric, thus under- or overestimating the information contained. Such practices may lead to worse inference and predictions compared to methods which are specifically designed for this purpose. We propose a new method for modelling ordinal predictors that applies in situations in which it is reasonable to assume their effects to be monotonic. The parameterization of such monotonic effects is realized in terms of a scale parameter b representing the direction and size of the effect and a simplex parameter ς modelling the normalized differences between categories. This ensures that predictions increase or decrease monotonically, while changes between adjacent categories may vary across categories. This formulation generalizes to interaction terms as well as multilevel structures. Monotonic effects may be applied not only to ordinal predictors, but also to other discrete variables for which a monotonic relationship is plausible. In simulation studies we show that the model is well calibrated and, if there is monotonicity present, exhibits predictive performance similar to or even better than other approaches designed to handle ordinal predictors. Using Stan, we developed a Bayesian estimation method for monotonic effects which allows us to incorporate prior information and to check the assumption of monotonicity. We have implemented this method in the R package *brms*, so that fitting monotonic effects in a fully Bayesian framework is now straightforward.

1. Introduction

Over the last few decades, a substantial amount of statistical research has been devoted to handling ordinal response variables in regression models, starting with the seminal paper of McCullagh (1980; for an overview, see also Agresti, 2010; Bürkner & Vuorre, 2019; Liu & Agresti, 2005; Tutz, 2011). In psychology, for instance, this kind of data is omnipresent in the form of Likert scale items, which are often treated as continuous out of convenience without ever testing this assumption (Liddell & Kruschke, 2017). With researchers realizing the importance of correctly modelling ordinal responses, the related models – often simply called *ordinal models* – are now increasingly applied in scientific practice. In the statistical language R (R Core Team, 2018), for instance, several packages are available to fit ordinal models, such as *ordinal* (Christensen, 2018), *VGAM* (Yee, Stoklosa, & Huggins, 2015), and *brms* (Bürkner, 2017, 2018), to name the perhaps most general ones.

*Correspondence should be addressed to Paul-Christian Bürkner, Department of Computer Science, Aalto University, Konemiehentie 2, 02150 Espoo, Finland (email: paul.buerkner@gmail.com).

Ordinal *predictors* seem to have received less attention in statistical research. In R, for instance, the standard treatment of ordinal predictors is still to compute orthogonal polynomials on their integer representations to model linear, quadratic, cubic, etc. terms of the predictors (Chambers & Hastie, 1992). We believe this approach to be suboptimal for various reasons, most notably because it assumes the ordinal categories to be equidistant, which is clearly an oversimplification, and because it yields parameter estimates we would consider hard to interpret.

The literature on ordinal predictors may be divided into three partially interconnected lines of research: the first is based on penalized regression/spline approaches specifically designed for ordinal predictors (Alvarez, Bailey, & Katz, 2011; Gertheiss, 2014; Gertheiss & Oehrlin, 2011; Gertheiss & Tutz, 2009; Gu, 2013); the second are categorical types of isotonic regression (Barlow, Bremner, Brunk, & Bartholomew, 1972; Robertson, Wright, & Dykstra, 1988); and the third are ordinal latent variable models (Jöreskog, 1994; Winship & Mare, 1984). We begin by explaining the penalized regression approach. The main idea of the method proposed by Gertheiss and Tutz (2009) is to penalize large differences between adjacent categories. This is done by imposing a penalty on the squared differences between the means of adjacent categories, that is, on $(\eta(x) - \eta(x-1))^2$, where x denotes values of the ordinal predictor and $\eta(x)$ denotes the predicted mean at category x . The penalty reflects the expectation that, if a predictor is ordinal, changes may happen smoothly and larger differences should thus be unlikely. This approach allows for a principled and flexible handling of ordinal predictors in a way closely related to regression splines (Gertheiss & Tutz, 2009; Gu, 2013). It also has a Bayesian interpretation in terms of priors on the category means (Gertheiss & Tutz, 2009). In the original version of this approach (used in Gertheiss, 2014; Gertheiss & Oehrlin, 2011; Gertheiss & Tutz, 2009), the direction of the changes remains unspecified and may vary across the range of the ordinal variable.

In many practical settings, we do often expect the changes between adjacent categories to be *monotonic*, that is, consistently negative or positive across the full range of the ordinal variable (e.g., Barlow *et al.*, 1972). For instance, subjective well-being may be monotonically related to measures of physical or psychological health, which we would typically assess via Likert scales and hence in an ordinal manner. If we have theoretical reasons to expect a monotonic relationship, we may want to incorporate this assumption into our model to improve accuracy of the parameter estimates and predictions, but of course also to test whether this assumption was justified in the first place. Even when monotonicity is justified, the size of the changes may still vary across ordinal categories by a substantial amount as ordinality does not contain information about the distance between categories.

The major line of statistical research which concerns itself with regression models subject to order constraints (i.e., monotonicity) is known as *isotonic regression*¹ (Barlow *et al.*, 1972; Robertson *et al.*, 1988). Depending on the research question and nature of the variable on which we want to impose a monotonicity constraint, different techniques may be more favourable. If the variable is essentially continuous, such as time intervals or the dose of a drug, we can use parametric functions which are known to be monotonic (e.g., the log or logistic function in simple cases) or use semi-parametric approaches such as monotonic splines (Gu, 2013; He & Shi, 1998; Helwig, 2017; Kelly & Rice, 1990; Lee,

¹ The term “isotonic” is mostly used synonymously with “monotonic” in the mathematical-statistical literature. We prefer the latter as we believe it to be understandable by a wider audience outside of mathematics.

1996; Leitenstorfer & Tutz, 2006; Pya & Wood, 2015; Ramsay, 1988; Wang & Small, 2015). If the variable under study is categorical, the monotonicity assumption reduces to an ordering constraint on the predicted category means. Using frequentist approaches, the latter case has been studied extensively in Barlow *et al.* (1972) and Robertson *et al.* (1988; see also Best & Chakravarti, 1990; Dykstra & Robertson, 1982; Lee, 1981; Rufibach, 2010; Wu, Woodroffe, & Mentz, 2001). Bayesian approaches to order constraint category means and testing of these constraints have been developed as well (e.g., Danaher, Roy, Chen, Mumford, & Schisterman, 2012; Klugkist & Mulder, 2008; Mulder & Raftery, 2019). For the purpose of studying ordinal predictors, we are primarily interested in the categorical type of isotonic regression although continuous types may provide useful predictions also for categorical predictors if they have a sufficient number of categories (e.g., see Helwig, 2017). Building on the penalized regression of Gertheiss and Tutz (2009) and Gu (2013), Helwig (2017) proposed to impose order constraints on the category means so that the implied relationship between response and ordinal predictor is monotonic. Combining the two approaches can lead to improved predictions compared to penalized or isotonic regression alone, provided that the true relationship is monotonic (Helwig, 2017).

The above-described approaches to modelling ordinal predictors, especially those which induce some regularization, have good theoretical and practical properties when it comes to predictive accuracy (e.g., Gertheiss & Tutz, 2009; Gu, 2013; Helwig, 2017). Furthermore, the parameter estimates are easy to interpret as they simply consist of the (regularized) response means for each ordinal predictor category. As such, they are conceptually closer to how categorical predictors, rather than continuous predictors, are handled in regression models. In contrast, in the present paper, we introduce a new monotonicity imposing parameterization for ordinal predictor terms which behaves much like a continuous predictor term. However, we do not make the assumption of equidistance of the predictor values, which is clearly unwarranted for ordinal variables. The proposed parameterization is designed to fit naturally into generalized linear modelling frameworks and their extensions. As such, it can be seamlessly combined with other types of predictor terms to model parameters of arbitrary response distributions, and may even be used within interactions or multilevel structures. To make this approach easy to remember, we simply call it *monotonic effects*, by which, of course, we do not wish to imply that this is the only possible way to impose monotonicity. As explained in detail in the next section, the estimated parameters have an intuitive meaning and are thus easy to interpret and communicate. In contrast to existing approaches, we work in a fully Bayesian framework for model specification and estimation, which increases the complexity of models in which monotonic effects can be incorporated and also allows us to specify prior distributions on the corresponding parameters. The latter may be used not only to incorporate additional subject-matter knowledge into the model that would otherwise remain unused, but also to regularize the model's predictions and make it robust against overfitting even in the absence of such specific knowledge.

The method proposed in the present paper, as well as other approaches discussed above, model ordinal predictors as *manifest* variables, that is, do not explicitly consider potential measurement error in these predictors. In contrast, in latent variable models, it is common to model ordinal variables as indicators of an underlying *latent* continuous variable, from which the observed ordinal variable originated via categorization (e.g., Finney & DiStefano, 2006; Jöreskog, 1994; Lei, 2009; Winship & Mare, 1984). Such models then estimate the relationship between this latent variable and the (manifest or latent) response variable. This way, latent ordinal models are able to take measurement error into

account and provide estimates of how the relationship between variables would have been if we had been able to directly observe the underlying true continuous construct. Importantly, this also implies a monotonic relationship between the manifest ordinal predictor and the response variable. A latent approach may be a reasonable modelling choice if there is substantial measurement error and/or we are interested in the (hypothetical) latent relationships between variables. In contrast, a manifest approach maybe a reasonable choice if the variables are measured very precisely or simply known by design (e.g., discrete points in time in a longitudinal study), if the main focus is on making predictions for new response values, and/or if the observable manifest relationships between variables are simply those which are of interest. It is beyond the scope of this paper to make a general point about manifest versus latent approaches for ordinal variables. However, we wish to point out that our proposed method treats ordinal predictors as manifest variables, as is the case for a lot of other prominent approaches (e.g., Gertheiss & Tutz, 2009; Gu, 2013; Klugkist & Mulder, 2008).

The structure of this paper is as follows. In Section 2 we will introduce monotonic effects as well as their mathematical foundation in detail. We continue by explaining a software implementation of monotonic effects in the R package *brms* (Bürkner, 2017, 2018) in Section 3, which supports a wide and growing range of Bayesian regression models. In Section 4 we perform a simulation study to investigate parameter recovery of monotonic effects and compare their performance to other approaches proposed in the literature. In Section 5 we discuss a case study dealing with measures of chronic widespread pain (Cieza *et al.*, 2004; Gertheiss, Hogger, Oberhauser, & Tutz, 2011), in which we make extensive use of monotonic effects. We conclude with a discussion in Section 6. Mathematical proofs concerning the properties of monotonic effects are presented in Appendix A and further simulation results are presented in Appendix B.

2. Monotonic effects

We will develop monotonic effects in the context of a distributional regression framework (Bürkner, 2018) in which the response y is distributed according to distribution D with P distributional parameters ψ_1, \dots, ψ_P . We write

$$y_n \sim D(\psi_{1n}, \psi_{2n}, \dots, \psi_{Pn})$$

to stress the dependence on the n th observation. The domain of each parameter ψ_p depends on the distribution D . For instance, the mean parameter of a normal distribution might take on all real values, while the probability parameter of a binomial distribution can only take on values in the interval $[0, 1]$. Each ψ_p ($1 \leq p \leq P$; with individual elements ψ_{pn}) may be predicted by a vector of predictor variables $\mathbf{X} = (\mathbf{x}_1, \dots, \mathbf{x}_K)$, where each variable \mathbf{x}_k is itself a vector of length N . To regress ψ_p on \mathbf{X} , we formulate ψ_p in terms of a generalized linear model (GLM). To reduce the burden of notation, we will drop the index p in the following as the GLM is formulated in the same way for all distributional parameters. We write $\psi = g(\eta)$, where g is the response function (i.e., inverse link function) and $\eta \in \mathbb{R}^N$ is a linear predictor term. Its n th element, η_n , may be written as.

$$\eta_n = \sum_{j=0}^J b_j f_j(X_n). \quad (1)$$

In equation (1), X_n denotes the vector (x_{1n}, \dots, x_{Kn}) of predictor values of the n th observation, the f_j are (possibly nonlinear) transformations of the predictor variables and

the b_j are the regression coefficients. Typically, $f_0 = 1$ is a constant function to include an intercept into the model. The notation above is a slightly non-standard formulation of a GLM (in fact we could speak of a generalized additive model in this context; Hastie, 2017). We use this notation in order to naturally generalize the framework to monotonic effects as explained in what follows.

A predictor variable which we want to model as monotonic must have discrete values in an ordered set, which are coded as integers. The integer value may represent, for instance, count data, discrete points in time, or categories of an ordinal variable. Since the latter is possibly the most relevant use case in psychology and related disciplines, in the following, we will concentrate on this example of an application for a monotonic predictor. We will refer to the values of such a variable as *predictor categories*. As opposed to the values of a continuous predictor, predictor categories should not be assumed equidistant with respect to their effect on the response variable. Instead, the distance between adjacent predictor categories is estimated from the data and may vary across categories.

Suppose we have an ordinal predictor \mathbf{x} which we want to model as having a monotonic effect. Ordinal variables contain no information about the distance between adjacent categories. Thus, without loss of generality, we can code the categories of \mathbf{x} so that the lowest possible category is 0,² and the largest is D . Since we start counting at 0, D is equal to the number of differences between two adjacent categories and also equal to the total number of categories minus 1. For any value $x \in \{0, \dots, D\}$ that \mathbf{x} can take on, we define.

$$\text{mo}: \{0, \dots, D\} \rightarrow [0, D], x \rightarrow \text{mo}(x, \zeta) = D \sum_{i=1}^x \zeta_i \quad (2)$$

and call it the *monotonic transform*. For notational convenience, we set $\sum_{i=1}^0 \zeta_i = 0$. The vector ζ is defined as a simplex, which means that it satisfies $\zeta_i \in [0, 1]$ and $\sum_{i=1}^D \zeta_i = 1$. By definition, the elements of ζ represent the normalized distances between consecutive predictor categories. As we can identify any set of $D + 1$ ordinal categories with $\{0, \dots, D\}$, the monotonic transform is invariant under ordinality preserving transformations of \mathbf{x} .

The additive increment of x_n (i.e., the n th value of \mathbf{x}) to η_n can be written as

$$b\text{mo}(x_n, \zeta) = bD \sum_{i=1}^{x_n} \zeta_i, \quad (3)$$

where b can take on any real value. In the above parameterization, b represents the size and the sign of the effect similar to an ordinary regression coefficient. That is, we do not have to specify the sign of the monotonic effect *a priori* but let the model find out itself if the effect is positive or negative, just as we do for coefficients in ordinary regression models. To explicitly bring monotonic effects into our GLM framework from equation (1) we can set $b_j = b$ and $f_j = \text{mo}(\cdot, \zeta)$. However, monotonic effects cannot be included in standard GLMs because the transformations f_j are not fully known *a priori* but contain the

² Note that this convention differs of the one customarily used in statistical software, where indices of vectors, matrices, etc. usually start at 1. However, starting at 0 simplifies the notation of monotonic effects and so we adopt this approach in the present paper.

parameter ζ , which needs to be estimated along with all other model parameters. As such, monotonic effects have some similarities with regression splines, a fact we will return to later on.

If the monotonic effect is used in a linear model, b can be interpreted as the expected average difference between two adjacent categories of \mathbf{x} , while ζ_i describes the expected difference between the categories i and $i - 1$ in the form of a proportion of the overall difference between lowest and highest categories. Thus, this parameterization has an intuitive interpretation while guaranteeing the monotonicity of the effect (see Section A.1 in Appendix A). As visualized in Figure 1, we can understand monotonic effects as implying a piecewise linear curve of which all components have the same sign. In a simple linear model, monotonic effects are equivalent to categorical isotonic regression (see Section A.2). A conceptual advantage of monotonic effects over isotonic regression – or other approaches working directly on the category means – is that the former outputs a single regression coefficient, b , which can directly be post-processed further. An example where this is useful are path models, in which regression coefficients are multiplied along the paths of interest, and monotonic effects can naturally be incorporated in such models.

Interaction terms including a monotonic predictor \mathbf{x} can be canonically written as.

$$bmo(x_n, \zeta)f(X_n). \quad (4)$$

where $f(\cdot)$ is an arbitrary function on the set of predictor variables \mathbf{X} , and may of course include further monotonic effects. In more complex predictor terms, monotonic effects of \mathbf{x} may also appear multiple times. As such, one modelling choice to be made is whether different simplex parameters related to \mathbf{x} should be the same or allowed to have different values. For example, a linear predictor term consisting of an intercept as well as the main effects and two-way interaction between a monotonic predictor \mathbf{x} and a continuous (or coded nominal) predictor \mathbf{z} could be formulated as




Figure 1. Visualization of a monotonic effect with $D + 1 = 4$ predictor categories. Parameters were set to $b = 100/3$ and $\zeta = (.6, .3, .1)$.

$$\eta_n = b_0 + b_1 z_n + b_2 \text{mo}(x_n, \zeta_2) + b_3 z_n \text{mo}(x_n, \zeta_3), \quad (5)$$

where ζ_2 and ζ_3 are two simplex parameters related to \mathbf{x} . If ζ_2 and ζ_3 are different, \mathbf{x} may not necessarily be conditionally monotonic for all values of \mathbf{z} (see Section A.4 for a counter-example). Rather the monotonicity being modelled depends on the chosen parameterization. For instance, if the predictor \mathbf{z} is dummy-coded as 0 and 1, representing the two categories of a dichotomous variable, the formulation above models the effect of \mathbf{x} being monotonic for category 0 as well as for the *change* between category 1 and 0. Conversely, when using cell mean coding rather than dummy coding for \mathbf{z} , the model assumes a different monotonic effect of \mathbf{x} for both categories of \mathbf{z} . In the latter case, \mathbf{x} is conditionally monotonic on \mathbf{z} . If we fix all simplex parameters corresponding to the same monotonic variable \mathbf{x} to the same value, conditional monotonicity is achieved in general (see Section A.3).

Proposition 1. *Let η be an arbitrary linear predictor term containing the monotonic predictor \mathbf{x} with the corresponding simplex parameter ζ being the same across all terms including \mathbf{x} . Then η is monotonic in \mathbf{x} conditionally on all possible combinations of all other predictor variables.*

While fixing all simplex parameters associated with \mathbf{x} to the same vector guarantees conditional monotonicity, it may be too restrictive for many common situations. For instance, if one wanted to model different monotonic effects for two groups, it would imply the shape (ζ) of the predictions being the same across groups with just their overall effect scale (b) being different. As explained in Section 2.2, in *brms* we make use of both parameterizations (varying and constant ζ) at different places in the package.

2.1. Monotonic effects in a Bayesian framework

The present paper describes monotonic effects as embedded in a fully Bayesian framework. We consider every statistical model a *Bayesian* model if it quantifies the uncertainty in all observed and unobserved variables (conventionally denoted as data and parameters, respectively) by means of probabilities. This is often expressed in terms of Bayes' theorem, which states that the posterior distribution $p(\theta)$ of the model parameters θ given the data y can be expressed in terms of the product of the likelihood $p(y|\theta)$ and the prior distribution $p(\theta)$ as well as a normalizing constant $p(y)$:

$$p(\theta|y) = \frac{p(y|\theta)p(\theta)}{p(y)}. \quad (6)$$

A thorough introduction to Bayesian statistics is outside the scope of the present paper. Instead, we refer to well-established textbooks such as McElreath (2016), Kruschke (2014), and Gelman *et al.* (2013).

With respect to monotonic effects, a fully Bayesian framework has two main implications. First, such a framework allows us to incorporate monotonic effects in a large class of regression models without the need to develop any model-specific estimators. Second, it implies that we can think of prior distributions for b and ζ . Such prior

distributions enable us to incorporate information, which does not come directly from data in terms of the likelihood contribution, such as expert knowledge or findings from previous studies.

Priors for b can be derived based on the *a priori* expectation regarding the *average difference* between adjacent categories. Any family of prior distributions typically applied to regression coefficients can be applied to b , as well. As a weakly informative prior for b , we can understand any location-scale distribution (such as a normal or Student t distribution) centred around zero and with a scale parameter large enough to allow for large but plausible average differences, while penalizing implausibly large differences. This scale will necessarily depend on the scale of the response distribution, the range of the monotonic predictor and the chosen link function (Gelman, Simpson, & Betancourt, 2017). Alternatively, one may use an improper flat prior that treats all real values as being equally likely *a priori*, in the hope that the data alone are sufficient to identify b . Importantly, when setting up a prior on b , we do not need to take into account the individual differences between adjacent categories since the latter are fully handled by the simplex parameter ζ .

Setting a prior on the simplex parameter ζ requires a different approach. A natural choice for a prior on simplex parameters is the Dirichlet distribution, a multivariate generalization of the beta distribution (Frigyik, Kapila, & Gupta, 2010). It is non-zero for all valid simplexes (i.e., for ζ with $\zeta_i \in (0, 1)$ and $\sum_{i=1}^D \zeta_i = 1$) and zero otherwise. The Dirichlet prior has a single parameter vector α of the same length as ζ . Its density is defined as

$$f(\zeta|\alpha) = \frac{1}{B(\alpha)} \prod_{i=1}^D \zeta_i^{\alpha_i-1}, \quad (7)$$

where $B(\alpha)$ is a normalizing constant (Balakrishnan, 2014). As the *a priori* expectation of ζ_i is given by $w_i = \mathbb{E}(\zeta_i) = \alpha_i/\alpha_0$, with $\alpha_0 = \sum_{i=1}^D \alpha_i$, higher values of α_i in comparison to the sum over α imply higher *a priori* values of ζ_i . Moreover, the higher the sum over α , the higher the certainty in each of the proportions w_i .

In the absence of any problem-specific information, a reasonable default prior on ζ would surely be one that assumed all differences between adjacent categories to be the same on average while being considerably uncertain about this expectation. Such a prior would imply, on average, a linear trend but with enough uncertainty to allow for all other possible monotonic trends as well. The Dirichlet prior with a constant $\alpha = 1$ puts equal probability on all valid simplexes and can thus be understood as the multivariate generalization of the uniform prior on simplexes. Since we have $w_i = 1/D$, this prior centres ζ around a linear trend with large uncertainty and thus appears to be a good default prior in the absence of any problem-specific information.

If the prior on b is centred around zero and the prior of ζ is centred around a linear trend, the implied joint prior of the monotonic effect is centred around zero, with potentially substantial uncertainty around it, depending how uncertain the priors on b and ζ are. That is, the data has to provide enough evidence for a non-zero effect in order to overcome the prior. The stronger the prior in favour of a zero effect, the more evidence we need from the data in order to be convinced of a non-zero effect. This property actually enables shrinkage priors for regression coefficient (e.g., Carvalho, Polson, & Scott, 2009; Piironen & Vehtari, 2017) to be applied to monotonic effects.

2.2. Regularizing larger changes between categories

In a Bayesian framework, larger differences between adjacent categories can naturally be penalized (i.e., made less likely *a priori*) by means of priors on b and ζ . Importantly, when we speak of penalizing larger differences, we do not mean making the overall functional form smoother. Since monotonic effects are piecewise linear, priors on b or ζ will not make them look smoother (unless the effect turns out to be exactly linear across all predictor categories). This is an important difference from other approaches such as monotonic regression splines and should be taken into account when interpreting the influence of priors on the parameter estimates obtained.

If we expect the total effect b to be small, we can use a zero-centred prior on b with comparatively small tails. For instance, if we expect b to be between -10 and 10 with probability 95% as well as higher probabilities for values closer to zero, we can use a $\text{Normal}(0, 5)$ prior. The logic behind this choice is straightforward as the normal distribution has approximately 95% probability between -2 and 2 standard deviations around its mean.

When it comes to the shape of the monotonic effect, we have to take a closer look at the prior on ζ . As discussed above, a constant vector α of the Dirichlet prior on ζ implies a linear trend in expectation. In other words, for constant α , the prior means of all changes ζ_i between adjacent categories are the same. The higher the sum over α , the higher the certainty in that expectation. Thus, if we expect a linear trend with some certainty, we assign all elements of α to the same value a . To get an intuition for what is a reasonable value for a , we may use the standard deviation of the elements ζ_i , which can be computed as (see Balakrishnan, 2014).

$$SD(\zeta_i) = \sqrt{\frac{w_i(1 - w_i)}{\alpha_0 + 1}}, \quad (8)$$

where $w_i = \alpha_i / \alpha_0$ is the expectation of the i th component. Although the standard deviation is an imperfect measure of variability for the Dirichlet distribution as the latter is not symmetric in general, we still believe the former to be helpful in better understanding the implications of one's chosen priors. For the default of $a = 1$ and a total of $D + 1 = 5$ categories, we get a rather large standard deviation of $SD(\zeta_i) = 0.19$. If we set, for example, $a = 5$, we get $SD(\zeta_i) = 0.09$ and thus much higher certainty in changes of equal size.

Of course, the process of increasing α on average works equally well even if we do not expect all changes to be the same *a priori*. For instance, if $D = 4$ and we expect a three times larger change between the first two categories than between all the other categories with some certainty, we may set $\alpha = (9, 3, 3, 3)$. As a result, we get $w_1 = 1/2$ and otherwise $w_i = 1/6$. As standard deviations, we get $SD(\zeta_1) = 0.11$ and $SD(\zeta_i) = 0.09$ otherwise.

Alternatively, and perhaps favourably, we can directly plot the marginals of the Dirichlet distribution. These marginal priors are known to be beta distributions with shape parameters $s_1 = \alpha_i$ and $s_2 = \alpha_0 - \alpha_i$ (Balakrishnan, 2014). For $\alpha = (9, 3, 3, 3)$, the marginal distributions of ζ are shown in Figure 2. All of the above approaches to better understand the Dirichlet prior have in common that they ignore the dependence between elements of ζ . More precisely, elements of ζ are always negatively correlated




Figure 2. Densities of marginal priors of ζ_1 and ζ_2 for $\alpha = (9, 3, 3, 3)$. The marginal priors of ζ_3 and ζ_4 are in this case identical to that of ζ_2 .

as an increase in one element needs to be reflected in a decrease in the other elements to satisfy the sum-to-one constraint (Balakrishnan, 2014):

$$\text{Cor}(\zeta_i, \zeta_j) = -\frac{w_i w_j}{\sqrt{w_i(1-w_i)w_j(1-w_j)}}. \quad (9)$$

A possible solution would be to plot the multivariate density of the Dirichlet prior, but this will become more difficult for higher-dimensional ζ (i.e., for variables with more than three categories) and so we do not illustrate this approach in the present paper.

3. Implementation in *brms*

The *brms* package (Bürkner, 2017, 2018) provides an interface to fit Bayesian generalized (non)linear (multilevel) regression models using Stan (Carpenter *et al.*, 2017; Stan Development Team, 2019), which is a C++ package for performing full Bayesian inference (see also <http://mc-stan.org/>). It supports a wide range of distributions, allowing users to fit (among others) linear, count data, survival, response times, ordinal, zero-inflated, and even self-defined mixture models all in a distributional multilevel context.

In *brms*, monotonic effects are fully integrated into the formula syntax, which builds on and extends standard R formula syntax as well as the multilevel formula syntax initially created for the *lme4* package (Bates, Mächler, Bolker, & Walker, 2015). Monotonic predictors can be used like any other predictor variable and, with respect to the formula syntax, behave like a numeric predictor. Suppose the response variable Y is predicted by a monotonic variable X and a non-monotonic variable Z (i.e., a continuous or categorical variable). Then the corresponding model formula is

$$Y \sim \text{mo}(x) + z$$

Modelling both main effects and interaction of X and Z can be achieved by

$$Y \sim \text{mo}(x) * z$$

Depending on whether Z is a continuous or categorical variable, this will imply a different predictor term, which is fully determined by and thus consistent with the basic R formula syntax. If Z is monotonic as well, then Z is simply replaced by $\text{mo}(Z)$. Note that for models including interactions with monotonic variables, *brms* will use *different* simplex parameters for different terms of the same monotonic variable (e.g., for the main effect of X and the interaction of X and Z). This results in much greater modelling flexibility as explained in the former section. The variable which should be modelled as monotonic may either be integer-valued or an ordered factor. In the latter case, the ordered factor will be transformed to an integer variable with the lowest factor level being identified with zero as described above.

An especially well-developed feature of *brms* is its multilevel formula syntax allowing us to model, for instance, hierarchically nested data structures such as multiple observations per person in a longitudinal study. Suppose we wanted to fit a monotonic effect per person in a multilevel model. Then we could specify this as follows:

$$Y \sim \text{mo}(x) + (\text{mo}(x) | \text{person})$$

The $\text{mo}(x)$ term outside the brackets denotes the *average* monotonic effect across persons, while the $(\text{mo}(x) | \text{person})$ term indicates that the *difference* between the individual monotonic effects per person and the average effect should be modelled as well (for more details on the *brms* formula syntax see Bürkner, 2018). For this parameterization to make sense in combination with monotonic effects, we treat the shape (i.e., the simplex parameter ζ) as constant across persons and only vary the size and direction of the effect (i.e., b) as varying across persons. This restricts the flexibility of the model but results in much more stable estimates and fewer convergence problems in particular if the number of observations per person (or more generally, per level of the grouping factor) is small.

4. Simulations

To verify the correctness of our implementation of monotonic effects and to compare them to other approaches for ordinal predictors, we performed a simulation study. All simulations were done in R (R Core Team, 2018) via the RStudio interface (RStudio Team, 2018). For data preparation and plotting we used packages from the tidyverse (Wickham, 2017) in particular *dipl* (Wickham, François, Henry, & Müller, 2019) and *ggplot2* (Wickham, 2016).

4.1. Parameter recovery

Before applying a statistical model in practice, we should first make sure that it is able to recover its own parameters (e.g., Cook, Gelman, & Rubin, 2006). This means that if data are simulated from the model under consideration (so that it is the true data generating model) we should, on average, be able to recover the true parameters of the model. What is more, our parameter estimates should have just the right amount of uncertainty so that we are neither overly certain nor overly uncertain about the location of the parameter. We may be tempted to just select a few parameter values to work as the ground truth, and evaluate parameter recovery on the basis of these. However, this is dangerous since we may (accidentally) select parameter values for which the algorithm works particularly well or particularly poorly (Talts, Betancourt, Simpson, Vehtari, & Gelman, 2018). A more robust approach is to sample the ground

truth from a distribution of ground truths in each simulation trial and then evaluate the set of estimates against the true distribution.

If an estimation algorithm for a given model succeeds in the procedure described above, we call the algorithm *well calibrated*. In Bayesian statistics, we could equivalently say that, given the prior and the likelihood, we are able to estimate the true posterior distribution of the parameters. This can be formally tested by means of *simulation-based calibration* (SBC; Talts *et al.*, 2018), a procedure which works as follows. First, sample true parameter values $\tilde{\theta}$ from the prior, $\tilde{\theta} \sim p(\theta)$. Second, sample data \tilde{y} from the likelihood, $\tilde{y} \sim p(y|\tilde{\theta})$. Third, using the algorithm which we want to validate, obtain L samples $\{\theta_1, \dots, \theta_L\}$ from the estimated posterior distribution, $\{\theta_1, \dots, \theta_L\} \sim p(\theta|\tilde{y})$. Fourth, for a quantity (or quantities) of interest $f(\theta)$, which can be computed on the basis of the posterior samples (e.g., the individual parameter estimates), count how many values in $\{f(\theta_1), \dots, f(\theta_L)\}$ fall below the true value $f(\tilde{\theta})$. We call this count the *rank statistic* and denote it by $r(\theta_{1:L}|\tilde{\theta}, f)$.

If the algorithm is well calibrated for a given model, $\tilde{\theta}$ and $\{\theta_1, \dots, \theta_L\}$ should follow the same distribution (Talts *et al.*, 2018). We can verify this by repeating the above steps multiple times (say, $T = 1,000$ times). For each repetition t , we compute the rank statistic $r_t(\theta|\tilde{\theta}, f)$. Afterwards, we create a histogram over $\{r_1(\theta_{1:L}|\tilde{\theta}, f), \dots, r_T(\theta_{1:L}|\tilde{\theta}, f)\}$ and investigate its shape. If the histogram is approximately uniform over $[0, L]$, the algorithm is well calibrated to the model. If it is skewed, the algorithm is biased. If the histogram is U-shaped or inverse-U-shaped, the estimated posterior distribution is narrower or wider, respectively, than the true posterior distribution. We may add confidence intervals to the histograms to indicate the range in which we would expect the bars to be for a well-calibrated quantity. This helps in differentiating actual estimation problems from random simulation noise. The SBC procedure needs to be adjusted slightly for use with autocorrelated samples such as those obtained by Markov chain Monte Carlo sampling. For more details see Talts *et al.* (2018).

To analyse the calibration of monotonic models using SBC in common settings, we performed a simulation study. We focused on normally distributed response variables

$$y \sim \text{Normal}(\mu, \sigma),$$

where the mean μ is regressed on some monotonic effects as detailed in the following, and σ is the residual standard deviation assumed constant across observations. This resembles a linear regression model except that the predictors were modelled as monotonic effects. We varied μ as containing either the main effect of a single monotonic predictor x ,

$$\mu = b_0 + b_1 \text{mo}(x, \zeta_x),$$

or the main effects of two monotonic predictors x and z plus their interaction,

$$\mu = b_0 + b_1 \text{mo}(x, \zeta_1) + b_2 \text{mo}(z, \zeta_2) + b_3 \text{mo}(x, \zeta_{31}) \text{mo}(z, \zeta_{32}).$$

Furthermore, the dimension of all simplex parameters was $D \in \{4, 10, 50\}$ so that the number of predictor categories of x and z took on values $D+1 \in \{4, 11, 51\}$, respectively. In each simulation trial, the values of x and z were sampled uniformly from the set of possible categories $\{0, \dots, D\}$. The number of observations took on values $N \in \{50, 200, 1,000\}$. As priors for the model parameters, we used $\text{Normal}(0, 1)$ distributions for all regression coefficients, uniform Dirichlet distributions of dimension D for all simplexes, and a truncated $\text{Normal}_+(0, 1)$ distribution for the residual standard

deviation σ . Further, regression coefficients were scaled to be independent of the number of predictor categories, that is, b_1 and b_2 were divided by D and b_3 was divided by D^2 . This ensures comparability of model predictions across different values of D . The monotonic models were fitted in Stan via the *brms* interface using 500 warm-up followed by 500 draws from the posterior obtained from a single Markov chain. For each of the $2 \times 3 \times 3 = 18$ conditions, the simulations were repeated $T = 1,000$ times.

For the sake of brevity, we only show the results of selected simulation conditions that are representative of the overall findings. A complete overview of all results is available on GitHub (<https://github.com/paul-buerkner/monotonic-effects-paper>). The SBC results for the monotonic main effect and interaction models for $N = 200$ and $D = 4$ are displayed in Figures 3 and 4, respectively. We clearly see that all model parameters are well calibrated under these conditions. Even if the number of parameters P becomes substantially larger than the number of observations N , the model is may be well calibrated as shown for the interaction model when $N = 50$ and $D = 50$ (see Figure 5).

However, this may not always be the case. In the interaction model for $N = 1,000$ and $D = 50$, we observe spikes in the histograms at very small and very large parameter values, in particular for the simplex parameters (see Figure 6). This indicates strong autocorrelation in the chains and thus convergence problems in the model (Talts *et al.*, 2018). A closer investigation of the fitted models revealed that most iterations exceeded the maximum tree depth (see Stan Development Team, 2019, for details). This indicates a highly complex posterior distribution which is hard to properly explore with the algorithm. For this model, the reason is the interaction term of two 50-dimensional simplex parameters, which the algorithm fails to explore efficiently (although model predictions are still accurate; see Section 4.2). Increasing the maximum tree depth can resolve this problem but increases the computation time noticeably.

Of course, monotonic effects can be applied in a lot of other modelling settings and so the present results provide no guarantee that they will be well calibrated in cases not studied in the present paper. Generally, we recommend building models specifically tuned to the study design, data, and subject-matter knowledge. These models should then be validated as a natural part of the research process using SBC or other validation procedures.




Figure 3. SBC results of the monotonic main effects model for $N = 200$ and $D = 4$. Facets show histograms of different model parameters whose names are taken from *brms*. Horizontal black lines indicate 99% confidence intervals under the assumption of correct calibration.




Figure 4. SBC results of the monotonic interaction model for $N = 200$ and $D = 4$. Facets show histograms of different model parameters whose names are taken from *brms*. Horizontal black lines indicate 99% confidence intervals under the assumption of correct calibration.




Figure 5. SBC results of the monotonic interaction model for $N = 50$ and $D = 50$. Facets show histograms of different model parameters whose names are taken from *brms*. For simplex parameters, only the first four elements are displayed. Horizontal black lines indicate 99% confidence intervals under the assumption of correct calibration.




Figure 6. SBC results of the monotonic interaction model for $N = 1,000$ and $D = 50$. Facets show histograms of different model parameters whose names are taken from *brms*. For simplex parameters, only the first four elements are displayed. Horizontal black lines indicate 99% confidence intervals under the assumption of correct calibration.

4.2. Comparison to other approaches

To compare the predictive performance of monotonic effects to alternative approaches, which can be used under the same circumstances, we performed another simulation study. We used the same simulation conditions as in Section 4.1 with one exception, described below. As underlying data-generating processes, we considered the main effects and interaction models described in Section 4.1 in three different variations: simplex values fixed to $1/D$, implying a linear relationship; simplex values sampled from a uniform Dirichlet distribution of dimension D , implying a nonlinear but monotonic relationship; and simplex values sampled from a uniform Dirichlet distribution of dimension D with approximately half of the values having a negative sign, implying a non-monotonic relationship.

As alternatives to the monotonic model (abbreviated as MO), we considered simple linear (LIN) and categorical (CAT) regression,³ isotonic regression (ISO; Barlow *et al.*, 1972; Robertson *et al.*, 1988), penalized ordinal regression (OS; Gertheiss & Tutz, 2009; Gu, 2013), penalized ordinal regression with monotonicity constraint (OSMO; Helwig, 2017), as well as linear and cubic spline models (LS and CS; e.g., Gu, 2013; Helwig, 2016). The latter two are primarily designed for continuous responses but may still perform reasonably well for linear relationships or a sufficiently large number of predictor categories. In what follows, we will refer to the different approaches using the abbreviations introduced above.

³ Here, categorical refers to treating the predictor(s) as categorical, not the response variable which was assumed to be normally distributed under all simulation conditions.

The MO models were fitted with *brms* using its default priors, that is, without considering the true priors used in the data-generating process. This avoids giving these models a possibly unfair advantage as in reality we are unlikely to be aware of the exact data-generating process. LIN and CAT models were fitted via the *lm* function, while ISO models were fitted via the *isoreg* function. All penalized regression/splines approaches were fitted in the *bigsplines* package (Helwig, 2018) using the *bigspline* and *bigssp* functions.

For the main effect models, simulation results are displayed in Figure 7 and 8 showing the models' root mean squared error (RMSE) under true linearity and monotonicity, respectively. From Figure 7 we see that, under true linearity, LIN performed consistently better than all other models, closely followed by CS and MO. Other penalized approaches had slightly but noticeably higher RMSEs, while unpenalized approaches such as CAT or ISO models had even higher RMSEs, in particular for larger D . From Figure 8 we see that, under true monotonicity, MO models exhibited predictive performance the same as or better than all other approaches, although the difference from CS, OS, and OSMO models was generally quite small. As expected, under true non-monotonicity, MO models performed worse than models without monotonicity assumption but similarly to other non-monotonicity-assuming models (see Figure B1 in Appendix B).

For the interaction models, simulation results are displayed in Figure 9 and 10 showing the models' RMSE under true linearity and monotonicity, respectively. We did not find




Figure 7. Simulation results for the main effects models under true linearity based on $T = 1,000$ simulation trials. Abbreviations: CAT = categorical model; CS = cubic spline model; D = number of categories minus 1; ISO = isotonic regression model; LIN = linear model; LS = linear spline model; MO = monotonic model; N = number of observations; OS = ordinal spline model; OSMO = ordinal monotonic spline model.




Figure 8. Simulation results for the main effects models under true monotonicity based on $T = 1,000$ simulation trials. LIN is not displayed as its RMSE is too large and thus obscures differences between other models. For the same reason, CAT is not displayed for $D = 50$. Abbreviations: CAT = categorical model; CS = cubic spline model; D = number of categories minus 1; ISO = isotonic regression model; LS = linear spline model; MO = monotonic model; N = number of observations; OS = ordinal spline model; OSMO = ordinal monotonic spline model.

implementations for interactions in ISO or OSMO models, which are thus not displayed in the figures. From Figure 9 we see that, under true linearity, LIN and CS models performed better than all other models, closely followed by MO, CS, and LS. For small N and large D , MO was even on par with LIN. From Figure 10 we see that, under true monotonicity, MO models had better predictive performance across all conditions than all other approaches. As expected, under true non-monotonicity, MO models performed worse than models without monotonicity assumption (see Figure B2 in Appendix B).

In summary, in our simulations, MO yielded predictions the same as or better than other penalized or unpenalized ordinal approaches if the monotonicity assumption was justified. Intuitively, one may expect that MO models tend to overfit the data in cases of small N and comparably large D , in particular for interaction models, as they have considerably more parameters than observations. However, as evident in our simulations, this is not actually what happens, although we found convergence issues under some of these conditions. The reason for this lies in the joint Dirichlet prior on the simplex parameters: if one particular element of ζ (i.e., one difference between two adjacent categories) is large, larger values of other elements are automatically penalized (i.e., made more unlikely) due to the sum-to-one constraint on ζ . The same property can be expressed in terms of the negative correlation between two distinct elements of ζ (see equation 9). This holds even if the Dirichlet prior is uniform over the set of possible simplexes, which is used as the default prior in *brms*.




Figure 9. Simulation results for the interaction models under true linearity based on $T = 1,000$ simulation trials. CAT is not displayed for $D = 50$ as its RMSE is too large and thus obscures differences between other models. ISO and OSMO are not displayed as they have no corresponding interaction model. Abbreviations: CAT = categorical model; CS = cubic spline model; D = number of categories minus 1; LIN = linear model; LS = linear spline model; MO = monotonic model; N = number of observations; OS = ordinal spline model.

There is also another aspect of the monotonic parameterization that can guard against overfitting. If the scale parameter b is close to 0, there is not much to learn about the corresponding simplex parameter ζ , which will thus have a posterior distribution close to its prior. Still, this uncertainty will not lead to overfitting as changes in ζ do not influence predictions as long as b is small. This is because the latter controls the overall effect size of the monotonic predictor, while ζ only controls the shape. In other words, the complexity of a monotonic predictor with an effect close to zero naturally reduces to the complexity of a simple linear predictor.

5. Case study: Measures of chronic widespread pain

To illustrate the application of monotonic effects in practice, we will reanalyse data used to validate measures of chronic widespread pain (CWP) from patients' point of view (Cieza *et al.*, 2004; Gertheiss *et al.*, 2011). There is no universally accepted definition of CWP, but 'it may be characterized by pain involving several regions of the body, which causes problems in functioning, psychological distress, poor quality of sleep or difficulties in daily life' (Gertheiss *et al.*, 2011, p. 378). The applied CWP measures stem from the international classification of functioning (ICF; World Health Organization, 2001) and are rated by clinical staff, not by patients themselves. Thus, it is important to validate which




Figure 10. Simulation results for the interaction models under true monotonicity based on $T = 1,000$ simulation trials. LIN is not displayed as its RMSE is too large and thus obscures differences between other models. For the same reason, CAT is not displayed for $D = 50$. ISO and OSMO are not displayed as they have no corresponding interaction model. Abbreviations: CAT = categorical model; CS = cubic spline model; D = number of categories minus 1; LS = Linear spline model; MO = monotonic model; N = number of observations; OS = ordinal spline model.

and to what degree CWP measures actually relate to subjective physical health in order to better understand their implications for patients' lives.

For each of 420 patients, the present data contain information on 67 CWP measures as well as a subjective measure of physical health based on the SF-36 questionnaire (Ware & Sherbourne, 1992). The data are freely available in the R package *ordPens* (Gertheiss, 2015) and are explained in detail in Gertheiss *et al.* (2011) and Cieza *et al.* (2004). In the data set, the variable of subjective physical health is called *phcs*, while the CWP measures are named according to their official ICF coding (see Gertheiss *et al.*, 2011, for explanation). Our fully reproducible analysis is available on OSF (<https://osf.io/kvrsg/>).

In the data set, the subjective physical health (variable *phcs*) ranges from 10.08 to 53.17 with a mean of 32.41 and a standard deviation of 8.17. For the purpose of this case study, we will predict *phcs* only by impairments in 'walking' (variable *d450*) and 'moving around' (variable *d450*), which were both measured on a five-point scale between 0 ('no problem') and 4 ('complete problem'). Both of these variables were strong predictors of *phcs* in the analysis of Gertheiss *et al.* (2011). The category labels of these variables suggest that their relationship with *phcs* will be monotonic. More specifically, we expect subjective physical health to decrease with an increase in impairments in walking or moving around or basically any other everyday functioning. Including more or even all of the 67 predictors would be possible as well in theory but barely sensible without principled variable selection techniques. Such techniques have yet to be developed for monotonic effects and are outside the scope of the present paper.




Figure 11. Effects of impairments in moving around on subjective physical health: (left) linear model; (middle) monotonic model; (right) categorical model.

We will start by predicting subjective physical health only by impairments in moving around. For the present example (and also more generally; see Section 4.2), the default priors of *brms* on monotonic effects work well in terms of sampling efficiency and convergence. However, for illustrative purposes, we still manually specify our own priors for each model even if priors are similar to the default ones. Based on knowledge about the outcome scale, it is unlikely that a one-point change in any WCP measure will imply a change in the subjective physical health by more than 5 points. We code this expectation as a $\text{Normal}(0, 2.5)$ prior on the scale parameters b . That way, $|b|$ will only exceed 2.5 and 5 outcome points with probabilities of roughly 32% and 5%, respectively. With regard to the shape of the effect of ‘moving around’, we have no particular prior expectations and thus assume a uniform Dirichlet prior as explained in Section 2.2, which is also the default in *brms*. In *brms*, we can specify the above priors by means of the following code:

```
library(brms)
prior_b <- prior(normal(0, 2.5), class = "b")
prior_s1 <- prior(dirichlet(1, 1, 1, 1), class = "simo",
                  coef = "mod4551")
```

We use class `simo` to refer to the simplex parameters of monotonic effects. The required coefficient name “mod4551” is constructed as `mo <variable><index>`, where `<index>` = 1 unless a single regression term contains multiple simplexes – which is only the case for interactions of monotonic effects. Finally, we fit the model in *brms* via

```
fit1 <- brm(phcs ~ mo(d455), data = cwp,
            prior = prior_b + prior_s1)
```

As illustrated in the middle of Figure 11, impairments in moving around show a strong negative relationship to subjective physical health. Moreover, this relationship is clearly (at least visually) nonlinear. Changes in the outcome are strongest between the first two categories and the third to fourth category. This impression is confirmed by the summary estimates of the regression parameters (see Table 1) as `simo[1]` and `simo[3]` have the largest estimates. For example, the estimate of `simo[1] = 0.38` implies that 38% of the total change in subjective physical health due to impairments in walking happens between the first two predictor categories. Further, the estimate of `slope = -2.70` implies that *on average* the subjective physical health *decreases* by 2.70 per increase in impairments in walking by one category.

Table 1. Summary of parameter estimates for impairments in moving around

	Estimate	l-95% CI	u-95% CI
Intercept	37.19	35.80	38.64
slope	−2.70	−3.34	−2.08
simo[1]	0.38	0.23	0.55
simo[2]	0.11	0.01	0.27
simo[3]	0.36	0.15	0.55
simo[4]	0.15	0.01	0.35

Note. simo = simplex parameter of the monotonic effect; Estimate = posterior mean; CI = credible interval based on quantiles.

Next, let us compare the monotonic model to a linear and an unordered categorical model, fitted as follows:

```
fit2 <- brm(phcs ~ d455, data = cwp, prior = prior_b)
fit3 <- brm(phcs ~ d455c, data = cwp, prior = prior_b)
```

The variable `d455c` denotes the categorical version of `d455` to which we applied sequential difference coding. Results are visualized on the left- and right-hand side of Figure 11. To compare models, we use approximate leave-one-out cross-validation (Vehtari, Gelman, & Gabry, 2017) together with corresponding information criteria and Akaike model weights (Vehtari *et al.*, 2017; Wagenmakers & Farrell, 2004):⁴

```
loo_compare(loo(fit1), loo(fit2), loo(fit3))
model_weights(fit1, fit2, fit3, weights = "loo")
```

As shown in Table 2, the monotonic models fits better than the categorical model followed by the linear model although the differences between the three models are not substantial. Looking more closely at the results, we see that the effective number of parameters is somewhat smaller for the monotonic model than for the categorical model: about the same difference as we see in the corresponding expected log posterior predictive density (ELPD) difference. Thus, the better predictive performance of the monotonic model is primarily driven by it being more parsimonious than the categorical model. Together, this provides evidence that the monotonicity assumption for the effect of predictor `d455` is justified by the data.

Next, we will use both impairments in walking (variable `d450`) and in moving around (variable `d455`) to predict subjective physical health. When specifying the Dirichlet prior for ‘walking’, we have to take into account that the highest category 4 (‘complete problem’) is actually not present in the data set. Thus, the corresponding prior requires a vector of reduced size.

```
prior_s2 <-
  prior(dirichlet(1, 1, 1), class = "simo",
        coef = "mod4501") +
  prior(dirichlet(1, 1, 1, 1), class = "simo",
        coef = "mod4551")
```

We fit the monotonic, linear, and categorical models as follows:

⁴In a Bayesian framework, models may be compared by various means, for instance, Bayes factors (Kass & Raftery, 1995), (approximate) cross-validation methods (Vehtari *et al.*, 2017), information criteria (Vehtari *et al.*, 2017; Watanabe, 2010) or stacking of posterior predictive distributions (Yao, Vehtari, Simpson, & Gelman, 2017). A discussion of the pros and cons of these various approaches is outside the scope of the present paper.

Table 2. Comparison of models fit1 to fit3 based on approximate leave-one-out crossvalidation

	ELPD-LOO	ELPD-Diff	SE-Diff	P-LOO	LOOIC	Akaike-Weight
fit1	-1,439.54	0.00	0.00	4.79	2,879.09	0.44
fit3	-1,439.85	-0.30	0.23	4.96	2,879.70	0.32
fit2	-1,440.14	-0.59	1.90	2.90	2,880.27	0.24

Note. ELPD-LOO = expected log posterior predictive density (higher is better); ELPD-DIFF = difference in ELPD values compared to the best model; LOOIC: leave-one-out information criterion (lower is better); P-LOO = effective number of model parameters (lower is better); Akaike-Weight = model weight based on the LOOIC values (higher is better); SE-DIFF = standard error of the ELPD difference.

```
fit4 <- brm(phcs ~ mo(d450) + mo(d455), data = cwp,  
            prior = prior_b + prior_s2)  
fit5 <- brm(phcs ~ d450 + d455, data = cwp,  
            prior = prior_b)  
fit6 <- brm(phcs ~ d450c + d455c, data = cwp,  
            prior = prior_b)
```

Conditional predictions of the three models are visualized in Figure 12. As visible on the right-hand side of Figure 12, the effect of moving around seems to be no longer monotonic when controlling for the effect of walking. Thus, we would expect the




Figure 12. Effects of impairments in walking and in moving around on subjective physical health: (left) linear model; (middle) monotonic model; (right) categorical model.

Table 3. Comparison of models fit4 to fit6 based on approximate leave-one-out crossvalidation

	ELPD-LOO	ELPD-Diff	SE-Diff	P-LOO	LOOIC	Akaike-Weight
fit6	-1,411.48	0.00	0.00	7.08	2,822.97	0.71
fit4	-1,412.45	-0.96	1.47	6.22	2,824.89	0.27
fit5	-1,415.01	-3.53	3.36	3.62	2,830.03	0.02

Note. ELPD-LOO = expected log posterior predictive density (higher is better); ELPD-DIFF = difference in ELPD values compared to the best model; LOOIC: leave-one-out information criterion (lower is better); P-LOO = effective number of model parameters (lower is better); Akaike-Weight = model weight based on the LOOIC values (higher is better); SE-DIFF = standard error of the ELPD difference.

categorical model to now show better predictions than the monotonic model. As can be seen in Table 3, this is indeed what happens as the categorical model has a higher ELPD value and corresponding model weight. That is, from a purely predictive perspective, we will likely prefer the categorical model. However, from a theoretical perspective, the situation may be different as it is more plausible that a change to worse in moving around always leads (in expectation) to a reduction in subjective physical health no matter what impairments in walking individuals may have. That is, even for impairments in walking held constant, the effect of impairments in moving around should still be monotonically decreasing. The fact that this is not strictly the case in the present data is clearly the result of the dependence structure between the two predictors as well as a large number of possible confounders that we have not accounted for in the present analysis. Just from the present data, it remains unclear how well the (non-) monotonicity will generalize to other samples or populations of impaired individuals.

6. Discussion

In the present paper, we proposed a new approach to including *monotonic effects* of ordinal predictors in regression models. The proposed parameterization not only ensures monotonicity but also naturally regularizes the model and its predictions even without the use of strong prior information. Thus, monotonic effects share important aspects with existing methods for modelling ordinal predictors. Moreover, monotonic effects nicely integrate into the framework of generalized linear regression and can even be used within multilevel models. By making an informed decision about the parameterization of interactions with monotonic effects, different kinds of monotonicity can be modelled depending on the research question and *a priori* information available. Monotonic effects are fully supported in the *brms* R package, which fits Bayesian regression models using Stan and provides an intuitive user interface based on widely known R formula syntax. Ordinal predictors are still mostly treated as either nominal or metric, thus under- or overstating the information contained. Monotonic effects avoid these problems but still allow for an intuitive interpretation of the parameters estimated. In summary, we think that monotonic effects provide a useful tool for handling of ordinal predictors in regression models in situations where the monotonicity assumption is justified.

One potential problem in the Bayesian estimation of monotonic effects is that elements of a simplex tend to be negatively correlated, sometimes rather strongly, thus making Markov chain Monte Carlo sampling more difficult (Hoffman & Gelman, 2014). However,

due to the advanced Hamiltonian Monte Carlo samplers implemented in Stan, which are designed to work well even for highly intercorrelated posteriors (Betancourt, Byrne, Livingstone, & Girolami, 2014; Hoffman & Gelman, 2014), these problems may be alleviated when fitting monotonic effects in Stan – either directly or indirectly through *brms*. Indeed, in the models estimated for the purpose of this paper, sampling efficiency and convergence were good and rarely worse than when using a purely linear approach. The only exceptions were monotonic interaction models with a very high number of predictor categories (i.e., 51 in our simulation). For that many predictor categories, it may be easier to fit (monotonic) splines or similar models which require fewer parameters.

With regard to predictions, monotonic effects performed very well under all simulation conditions where the monotonicity assumption was justified, even those where convergence was an issue. More precisely, monotonic effects made predictions similar to or better than other penalized ordinal approaches such as ordinal splines and much better predictions than unpenalized approaches such as standard isotonic regression or approaches treating the predictors as categorical. This nicely illustrates the advantages of a fully Bayesian approach, where joint priors, even weakly informative ones, can regularize the model parameters and ultimately lead to improved predictions. It has to be noted that all penalized/regularized approaches generally performed well in our simulations and differences between them were comparably small (which we believe is a good sign for the validity of these methods in general). Where monotonic effects differ from previously proposed approaches is in their conceptualization as generalizations of continuous predictor terms, rather than as (penalized) categorical predictor means. This allows for the intuitive interpretation of the scale parameter as an ordinary regression coefficient, except that we do not assume the shape of the relationship as linear, but more generally as monotonic. It is this clear separation between the strength of the relationship and its shape that, in our opinion, makes monotonic effects very appealing for interpretation and communication.

This separation is especially advantageous when dealing with interactions of ordinal predictors. If we used any type of categorical coding (e.g., dummy coding) for the interaction of two ordinal predictors, the number of regression coefficients would increase quadratically with the number of predictor categories, which complicates interpretation. When working with monotonic effects, by contrast, we would only have three related regression coefficients (two for the main effects and one for the interaction), which would essentially have the same interpretative complexity as a linear model with the ordinal predictors treated as continuous. Of course, interpreting the shape parameters of the main effects and interactions will again increase the complexity to a level similar to what is implied by categorical coding. However, often we may not be interested in the exact shape of the effect, in which case it would simply be sufficient to know that the shape has been taken into account by the model. Of course, there is nothing wrong with directly reporting and interpreting the estimated category means (or their contrasts), which, depending on one's preferences and the overall complexity of the model, may also be seen as more intuitive (see, for example, Barlow *et al.*, 1972; Danaher *et al.*, 2012; Gertheiss & Tutz, 2009; Klugkist & Mulder, 2008; Mulder & Raftery, 2019, for related approaches).

Regardless of what formulation one chooses, the assumption of monotonicity is critical and needs to be theoretically justified and/or statistically investigated. A general approach to the latter is to fit one model with and one without monotonicity constraint and then compare the two models using one's preferred criteria of model fit. From a Bayesian

perspective, we could use, for example, Bayes factors or cross-validation procedures/information criteria, depending on whether we aim to penalize prior or posterior complexity, respectively (Gelman *et al.*, 2013; Hoijtink, Klugkist, & Boelen, 2008). In our case study, we applied approximate leave-one-out cross-validation for this purpose, but other criteria would have been possible as well (see, for example, Mulder & Raftery, 2019).

As pointed out earlier, one important distinction within the class of models dealing with ordinal predictors is the assumption about the generative process of these variables. In our monotonic effects approach, as well as in a lot of other prominent approaches (e.g., Gertheiss & Tutz, 2009; Gu, 2013; Klugkist & Mulder, 2008), we treat ordinal predictors as manifest variables. Specifically, we do not make the assumption that the observations originate from the categorization of a latent continuous variable as is commonly the case in latent variable models (e.g., Finney & DiStefano, 2006; Jöreskog, 1994; Lei, 2009; Winship & Mare, 1984). Such a latent variable assumption may be sensible if the ordinal predictor represents the chosen categories of a Likert item, for instance, with the intention of measuring a latent psychological construct, but would certainly not be sensible in some other cases, for instance, if the categories were known discrete points in time. Thus, both approaches seem valid in our opinion and are called for in different settings and research questions. We note again that our proposed approach is targeted at data settings where modelling manifest observations is desired, and that if researchers desire to examine a latent relationship, they are advised to use other methods.

Although our primary focus was the use of monotonic effects for modelling strictly ordinal predictors, we wish to point out that they may be applied to other kinds of discrete variables as well. Such variables may represent, for instance, count data or discrete points in time. As an example of the former, we can think of participants solving a sequence of figural analogy tasks with the value of interest being the number of tasks solved correctly. This count variable could then be used as a predictor for a general intelligence score. It is plausible to assume the number of correctly solved items to be monotonically related to general intelligence and so the application of a monotonic effect appears reasonable. As an example of the latter, we could think of a longitudinal study with few measurement points. If the outcome were a skill gradually acquired over time, we would expect time to be monotonically related to it. Of course, time may also be modelled as continuous, but, for very few time points, using a monotonic effect may be a more reliable solution without strong assumptions outside of monotonicity.

For simple cases such as regression models with only a single monotonic effect and normally distributed errors, maximum likelihood estimators can be developed as well (Barlow *et al.*, 1972; Robertson *et al.*, 1988). As we prefer a fully Bayesian approach to statistical modelling, we did not pursue this line of investigation more deeply. However, we still believe that developing frequentist estimators and corresponding uncertainty estimates for more complex monotonic models including, for instance, interactions or multilevel structure, may be a worthwhile endeavour for future research. Finally, we wish to note that the general idea of monotonic effects should also generalize to continuous data. In this case, the sum in the definition of monotonic effects becomes an integral and ζ a non-negative function to be integrated over. A similar idea is used in I-splines (i.e., integral splines) whose basis functions represent integrals over the non-negative basis functions of another spline (Ramsay, 1988). In future research, it may thus be worthwhile to study continuous versions of monotonic effects and to relate them to existing methods (e.g., Pya & Wood, 2015; Ramsay, 1988) that ensure monotonicity in the continuous case.

Acknowledgements

We would like to thank Marie Beisemann, Andrew Rutherford, and three anonymous reviewers for valuable comments on earlier versions of this paper.

References

- Agresti, A. (2010). *Analysis of ordinal categorical data*. Chichester, UK: John Wiley & Sons.
- Alvarez, R. M., Bailey, D., & Katz, J. N. (2011). An empirical Bayes approach to estimating ordinal treatment effects. *Political Analysis*, 19(1), 20–31. <https://doi.org/10.1093/pan/mpq033>
- Balakrishnan, N. (2014). *Continuous multivariate distributions*. Wiley StatsRef: Statistics Reference Online. <https://doi.org/10.1002/9781118445112.stat01249>
- Barlow, R. E., Bremner, J. M., Brunk, H. D., & Bartholomew, D. J. (1972). *Statistical inference under order restrictions: The theory and application of isotonic regression*. London, UK: John Wiley & Sons.
- Bates, D., Mächler, M., Bolker, B., & Walker, S. (2015). Fitting linear mixed-effects models using lme4. *Journal of Statistical Software*, 67(1), 1–48. <https://doi.org/10.18637/jss.v067.i01>
- Best, M. J., & Chakravarti, N. (1990). Active set algorithms for isotonic regression: A unifying framework. *Mathematical Programming*, 47(1–3), 425–439. <https://doi.org/10.1007/BF01580873>
- Betancourt, M., Byrne, S., Livingstone, S., & Girolami, M. (2014). *The geometric foundations of Hamiltonian Monte Carlo*, Preprint, arXiv:1410.5110.
- Bürkner, P.-C. (2017). brms: An R package for Bayesian multilevel models using Stan. *Journal of Statistical Software*, 80(1), 1–28. <https://doi.org/10.18637/jss.v080.i01>
- Bürkner, P.-C. (2018). Advanced Bayesian multilevel modeling with the R package brms. *R Journal*, 10(1), 395–411. <https://doi.org/10.32614/RJ-2018-017>
- Bürkner, P. C., & Vuorre, M. (2019). Ordinal regression models in psychology: A tutorial. *Advances in Methods and Practices in Psychological Science*. Advance online publication. <https://doi.org/10.1177/2515245918823199>
- Carpenter, B., Gelman, A., Hoffman, M., Lee, D., Goodrich, B., Betancourt, M., . . . Ridell, A. (2017). Stan: A probabilistic programming language. *Journal of Statistical Software*, 76(1), 1–32. <https://doi.org/10.18637/jss.v076.i01>
- Carvalho, C. M., Polson, N. G., & Scott, J. G. (2009). Handling sparsity via the horseshoe. In D. A. Van Dyk, & M. Welling (Eds.), *Proceedings of the Twelfth International Conference on Artificial Intelligence and Statistics (JMLR Proceedings 5*, pp. 73–80).
- Chambers, J. M., & Hastie, T. J. (1992). *Statistical models in S*. Pacific Grove, CA: Wadsworth & Brooks/Cole Advanced Books & Software.
- Christensen, R. H. B. (2018). *ordinal – Regression models for ordinal data*. Retrieved from <http://www.cran.r-project.org/package=ordinal/>
- Cieza, A., Stucki, G., Weigl, M., Kullmann, L., Stoll, T., Kamen, L., . . . Walsh, N. (2004). ICF core sets for chronic widespread pain. *Journal of Rehabilitation Medicine*, 36(1), 63–68. <https://doi.org/10.1080/16501960410016046>
- Cook, S. R., Gelman, A., & Rubin, D. B. (2006). Validation of software for Bayesian models using posterior quantiles. *Journal of Computational and Graphical Statistics*, 15, 675–692. <https://doi.org/10.1198/106186006X136976>
- Danaher, M. R., Roy, A., Chen, Z., Mumford, S. L., & Schisterman, E. F. (2012). Minkowski-Weyl priors for models with parameter constraints: An analysis of the biocycle study. *Journal of the American Statistical Association*, 107, 1395–1409. <https://doi.org/10.1080/01621459.2012.712414>
- Dykstra, R. L., & Robertson, T. (1982). An algorithm for isotonic regression for two or more independent variables. *Annals of Statistics*, 10, 708–716. <https://doi.org/10.1080/01621459.2012.712414>
- Finney, S. J., & DiStefano, C. (2006). Nonnormal and categorical data in structural equation modeling. In G. R. Hancock & R. O. Mueller (Eds.), *Structural equation modeling: A second course*. Greenwich, CT: Information Age Publishing.

- Frigyik, B. A., Kapila, A., & Gupta, M. R. (2010). *Introduction to the Dirichlet distribution and related processes (UWEE Tech Report Series UWEETR-2010-0006)*. Seattle, WA: Department of Electrical Engineering, University of Washington.
- Gelman, A., Carlin, J. B., Stern, H. S., Dunson, D. B., Vehtari, A., & Rubin, D. B. (2013). *Bayesian data analysis* (3rd ed.). Boca Raton, FL: Chapman and Hall/CRC.
- Gelman, A., Simpson, D., & Betancourt, M. (2017). The prior can often only be understood in the context of the likelihood. *Entropy*, 19, 555–567. <https://doi.org/10.3390/e19100555>
- Gertheiss, J. (2014). ANOVA for factors with ordered levels. *Journal of Agricultural, Biological, and Environmental Statistics*, 19(2), 258–277. <https://doi.org/10.1080/01621459.2012.712414>
- Gertheiss, J. (2015). *ordPens: Selection and/or smoothing of ordinal predictors*. Retrieved from <https://CRAN.R-project.org/package=ordPens>
- Gertheiss, J., Hogger, S., Oberhauser, C., & Tutz, G. (2011). Selection of ordinally scaled independent variables with applications to international classification of functioning core sets. *Applied Statistics*, 60(3), 377–395. <https://doi.org/10.1111/j.1467-9876.2010.00753.x>
- Gertheiss, J., & Oehrlin, F. (2011). Testing linearity and relevance of ordinal predictors. *Electronic Journal of Statistics*, 5, 1935–1959. <https://doi.org/10.1214/11-EJS661>
- Gertheiss, J., & Tutz, G. (2009). Penalized regression with ordinal predictors. *International Statistical Review*, 77(3), 345–365. <https://doi.org/10.1111/j.1751-5823.2009.00088.x>
- Gu, C. (2013). *Smoothing spline ANOVA models*. New York, NY: Springer.
- Hastie, T. J. (2017). Generalized additive models. In T. J. Hastie (Ed.), *Statistical models in S* (pp. 249–307). Abingdon, UK: Routledge.
- He, X., & Shi, P. (1998). Monotone B-spline smoothing. *Journal of the American Statistical Association*, 93, 643–650. <https://doi.org/10.2307/2670115>
- Helwig, N. E. (2016). Efficient estimation of variance components in nonparametric mixed-effects models with large samples. *Statistics and Computing*, 26, 1319–1336. <https://doi.org/10.1007/s11222-015-9610-5>
- Helwig, N. E. (2017). Regression with ordered predictors via ordinal smoothing splines. *Frontiers in Applied Mathematics and Statistics*, 3, 15. <https://doi.org/10.3389/fams.2017.00015>
- Helwig, N. E. (2018). *Bigsplines: Smoothing splines for large samples*. Retrieved from <https://CRAN.R-project.org/package=bigsplines>
- Hoffman, M., & Gelman, A. (2014). The No-U-Turn Sampler: Adaptively setting path lengths in Hamiltonian Monte Carlo. *Journal of Machine Learning Research*, 15(1), 1593–1623.
- Hoijtink, H., Klugkist, I., & Boelen, P. (2008). *Bayesian evaluation of informative hypotheses*. New York, NY: Springer.
- Jöreskog, K. G. (1994). Structural equation modeling with ordinal variables. In T. W. Anderson, K. T. Fang & I. Olkin (Eds.), *Multivariate analysis and its applications (IMS Lecture Notes-Monograph Series (Vol. 24, pp. 297–310)*. Hayward, CA: Institute of Mathematical Statistics.
- Kass, R. E., & Raftery, A. E. (1995). Bayes factors. *Journal of the American Statistical Association*, 90, 773–795. <https://doi.org/10.1080/01621459.1995.10476572>
- Kelly, C., & Rice, J. (1990). Monotone smoothing with application to dose-response curves and the assessment of synergism. *Biometrics*, 46, 1071–1085. <https://doi.org/10.2307/2532449>
- Klugkist, I., & Mulder, J. (2008). Bayesian estimation for inequality constrained analysis of variance. In H. Hoijtink, I. Klugkist & P. A. Boelen (Eds.), *Bayesian evaluation of informative hypotheses* (pp. 27–52). New York, NY: Springer.
- Kruschke, J. K. (2014). *Doing Bayesian data analysis: A tutorial introduction with R* (2nd ed.). Boston, MA: Academic Press.
- Lee, C.-I. C. (1981). The quadratic loss of isotonic regression under normality. *Annals of Statistics*, 9, 686–688. <https://doi.org/10.2307/2532449>
- Lee, C.-I. C. (1996). On estimation for monotone dose-response curves. *Journal of the American Statistical Association*, 91, 1110–1119. <https://doi.org/10.2307/2291730>
- Lei, P.-W. (2009). Evaluating estimation methods for ordinal data in structural equation modeling. *Quality and Quantity*, 43(3), 495. <https://doi.org/10.1007/s11135-007-9133-z>

- Leitenstorfer, F., & Tutz, G. (2006). Generalized monotonic regression based on B-splines with an application to air pollution data. *Biostatistics*, 8, 654–673. <https://doi.org/10.1093/biostatistics/kxl036>
- Liddell, T., & Kruschke, J. K. (2017). *Analyzing ordinal data with metric models: What could possibly go wrong?* Preprint, Open Science Framework. <https://doi.org/10.17605/OSF.IO/9H3ET>
- Liu, I., & Agresti, A. (2005). The analysis of ordered categorical data: An overview and a survey of recent developments. *Test*, 14(1), 1–73. <https://doi.org/10.1007/BF02595397>
- McCullagh, P. (1980). Regression models for ordinal data. *Journal of the Royal Statistical Society, Series B*, 42(2), 109–142. <https://doi.org/10.1111/j.2517-6161.1980.tb01109.x>
- McElreath, R. (2016). *Statistical rethinking: A Bayesian course with examples in R and Stan*. Boca Raton, FL: CRC Press.
- Mulder, J., & Raftery, A. E. (2019). BIC extensions for order-constrained model selection. *Sociological Methods and Research*. Advance online publication. <https://doi.org/10.1177/0049124119882459>
- Piironen, J., & Vehtari, A. (2017). Sparsity information and regularization in the horseshoe and other shrinkage priors. *Electronic Journal of Statistics*, 11, 5018–5051. <https://doi.org/10.1214/17-EJS1337SI>
- Pya, N., & Wood, S. N. (2015). Shape constrained additive models. *Statistics and Computing*, 25, 543–559. <https://doi.org/10.1007/s11222-013-9448-7>
- R Core Team. (2018). *R: A language and environment for statistical computing*. Vienna, Austria: R Foundation for Statistical Computing.
- Ramsay, J. O. (1988). Monotone regression splines in action. *Statistical Science*, 3(4), 425–441. <https://doi.org/10.1214/ss/1177012761>
- Robertson, T., Wright, F. T., & Dykstra, R. L. (1988). *Order restricted statistical inference*. Chichester, UK: John Wiley & Sons.
- RStudio Team. (2018). *RStudio: Integrated development for R [Computer software]*. Boston, MA: RStudio.
- Rufibach, K. (2010). An active set algorithm to estimate parameters in generalized linear models with ordered predictors. *Computational Statistics and Data Analysis*, 54, 1442–1456. <https://doi.org/10.1016/j.csda.2010.01.014>
- Stan Development Team. (2019). *Stan modeling language: User's guide and reference manual*. Retrieved from <http://mc-stan.org/manual.html>
- Talts, S., Betancourt, M., Simpson, D., Vehtari, A., & Gelman, A. (2018). *Validating Bayesian inference algorithms with simulation-based calibration*. Preprint, arXiv:1804.06788.
- Tutz, G. (2011). *Regression for categorical data*. Cambridge, UK: Cambridge University Press.
- Vehtari, A., Gelman, A., & Gabry, J. (2017). Practical Bayesian model evaluation using leave-one-out cross-validation and WAIC. *Statistics and Computing*, 27, 1413–1432. <https://doi.org/10.1007/s11222-016-9696-4>
- Wagenmakers, E.-J., & Farrell, S. (2004). AIC model selection using Akaike weights. *Psychonomic Bulletin and Review*, 11(1), 192–196. <https://doi.org/10.3758/BF03206482>
- Wang, W., & Small, D. S. (2015). Monotone B-spline smoothing for a generalized linear model response. *American Statistician*, 69(1), 28–33. <https://doi.org/10.1080/00031305.2014.969445>
- Ware, J. E., & Sherbourne, C. D. (1992). The MOS 36-item short-form health survey (SF-36): I. Conceptual framework and item selection. *Medical Care*, 30, 473–483.
- Watanabe, S. (2010). Asymptotic equivalence of Bayes cross validation and widely applicable information criterion in singular learning theory. *Journal of Machine Learning Research*, 11, 3571–3594.
- Wickham, H. (2016). *ggplot2: Elegant graphics for data analysis* (2nd ed.). Cham, Switzerland: Springer.
- Wickham, H. (2017). *Tidyverse: Easily install and load the 'tidyverse'*. Retrieved from <https://CRAN.R-project.org/package=tidyverse>
- Wickham, H., François, R., Henry, L., & Müller, K. (2019). *Dplyr: A grammar of data manipulation*. Retrieved from <https://CRAN.R-project.org/package=dplyr>

- Winship, C., & Mare, R. D. (1984). Regression models with ordinal variables. *American Sociological Review*, 49(4), 512–525. <https://doi.org/10.2307/2095465>
- World Health Organization (2001). *International classification of functioning disability and health: ICF*. Geneva, Switzerland: World Health Organization.
- Wu, W. B., Woodroffe, M., & Mentz, G. (2001). Isotonic regression: Another look at the changepoint problem. *Biometrika*, 88, 793–804. <https://doi.org/10.1093/biomet/88.3.793>
- Yao, Y., Vehtari, A., Simpson, D., & Gelman, A. (2017). *Using stacking to average Bayesian predictive distributions*. Preprint, arXiv:1704.02030.
- Yee, T. W., Stoklosa, J., & Huggins, R. M. (2015). The VGAM package for capture–recapture data using the conditional likelihood. *Journal of Statistical Software*, 65, 1–33. <https://doi.org/10.18637/jss.v065.i05>

Received 17 July 2019; revised version received 2 October 2019

Appendix A: Mathematical proofs

A.1. Monotonicity

For all values x between 0 and $D-1$, we have

$$bmo(x+1, \zeta) - bmo(x, \zeta) = bD \sum_{i=1}^{x+1} \zeta_i - bD \sum_{i=1}^x \zeta_i = bD \zeta_{x+1}. \quad (A1)$$

Since $D > 0$ and $\zeta_{x+1} > 0$, the linear predictor $\eta(x)$ is monotonically increasing if $b \geq 0$ and monotonically decreasing if $b \leq 0$.

A.2. Equivalence to categorical isotonic regression

Consider a simple linear model of a continuous response y regressed on a categorical predictor x with categories $j \in \{0, 1, \dots, D\}$. Further, let μ_j be the group mean of category j with respect to the response variable. Then the model for observation n can be written as

$$y_n = \mu_{x_n} + e_n, \quad (A2)$$

where the e_n are errors of the regression. In categorical isotonic regression, we estimate $\mu = (\mu_0, 1, \dots, \mu_C)$ under the order-constraint $\mu_0 \leq \mu_1 \leq \dots \leq \mu_C$ or $\mu_0 \geq \mu_1 \geq \dots \geq \mu_C$. Using a monotonic effect, we write

$$y_n = b_0 + b_1 D \sum_{i=1}^{x_n} \zeta_i + e_n. \quad (A3)$$

Hence, we can identify μ_0 with b_0 and μ_j with $b_0 + b_1 D \sum_{i=1}^j \zeta_i$ for $j > 0$. This identification is bijective within the set of order-constraint μ .

A.3. Proof of proposition 1

Under the stated assumptions, we can, without loss of generality, write the linear predictor $\eta = \eta(x)$ as

$$\eta(x) = b_0 + \sum_{k=1}^K b_k D_k \sum_{i=1}^x \zeta_i = b_0 + \left(\sum_{k=1}^K b_k D_k \right) \left(\sum_{i=1}^x \zeta_i \right). \quad (A4)$$

Since all other predictors have been fixed to some constants, their contribution to η can be absorbed by the intercept b_0 and the regression coefficients b_1, \dots, b_K which are all

related to x . If we define $b = \sum_{i=1}^K b_i D_i$ we see that $\eta(x)$ is monotonic in x with the sign of the effect determined by the sign of b .

A.4. Counter-example to conditional monotonicity for varying simplex parameters

Consider the situation shown in Figure A1, where quite clearly the effect of \mathbf{x} is monotonic for group G , but non-monotonic for group H . Suppose further that we named the grouping variable \mathbf{z} and applied dummy coding such that $G = 0$ and $H = 1$. Using different simplex parameters for the main effect of \mathbf{x} and the interaction effects between \mathbf{x} and \mathbf{z} , the linear predictor reads as

$$\eta(x, z) = b_0 + b_1 z + b_2 \text{mo}(x, \zeta_2) + b_3 z \text{mo}(x, \zeta_3). \quad (\text{A5})$$

Clearly, $b_0 = 0$. For group G this implies $\eta(x, 0) = b_2 \text{mo}(x, \zeta_2)$ so that $b_2 = 50$ as well as $\zeta_2 = (.8, .2)$ are completely defined by the curve of group G . For group H , we have

$$\eta(x, 1) = b_0 + b_1 + b_2 \text{mo}(x, \zeta_2) + b_3 \text{mo}(x, \zeta_3). \quad (\text{A6})$$

As the curve of group H starts at the origin, we have $b_1 = 0$. Due to the chosen parameterization of \mathbf{z} , the term $b_3 \text{mo}(x, \zeta_3)$ models the *difference* between in the effect of \mathbf{x} between the two groups, which is visualized as a dashed line in Figure A1 and is clearly monotonic. Consequently, we have $b_3 = 30$ and $\zeta_3 = (\frac{1}{6}, \frac{5}{6})$. Although the assumptions of the monotonic effects are fully met, the effect of \mathbf{x} in group H is non-monotonic. Thus, \mathbf{x} is not conditionally monotonic given \mathbf{z} .




Figure A1. Counter-example to the conditional monotonicity for varying simplex parameters. The dashed line shows the difference between the groups G and H as a function of \mathbf{x} .

Appendix B: Additional simulation results




Figure B1. Simulation results for the main effects models under true non-monotonicity based on $T = 1,000$ simulation trials. LIN is not displayed as its RMSE is too large and thus obscures differences between other models. Abbreviations: CAT = categorical model; CS = cubic spline model; D = number of categories minus 1; ISO = isotonic regression model; LS = linear spline model; MO = monotonic model; N = number of observations; OS = ordinal spline model; OSMO = ordinal monotonic spline model.




Figure B2. Simulation results for the interaction models under true non-monotonicity based on $T = 1,000$ simulation trials. LIN is not displayed as its RMSE is too large and thus obscures differences between other models. For the same reason, CAT is not displayed for $D = 50$. ISO and OSMO are not displayed as they have no corresponding interaction model. Abbreviations: CAT categorical model; CS = cubic spline model; D = number of categories minus 1; LS = linear spline model; MO = monotonic model; N = number of observations; OS = ordinal spline model.