

MATH 115AH: Week 1 (September 25 – 30, 2026)

Lecture notes for the first week of MATH 115AH (Linear Algebra, honors), Fall 2026: vector spaces and matrices, subspaces, fields and subfields, linear combinations and span, linear independence, bases, the Replacement Theorem and dimension, and a first look at linear maps.

1 September 25, 2026: Vector Spaces

1.1 The Example of $\mathbb{R}^2$

Example 1.1. $\mathbb{R}^2 = \left\{ \begin{bmatrix} x \\ y \end{bmatrix} : x \in \mathbb{R}, y \in \mathbb{R} \right\}$, with

$$\begin{bmatrix} x_1 \\ y_1 \end{bmatrix} + \begin{bmatrix} x_2 \\ y_2 \end{bmatrix} = \begin{bmatrix} x_1 + x_2 \\ y_1 + y_2 \end{bmatrix} \quad (1)$$

and, for $a \in \mathbb{R}$,

$$a \begin{bmatrix} x \\ y \end{bmatrix} = \begin{bmatrix} ax \\ ay \end{bmatrix}. \quad (2)$$

[Sketch: the vector (x, y) drawn as an arrow from the origin in the plane, with its negative $(-x, -y)$.]

1.2 Definition of a Vector Space

Definition 1.2. A *vector space* (linear space) V over F ($\mathbb{R}$ or $\mathbb{C}$) is a set with 2 operations, addition and scalar multiplication, $(V, +, \cdot; 0)$, with a distinguished element 0 , such that $\forall x, y \in V$, $\exists! x + y \in V$, and $\forall a \in F, x \in V, \exists! ax \in V$, satisfying

- (i) $\forall x, y \in V, x + y = y + x$;
- (ii) $\forall x, y, z \in V, (x + y) + z = x + (y + z)$;
- (iii) $\forall x \in V, x + 0 = 0 + x = x$;
- (iv) $\forall x \in V, \exists! y \in V$ s.t. $x + y = 0$;
- (v) $\forall x \in V, 1 \cdot x = x$ ($1 \in F$);
- (vi) $\forall a, b \in F, x \in V, a(bx) = (ab)x$;
- (vii) $\forall a \in F, x, y \in V, a(x + y) = ax + ay$;
- (viii) $\forall a, b \in F, x \in V, (a + b)x = ax + bx$.

Example 1.3. $\mathbb{R}$ is a vector space over $\mathbb{R}$. $\mathbb{C}$ is a vector space over $\mathbb{C}$.

Elements in V are *vectors*; elements in F are *scalars*.

Example 1.4. F^n , $n \geq 1$ integer:

$$F^n = \left\{ \begin{pmatrix} x_1 \\ \vdots \\ x_n \end{pmatrix} : x_1, \dots, x_n \in F \right\}. \quad (3)$$

For $\vec{x}, \vec{y} \in F^n$ and $\forall a \in F$,

$$\vec{x} + \vec{y} = \begin{pmatrix} x_1 + y_1 \\ \vdots \\ x_n + y_n \end{pmatrix}, \quad a\vec{x} = \begin{pmatrix} ax_1 \\ \vdots \\ ax_n \end{pmatrix}. \quad (4)$$

Remark. “0 and negation are unique.”

1.3 Matrices

Example 1.5 (Matrices). Let $m, n \geq 1$ be integers. An $m \times n$ matrix with entries $\in F$ is

$$\begin{bmatrix} a_{11} & \cdots & a_{1n} \\ \vdots & \ddots & \vdots \\ a_{m1} & \cdots & a_{mn} \end{bmatrix}. \quad (5)$$

An $n \times 1$ matrix is a column vector. For example,

$$\begin{bmatrix} 2 + i & 2 \\ 3 + 2\sqrt{3} & 4 \end{bmatrix} \quad (6)$$

is 2×2 in $\mathbb{C}$. Then $M_{m \times n}(F) := \{m \times n \text{ matrices}/F\}$ is a vector space, with addition

$$A + B = \begin{bmatrix} a_{11} + b_{11} & \cdots & a_{1n} + b_{1n} \\ \vdots & \ddots & \vdots \\ a_{m1} + b_{m1} & \cdots & a_{mn} + b_{mn} \end{bmatrix} \quad (7)$$

and scalar multiplication

$$c \cdot A = \begin{bmatrix} ca_{11} & \cdots & ca_{1n} \\ \vdots & \ddots & \vdots \\ ca_{m1} & \cdots & ca_{mn} \end{bmatrix}. \quad (8)$$

How to relate to equations: from

$$\begin{cases} ax + by = c \\ dx + ey = f \end{cases} \quad (9)$$

we can get

$$\begin{bmatrix} a & b \\ d & e \end{bmatrix} \begin{bmatrix} x \\ y \end{bmatrix} = \begin{bmatrix} c \\ f \end{bmatrix}. \quad (10)$$

1.4 Cancellation Law

Theorem 1.6 (Cancellation law for addition). *Assume $x, y, z \in V$. If $x + z = y + z$, then $x = y$.*

Proof. For $z \in V$, there exists $v \in V$ s.t. $z + v = 0$. Then,

$$(x + z) + v = (y + z) + v \quad (11)$$

$$x + (z + v) = y + (z + v) \quad (12)$$

$$x = y. \quad (13)$$

□

1.5 Properties of Zero and Negatives

Theorem 1.7. *For the number $0 \in F$,*

1. $0 \cdot v = 0, \forall v \in V$;
2. $(-a)x = -(ax) = a(-x), \forall a \in F, \forall x \in V$;
3. $a \cdot 0 = 0, \forall a \in F$.

Proof. (1) By the distributive law,

$$0 \cdot v + 0 \cdot v = (0 + 0)v = 0v \quad (14)$$

hence $0 \cdot v = 0$ (by cancellation).

(2) Using the distributive law, then (1),

$$(-a)x + ax = (-a + a)x = 0x = 0, \quad (15)$$

and using (3),

$$a(-x) + ax = a(-x + x) = a \cdot 0 = 0. \quad (16)$$

(3)

$$a \cdot 0 + a \cdot 0 = 0(a + a) = 0 \cdot 2a = 0a = 0. \quad (17)$$

□

1.6 Back to F

Back to F : $\in \mathbb{R}$ or $\mathbb{C}$; also $\mathbb{Q}$, and $\mathbb{Z}/p\mathbb{Z}$. For $\mathbb{Z}/p\mathbb{Z}$: $\forall a \in \mathbb{Z}, \exists$

$$a \cdot u + p \cdot v = 1 \quad (18)$$

for some $u, v \in \mathbb{Z}$ (since $(a, p) = 1$). Then, mod p ,

$$\bar{a} \cdot \bar{u} = \bar{1}. \quad (19)$$

Also,

$$\underbrace{\bar{1} + \cdots + \bar{1}}_p = \bar{0}. \quad (20)$$

2 September 28, 2026: Subspaces

2.1 Recall

Recall $(V, +, \cdot)$ over $F = \mathbb{R}, \mathbb{C}$. V is a set, $0 \in V$,

$$+ : V \times V \rightarrow V \quad (21)$$

$$\cdot : F \times V \rightarrow V \quad (22)$$

s.t. $\forall x, y \in V: x + y = y + x; (x + y) + z = x + (y + z); 0 + x = x = x + 0; \exists! x' \in V$ s.t. $x + x' = 0$;
and $\forall a, b \in F, x, y \in V$,

$$(a + b)(x + y) = a \cdot x + a \cdot y + b \cdot x + b \cdot y. \quad (23)$$

2.2 Subspaces and the Subspace Test

Definition 2.1. A subset W of V is called a *subspace* if W itself is a vector space over F with respect to $+|_W, \cdot|_W$.

Remark (Digest). Let $(V, +, \cdot; 0)$ and $W \subseteq V$. Then W is a subspace of V if

1. $\bar{0} \in W$;
2. W is closed under addition;
3. W is closed under scalar multiplication.

Example 2.2. $V = \mathbb{R}^2 = \left\{ \begin{bmatrix} x_1 \\ x_2 \end{bmatrix} : x_1, x_2 \in \mathbb{R} \right\}$. Then $W = \left\{ \begin{bmatrix} x_1 \\ 0 \end{bmatrix} : x_1 \in \mathbb{R} \right\}$ is a subspace ($\checkmark$),
but $W = \left\{ \begin{bmatrix} x_1 \\ 1 \end{bmatrix} : x_1 \in \mathbb{R} \right\}$ is not ($\times$).

Proof. ($\Downarrow$) Since W is a vector space over F w.r.t. $+|_W$ and $\cdot|_W$, conditions 2 and 3 are given true, and $\exists \bar{0}_W \in W$ s.t.

$$\bar{0}_W + x = x, \quad \forall x \in W. \quad (24)$$

But

$$\bar{0} + x = x \quad \forall x \in W \subseteq V, \quad (25)$$

so

$$\bar{0} + x = x = \bar{0}_W + x \quad (26)$$

$$\bar{0} = \bar{0}_W. \quad (27)$$

($\uparrow$) $\forall x, y \in W, a, b \in F$:

$$x + y = y + x, \quad (x + y) + z = x + (y + z) \quad (28)$$

$$x + \bar{0} = x \quad (29)$$

$$\exists (-x) \in V \text{ s.t. } x + (-x) = \bar{0}, \quad -x = -1 \cdot x \in W \text{ (by 3)} \quad (30)$$

$$1 \cdot x = x, \quad (a + b)(x + y) = ax + ay + bx + by \quad (31)$$

$$(ab)x = a(bx). \quad (32)$$

Therefore $\forall a, b \in F, x, y \in W \Rightarrow ax + by \in W$. $\square$

Example 2.3.

$$\mathbb{R}[x] = \{ a_0 + a_1x + \cdots + a_nx^n : a_0, \dots, a_n \in \mathbb{R} \} \quad (33)$$

$$\mathbb{R}[x]_{<m} = \{ \text{polynomials of degree } < m \} \quad (34)$$

2.3 Transpose and Trace Examples

Example 2.4. $V = M_{m \times n}(F)$. Define the *transpose* of A to be

$$A^t = \begin{bmatrix} a_{11} & \cdots & a_{m1} \\ \vdots & \ddots & \vdots \\ a_{1n} & \cdots & a_{mn} \end{bmatrix}. \quad (35)$$

(i) $m = n$:

$$W_1 = \{ A \in M_{m \times n}(F) : A = A^t \} \quad (36)$$

$$W_2 = \{ A \in M_{m \times n}(F) : a_{ij} = 0, \forall i > j \} \quad (37)$$

$$W_3 = W_2^t \quad (38)$$

$$W_4 = \{ \text{diagonal matrices} \} \quad (39)$$

$$W_4 = W_1 \cap W_2 \text{ and } W_1 \cap W_3 \text{ and } W_2 \cap W_3 \quad (40)$$

[Sketch: W_2 drawn as a square matrix with entries on and above the diagonal and 0 below it.]

For $m = n$: $\text{tr}(A) = a_{11} + a_{22} + \cdots + a_{nn}$, and

$$W_5 = \{ A \in M_{n \times n}(F) : \text{tr}(A) = 0 \}. \quad (41)$$

2.4 Intersections of Subspaces

Theorem 2.5. *Any intersection of subspaces of V is a subspace of V .*

Proof. Let $\mathcal{C} = \{ \text{subspaces of } V \}$ and

$$W_0 := \bigcap_{W \in \mathcal{C}} W. \quad (42)$$

$\bar{0} \in W$ for any $W \in \mathcal{C}$, so $\bar{0} \in \bigcap_{W \in \mathcal{C}} W = W_0$. $\forall a, b \in F, \forall x, y \in W_0 \subseteq W, ax + by \in W$ for any $W \in \mathcal{C}$, hence

$$ax + by \in \bigcap_{W \in \mathcal{C}} W = W_0. \quad (43)$$

□

Remark. The union of a subspace is only a subspace if one subspace is contained in the other.

3 September 29, 2026: Fields and Subfields

3.1 Basic Properties of Fields

Proposition 3.1. *Let F be a field. $\forall a \in F, 0 \cdot a = 0$.*

Proof. First,

$$0 \cdot a + a = (0 + 1) \cdot a = 1 \cdot a = a \quad (44)$$

then,

$$0 = a + (-a) = (0 \cdot a + a) + (-a) \quad (45)$$

$$= 0 \cdot a + (a + (-a)) \quad (46)$$

$$= 0 \cdot a + 0 = 0 \cdot a. \quad (47)$$

□

Proposition 3.2. *Let F be a field. If $a, b \in F$ satisfy $a \cdot b = 0$, then $a = 0$ or $b = 0$.*

Proof. Assume $a \neq 0$. Then

$$b = 1 \cdot b = (a^{-1} \cdot a) \cdot b = a^{-1} \cdot (a \cdot b) \quad (48)$$

$$= a^{-1} \cdot 0 = 0. \quad (49)$$

□

3.2 Subfields

Definition 3.3. Let F be a field. A subset $E \subseteq F$ is called a *subfield* if

- (i) $0, 1 \in E$;
- (ii) $\forall a, b \in E, a + b \in E, a \cdot b \in E$;
- (iii) $\forall a \in E, -a \in E$;
- (iv) $\forall a \in E, a \neq 0, a^{-1} \in E$.

Let E be a subfield of F . By (ii) we get operations $+, \cdot : E \times E \rightarrow E$.

Exercise: show these operations make E into a field.

Example 3.4. $\mathbb{Q}(\sqrt{2}) = \{ a + b\sqrt{2} : a, b \in \mathbb{Q} \} \subseteq \mathbb{R}$ is a subfield of $\mathbb{R}$.

Proof. $0, 1 \in \mathbb{Q}(\sqrt{2})$. Clear: closure under $+$, closure under additive inverse. For $a, b, c, d \in \mathbb{Q}$,

$$(a + b\sqrt{2})(c + d\sqrt{2}) = ac + 2bd + (ad + bc)\sqrt{2} \in \mathbb{Q}(\sqrt{2}). \quad (50)$$

Say $a, b \in \mathbb{Q}$ s.t. $a + b\sqrt{2} \neq 0$; then a, b are not both 0, so $a - b\sqrt{2} \neq 0$. Then,

$$\frac{1}{a + b\sqrt{2}} = \frac{a - b\sqrt{2}}{(a - b\sqrt{2})(a + b\sqrt{2})} \quad (51)$$

$$= \frac{a}{a^2 - 2b^2} + \frac{(-b)}{a^2 - 2b^2}\sqrt{2} \in \mathbb{Q}(\sqrt{2}). \quad (52)$$

□

4 September 30, 2026 (Part 1): Linear Combinations and Span

4.1 Linear Combinations

Let $(V = \text{vector space}/F, +, \cdot; \bar{0})$. Given $v_1, \dots, v_n \in V$ and $a_1, \dots, a_n \in F$, then we get

$$a_1v_1 + a_2v_2 + \dots + a_nv_n \in V. \quad (53)$$

Definition 4.1. Let S be a subset of V ($S \subseteq V$). A vector $v \in V$ is called a *linear combination of vectors in S* if $\exists u_1, \dots, u_n \in S, a_1, \dots, a_n \in F$ s.t.

$$v = a_1u_1 + \dots + a_nu_n. \quad (54)$$

Remark (Digest). If $S = \emptyset$, then no vector is a linear combination of vectors in S .

(i) Assume $S \neq \emptyset$; then $\bar{0} = 0 \cdot u$ for any $u \in S$, so $\bar{0}$ is a linear combination.

(ii) If v, v' are linear combinations of vectors $\in S$ and $c, c' \in F$, what about $cv + c'v'$?

For (ii), say

$$v = a_1u_1 + \dots + a_nu_n \quad (55)$$

$$v' = a'_1u'_1 + \dots + a'_mu'_m \quad (56)$$

then consider $\{u_1, \dots, u_n, u'_1, \dots, u'_m\} = \{u_1, \dots, u_N\}$, $N \leq n + m$. Then

$$cv + c'v' = (ca_1 + c'a'_1)u_1 + \dots + (ca_N + c'a'_N)u_N \quad (57)$$

is a linear combination of vectors in S .

4.2 Span Is a Subspace

Theorem 4.2 (Theorem/Definition). *Define $\text{Span}(S)$ as the set of all linear combinations of vectors in S . Assume $S \neq \emptyset$. Then $\text{Span}(S)$ is a subspace of V .*

5 September 30, 2026 (Part 2): Span and Linear Independence

5.1 Examples of Span

Example 5.1. In $\mathbb{R}^3$, $u_1 = (1, 0, 1)$, $u_2 = (0, 1, 1)$:

$$3u_1 - 2u_2 = (3, -2, 1) \in \text{Span}(\{u_1, u_2\}). \quad (58)$$

Example 5.2. In $\mathbb{R}^2$: [Sketch: two non-parallel vectors u_1, u_2 drawn from the origin.] $\text{Span}(\{u_1, u_2\}) = \mathbb{R}^2$.

Example 5.3. In F^n , let

$$u_1 = (x_1^{(1)}, \dots, x_n^{(1)}), \quad u_2 = (x_1^{(2)}, \dots, x_n^{(2)}), \quad u_3 = (x_1^{(3)}, \dots, x_n^{(3)}). \quad (59)$$

Given $v = (y_1, \dots, y_n) \in F^n$, how to check if $v \in \text{Span}(\{u_1, u_2, u_3\})$? Equivalently, $\exists a, b, c \in F$ s.t. $v = au_1 + bu_2 + cu_3$. Then, write

$$v = \begin{bmatrix} u_1 & u_2 & u_3 \end{bmatrix} \begin{bmatrix} a \\ b \\ c \end{bmatrix} = au_1 + bu_2 + cu_3 \quad (60)$$

$$\begin{bmatrix} x_1^{(1)} & x_1^{(2)} & x_1^{(3)} \\ \vdots & \vdots & \vdots \\ x_n^{(1)} & x_n^{(2)} & x_n^{(3)} \end{bmatrix} \begin{bmatrix} a \\ b \\ c \end{bmatrix} = \begin{bmatrix} y_1 \\ \vdots \\ y_n \end{bmatrix}. \quad (61)$$

Definition 5.4. We say that S generates V if $V = \text{span}(S)$.

Example 5.5. In $\mathbb{R}^2$, $\{u_1, u_2\}$ generates $\mathbb{R}^2$. [Sketch: u_1 along the horizontal axis and u_2 along the vertical axis.]

Example 5.6. In F^n , let

$$e_1 = (1, 0, \dots, 0) \quad (62)$$

$$e_2 = (0, 1, \dots, 0) \quad (63)$$

$$e_n = (0, 0, \dots, 1). \quad (64)$$

Then $F^n = \text{span}(\{e_1, e_2, \dots, e_n\})$, since any $v = (x_1, \dots, x_n) \in V$ is $x_1e_1 + \dots + x_ne_n$.

5.2 Linear Independence

Definition 5.7 (“redundancy”). A subset $S \subseteq V$ is called *linearly independent* if $\forall u_1, \dots, u_n \in S$ and $a_1, \dots, a_n \in F$, if we have

$$a_1u_1 + \dots + a_nu_n = 0 \quad (65)$$

then $a_1 = \cdots = a_n = 0$.

S is *linearly dependent* if $\exists u'_1, \dots, u'_n \in S$ and $a'_1, \dots, a'_n \in F$ not all 0 s.t. $a'_1 u'_1 + \cdots + a'_n u'_n = 0$.

Example 5.8. In F^n , $S = \{e_1, e_2, \dots, e_n\}$. If $a_1, \dots, a_n \in F$ s.t. $a_1 e_1 + \cdots + a_n e_n = 0$, then $(a_1, \dots, a_n) = (0, \dots, 0)$, so $a_1 = \cdots = a_n = 0$: l.i.

Example 5.9. In $\mathbb{R}^3$, $S = \{e_1, e_2, e_1 + 2e_3\}$. If $a_1, a_2, a_3 \in \mathbb{R}$ s.t.

$$a_1 e_1 + a_2 e_2 + a_3 (e_1 + 2e_3) = 0 \quad (66)$$

$$(a_1, 0, 0) + (0, a_2, 0) + (a_3, 0, 2a_3) = 0 \quad (67)$$

$$(a_1 + a_3, a_2, 2a_3) = (0, 0, 0) \quad (68)$$

hence $a_1 = a_2 = a_3 = 0$: l.i. Or,

$$(a_1 + a_3)e_1 + a_2 e_2 + 2a_3 e_3 = 0 \quad (69)$$

$$a_1 + a_3 = a_2 = 2a_3 = 0 \quad (70)$$

hence $a_1 = a_2 = a_3 = 0$.

HW: V ; $S = \{v_1, v_2, v_3\}$ l.i. Prove: $v_1, v_2, v_1 + 2v_3$ l.i.

6 September 30, 2026 (Part 3): Bases

6.1 Proof of the HW Statement

Proof. If $a_1 v_1 + a_2 v_2 + a_3 (v_1 + 2v_3) = 0$, then

$$(a_1 + a_3)v_1 + a_2 v_2 + 2a_3 v_3 = 0 \quad (71)$$

and by l.i., $a_1 + a_3 = a_2 = 2a_3 = 0$, hence $a_1 = a_2 = a_3 = 0$. $\square$

6.2 Removing a Dependent Vector

Example 6.1. In $\mathbb{R}^4$,

$$u_1 = (1, 3, -4, 2), \quad u_2 = (2, 2, -4, 0), \quad u_3 = (1, -3, 2, -4). \quad (72)$$

Then

$$4u_1 - 3u_2 + 2u_3 = 0 \quad (73)$$

$$u_3 = -\frac{1}{2}(4u_1 - 3u_2) \quad (74)$$

hence $\text{Span}(\{u_1, u_2, u_3\}) = \text{span}(\{u_1, u_2\})$.

6.3 Adding a Vector to a Linearly Independent Set

Theorem 6.2. Let $\emptyset \neq S \subseteq V$ be l.i., and $v \in V$ s.t. $v \notin S$. Then $S \cup \{v\}$ is l.i. $\iff v \notin \text{span}(S)$.

Proof. ($\Rightarrow$) Assume that $v \in \text{span}(S)$. Then $v = a_1u_1 + \cdots + a_nu_n$ for $a_1, \dots, a_n \in F$ and $u_1, \dots, u_n \in S$. Then

$$a_1u_1 + \cdots + a_nu_n + (-1) \cdot v = 0 \quad (75)$$

so $S \cup \{v\}$ is l.d., a contradiction.

($\Leftarrow$) Assume

$$a_1u_1 + \cdots + a_nu_n + a_{n+1}v = 0 \quad (76)$$

for $a_1, \dots, a_{n+1} \in F$ and $u_1, \dots, u_n \in S$. If $a_{n+1} = 0$, then $a_1u_1 + \cdots + a_nu_n = 0$, and since S is l.i., $a_1 = \cdots = a_n = 0$. If $a_{n+1} \neq 0$, then

$$v = -\frac{1}{a_{n+1}}(a_1u_1 + \cdots + a_nu_n) \in \text{span}(S), \quad (77)$$

a contradiction. □

Recall: $(V, +, \cdot)/F$, $S \subseteq V$,

$$\text{span}(S) = \{ \sum a_iu_i : u_i \in S \}, \quad \text{span}(\emptyset) = \{0\}. \quad (78)$$

$\{v_1, \dots, v_n\} \subseteq V$ is l.i. iff $\forall a_1, \dots, a_n \in F, \sum_{i=1}^n a_i v_i = 0 \Rightarrow a_1 = \cdots = a_n = 0$.

Theorem 6.3. Let $S \subseteq V$ be l.i., $v \in V$. Then $S \cup \{v\}$ is l.i. $\iff v \notin \text{span}(S)$.

6.4 Bases

Definition 6.4. A subset $\beta \subseteq V$ is called a *basis* for V if

- (i) β is l.i.;
- (ii) V is the $\text{span}(\beta)$.

Remark (Digest). In $\mathbb{R}^{\text{[illegible]}}$: $\{e_1, e_2\} \checkmark$, $\{e_1, e_3\} \checkmark$, $\{e_1, e_2, e_3\} \times$.

Example 6.5. F^n : $\{e_1, \dots, e_n\}$, the *standard basis*.

Example 6.6. $M_{m \times n}(F)$: $\{E_{ij} : 1 \leq i \leq m, 1 \leq j \leq n\}$, where E_{ij} is the $m \times n$ matrix s.t. the (i, j) entry is 1 and the rest are 0.

Example 6.7. $\{\bar{0}\}$ is NOT l.i., since $1 \cdot \bar{0} = \bar{0}$. $\emptyset$ is a basis of $\{\bar{0}\}$.

6.5 Unique Representation

Theorem 6.8. For distinct vectors $u_1, \dots, u_n \in V$, the set $\beta = \{u_1, \dots, u_n\}$ is a basis of $V \iff \forall v \in V$ there is a unique expression of v as a combination of $u_1, \dots, u_n$, i.e. $\forall v \in V, \exists! a_1, \dots, a_n \in F$ s.t.

$$v = \sum_{i=1}^n a_i u_i. \quad (79)$$

Proof. ($\Downarrow$) $V = \text{span}(\beta)$, so $v = \sum_{i=1}^n a_i u_i$ for some $a_i \in F$. For uniqueness, assume

$$v = \sum a_i u_i = \sum a'_i u_i \quad (80)$$

then

$$\sum (a_i - a'_i) u_i = \bar{0} \quad (81)$$

and by l.i., $a_i - a'_i = 0$, i.e. $a_i = a'_i$.

($\Uparrow$) $V = \text{span}(\beta)$ $\checkmark$, and $\bar{0} = 0 \cdot u_1 + \dots + 0 \cdot u_n$. Assume $a_1 u_1 + \dots + a_n u_n = \bar{0}$. Uniqueness of the expression of $\bar{0}$ implies that $a_1 = 0, a_2 = 0, \dots, a_n = 0$. So β is l.i. $\square$

6.6 A Basis from a Finite Generating Set

Theorem 6.9. If S is a finite set that generates V , a subset of S is a basis for V .

Proof. Let $S = \{u_1, \dots, u_n, \dots, u_N\}$, $V = \text{span}(S)$.

- (i) S_1 : is $\{u_1\}$ l.i., i.e. is $u_1 \neq 0$? Keep if l.i.
- (ii) S_2 : is $\{u_1, u_2\}$ l.i.? Keep u_2 if yes.
- (iii) S_3 : is $\{u_1, u_2, u_3\}$ l.i.? Keep u_3 if yes.

Continuing, we get $S_N \subseteq S$, with $\text{span}(S_1) = \text{span}(\{u_1\})$, $\text{span}(S_2) = \text{span}(\{u_1, u_2\})$, $\dots$, s.t.

$$\text{span}(S_N) = \text{span}(\{u_1, \dots, u_N\}) = \text{span}(S) = V, \quad (82)$$

and S_N is l.i. by our construction. So S_N is a basis for V . $\square$

6.7 The Replacement Theorem and Dimension

Theorem 6.10 (Replacement Theorem (*)). Let $S \subseteq V$ s.t. $V = \text{Span}(S)$, $|S| = n$. Let $L \subseteq V$ be l.i., $m = |L|$. Then $|L| \leq n$, and $\exists n - m$ vectors H s.t. $\text{span}(L \cup H) = V$.

Remark (Upshot). If V has a basis consisting of n vectors, then any basis for V has n vectors. So $n := \dim(V)$ is intrinsic.

Proof of upshot assuming the theorem. Let β be a basis for V with n vectors, and let β' be a basis for V . Applying $(*)$ to $S = \beta$ and $L = \beta'$,

$$|\beta'| \leq |\beta| = n. \quad (83)$$

Applying $(*)$ to $S = \beta'$ and $L = \beta$,

$$|\beta| \leq |\beta'| \quad (84)$$

hence $|\beta'| = |\beta|$. $\square$

Proof of $()$.* Let $L = \{v_1, \dots, v_m\}$.

Base step: $m = 0$, we can take $H = S$.

Induction: assume the theorem is proved for $|L'| = 1, \dots, m-1$ (so $m-1 \leq n$). Now for L , consider $L' = \{v_1, \dots, v_{m-1}\}$. The induction hypothesis gives $m-1 \leq n$ and $\exists H_{m-1} = \{w_1, \dots, w_{n-(m-1)}\}$ s.t. $V = \text{span}(L' \cup H_{m-1})$. So

$$v_m = \sum_{i=1}^{m-1} a_i v_i + \sum_{j=1}^{n-(m-1)} b_j w_j. \quad (\dagger)$$

If $n - (m-1) = 0$, then $v_m \in \text{span}(v_1, \dots, v_{m-1})$, a contradiction to L being l.i. So $n - (m-1) \geq 1$, hence $m \leq n$.

At least one $b_j \neq 0$ (by $(\dagger)$). Say $b_1 \neq 0$ (up to renumbering); then $(\dagger)$ implies

$$w_1 = \frac{1}{b_1} v_m - \sum_{i=1}^{m-1} \frac{a_i}{b_1} v_i - \sum_{j=2}^{n-(m-1)} \frac{b_j}{b_1} w_j. \quad (85)$$

Say we can take

$$H = \{w_2, \dots, w_{n-(m-1)}\} \quad (86)$$

with $|H| = n - m$ and $V = \text{span}(L \cup H)$. $\square$

Theorem 6.11. *Let $W \subseteq V$ be a subspace. Then $\dim(W) \leq \dim(V)$, and $\dim W = \dim V$ if and only if $W = V$.*

6.8 Linear Maps

Definition 6.12 (Maps). Let V, W be vector spaces over F . $T : V \rightarrow W$ is a *linear transformation* if

$$T(c_1x + c_2y) = c_1T(x) + c_2T(y) \quad \forall c_1, c_2 \in F, x, y \in V. \quad (87)$$

Equivalently, $T(x + y) = T(x) + T(y)$ and $T(cx) = cT(x)$.

Example 6.13. 1. $T(\bar{0}) = \bar{0}$.

2. $T : \mathbb{R}^3 \rightarrow \mathbb{R}^2, \begin{pmatrix} x_1 \\ x_2 \\ x_3 \end{pmatrix} \mapsto \begin{pmatrix} 2x_1 - x_3 \\ 4x_2 + x_3 \end{pmatrix}.$

4. $T : \mathbb{R}^2 \rightarrow \mathbb{R}^2$, rotation by θ .
5. $T : \mathbb{R}^n \rightarrow \mathbb{R}^m$, projection to the first m elements.
6. $T : M_{m \times n}(F) \rightarrow M_{n \times m}(F)$, $A \mapsto A^t$:

$$T(c_1A + c_2B) = (c_1A + c_2B)^t = (c_1A)^t + (c_2B)^t \quad (88)$$

$$= T(c_1A) + T(c_2B). \quad (89)$$

7. $T : \mathbb{R} \rightarrow \mathbb{R}^2$, $x \mapsto \begin{bmatrix} x \\ 2x \end{bmatrix}$.

Example 6.14 (*). For $A \in M_{m \times n}(\mathbb{R})$, $T_A : \mathbb{R}^n \rightarrow \mathbb{R}^m$, $x \mapsto Ax$.