

Training on *Mycobacterium tuberculosis* drug susceptibility testing (first and second line LJ DST)

Module 2: Epidemiology of Tuberculosis

Venue:

Presenter:

Date:

Module Outline

- Introduction
- Learning Objectives
- Course Content
 - () What is TB and its causative agent.
 - () Natural history of TB
 - () How is TB transmitted
 - () Risk factors associated with Tuberculosis Infection
 - () Global and national burden Tuberculosis infection
 - () Organization of TB laboratory services
- Assessment
- Summary
- References

Introduction

- Tuberculosis (TB) is a communicable disease that is a major cause of ill health and one of the leading causes of death worldwide.
- Until the coronavirus (COVID-19) pandemic, TB was the leading cause of death from a single infectious agent, ranking above HIV/AIDS.

Learning Objectives


At the end of this presentation , the learner should be able to:

- Describe the natural history of TB disease
- Understand the underlying risk factors to TB infection
- Appreciate the current WHO Global and national TB burden

What is TB

- Main cause of TB: MTB bacteria
- But, not all MTB infected people are developing to “Active TB”
- TB is an infectious disease that affects mainly the lungs (pulmonary TB, or PTB) but can also attack any part of the body (extra pulmonary TB, or EPTB)

Natural History of TB



ACTIVE TB

Active infection- person excreting tubercle bacilli in sputum, and has other signs and symptoms of TB such as cough > 2 weeks, fever, loss of weight

📖 **A “Presumptive TB Case”**- a person presenting with cough for more than 2 weeks

📖 **A “TB case”**- A person who has been diagnosed by a clinician or confirmed bacteriologically as having TB

Group Exercise


Describe 5 risk factors that can pre-dispose an individual to TB infection.

Causative agent of TB

- *Mycobacterium tuberculosis*
- *Mycobacterium bovis*

GLOBAL TB BURDEN

Estimated TB incidence rates, 2020



Countries in the three global lists of high-burden countries for TB, HIV-associated TB and MDR/RR-TB to be used by WHO in the period 2021–2025.

The red square indicates that a country is in a list.


COUNTRY	TB	TB/HIV	MDR/RR-TB
Angola	■		■
Azerbaijan			■
Bangladesh	■		■
Belarus			■
Botswana		■	
Brazil	■	■	
Cameroon		■	
Central African Republic	■	■	
China	■	■	■
Congo	■	■	
Democratic People's Republic of Korea	■		■
Democratic Republic of the Congo	■	■	■
Eswatini		■	
Ethiopia	■	■	
Gabon	■	■	
Guinea		■	
Guinea-Bissau		■	
India	■	■	■
Indonesia	■	■	■

Kazakhstan			■
Kenya	■	■	
Kyrgyzstan			■
Lesotho	■	■	
Liberia	■	■	
Malawi		■	
Mongolia	■		■
Mozambique	■	■	■
Myanmar	■	■	■
Namibia	■	■	
Nepal			■
Nigeria	■	■	■
Pakistan	■		■
Papua New Guinea	■		■
Peru			■
Philippines	■	■	■
Republic of Moldova			■
Russian Federation		■	■
Sierra Leone	■		
Somalia			■
South Africa	■	■	■
Tajikistan			■
Thailand	■	■	
Uganda	■	■	
Ukraine			■
United Republic of Tanzania	■	■	
Uzbekistan			■
Viet Nam	■		■
Zambia	■	■	■
Zimbabwe		■	■



MDR/RR TB GLOBAL SITUATION

Percentage of new TB cases with MDR/RR-TB^a




^a Percentages are based on the most recent data point for countries with representative data from 2004 to 2019. Model-based estimates for countries with data before 2004 are not shown. MDR-TB is a subset of RR-TB.




MDR/RR-TB GLOBAL BURDEN

Percentage of MDR/RR-TB cases tested for susceptibility to second-line drugs, 2018



USE OF BDQ FOR MDR/XDR-TB: GLOBAL SITUATION


Countries that used bedaquiline for the treatment of MDR/XDR-TB as part of expanded access, compassionate use or under normal programmatic conditions by the end of 2018



TB BURDEN IN UGANDA

- Estimated TB incidence (2016): 234 cases/100,000 population
- Estimated TB Prevalence (2016): 253 cases/100,000 population

TB Laboratory Network



Peripheral laboratories

- Are located within a general dispensary, clinic or hospital
- Have limited services of TB diagnosis that may include
 - ↻ Sputum specimen collection
 - ↻ Sputum-smear microscopy
 - ↻ GeneXpert MTB/RIF testing
- Should participate in External Quality Assurance (EQA) programmes

Intermediate laboratories..1

Are in regional or large hospitals

🌐 Have expanded services for TB diagnosis that may include

- Sputum specimen collection
- Sputum-smear microscopy
- GeneXpert MTB/RIF testing
- Culture and identification of *M. tuberculosis*
- LPA

Intermediate Laboratories..2

🌐 Provide support for peripheral laboratories in terms of;

- Supply of reagents and materials
- Offer training, supervision, EQA of sputum-smear microscopy and GeneXpert MTB/RIF testing.

National/Central Laboratory ..1

Central laboratories:

- Are at the country, provincial or state level
- Provide comprehensive services for TB diagnosis that may include
 - Sputum specimen collection
 - Sputum-smear microscopy
 - Xpert MTB/RIF testing
 - LPA
 - Culture and identification of *M. tuberculosis*
 - DST for first-line and second-line anti-TB agents

National/Central Laboratories..2

Provide support for the laboratory network

- Organizing and participating in training, providing supervision and EQA of sputum-smear microscopy, GeneXpert MTB/RIF testing and culture; offering advice on procurement
- Engage in other activities
 - Participate in operational research, drug-resistance surveillance

Assessment

- What is TB and how it is transmitted?
- What is the difference between Active and Latent TB?
- Give five risk factors that predispose an individual to TB?
- What is the role of TB laboratories?

Summary

- TB is an infectious disease that mainly affects the lungs but can affect any part of the body.
- Being HIV-positive, smoking, DM, malnourished, pregnant increases the risk of developing TB disease: people coinfecting with HIV and TB have a 10% annual risk of developing active TB.
- The TB laboratory network plays a critical role in TB control, and is generally organized into 3 levels: central, intermediate and peripheral. Each level has well defined technical or managerial tasks, or both.



References

- GLI TB training package
<http://www.stoptb.org/wg/gli/trainingpackages.asp>

Acknowledgement

