Proton-to-Electron Mass Ratio from Closure Geometry and Graph Invariants

Lee Smart

Institute of Vibrational Field Dynamics

contact@vibrationalfielddynamics.org

@vfd_org

April 2026

Status: Preprint (not peer-reviewed)

Abstract

We develop a structural model of particles and mass within the deterministic crystallisation framework. Particles are identified with stable closure classes — equivalence classes of bounded field configurations on a φ-structured manifold, characterised by discrete shell support, winding number, and symmetry sector. Five explicit assignment rules constrain admissible closure classes and single out the electron as the minimal boundary closure ($\{1\}$) and the proton as the minimal interior composite ($\{2, 3, 4\}$) within the stated rule system. A combinatorial class invariant yields $\Delta C = 15$ with zero fitted continuous parameters. The result still depends on explicit discrete structural assumptions, which are stated separately (Section 4). Metric-induced class energies show that additive composition is insufficient for mass, requiring a nonlinear law. A closure-graph analysis then yields a three-order mass law,

$$\frac{m_p}{m_e} = \varphi^{\Delta C + \Delta(|E|/|V|) - \Delta\left(\mathrm{Var(deg)}\,\frac{|E|}{|V|^2}\right)} = \varphi^{1265/81},$$

where $|E|$ and $|V|$ denote the edge and vertex counts of the closure graph. The first-order graph term is derived within the present framework, conditional on the shell assignments, from the per-vertex normalised trace of the closure-graph Laplacian, and the second-order term from degree variance, with a self-consistent normalisation that is uniquely admissible within the tested power-law family under the stated selection criteria. This gives $m_p/m_e \approx 1835.8$, in strong structural agreement with the observed value 1836.15 at the $\sim 2 \times 10^{-4}$ relative level. The agreement is structural rather than metrological: the experimental ratio is known to much higher precision than the framework currently reproduces. The neutron-proton mass splitting is compatible with the architecture at the correct scale but not yet derived. Heavier lepton masses (muon, tau) are not reproduced under the current rule set, identifying lepton-generation structure as the primary limitation. The treatment is presented as a falsifiable structural hypothesis, not as a first-principles derivation or a general mass theory.

1 Introduction

The Standard Model classifies particles by quantum numbers and predicts their interactions, but particle masses are free parameters. The question of why there exist discrete particle types with specific mass hierarchies remains open.

A companion paper [Smart, 2026a] introduced the crystallisation operator: a deterministic mechanism for state selection in which unresolved configurations converge to stable attractors of a closure functional. The present paper asks two questions: what are those attractors, and what mass do they carry?

We develop a structural model in which particles are *closure classes* — discrete equivalence classes of bounded field configurations on a φ-structured manifold — and mass emerges from the geometric and graph-theoretic structure of these classes. The development proceeds through six layers:

1. Closure classes as particle identities (Sections 2–6).
2. Manifold geometry and class energy (Sections 7–8).
3. The nonlinear mass-map problem (Section 9).
4. Structural bounds on the mass ratio (Section 10).
5. First-order graph composition (Sections 11–12).
6. Second-order degree-variance refinement (Section 13).

Each layer is obtained from the preceding one within the present framework, conditional on the structural assumptions stated explicitly in Section 4. The absence of fitted continuous parameters does not imply absence of structural assumptions; the framework relies on a discrete set of structural choices whose physical justification remains part of the programme (see Section 4).

2 Particles as Solutions to Constraint Systems

We propose that particles are stable solutions of constraint systems. A particle exists because a specific combination of constraints — closure, boundedness, symmetry, topology, stability — admits a stable bounded field configuration. The discreteness of the particle spectrum reflects the discreteness of admissible constraint solutions.

3 Closure Classes

Definition 1 (Closure Class)**.** *A closure class $\mathcal{A}_{S,w,\sigma}$ is the set of all bounded, normalisable field configurations on $\mathcal{M}$ with shell support S, winding number w, and symmetry sector σ, satisfying exact phase closure and stability.*

Within the present framework, the closure class serves as the particle identity. The labels (S, w, σ) are the structural analogue of quantum numbers.

4 Scope and Structural Assumptions

The framework rests on the following structural assumptions, which are not derived from the Standard Model:

1. The φ-structured manifold $\mathcal{M}$ with nested shells at scales φ^{-n} is a modelling assumption. The choice of φ is motivated by its unique self-similar nesting property ($\varphi^{-1} = \varphi - 1$); the framework does not claim φ is the only possible geometric substrate. Whether this structure is realised in nature is an empirical question.

2. Assignment rules R1–R5 (Section 5) are structural constraints motivated by the closure framework, not derived from a fundamental Lagrangian.

3. The closure-graph construction and its Laplacian are exact given the shell assignments, but the shell assignments themselves rest on R1–R5.

4. The self-consistent normalisation of the second-order term is selected within a tested family, not derived from a deeper axiom.

5. The closure graph used in the present paper is the shell-support adjacency graph with one vertex per occupied shell. Alternative graph encodings are possible and are not analysed here.

The framework should therefore be understood as a *candidate geometric substrate* for particle classification and mass, not as an established physical theory. Its value lies in producing a mass law with zero fitted continuous parameters with strong structural agreement from a small number of explicit assumptions.

5 Assignment Rules and Class Enumeration

Five rules constrain admissible closure classes:

Definition 2 (R1–R5)**.** ***R1*** *(Minimal Support): minimal charged closure occupies the smallest shell set; composites require $|S| \geq 3$; shell 1 reserved for minimal charged.* ***R2*** *(Boundary/Interior): shell 1 is boundary; baryons occupy interior only, starting at shell 2.* ***R3*** *(Complexity Threshold): $C(\mathcal{A}) \geq -1$ for leptons, $C(\mathcal{A}) \geq 5$ for baryons.* ***R4*** *(Symmetry/Topology): contiguous shells; winding restrictions per sector.* ***R5*** *(Connectivity): all shells form a connected subgraph.*

Table 1: Closure-class admissibility under R1–R5.

Candidate	Shells	C	All Pass?	Rejection
Electron	$\{1\}$	-1	**Yes**	—
Electron (ext.)	$\{1, 2\}$	-2	No	Fails R1, R2, R3
Baryon (low)	$\{1, 2, 3\}$	-1	No	Fails R1, R2, R3
Proton	$\{2, 3, 4\}$	14	**Yes**	—
Baryon (high)	$\{3, 4, 5\}$	47	No	Fails R2

Proposition 3 (Unique Minimality)**.** *Under R1–R5, the electron assignment* $\{1\}$ *and proton assignment* $\{2, 3, 4\}$ *are uniquely determined as the minimal admissible classes in their respective sectors.*

The neutron shares the proton's shell support $\{2, 3, 4\}$ in the baryon-neutral symmetry sector (R4).

6 Combinatorial Class Invariant

Definition 4 (Combinatorial Complexity)**.** $C(\mathcal{A}) = \prod_i n_i - \sum_i n_i - 1$ *for shell support* $\{n_1, \ldots, n_s\}$.

Theorem 5 (Inter-Class Difference)**.** $\Delta C = C(\mathcal{A}_p) - C(\mathcal{A}_e) = 14 - (-1) = 15$. *This is a fixed integer determined by R1–R5 with zero fitted continuous parameters.*

7 φ-Structured Manifold Geometry

The manifold $\mathcal{M}$ consists of nested shells at scales $r_n = r_0\, \varphi^{-n}$. A field configuration is:

$$\Psi(\mathbf{x}, t) = \sum_{n \in S} a_n\, \varphi^{-n}\, e^{i(\omega_n t + \theta_n)}\, \chi_n(\mathbf{x}) \tag{1}$$

8 Metric, Measure, and Class Energy

The minimal radial metric assigns spectral weight $\lambda(n) = \varphi^{2n}$ to shell n. The class energy under additive summation is $E_{\text{sum}}(\mathcal{A}) = \sum_{n \in S} \varphi^{2n}$, giving $E_p/E_e \approx 27.4$.

Remark 6 (Diagnostic Result)**.** *The additive class-energy ratio is of order* 10^1*, while the observed mass ratio is of order* 10^3*. A nonlinear composition law is required.*

9 The Nonlinear Mass-Map Problem

Seven candidate nonlinear map families were evaluated. The tested families were required to satisfy three criteria: invariance under shell relabelling within a fixed support, zero fitted continuous parameters, and monotone growth with combinatorial complexity. Within this restricted class, two produce φ-power ratios:

Combinatorial map: $m_p/m_e = \varphi^{15} \approx 1364$.

Multiplicative spectral product: $m_p/m_e = \varphi^{16} \approx 2207$.

10 Structural Bounds

Proposition 7 (Bracketing)**.** $\varphi^{15} < m_p/m_e < \varphi^{16}$. *Both bounds are derived within the framework with zero fitted continuous parameters. The observed ratio corresponds to* $\varphi^{15.618}$*, lying strictly between.*

11 Closure-Graph Formulation

The shell support S induces a *closure graph* $G(S)$: vertices are occupied shells, edges are adjacency links between consecutive shells. We write $|V|$ and $|E|$ for the vertex and edge counts of G (not to be confused with the energy functional E_{sum} of Section 8). For the electron: $|V| = 1$, $|E| = 0$. For the proton: $|V| = 3$, $|E| = 2$.

In the present paper, the closure graph is the shell-support adjacency graph with one vertex per occupied shell; no multiplicity weighting by shell index is used at this stage.

The edge-to-vertex ratio $|E|/|V|$ equals $\text{tr}(L)/(2|V|)$, where L is the unnormalised graph Laplacian. For a simple unweighted graph, $\text{tr}(L) = 2|E|$, so the per-vertex normalised trace equals $|E|/|V|$. This is an exact operator identity.

12 First-Order Graph Composition

The present framework adopts a leading-order factorisation principle in which distinct closure-class sectors contribute multiplicatively to mass. Since the combinatorial invariant C (depending on shell *indices*) and the graph term $|E|/|V|$ (depending on shell *adjacency*) act on distinct structural aspects of the closure class at leading order, the factorisation gives:

$$m(\mathcal{A}) = \Lambda\, \varphi^{C(\mathcal{A})} \cdot \varphi^{|E(\mathcal{A})|/|V(\mathcal{A})|} \tag{2}$$

The exponents add because independent factors multiply. This yields the first-order law:

$$\frac{m_p}{m_e} = \varphi^{\Delta C + \Delta(|E|/|V|)} = \varphi^{15+2/3} = \varphi^{47/3} \approx 1880 \quad \text{(error: 2.4\%)} \tag{3}$$

13 Second-Order Degree-Variance Refinement

13.1 Source

The first-order graph term uses the *first moment* (mean) of the degree distribution. The leading second-order correction considered here uses the *second central moment* (variance):

$$\delta_2(\mathcal{A}) = -\frac{\text{Var}(\deg(\mathcal{A})) \;\cdot\; |E(\mathcal{A})|}{|V(\mathcal{A})|^2}$$

The negative sign is consistent with the interpretation that degree heterogeneity (irregular connectivity) reduces effective coupling. The electron ($\text{Var} = 0$) receives no correction. The proton ($\deg = [1, 2, 1]$, $\text{Var} = 2/9$) receives $\delta_2 = -(2/9)(2/9) = -4/81$.

13.2 Normalisation

Within the tested power-law family $N = |V|^a|E|^b$, the normalisation $|V|^2/|E|$ is the only member satisfying both the self-consistency criterion (each order normalised by the previous) and the empirical screening criterion of sub-1% agreement. It equals $|V|/g_1$, where $g_1 = |E|/|V|$ is the first-order graph term. This is the *self-consistent normalisation*: each order of the expansion is normalised relative to the previous order.

13.3 Three-Order Mass Law

Proposition 8 (Three-Order Mass Law)**.** *Within the present framework, the mass ratio between proton and electron closure classes is:*

$$\frac{m_p}{m_e} = \varphi^{\Delta C + \Delta(|E|/|V|) - \Delta\left(\mathrm{Var(deg)}\,\frac{|E|}{|V|^2}\right)} = \varphi^{15 + 2/3 - 4/81} = \varphi^{1265/81} \tag{4}$$

where $1265/81$ *is in lowest terms (*$\gcd(1265, 81) = 1$*). Numerically:*

$$\varphi^{1265/81} \approx 1835.8 \qquad (\textit{observed: } 1836.15, \quad \textit{relative error: } 2 \times 10^{-4})$$

No continuous parameters are fitted. The result depends on the discrete structural choices detailed in Section 4.

Table 2: Three-order mass law: cumulative accuracy.

Order	Term	Exponent	Computed m_p/m_e	Error
0th	$\varphi^{\Delta C}$	15	1364	25.7%
1st	$+\,\Delta(\lvert E\rvert/\lvert V\rvert)$	$15 + 2/3 = 47/3$	1880	2.4%
2nd	$\boldsymbol{-\,\Delta(\mathrm{Var(deg)}\,\lvert E\rvert/\lvert V\rvert^2)}$	$\mathbf{47/3 - 4/81 = 1265/81}$	**1835.8**	$\mathbf{2 \times 10^{-4}}$

14 Operator Origin and Factorisation

The three-order law is organised by a factorisation ansatz in which mass contributions from distinct closure-class sectors multiply at leading order.

- **Set sector** (φ^C): depends on shell *indices* (which shells). Derived from shell-support combinatorics.
- **Graph sector** ($\varphi^{|E|/|V|}$): depends on shell *adjacency* (how connected). Derived within the framework from $\mathrm{tr}(L)/(2|V|)$.
- **Variance sector** ($\varphi^{-\mathrm{Var(deg)}|E|/|V|^2}$): depends on degree *heterogeneity*. Self-consistently normalised by the first-order graph term. Structurally selected within the tested power-law family.

These sectors act on independent aspects of the closure class at leading order. Independent sectors multiply at the mass level, yielding additive exponents. Subleading sector couplings may exist but are not included in the present treatment. The factorised form is:

$$m(\mathcal{A}) = \Lambda \;\cdot\; \varphi^{C(\mathcal{A})} \;\cdot\; \varphi^{\mathrm{tr}(L_{\mathcal{A}})/(2|V_{\mathcal{A}}|)} \;\cdot\; \varphi^{-\mathrm{Var}(\deg_{\mathcal{A}})\,|E_{\mathcal{A}}|/|V_{\mathcal{A}}|^2} \tag{5}$$

Scope of the Derived Claim

The main derived claim of this paper is the proton/electron mass ratio within the stated closure framework. The framework assumptions include the φ-structured shell manifold, the assignment rules R1–R5, the combinatorial invariant, the shell-support closure graph, and the order-by-order factorisation ansatz. All numerical agreement claims should be understood relative to those assumptions.

15 Status Summary

Table 3: Claim status map.

Claim	Status	Basis
Closure-class definition	POSTULATED	Structural framework
Assignment rules R1–R5	POSTULATED	Closure constraints
Combinatorial invariant $C(\mathcal{A})$	POSTULATED	Combinatorial definition
Unique minimality of $\{1\}$, $\{2,3,4\}$	DERIVED	Follows from R1–R5
$\Delta C = 15$	DERIVED	Arithmetic from assignments
Additive energy insufficient	DERIVED	Direct computation
Bounds φ^{15}–φ^{16}	DERIVED	From framework
Graph Laplacian term	DERIVED	Operator identity $\mathrm{tr}(L)/(2\|V\|)$
Degree-variance source	DERIVED	Second central moment
Normalisation $\|V\|^2/\|E\|$	STRUCTURALLY SELECTED	Unique in tested power-law family
Three-order exponent 1265/81	DERIVED (within framework)	From above
Numerical agreement ≈ 1835.8	NUMERICAL OBSERVATION	2×10^{-4} vs. 1836.15
Neutron-proton splitting	CONDITIONAL	Compatible at correct scale; not d
Muon/tau generations	Not reproduced	Rule extension needed

16 First Validation Extensions

16.1 Neutron-Proton Splitting

The neutron and proton share the closure class $\{2,3,4\}$ and are therefore identical at all three orders of the mass law. The observed neutron-proton mass difference $\Delta m_{np} = 1.293\,\mathrm{MeV}/c^2$ ($m_n/m_p = 1.00138$) corresponds to an exponent correction $\delta_\sigma \approx 0.00286$. This is of the correct order of magnitude for a third-order or symmetry-sector correction within the existing architecture: candidate structural quantities at this scale include $\mathrm{Var(deg)}/|V|^3 \approx 0.008$ and $1/C^2 \approx 0.005$. The framework naturally leaves room for a correction of the observed order of magnitude as an additional multiplicative factor φ^{δ_σ}, entering through the distinction between charged and neutral baryon symmetry sectors. However, no specific correction form has been derived; the splitting is *compatible with* the framework but *not yet derived from* it.

16.2 Heavier Lepton Tests

The muon and tau were tested as tentative closure classes under extensions of R1–R5. The assignment $\mu \to \{1,2\}$ violates R3 ($C = -2 < -1$) and yields $m_\mu/m_e \approx 0.8$, orders of magnitude below the observed 206.8. The assignment $\tau \to \{1,2,3\}$ satisfies R3 ($C = -1$) but yields $m_\tau/m_e \approx 1.3$, far from the observed 3477.

Both failures arise from the same structural issue: the current shell-support model assigns small or negative combinatorial complexity to multi-shell leptonic states, producing mass ratios near unity rather than the observed 10^2–10^3 hierarchy. The lepton generation problem requires either an extension of the rule set (e.g. generation-dependent

winding, higher-dimensional shell structure, or an entirely different assignment mechanism for heavier leptons) or a fundamental revision of how leptonic mass scales with shell support.

16.3 Scope of Current Results

Table 4: Validation results under the current framework.

Ratio	Assignment	Computed	Observed	Verdict
m_p/m_e	$\{2,3,4\}$ vs $\{1\}$	1835.8	1836.15	Structural agreement (2×10^{-4})
m_n/m_p	same class + δ_σ	(compatible)	1.00138	Compatible at correct scale
m_μ/m_e	$\{1,2\}$ vs $\{1\}$	0.8	206.8	Fails (rule violation; wrong scale)
m_τ/m_e	$\{1,2,3\}$ vs $\{1\}$	1.3	3477	Fails (wrong scale)

The proton-to-electron ratio is the framework's strongest result. The neutron-proton splitting is compatible but not derived. The muon and tau are not reproduced, representing a genuine limitation that constrains the scope of the current claim. The present result should be interpreted as evidence of structural viability for the baryon-lepton mass hierarchy within the stated framework, not as a general mass theory. Broader validation — particularly extension to lepton generations — is required to exclude the possibility of structural coincidence in the proton-to-electron case.

17 Discussion and Limitations

Derived within the framework:

- combinatorial invariant $\Delta C = 15$
- graph Laplacian term $\mathrm{tr}(L)/(2|V|)$
- degree-variance source for the second-order term

Structurally selected within a tested family:

- self-consistent normalisation $|V|^2/|E|$ (unique within the power-law family $|V|^a|E|^b$)

Validated:

- neutron-proton splitting: compatible at the correct scale (not derived)

Not reproduced:

- muon and tau masses: current rule set does not generate heavier lepton generations

Open:

- deeper axiomatic justification of the self-consistency normalisation principle
- derivation (not just compatibility) of neutron-proton splitting
- extension of rule set for lepton generations

- physical realisation of the φ-manifold

What this paper does not claim: First-principles mass derivation in the strong physics sense. The normalisation principle is structurally selected, not axiomatically derived. The φ-structured manifold is a modelling assumption. The framework does not replace the Standard Model.

The agreement obtained here is structural rather than metrological: the computed ratio is accurate at the $\sim 10^{-4}$ relative level, whereas the experimental ratio $m_p/m_e = 1836.15267343(11)$ is known to a relative precision of $\sim 6 \times 10^{-11}$. The framework does not approach experimental precision; it provides structural agreement at a level that is not attributable to adjustment of a fitted continuous parameter, but does not yet constitute a rigorous derivation.

18 Conclusion

Particles correspond to stable closure classes on a φ-structured manifold. Five assignment rules single out the electron and proton as minimal admissible classes within the stated rule system, yielding the combinatorial invariant $\Delta C = 15$ with zero fitted continuous parameters. Metric-induced energies show that mass cannot arise from additive shell energy alone and instead requires nonlinear composition. The resulting mass law factorises into three structural sectors: a combinatorial sector φ^C, a graph-connectivity sector $\varphi^{|E|/|V|}$, and a second-order degree-variance refinement $\varphi^{-\mathrm{Var(deg)}\,|E|/|V|^2}$. For the proton-to-electron pair this yields the rational exponent 1265/81 and a computed ratio $m_p/m_e \approx 1835.8$, in structural agreement with the observed value at the $\sim 2 \times 10^{-4}$ level.

The leading proton/electron mass law is established within the present framework through second order. First validation tests show that the neutron-proton splitting is compatible with the architecture at the correct scale, but heavier lepton masses (muon, tau) are not reproduced under the current rule set, identifying lepton-generation structure as the primary open problem. What remains is not the existence of the leading law, but the deeper axiomatic justification of the second-order normalisation, the derivation of neutron-proton splitting, and the extension of the rule set to accommodate lepton generations. The framework is falsifiable: the muon and tau failures already constrain the scope of the current claim, and broader validation is required before the proton-to-electron success can be interpreted as more than a structurally interesting result within the present framework.

Subsequent work [Smart, 2026b] suggests that the spectral structure required for lepton generation may be native to the vertex geometry of the 600-cell. This does not affect the derivation presented here, which is self-contained at the baryon-lepton level.

A Discarded Metric and Map Families

Five metric families were evaluated. Best additive ratio: ~ 376 under weighted energy. Seven nonlinear map families were tested; only the multiplicative product (exponent 16) and combinatorial map (exponent 15) remain viable as bounds. "Viable" here means producing a proton/electron exponent in the observed bracket $[\varphi^{15}, \varphi^{16}]$ without introducing fitted continuous parameters. Twelve graph rules were tested; only the edge-to-vertex ratio produces an exponent in the correct range.

B Normalisation Selection within the Power-Law Family

The self-consistency criterion requires the n-th correction to be normalised by a quantity constructed from the $(n-1)$-th sector, so that each order is scaled relative to the preceding one rather than introduced independently.

Within the power-law family $N = |V|^a|E|^b$, the normalisation $|V|^2/|E|$ (corresponding to $a = 2$, $b = -1$) is the only member satisfying both this self-consistency criterion and the empirical screening criterion of sub-1% agreement on the proton/electron ratio. The next-best alternative ($|V|^2$, corresponding to $a = 2$, $b = 0$) gives 1.17% error. The selected normalisation equals $|V|/g_1$, where $g_1 = |E|/|V|$ is the first-order graph term, establishing a self-referential structure in which each order of the expansion is normalised by the previous one.

References

Lee Smart. Crystallisation dynamics as a deterministic alternative to collapse in open quantum systems. 2026a. Preprint, Institute of Vibrational Field Dynamics.

Lee Smart. Lepton generations and the missing mass operator in deterministic field closure. 2026b. Companion paper. Preprint, Institute of Vibrational Field Dynamics.