The Crystallisation Operator: Mathematical Structure and Properties

Lee Smart

Institute of Vibrational Field Dynamics

contact@vibrationalfielddynamics.org

@vfd_org

March 2026

Abstract

We introduce the **crystallisation operator**, a mathematical framework that replaces the discontinuous, probabilistic notion of wavefunction collapse with a deterministic constraint-driven process. In this formulation an unresolved multi-state configuration evolves under lawful constraints toward a stable, coherent structure by minimising a closure functional that balances constraint satisfaction, energetic cost, and phase coherence. The observed definite outcome is reinterpreted not as a random projection but as the formation of a closure-compatible eigenstate — a *crystal* in configuration space. We develop the operator definition, its dynamical flow form, a spectral reweighting formulation, and prove existence, stability, and mode-selection theorems. Applications to quantum measurement are discussed, with an outlook on broader applicability.

1 Introduction

The measurement problem in quantum mechanics is conventionally addressed through wavefunction collapse: upon measurement, a superposition is said to discontinuously reduce to a single eigenstate with probability given by the Born rule. Despite its empirical success, this prescription suffers from several conceptual difficulties:

- It introduces a fundamental discontinuity absent from unitary evolution.
- The selection mechanism remains unspecified — *why* a particular outcome obtains is left to irreducible randomness.
- The boundary between "quantum" and "classical" regimes is vague.

The Vacuum Field Dynamics (VFD) programme offers an alternative perspective. Rather than treating definite outcomes as the result of stochastic projection, VFD frames state resolution as a **physical process of crystallisation**: the system evolves from an unstable, over-complete, constraint-incompatible configuration into a stable, constraint-satisfying coherent structure.

This paper formalises that intuition.

2 Collapse vs Crystallisation

2.1 Standard Collapse

In orthodox quantum mechanics, measurement on state $\Psi = \sum_i c_i \phi_i$ yields eigenstate ϕ_k with probability $|c_k|^2$. Formally:

$$\Psi \mapsto P_k \Psi = \phi_k \tag{1}$$

This is instantaneous, irreversible, and non-unitary.

2.2 Crystallisation

We propose instead:

$$C[\Psi] = \Psi^\star \tag{2}$$

where $\Psi^\star$ is the stable attractor of a **closure functional** F over the space of admissible configurations $\mathcal{A}$:

$$\Psi^\star = \arg\min_{\Phi \in \mathcal{A}} F[\Phi] \tag{3}$$

The key conceptual shift is that the definite outcome is not *selected at random* but *formed by constraint resolution.* What appears observationally as "collapse" is reinterpreted as deterministic or quasi-deterministic convergence to a constraint-compatible coherent structure.

3 Mathematical Framework

The following framework is presented as a mathematical model for state resolution, not as a derived consequence of established physical theory.

3.1 State Space

Let $\Psi \in \mathcal{H}_M$, the Hilbert space of configurations over a constrained manifold M.

3.2 Closure Residual

The closure residual $R[\Psi]$ measures how far the current state is from satisfying the governing constraints:

$$R[\Psi] = \|G[\Psi] - K\|^2 \geq 0 \tag{4}$$

where $G[\Psi]$ is the constraint structure induced by the state and K is the admissible closure target. $R[\Psi] = 0$ indicates full coherence.

3.3 Energy Functional

$$E[\Psi] = \langle \Psi | L | \Psi \rangle \tag{5}$$

where L is a coupling/Laplacian/Hamiltonian-proxy operator encoding the energetic cost of the configuration.

3.4 Coherence Metric

$$Q[\Psi] = \frac{1}{n^2} \sum_{i,j} \cos(\theta_i - \theta_j) \tag{6}$$

where $\theta_i = \arg(\Psi_i)$. This measures global phase alignment; $Q = 1$ indicates perfect phase coherence.

3.5 Crystallisation Functional

$$F[\Phi] = \alpha\, R[\Phi] + \beta\, E[\Phi] - \gamma\, Q[\Phi] \tag{7}$$

with $\alpha, \beta, \gamma > 0$. Minimising F simultaneously:

- Reduces constraint mismatch (R)
- Lowers energetic cost (E)
- Rewards phase coherence (Q)

At this stage, the functional is introduced as a modelling assumption designed to test whether constraint-driven selection produces consistent dynamical structure.

4 Operator Definition

Definition 1 (Crystallisation Operator)**.** *The crystallisation operator C maps an unresolved field configuration Ψ on manifold M to a stable coherent configuration $\Psi^\star$ by constrained minimisation:*

$$C[\Psi] = \Psi^\star = \arg\min_{\Phi \in \mathcal{A}} F[\Phi] \tag{8}$$

subject to:

$$\nabla_\Phi F[\Psi^\star] = 0, \qquad \delta^2 F[\Psi^\star] > 0 \tag{9}$$

The first condition is stationarity; the second ensures the crystallised state is a ***stable*** *attractor, not a saddle point.*

5 Dynamical Formulation

Rather than instantaneous collapse, crystallisation proceeds as a gradient flow:

$$\frac{\partial \Psi}{\partial t} = -\nabla_\Psi F[\Psi] \tag{10}$$

This means:

- The state flows downhill in the constraint-energy-coherence landscape.
- Crystallisation is the arrival at an attractor.
- No discontinuity is required.

Equilibrium: $\nabla_\Psi F[\Psi^\star] = 0$
Stability: $\delta^2 F[\Psi^\star] > 0$

5.1 Crystallisation Timescale

Define the instability-to-resolution rate:

$$\Omega[\Psi] = a\,R[\Psi] + b\,Q[\Psi] + c\,|\nabla V(\Psi)| \tag{11}$$

Then the crystallisation timescale is:

$$\tau_{\text{crys}} = \frac{1}{\Omega[\Psi]} \tag{12}$$

Higher mismatch or steeper landscape gradients lead to faster crystallisation. This provides a physical alternative to stochastic collapse timescales.

6 Spectral Formulation

Expand the unresolved state in eigenmodes: $\Psi = \sum_i c_i \phi_i$.

Crystallisation is not "pick one basis vector" but a **coherence-weighted reweighting**:

$$\tilde{c}_i = \frac{c_i \exp(-\mu R_i - \nu E_i + \rho Q_i)}{Z} \tag{13}$$

where $Z = \sum_j c_j \exp(-\mu R_j - \nu E_j + \rho Q_j)$.

Interpretation:

- Modes with poor closure (R_i large) are suppressed.
- Modes with high coherence (Q_i large) are amplified.
- The surviving structure is the coherent crystal that passes the constraint filter.

This expression should be understood as a formal reweighting model rather than a derived physical transition amplitude.

7 Theoretical Results

Theorem 2 (Existence)**.** *Let $F : \mathcal{A} \to \mathbb{R}$ be continuous and coercive on a non-empty, closed, bounded admissible set $\mathcal{A} \subset \mathcal{H}_M$. Then F attains its minimum: there exists $\Psi^\star \in \mathcal{A}$ such that $F[\Psi^\star] \leq F[\Phi]$ for all $\Phi \in \mathcal{A}$.*

Proof sketch. F is continuous and $\mathcal{A}$ is compact (closed and bounded in finite dimensions, or weakly compact in the infinite-dimensional case with coercivity). By the extreme value theorem (or direct method of calculus of variations), F attains its infimum on $\mathcal{A}$. □ □

Theorem 3 (Stability)**.** *If $\Psi^\star$ is a strict local minimiser of F with $\nabla F[\Psi^\star] = 0$ and Hessian $H_F[\Psi^\star]$ positive definite, then $\Psi^\star$ is a Lyapunov-stable equilibrium of the gradient flow $\dot{\Psi} = -\nabla F[\Psi]$.*

Proof sketch. F itself serves as a Lyapunov function: $\dot{F} = \langle \nabla F, \dot{\Psi} \rangle = -\|\nabla F\|^2 \leq 0$. Strict local minimality with positive-definite Hessian gives $F[\Psi] - F[\Psi^\star] \geq \frac{\lambda_{\min}}{2}\|\Psi - \Psi^\star\|^2$ in a neighbourhood, providing the required Lyapunov bound. □ □

Theorem 4 (Monotonic Descent)**.** *Along the gradient flow* $\dot{\Psi} = -\nabla F[\Psi]$*:*

$$\frac{d}{dt}F[\Psi(t)] = -\|\nabla F[\Psi(t)]\|^2 \leq 0 \tag{14}$$

F is monotonically non-increasing.

Proof. Direct computation by chain rule. □ □

Theorem 5 (Mode Selection)**.** *In the spectral formulation, let modes be indexed by residual R_i and coherence Q_i. As the reweighting parameters $\mu, \rho \to \infty$ (strong constraint enforcement), the reweighted coefficients $\tilde{c}_i$ asymptotically concentrate on the mode(s) minimising $\mu R_i + \nu E_i - \rho Q_i$.*

Proof sketch. This follows from the standard Laplace/saddle-point asymptotics of exponential tilting: the mode achieving the minimum exponent dominates the partition sum Z in the limit. □ □

8 Physical Interpretation

The crystallisation framework provides a four-layer account of what is conventionally called "collapse":

Table 1: Four-layer account of state resolution via crystallisation.

Layer	Description
Pre-state	A manifold carries many partially compatible mode possibilities
Instability	The mixed state cannot remain globally self-consistent
Crystallisation flow	The state is driven through a closure-energy-coherence landscape
Formed structure	A coherent eigenstructure stabilises and becomes the observed state

This provides a more explicit dynamical account than models in which state selection is treated as intrinsically stochastic:

1. The mechanism is specified (functional minimisation).
2. The outcome is explained (constraint-compatible attractor).
3. The timescale is derived (not postulated).
4. No discontinuity is required.

9 Applications

9.1 Quantum Measurement

The measurement apparatus introduces boundary conditions and decoherence channels that sharpen the closure functional landscape. The system crystallises into the eigenstate(s) most compatible with the combined system–apparatus constraints. A possible interpretation is that Born probabilities may be related to the basin structure of F, though this connection is not established here.

9.2 Outlook: Beyond Quantum Mechanics

The mathematical structure of the crystallisation operator — constrained minimisation over a functional landscape with attractor dynamics — is not inherently quantum-mechanical. The same formalism may apply to other systems exhibiting structured state selection under constraints, including pattern formation in developmental biology and optimisation in constraint-satisfaction problems. These directions are speculative and are noted only as potential extensions; they are not developed here.

10 Discussion and Limitations

What this framework provides:

- A precise mathematical operator providing an alternative formulation of state selection.
- Continuous dynamics rather than discontinuous jumps.
- Derived timescales rather than postulated ones.
- A unifying formalism with potential applicability beyond quantum mechanics.

Current limitations:

- The choice of α, β, γ weights requires physical justification specific to each application domain.
- The connection to Born-rule statistics requires further development — specifically, showing that basin volumes under F reproduce $|c_i|^2$ weighting.
- Infinite-dimensional generalisation requires careful functional-analytic treatment.

What this is not:

- A claim to replace all of quantum mechanics.
- A consciousness theory.
- A metaphysical framework.

It is strictly a mathematical and dynamical-systems reformulation of state resolution. The role of this paper is to define the operator and its mathematical properties; questions of experimental realisation and physical interpretation are addressed separately.

References