
ALGEBRA-SUMMER

Vincent Ferrara

Contents

0.1	Intro	3
1	Groups	5
1.1	Dihedral Groups	5

0.1 Intro

Theorem 0.1.1. (Division Algorithm) Suppose $a, b \in \mathbb{Z}$ s.t. $b > 0$. Then there are unique $q, r \in \mathbb{Z}$ s.t. $a = bq + r$ and $0 \leq r < b$

Proof. Let $a, b \in \mathbb{Z}$ s.t. $b > 0$. Consider $A = \{a - bx \mid x \in \mathbb{Z}\} \cap \mathbb{N}$

By the Well Ordering Principle since A has all non-negative terms it has a minimum lets call this r and call the associated $x \in \mathbb{Z}$, q .

Thus, $r = a - bq \implies a = bq + r$.

For contradiction, assume $r \geq b$.

$\implies r = b + m$ s.t. $m \geq 0$

$\implies b + m = a - bq \implies m = a - b(q + 1) \in A \implies m < r = \text{minimum} \implies \Leftarrow$ Thus $0 \leq r < b$

Fix $q_1, q_2, r_1, r_2 \in \mathbb{Z}$ s.t. $a = bq_1 + r_1 = bq_2 + r_2$ and $0 \leq r_1, r_2 < b$. WLOG let $r_1 \geq r_2$

$\implies bq_1 + r_1 = bq_2 + r_2 \implies r_1 - r_2 = b(q_2 - q_1) \implies r_1 - r_2$ is a multiple of b

Since $0 \leq r_1, r_2 < b \implies 0 \leq r_1 - r_2 < b$

Since $r_1 - r_2$ is strictly less than b the only multiple of b it could be is 0 thus $r_1 = r_2$

$\implies q_1 = q_2$ □

Lemma 0.1.2. (Bezout's Lemma) Suppose $a, b \in \mathbb{Z} \setminus \{0\}$. Then there are $x, y \in \mathbb{Z}$ s.t. $ax + by = \gcd(a, b)$.

Proof. Let $a, b \in \mathbb{Z} \setminus \{0\}$. Consider $A = \{ax + by \mid x, y \in \mathbb{Z}\} \cap \mathbb{N}$

By the Well Ordering Principle since A has all non-negative terms it has a minimum lets call this d and call the associated $x, y \in \mathbb{Z}$, u and v .

By Theorem 0.1.1 $\exists q, r \in \mathbb{Z}$ s.t. $a = dq + r$ and $0 \leq r < d$

Case 1: $r = 0 \implies d \mid a$

Case 2: $0 < r < d \implies r = a - dq \implies r = a - (au + bv)q = a(1 - uq) + b(-vq) \in A$

$\implies r < d = \text{minimum} \implies \Leftarrow$

Thus $d \mid a$ and similarly $d \mid b$

Let $c \in \mathbb{N}$ s.t. $c \mid a$ and $c \mid b$


$\implies a = cm$ and $b = cn$ for $m, n \in \mathbb{Z}$

$d = au + bv = c(mu) + c(mv) = c(mu + mv) \implies c \mid d \implies d = \gcd(a, b)$ □


Algorithm 0.1.3. (Euclidean Algorithm)

Find $\gcd(272, 571)$ and $x, y \in \mathbb{Z}$ s.t. $272x + 571y = \gcd(272, 571)$

$$571 = 272 \cdot 2 + 27 \quad (1)$$



$$272 = 27 \cdot 10 + 2 \quad (2)$$



$$27 = 2 \cdot 13 + 1 \quad (3)$$

Last term before remainder is zero is gcd

$$\gcd(272, 571) = 1$$

$$2 = 272 - 27 \cdot 10$$

Use equation (2)

$$= 272(21) - 571(10)$$

$$1 = (571 - 272 \cdot 2) - 13(272(21) - 571)$$

Use equation (3) with (2)

$$= 571(131) + 272(-275)$$

1 Groups

Definition 1.0.1. A group is a set G together with a binary operation $*$: $G \times G \rightarrow G$ s.t.

- **(Closure)** For all $a, b \in G$ we have $a * b \in G$
- **(Identity)** There is $e \in G$ s.t. for all $a \in G$ $a * e = e * a = a$
- **(Inverses)** For all $a \in G$ there is an $a^{-1} \in G$ s.t. $a * a^{-1} = a^{-1} * a = e$
- **(Associativity)** For all $a, b, c \in G$ we have $a * (b * c) = (a * b) * c$

Definition 1.0.2. Given a group G and $g \in G$ we say the order of g is $n \in \mathbb{N}$ if n is the minimum value s.t. $g^n = g * g * \dots * g = e$

Write as: $\text{ord}(g) = |g| = n$

1.1 Dihedral Groups