

ECE750T-28:
Computer-aided Reasoning for Software Engineering

Lecture 6: First Order Logic
Syntax and Semantics

Vijay Ganesh
(Original notes from Isil Dillig)

Overview

- ▶ So far: Automated reasoning in propositional logic.

Overview

- ▶ So far: Automated reasoning in propositional logic.
- ▶ Propositional logic is simple and easy to automate, but not very expressive

Overview

- ▶ So far: Automated reasoning in propositional logic.
- ▶ Propositional logic is simple and easy to automate, but not very expressive
- ▶ **Today:** First order logic, also known as relational logic, predicate logic, or first-order predicate calculus

Overview

- ▶ So far: Automated reasoning in propositional logic.
- ▶ Propositional logic is simple and easy to automate, but not very expressive
- ▶ **Today:** First order logic, also known as relational logic, predicate logic, or first-order predicate calculus
- ▶ Much richer and more expressive, but does not admit completely automated reasoning (more on this later)

The Plan

- ▶ Syntax and informal semantics (review, today's lecture)

The Plan

- ▶ Syntax and informal semantics (review, today's lecture)
- ▶ Formal semantics, model theory (today's lecture)

The Plan

- ▶ Syntax and informal semantics (review, today's lecture)
- ▶ Formal semantics, model theory (today's lecture)
- ▶ Semantic argument method for FOL and properties (next lecture)

The Plan

- ▶ Syntax and informal semantics (review, today's lecture)
- ▶ Formal semantics, model theory (today's lecture)
- ▶ Semantic argument method for FOL and properties (next lecture)
- ▶ Unification, clausal form (third lecture)

The Plan

- ▶ Syntax and informal semantics (review, today's lecture)
- ▶ Formal semantics, model theory (today's lecture)
- ▶ Semantic argument method for FOL and properties (next lecture)
- ▶ Unification, clausal form (third lecture)
- ▶ Resolution and first-order theorem proving (fourth lecture)

Constants in First-Order Logic

- ▶ In propositional logic, we had two constants \top and \perp

Constants in First-Order Logic

- ▶ In propositional logic, we had two constants \top and \perp
- ▶ In first order logic, constants are more involved.

Constants in First-Order Logic

- ▶ In propositional logic, we had two constants \top and \perp
- ▶ In first order logic, constants are more involved.
- ▶ Three kinds of constants:
 1. object constants
 2. function constants
 3. relation constants

Object Constants

- ▶ **Object constants** refer to objects in a universe of discourse.

Object Constants

- ▶ **Object constants** refer to objects in a universe of discourse.
- ▶ Objects can be anything we want to say something about: integers, people, real numbers, geometric shapes, ...

Object Constants

- ▶ **Object constants** refer to objects in a universe of discourse.
- ▶ Objects can be anything we want to say something about: integers, people, real numbers, geometric shapes, ...
- ▶ **Example:** If our universe of discourse is people, object constants can be *jack, jane, joe, ...*

Object Constants

- ▶ **Object constants** refer to objects in a universe of discourse.
- ▶ Objects can be anything we want to say something about: integers, people, real numbers, geometric shapes, ...
- ▶ **Example:** If our universe of discourse is people, object constants can be *jack, jane, joe, ...*
- ▶ As a convention, we will use letters starting with $a - t$ or digits to denote object constants.

Object Constants

- ▶ **Object constants** refer to objects in a universe of discourse.
- ▶ Objects can be anything we want to say something about: integers, people, real numbers, geometric shapes, ...
- ▶ **Example:** If our universe of discourse is people, object constants can be *jack, jane, joe, ...*
- ▶ As a convention, we will use letters starting with $a - t$ or digits to denote object constants.
- ▶ Example: *a, art, beth, 1* etc. refer to object constants.

Function Constants

- ▶ **Function constants** refer to functions

Function Constants

- ▶ **Function constants** refer to functions
- ▶ Examples: *mother*, *age*, *plus*, *times*, ...

Function Constants

- ▶ **Function constants** refer to functions
- ▶ Examples: *mother*, *age*, *plus*, *times*, ...
- ▶ Each function constant has an associated **arity** indicating its number of arguments

Function Constants

- ▶ **Function constants** refer to functions
- ▶ Examples: *mother*, *age*, *plus*, *times*, ...
- ▶ Each function constant has an associated **arity** indicating its number of arguments
- ▶ Example: *mother* has arity 1 (unary), *times* has arity 2 (binary) etc.

Function Constants

- ▶ **Function constants** refer to functions
- ▶ Examples: *mother*, *age*, *plus*, *times*, ...
- ▶ Each function constant has an associated **arity** indicating its number of arguments
- ▶ Example: *mother* has arity 1 (unary), *times* has arity 2 (binary) etc.
- ▶ An object constant is really a special case of a function constant with arity 0

Relation Constants

- ▶ **Relation constants** refer to relations between or properties of objects

Relation Constants

- ▶ **Relation constants** refer to relations between or properties of objects
- ▶ Example: *loves*, *betterthan*, *ishappy*, ...

Relation Constants

- ▶ **Relation constants** refer to relations between or properties of objects
- ▶ Example: *loves*, *betterthan*, *ishappy*, ...
- ▶ Each relation constant also has an associated arity

Relation Constants

- ▶ **Relation constants** refer to relations between or properties of objects
- ▶ Example: *loves*, *betterthan*, *ishappy*, ...
- ▶ Each relation constant also has an associated arity
- ▶ Example: *loves* has arity 2, *ishappy* has arity 1 etc.

Terms

- ▶ A set of object, function, and relation constants C , F , R specifies a **first-order language**, written $L(C, F, R)$

Terms

- ▶ A set of object, function, and relation constants C, F, R specifies a **first-order language**, written $L(C, F, R)$
- ▶ C, F, R form the **signature** of the language.

Terms

- ▶ A set of object, function, and relation constants C, F, R specifies a **first-order language**, written $L(C, F, R)$
- ▶ C, F, R form the **signature** of the language.
- ▶ Terms t for a first order language are generated using C, F

Terms

- ▶ A set of object, function, and relation constants C, F, R specifies a **first-order language**, written $L(C, F, R)$
- ▶ C, F, R form the **signature** of the language.
- ▶ Terms t for a first order language are generated using C, F
- ▶ **Basic terms**: Any object constant in C and **variables**, denoted x, y, z, \dots

Terms

- ▶ A set of object, function, and relation constants C, F, R specifies a **first-order language**, written $L(C, F, R)$
- ▶ C, F, R form the **signature** of the language.
- ▶ Terms t for a first order language are generated using C, F
- ▶ **Basic terms**: Any object constant in C and **variables**, denoted x, y, z, \dots
- ▶ **Composite terms**: $f(t_1, \dots, t_k)$ where $f \in F$ is function of arity k , and t_1, \dots, t_k are terms

Terms

- ▶ A set of object, function, and relation constants C, F, R specifies a **first-order language**, written $L(C, F, R)$
- ▶ C, F, R form the **signature** of the language.
- ▶ Terms t for a first order language are generated using C, F
- ▶ **Basic terms**: Any object constant in C and **variables**, denoted x, y, z, \dots
- ▶ **Composite terms**: $f(t_1, \dots, t_k)$ where $f \in F$ is function of arity k , and t_1, \dots, t_k are terms
- ▶ Examples: $mary, x, sister(mary), price(x, macys), age(mother(y)), \dots$

Formulas

- **Formulas** of $L(C, F, R)$ are formed using the terms of this language, relation constants R , logical connectives $\neg, \wedge, \vee, \dots$, and quantifiers \forall, \exists .

Formulas

- ▶ **Formulas** of $L(C, F, R)$ are formed using the terms of this language, relation constants R , logical connectives $\neg, \wedge, \vee, \dots$, and quantifiers \forall, \exists .
- ▶ **Atomic formula (predicate)**: Expression $p(t_1, \dots, t_n)$ where $p \in R$ (of arity n), and t_1, \dots, t_n are terms of $L(C, F, R)$

Formulas

- ▶ **Formulas** of $L(C, F, R)$ are formed using the terms of this language, relation constants R , logical connectives $\neg, \wedge, \vee, \dots$, and quantifiers \forall, \exists .
- ▶ **Atomic formula (predicate)**: Expression $p(t_1, \dots, t_n)$ where $p \in R$ (of arity n), and t_1, \dots, t_n are terms of $L(C, F, R)$
- ▶ If F_1 and F_2 are formulas, then so is $F_1 \star F_2$ where \star is any binary connective

Formulas

- ▶ **Formulas** of $L(C, F, R)$ are formed using the terms of this language, relation constants R , logical connectives $\neg, \wedge, \vee, \dots$, and quantifiers \forall, \exists .
- ▶ **Atomic formula (predicate)**: Expression $p(t_1, \dots, t_n)$ where $p \in R$ (of arity n), and t_1, \dots, t_n are terms of $L(C, F, R)$
- ▶ If F_1 and F_2 are formulas, then so is $F_1 \star F_2$ where \star is any binary connective
- ▶ If F is a formula, then so is $\neg F$

Formulas

- ▶ **Formulas** of $L(C, F, R)$ are formed using the terms of this language, relation constants R , logical connectives $\neg, \wedge, \vee, \dots$, and quantifiers \forall, \exists .
- ▶ **Atomic formula (predicate)**: Expression $p(t_1, \dots, t_n)$ where $p \in R$ (of arity n), and t_1, \dots, t_n are terms of $L(C, F, R)$
- ▶ If F_1 and F_2 are formulas, then so is $F_1 \star F_2$ where \star is any binary connective
- ▶ If F is a formula, then so is $\neg F$
- ▶ If F is a formula and x a variable, so are $\forall x.F$ (asserts facts about *all* objects) and $\exists x.F$ (asserts facts about *some* object)

Important Reminder

- ▶ Predicates (e.g., $p(x)$) and function terms (e.g., $f(x)$) look similar, but they are very different!

Important Reminder

- ▶ Predicates (e.g., $p(x)$) and function terms (e.g., $f(x)$) look similar, but they are very different!
- ▶ Function terms can be nested within each other and inside relation constants: $f(f(x)), p(f(x)), \dots$

Important Reminder

- ▶ Predicates (e.g., $p(x)$) and function terms (e.g., $f(x)$) look similar, but they are very different!
- ▶ Function terms can be nested within each other and inside relation constants: $f(f(x)), p(f(x)), \dots$
- ▶ Predicates such as $p(x)$ cannot be nested within function terms or other predicates!

Important Reminder

- ▶ Predicates (e.g., $p(x)$) and function terms (e.g., $f(x)$) look similar, but they are very different!
- ▶ Function terms can be nested within each other and inside relation constants: $f(f(x)), p(f(x)), \dots$
- ▶ Predicates such as $p(x)$ cannot be nested within function terms or other predicates!
- ▶ $f(p(x)), p(p(x))$ etc. not valid in FOL!

Quantifiers and Scoping

- ▶ The subformula embedded inside a quantifier is called the **scope** of that quantifier.

Quantifiers and Scoping

- ▶ The subformula embedded inside a quantifier is called the **scope** of that quantifier.
- ▶ Example: $\forall y.((\forall x.p(x)) \rightarrow q(x, y))$

Quantifiers and Scoping

- ▶ The subformula embedded inside a quantifier is called the **scope** of that quantifier.
- ▶ Example: $\forall y.((\forall x.p(x)) \rightarrow q(x, y))$
- ▶ An occurrence of a variable is **bound** if it is in the scope of some quantifier.

Quantifiers and Scoping

- ▶ The subformula embedded inside a quantifier is called the **scope** of that quantifier.
- ▶ Example: $\forall y.((\forall x.p(x)) \rightarrow q(x, y))$
- ▶ An occurrence of a variable is **bound** if it is in the scope of some quantifier.
- ▶ An occurrence of a variable is **free** if it is not in the scope of any quantifiers.

Free vs. Bound Variable Example

- ▶ Consider the formula:

$$\forall y.((\forall x.p(x)) \rightarrow q(x, y))$$

Free vs. Bound Variable Example

- ▶ Consider the formula:

$$\forall y.((\forall x.p(x)) \rightarrow q(x, y))$$

- ▶ Is variable y bound or free?

Free vs. Bound Variable Example

- ▶ Consider the formula:

$$\forall y.((\forall x.p(x)) \rightarrow q(x, y))$$

- ▶ Is variable y bound or free? **bound**

Free vs. Bound Variable Example

- ▶ Consider the formula:

$$\forall y.((\forall x.p(x)) \rightarrow q(x, y))$$

- ▶ Is variable y bound or free? **bound**
- ▶ Is first occurrence of x bound or free?

Free vs. Bound Variable Example

- ▶ Consider the formula:

$$\forall y.((\forall x.p(x)) \rightarrow q(x, y))$$

- ▶ Is variable y bound or free? **bound**
- ▶ Is first occurrence of x bound or free? **bound**

Free vs. Bound Variable Example

- ▶ Consider the formula:

$$\forall y.((\forall x.p(x)) \rightarrow q(x, y))$$

- ▶ Is variable y bound or free? **bound**
- ▶ Is first occurrence of x bound or free? **bound**
- ▶ What about second occurrence of x ?

Free vs. Bound Variable Example

- ▶ Consider the formula:

$$\forall y.((\forall x.p(x)) \rightarrow q(x, y))$$

- ▶ Is variable y bound or free? **bound**
- ▶ Is first occurrence of x bound or free? **bound**
- ▶ What about second occurrence of x ? **free**

Closed, Open, and Ground Formulas

- ▶ A formula with no free variables is called a **closed** formula.

Closed, Open, and Ground Formulas

- ▶ A formula with no free variables is called a **closed** formula.
- ▶ A closed formula is also called a **sentence**.

Closed, Open, and Ground Formulas

- ▶ A formula with no free variables is called a **closed** formula.
- ▶ A closed formula is also called a **sentence**.
- ▶ A formula containing free variables is said to be **open**.

Closed, Open, and Ground Formulas

- ▶ A formula with no free variables is called a **closed** formula.
- ▶ A closed formula is also called a **sentence**.
- ▶ A formula containing free variables is said to be **open**.
- ▶ Example: Is the formula $\forall y.((\forall x.p(x)) \rightarrow q(x, y))$ closed or open?

Closed, Open, and Ground Formulas

- ▶ A formula with no free variables is called a **closed** formula.
- ▶ A closed formula is also called a **sentence**.
- ▶ A formula containing free variables is said to be **open**.
- ▶ Example: Is the formula $\forall y.((\forall x.p(x)) \rightarrow q(x, y))$ closed or open? **open**

Closed, Open, and Ground Formulas

- ▶ A formula with no free variables is called a **closed** formula.
- ▶ A closed formula is also called a **sentence**.
- ▶ A formula containing free variables is said to be **open**.
- ▶ Example: Is the formula $\forall y.((\forall x.p(x)) \rightarrow q(x, y))$ closed or open? **open**
- ▶ Is the formula $\forall y.((\forall x.p(x)) \rightarrow (\exists x.q(x, y)))$ a sentence?

Closed, Open, and Ground Formulas

- ▶ A formula with no free variables is called a **closed** formula.
- ▶ A closed formula is also called a **sentence**.
- ▶ A formula containing free variables is said to be **open**.
- ▶ Example: Is the formula $\forall y.((\forall x.p(x)) \rightarrow q(x, y))$ closed or open? **open**
- ▶ Is the formula $\forall y.((\forall x.p(x)) \rightarrow (\exists x.q(x, y)))$ a sentence? **yes**

Closed, Open, and Ground Formulas

- ▶ A formula with no free variables is called a **closed** formula.
- ▶ A closed formula is also called a **sentence**.
- ▶ A formula containing free variables is said to be **open**.
- ▶ Example: Is the formula $\forall y.((\forall x.p(x)) \rightarrow q(x, y))$ closed or open? **open**
- ▶ Is the formula $\forall y.((\forall x.p(x)) \rightarrow (\exists x.q(x, y)))$ a sentence? **yes**
- ▶ A formula is said to be **ground** if it contains no variables.

Closed, Open, and Ground Formulas

- ▶ A formula with no free variables is called a **closed** formula.
- ▶ A closed formula is also called a **sentence**.
- ▶ A formula containing free variables is said to be **open**.
- ▶ Example: Is the formula $\forall y.((\forall x.p(x)) \rightarrow q(x, y))$ closed or open? **open**
- ▶ Is the formula $\forall y.((\forall x.p(x)) \rightarrow (\exists x.q(x, y)))$ a sentence? **yes**
- ▶ A formula is said to be **ground** if it contains no variables.
- ▶ Example: $p(a, f(b)) \rightarrow q(c)$ is ground.

Closed, Open, and Ground Formulas

- ▶ A formula with no free variables is called a **closed** formula.
- ▶ A closed formula is also called a **sentence**.
- ▶ A formula containing free variables is said to be **open**.
- ▶ Example: Is the formula $\forall y.((\forall x.p(x)) \rightarrow q(x, y))$ closed or open? **open**
- ▶ Is the formula $\forall y.((\forall x.p(x)) \rightarrow (\exists x.q(x, y)))$ a sentence? **yes**
- ▶ A formula is said to be **ground** if it contains no variables.
- ▶ Example: $p(a, f(b)) \rightarrow q(c)$ is ground.
- ▶ Is $\forall x.p(x)$ ground?

Closed, Open, and Ground Formulas

- ▶ A formula with no free variables is called a **closed** formula.
- ▶ A closed formula is also called a **sentence**.
- ▶ A formula containing free variables is said to be **open**.
- ▶ Example: Is the formula $\forall y.((\forall x.p(x)) \rightarrow q(x, y))$ closed or open? **open**
- ▶ Is the formula $\forall y.((\forall x.p(x)) \rightarrow (\exists x.q(x, y)))$ a sentence? **yes**
- ▶ A formula is said to be **ground** if it contains no variables.
- ▶ Example: $p(a, f(b)) \rightarrow q(c)$ is ground.
- ▶ Is $\forall x.p(x)$ ground? **no**

A Few Examples

- ▶ To give some intuition about meaning of formula in FOL, some examples:

A Few Examples

- ▶ To give some intuition about meaning of formula in FOL, some examples:
- ▶ Given a relation constant **friend** of arity 2, how do we express the following in FOL?

A Few Examples

- ▶ To give some intuition about meaning of formula in FOL, some examples:
- ▶ Given a relation constant **friend** of arity 2, how do we express the following in FOL?
- ▶ *“Everyone has a friend”*

A Few Examples

- ▶ To give some intuition about meaning of formula in FOL, some examples:
- ▶ Given a relation constant **friend** of arity 2, how do we express the following in FOL?
- ▶ *"Everyone has a friend"*
- ▶ $\forall x. \exists y. \text{friend}(x, y)$

A Few Examples

- ▶ To give some intuition about meaning of formula in FOL, some examples:
- ▶ Given a relation constant **friend** of arity 2, how do we express the following in FOL?
- ▶ *"Everyone has a friend"*
- ▶ $\forall x. \exists y. \text{friend}(x, y)$
- ▶ *"There is someone who is friends with everyone"*

A Few Examples

- ▶ To give some intuition about meaning of formula in FOL, some examples:
- ▶ Given a relation constant **friend** of arity 2, how do we express the following in FOL?
- ▶ *"Everyone has a friend"*
- ▶ $\forall x. \exists y. \text{friend}(x, y)$
- ▶ *"There is someone who is friends with everyone"*
- ▶ $\exists x. \forall y. \text{friend}(x, y)$

A Few Examples

- ▶ To give some intuition about meaning of formula in FOL, some examples:
- ▶ Given a relation constant **friend** of arity 2, how do we express the following in FOL?
- ▶ *"Everyone has a friend"*
- ▶ $\forall x. \exists y. \text{friend}(x, y)$
- ▶ *"There is someone who is friends with everyone"*
- ▶ $\exists x. \forall y. \text{friend}(x, y)$
- ▶ If you flip the quantifiers, completely different meaning!

More Friendship Examples

- ▶ Given unary relation constants `student` and `atWM` and binary relation constant `friend`, how do we say the following in FOL?

More Friendship Examples

- ▶ Given unary relation constants `student` and `atWM` and binary relation constant `friend`, how do we say the following in FOL?
- ▶ *"Every student has a friend."*

More Friendship Examples

- ▶ Given unary relation constants `student` and `atWM` and binary relation constant `friend`, how do we say the following in FOL?
- ▶ *"Every student has a friend."*
- ▶ $\forall x. \text{student}(x) \rightarrow \exists y. \text{friend}(x, y)$

More Friendship Examples

- ▶ Given unary relation constants `student` and `atWM` and binary relation constant `friend`, how do we say the following in FOL?
- ▶ *"Every student has a friend."*
- ▶ $\forall x. \text{student}(x) \rightarrow \exists y. \text{friend}(x, y)$
- ▶ *"No student has a friend unless he/she is at William & Mary."*

More Friendship Examples

- ▶ Given unary relation constants `student` and `atWM` and binary relation constant `friend`, how do we say the following in FOL?
- ▶ *"Every student has a friend."*
- ▶ $\forall x. \text{student}(x) \rightarrow \exists y. \text{friend}(x, y)$
- ▶ *"No student has a friend unless he/she is at William & Mary."*
- ▶ $\forall x. ((\text{student}(x) \wedge \neg \text{atWM}(x)) \rightarrow \neg \exists y. \text{friend}(x, y))$

Even More Friendship Examples

- ▶ “William & Mary students only have friends at William & Mary.”

Even More Friendship Examples

- ▶ “William & Mary students only have friends at William & Mary.”
- ▶ $\forall x. \forall y. (student(x) \wedge atWM(x) \wedge friends(x, y)) \rightarrow atWM(y)$

Even More Friendship Examples

- ▶ “William & Mary students only have friends at William & Mary.”

- ▶ $\forall x. \forall y. (student(x) \wedge atWM(x) \wedge friends(x, y)) \rightarrow atWM(y)$

- ▶ Another way of saying this:

$$\forall x. (student(x) \wedge atWM(x)) \rightarrow (\forall y. (friends(x, y) \rightarrow atWM(y)))$$

Even More Friendship Examples

- ▶ “William & Mary students only have friends at William & Mary.”

- ▶ $\forall x. \forall y. (student(x) \wedge atWM(x) \wedge friends(x, y)) \rightarrow atWM(y)$

- ▶ Another way of saying this:

$$\forall x. (student(x) \wedge atWM(x)) \rightarrow (\forall y. (friends(x, y) \rightarrow atWM(y)))$$

- ▶ These two formulas are actually semantically equivalent

More Friendship Examples

- ▶ “William & Mary students are all friends with each other.”

More Friendship Examples

- ▶ “William & Mary students are all friends with each other.”

$$\forall x. \forall y. \quad ((\textit{student}(x) \wedge \textit{atWM}(x) \wedge \textit{student}(y) \wedge \textit{atWM}(y)) \\ \rightarrow \textit{friends}(x, y))$$

More Friendship Examples

- ▶ “William & Mary students are all friends with each other.”

$$\forall x. \forall y. \quad ((student(x) \wedge atWM(x) \wedge student(y) \wedge atWM(y)) \\ \rightarrow friends(x, y))$$

- ▶ “Every William & Mary student has friends outside of their own college”

More Friendship Examples

- ▶ “William & Mary students are all friends with each other.”

$$\forall x. \forall y. \quad ((student(x) \wedge atWM(x) \wedge student(y) \wedge atWM(y)) \rightarrow friends(x, y))$$

- ▶ “Every William & Mary student has friends outside of their own college”

$$\forall x. \quad ((atWM(x) \wedge student(x)) \rightarrow \exists y. (friends(x, y) \wedge \neg atWM(y)))$$

Mathematical Theorems in FOL

- **Fermat's Last Theorem:** No three positive integers a, b, c satisfy the equation $a^n + b^n = c^n$ for any integer n greater than 2.

Mathematical Theorems in FOL

- ▶ **Fermat's Last Theorem:** No three positive integers a, b, c satisfy the equation $a^n + b^n = c^n$ for any integer n greater than 2.
- ▶ **Aside:** French (amateur) mathematician Pierre de Fermat wrote this in 1637 in the margin of Arithmetica, and claimed his proof is too large to fit in the margin.

Mathematical Theorems in FOL

- ▶ **Fermat's Last Theorem:** No three positive integers a, b, c satisfy the equation $a^n + b^n = c^n$ for any integer n greater than 2.
- ▶ **Aside:** French (amateur) mathematician Pierre de Fermat wrote this in 1637 in the margin of Arithmetica, and claimed his proof is too large to fit in the margin.
- ▶ Could not be proved until British mathematician Andrew Wiles proved in 1995 using properties of elliptical curves.

Mathematical Theorems in FOL

- ▶ **Fermat's Last Theorem:** No three positive integers a, b, c satisfy the equation $a^n + b^n = c^n$ for any integer n greater than 2.
- ▶ **Aside:** French (amateur) mathematician Pierre de Fermat wrote this in 1637 in the margin of Arithmetica, and claimed his proof is too large to fit in the margin.
- ▶ Could not be proved until British mathematician Andrew Wiles proved in 1995 using properties of elliptical curves.
- ▶ How do we express Fermat's last theorem in FOL (given a function constant \wedge)?

Mathematical Theorems in FOL

- ▶ **Fermat's Last Theorem:** No three positive integers a, b, c satisfy the equation $a^n + b^n = c^n$ for any integer n greater than 2.
- ▶ **Aside:** French (amateur) mathematician Pierre de Fermat wrote this in 1637 in the margin of Arithmetica, and claimed his proof is too large to fit in the margin.
- ▶ Could not be proved until British mathematician Andrew Wiles proved in 1995 using properties of elliptical curves.
- ▶ How do we express Fermat's last theorem in FOL (given a function constant \wedge)?

$$\forall n. n > 2 \rightarrow \neg \exists a, b, c. a > 0 \wedge b > 0 \wedge c > 0 \wedge a^n + b^n = c^n$$

One Last Example

- ▶ Given binary relation constant `friend`, how do we say this in FOL?

One Last Example

- ▶ Given binary relation constant **friend**, how do we say this in FOL?
- ▶ *“Every pair of friends has something in common”*

One Last Example

- ▶ Given binary relation constant **friend**, how do we say this in FOL?
- ▶ *“Every pair of friends has something in common”*
- ▶ This cannot be expressed in FOL because it requires quantification over relation constants!

One Last Example

- ▶ Given binary relation constant **friend**, how do we say this in FOL?
- ▶ *“Every pair of friends has something in common”*
- ▶ This cannot be expressed in FOL because it requires quantification over relation constants!
- ▶ But it can, however, be expressed in **second-order logic**:

$$\forall x, y. \text{friend}(x, y) \rightarrow \exists p. p(x) \wedge p(y)$$

Semantics of First Order Logic

- ▶ In propositional logic, the concepts of interpretation, satisfiability, validity were all straightforward.

Semantics of First Order Logic

- ▶ In propositional logic, the concepts of interpretation, satisfiability, validity were all straightforward.
- ▶ In FOL, these concepts are a bit more involved . . .

Semantics of First Order Logic

- ▶ In propositional logic, the concepts of interpretation, satisfiability, validity were all straightforward.
- ▶ In FOL, these concepts are a bit more involved . . .
- ▶ To give semantics to FOL, we need to talk about a **universe of discourse** (also sometimes called just “universe” or “domain”)

Universe of Discourse

- ▶ A **universe of discourse** is a *non-empty* set of objects about which we want to say something

Universe of Discourse

- ▶ A **universe of discourse** is a *non-empty* set of objects about which we want to say something
- ▶ Universe of discourse can be finite, countably infinite, or uncountably infinite, but not empty

Universe of Discourse

- ▶ A **universe of discourse** is a *non-empty* set of objects about which we want to say something
- ▶ Universe of discourse can be finite, countably infinite, or uncountably infinite, but not empty
- ▶ Example universes:

Universe of Discourse

- ▶ A **universe of discourse** is a *non-empty* set of objects about which we want to say something
- ▶ Universe of discourse can be finite, countably infinite, or uncountably infinite, but not empty
- ▶ Example universes:
 - ▶ Set of non-negative integers:

Universe of Discourse

- ▶ A **universe of discourse** is a *non-empty* set of objects about which we want to say something
- ▶ Universe of discourse can be finite, countably infinite, or uncountably infinite, but not empty
- ▶ Example universes:
 - ▶ Set of non-negative integers: countably infinite

Universe of Discourse

- ▶ A **universe of discourse** is a *non-empty* set of objects about which we want to say something
- ▶ Universe of discourse can be finite, countably infinite, or uncountably infinite, but not empty
- ▶ Example universes:
 - ▶ Set of non-negative integers: countably infinite
 - ▶ Set of real numbers:

Universe of Discourse

- ▶ A **universe of discourse** is a *non-empty* set of objects about which we want to say something
- ▶ Universe of discourse can be finite, countably infinite, or uncountably infinite, but not empty
- ▶ Example universes:
 - ▶ Set of non-negative integers: countably infinite
 - ▶ Set of real numbers: uncountably infinite

Universe of Discourse

- ▶ A **universe of discourse** is a *non-empty* set of objects about which we want to say something
- ▶ Universe of discourse can be finite, countably infinite, or uncountably infinite, but not empty
- ▶ Example universes:
 - ▶ Set of non-negative integers: countably infinite
 - ▶ Set of real numbers: uncountably infinite
 - ▶ The set of suits in playing cards $\{\clubsuit, \diamond, \heartsuit, \spadesuit\}$: finite

Universe of Discourse

- ▶ A **universe of discourse** is a *non-empty* set of objects about which we want to say something
- ▶ Universe of discourse can be finite, countably infinite, or uncountably infinite, but not empty
- ▶ Example universes:
 - ▶ Set of non-negative integers: countably infinite
 - ▶ Set of real numbers: uncountably infinite
 - ▶ The set of suits in playing cards $\{\clubsuit, \diamond, \heartsuit, \spadesuit\}$: finite
 - ▶ Students in this class: finite

First-Order Interpretations

- ▶ An **interpretation** for a first order language $L(C, F, R)$ is a mapping I from C, F, R to objects universe of discourse U

First-Order Interpretations

- ▶ An **interpretation** for a first order language $L(C, F, R)$ is a mapping I from C, F, R to objects universe of discourse U
- ▶ I maps every $c \in C$ to some member of U : $I(c) \in U$

First-Order Interpretations

- ▶ An **interpretation** for a first order language $L(C, F, R)$ is a mapping I from C, F, R to objects universe of discourse U
- ▶ I maps every $c \in C$ to some member of U : $I(c) \in U$
- ▶ I maps every n -ary function constant $f \in F$ to an n -ary function $f^I : U^n \rightarrow U$

First-Order Interpretations

- ▶ An **interpretation** for a first order language $L(C, F, R)$ is a mapping I from C, F, R to objects universe of discourse U
- ▶ I maps every $c \in C$ to some member of U : $I(c) \in U$
- ▶ I maps every n -ary function constant $f \in F$ to an n -ary function $f^I : U^n \rightarrow U$
- ▶ I maps every n -ary relation constant $p \in R$ to an n -ary relation p^I such that $p^I \subseteq U^n$

First-Order Interpretations

- ▶ An **interpretation** for a first order language $L(C, F, R)$ is a mapping I from C, F, R to objects universe of discourse U
- ▶ I maps every $c \in C$ to some member of U : $I(c) \in U$
- ▶ I maps every n -ary function constant $f \in F$ to an n -ary function $f^I : U^n \rightarrow U$
- ▶ I maps every n -ary relation constant $p \in R$ to an n -ary relation p^I such that $p^I \subseteq U^n$
- ▶ **Observe:** A first-order interpretation does not talk about variables (only constants)

An Example

- ▶ Consider the first order language containing object constants $\{a, b, c\}$, binary function constant f , and ternary relation constant r .

An Example

- ▶ Consider the first order language containing object constants $\{a, b, c\}$, binary function constant f , and ternary relation constant r .
- ▶ Universe of discourse: $U = \{1, 2, 3\}$

An Example

- ▶ Consider the first order language containing object constants $\{a, b, c\}$, binary function constant f , and ternary relation constant r .
- ▶ Universe of discourse: $U = \{1, 2, 3\}$
- ▶ Possible interpretation I :

$$I(a) = 1, I(b) = 2, I(c) = 2$$

An Example

- ▶ Consider the first order language containing object constants $\{a, b, c\}$, binary function constant f , and ternary relation constant r .
- ▶ Universe of discourse: $U = \{1, 2, 3\}$
- ▶ Possible interpretation I :

$$I(a) = 1, I(b) = 2, I(c) = 2$$

$$I(f) = \{1 \rightarrow 2, 2 \rightarrow 3, 3 \rightarrow 3\}$$

An Example

- ▶ Consider the first order language containing object constants $\{a, b, c\}$, binary function constant f , and ternary relation constant r .
- ▶ Universe of discourse: $U = \{1, 2, 3\}$
- ▶ Possible interpretation I :

$$I(a) = 1, I(b) = 2, I(c) = 2$$

$$I(f) = \{1 \rightarrow 2, 2 \rightarrow 3, 3 \rightarrow 3\}$$

$$I(r) = \{\langle 1, 2, 1 \rangle, \langle 2, 2, 1 \rangle\}$$

An Example

- ▶ Consider the first order language containing object constants $\{a, b, c\}$, binary function constant f , and ternary relation constant r .
- ▶ Universe of discourse: $U = \{1, 2, 3\}$
- ▶ Possible interpretation I :

$$I(a) = 1, I(b) = 2, I(c) = 2$$

$$I(f) = \{1 \rightarrow 2, 2 \rightarrow 3, 3 \rightarrow 3\}$$

$$I(r) = \{\langle 1, 2, 1 \rangle, \langle 2, 2, 1 \rangle\}$$

- ▶ **Observe:** Different object constants do not have to map to distinct objects in U !

Another Example

- ▶ Consider the universe of discourse $U = \{\square, \triangle\}$

Another Example

- ▶ Consider the universe of discourse $U = \{\square, \triangle\}$
- ▶ For an object constant a , what are the possible interpretations?

Another Example

- ▶ Consider the universe of discourse $U = \{\square, \triangle\}$
- ▶ For an object constant a , what are the possible interpretations? $I(a) = \square$
or $I(a) = \triangle$

Another Example

- ▶ Consider the universe of discourse $U = \{\square, \triangle\}$
- ▶ For an object constant a , what are the possible interpretations? $I(a) = \square$
or $I(a) = \triangle$
- ▶ For a unary function constant f , what are the possible interpretations?

Another Example

- ▶ Consider the universe of discourse $U = \{\square, \triangle\}$
- ▶ For an object constant a , what are the possible interpretations? $I(a) = \square$
or $I(a) = \triangle$
- ▶ For a unary function constant f , what are the possible interpretations?
- ▶ $I(f) = \{\square \mapsto \square, \triangle \mapsto \square\}$, or $I(f) = \{\square \mapsto \square, \triangle \mapsto \triangle\}$, or
 $I(f) = \{\square \mapsto \triangle, \triangle \mapsto \square\}$ or $I(f) = \{\square \mapsto \triangle, \triangle \mapsto \triangle\}$

Another Example

- ▶ Consider the universe of discourse $U = \{\square, \triangle\}$
- ▶ For an object constant a , what are the possible interpretations? $I(a) = \square$
or $I(a) = \triangle$
- ▶ For a unary function constant f , what are the possible interpretations?
- ▶ $I(f) = \{\square \mapsto \square, \triangle \mapsto \square\}$, or $I(f) = \{\square \mapsto \square, \triangle \mapsto \triangle\}$, or
 $I(f) = \{\square \mapsto \triangle, \triangle \mapsto \square\}$ or $I(f) = \{\square \mapsto \triangle, \triangle \mapsto \triangle\}$
- ▶ For a unary relation constant r , what are all the possible interpretations?

Another Example

- ▶ Consider the universe of discourse $U = \{\square, \triangle\}$
- ▶ For an object constant a , what are the possible interpretations? $I(a) = \square$
or $I(a) = \triangle$
- ▶ For a unary function constant f , what are the possible interpretations?
- ▶ $I(f) = \{\square \mapsto \square, \triangle \mapsto \square\}$, or $I(f) = \{\square \mapsto \square, \triangle \mapsto \triangle\}$, or
 $I(f) = \{\square \mapsto \triangle, \triangle \mapsto \square\}$ or $I(f) = \{\square \mapsto \triangle, \triangle \mapsto \triangle\}$
- ▶ For a unary relation constant r , what are all the possible interpretations?
- ▶ $I(r) = \{\}$, $I(r) = \{\langle \square \rangle\}$, or $I(r) = \{\langle \triangle \rangle\}$, or $I(r) = \{\langle \square \rangle, \langle \triangle \rangle\}$

First-Order Structures and Variable Assignments

- ▶ A **structure** $S = \langle U, I \rangle$ for a first order language consists of a universe of discourse of U and an interpretation I .

First-Order Structures and Variable Assignments

- ▶ A **structure** $S = \langle U, I \rangle$ for a first order language consists of a universe of discourse of U and an interpretation I .
- ▶ This is sometimes also called an **algebra**.

First-Order Structures and Variable Assignments

- ▶ A **structure** $S = \langle U, I \rangle$ for a first order language consists of a universe of discourse of U and an interpretation I .
- ▶ This is sometimes also called an **algebra**.
- ▶ A **variable assignment** σ (or assignment) to a FOL formula ϕ in a structure $S = \langle U, I \rangle$ is a mapping from variables in ϕ to an element of U .

First-Order Structures and Variable Assignments

- ▶ A **structure** $S = \langle U, I \rangle$ for a first order language consists of a universe of discourse of U and an interpretation I .
- ▶ This is sometimes also called an **algebra**.
- ▶ A **variable assignment** σ (or assignment) to a FOL formula ϕ in a structure $S = \langle U, I \rangle$ is a mapping from variables in ϕ to an element of U .
- ▶ **Example:** Given $U = \{\square, \triangle\}$, a possible variable assignment for x :
 $\sigma(x) = \triangle$

First-Order Structures and Variable Assignments

- ▶ A **structure** $S = \langle U, I \rangle$ for a first order language consists of a universe of discourse of U and an interpretation I .
- ▶ This is sometimes also called an **algebra**.
- ▶ A **variable assignment** σ (or assignment) to a FOL formula ϕ in a structure $S = \langle U, I \rangle$ is a mapping from variables in ϕ to an element of U .
- ▶ **Example:** Given $U = \{\square, \triangle\}$, a possible variable assignment for x :
 $\sigma(x) = \triangle$
- ▶ **Observe:** σ does not map variables to object constants but to objects in U !

Evaluation of Terms

- ▶ We define how to evaluate a term t under interpretation I and assignment σ , written $\langle I, \sigma \rangle(t)$

Evaluation of Terms

- ▶ We define how to evaluate a term t under interpretation I and assignment σ , written $\langle I, \sigma \rangle(t)$
- ▶ **Object constants:** $\langle I, \sigma \rangle(a) = I(a)$

Evaluation of Terms

- ▶ We define how to evaluate a term t under interpretation I and assignment σ , written $\langle I, \sigma \rangle(t)$
- ▶ Object constants: $\langle I, \sigma \rangle(a) = I(a)$
- ▶ Variable terms: $\langle I, \sigma \rangle(v) = \sigma(v)$

Evaluation of Terms

- ▶ We define how to evaluate a term t under interpretation I and assignment σ , written $\langle I, \sigma \rangle(t)$
- ▶ Object constants: $\langle I, \sigma \rangle(a) = I(a)$
- ▶ Variable terms: $\langle I, \sigma \rangle(v) = \sigma(v)$
- ▶ Function terms:

$$\langle I, \sigma \rangle(f(t_1, \dots, t_k)) = I(f)(\langle I, \sigma \rangle(t_1), \dots, \langle I, \sigma \rangle(t_k))$$

Example: Evaluation of Terms

- ▶ Consider a first-order language containing object constants a, b and binary function f

Example: Evaluation of Terms

- ▶ Consider a first-order language containing object constants a, b and binary function f
- ▶ Consider universe $\{1, 2\}$ and interpretation I :

$$\begin{aligned} I(a) &= 1 \quad I(b) = 2 \\ I(f) &= \{\langle 1, 1 \rangle \mapsto 2, \langle 1, 2 \rangle \mapsto 2, \langle 2, 1 \rangle \mapsto 1, \langle 2, 2 \rangle \mapsto 1\} \end{aligned}$$

Example: Evaluation of Terms

- ▶ Consider a first-order language containing object constants a, b and binary function f
- ▶ Consider universe $\{1, 2\}$ and interpretation I :

$$\begin{aligned} I(a) &= 1 \quad I(b) = 2 \\ I(f) &= \{\langle 1, 1 \rangle \mapsto 2, \langle 1, 2 \rangle \mapsto 2, \langle 2, 1 \rangle \mapsto 1, \langle 2, 2 \rangle \mapsto 1\} \end{aligned}$$

- ▶ Consider variable assignment $\sigma : \{x \mapsto 2, y \mapsto 1\}$

Example: Evaluation of Terms

- ▶ Consider a first-order language containing object constants a, b and binary function f
- ▶ Consider universe $\{1, 2\}$ and interpretation I :

$$\begin{aligned} I(a) &= 1 \quad I(b) = 2 \\ I(f) &= \{\langle 1, 1 \rangle \mapsto 2, \langle 1, 2 \rangle \mapsto 2, \langle 2, 1 \rangle \mapsto 1, \langle 2, 2 \rangle \mapsto 1\} \end{aligned}$$

- ▶ Consider variable assignment $\sigma : \{x \mapsto 2, y \mapsto 1\}$
- ▶ Under I and σ , what do these terms evaluate to?

$$f(a, y) \quad =$$

Example: Evaluation of Terms

- ▶ Consider a first-order language containing object constants a, b and binary function f
- ▶ Consider universe $\{1, 2\}$ and interpretation I :

$$\begin{aligned} I(a) &= 1 \quad I(b) = 2 \\ I(f) &= \{\langle 1, 1 \rangle \mapsto 2, \langle 1, 2 \rangle \mapsto 2, \langle 2, 1 \rangle \mapsto 1, \langle 2, 2 \rangle \mapsto 1\} \end{aligned}$$

- ▶ Consider variable assignment $\sigma : \{x \mapsto 2, y \mapsto 1\}$
- ▶ Under I and σ , what do these terms evaluate to?

$$f(a, y) \quad = \quad 2$$

Example: Evaluation of Terms

- ▶ Consider a first-order language containing object constants a, b and binary function f
- ▶ Consider universe $\{1, 2\}$ and interpretation I :

$$\begin{aligned} I(a) &= 1 & I(b) &= 2 \\ I(f) &= \{\langle 1, 1 \rangle \mapsto 2, \langle 1, 2 \rangle \mapsto 2, \langle 2, 1 \rangle \mapsto 1, \langle 2, 2 \rangle \mapsto 1\} \end{aligned}$$

- ▶ Consider variable assignment $\sigma : \{x \mapsto 2, y \mapsto 1\}$
- ▶ Under I and σ , what do these terms evaluate to?

$$\begin{array}{rcl} f(a, y) & = & 2 \\ f(x, b) & = & \end{array}$$

Example: Evaluation of Terms

- ▶ Consider a first-order language containing object constants a, b and binary function f
- ▶ Consider universe $\{1, 2\}$ and interpretation I :

$$\begin{aligned} I(a) &= 1 & I(b) &= 2 \\ I(f) &= \{\langle 1, 1 \rangle \mapsto 2, \langle 1, 2 \rangle \mapsto 2, \langle 2, 1 \rangle \mapsto 1, \langle 2, 2 \rangle \mapsto 1\} \end{aligned}$$

- ▶ Consider variable assignment $\sigma : \{x \mapsto 2, y \mapsto 1\}$
- ▶ Under I and σ , what do these terms evaluate to?

$$\begin{aligned} f(a, y) &= 2 \\ f(x, b) &= 1 \end{aligned}$$

Example: Evaluation of Terms

- ▶ Consider a first-order language containing object constants a, b and binary function f
- ▶ Consider universe $\{1, 2\}$ and interpretation I :

$$\begin{aligned} I(a) &= 1 & I(b) &= 2 \\ I(f) &= \{ \langle 1, 1 \rangle \mapsto 2, \langle 1, 2 \rangle \mapsto 2, \langle 2, 1 \rangle \mapsto 1, \langle 2, 2 \rangle \mapsto 1 \} \end{aligned}$$

- ▶ Consider variable assignment $\sigma : \{x \mapsto 2, y \mapsto 1\}$
- ▶ Under I and σ , what do these terms evaluate to?

$$\begin{aligned} f(a, y) &= 2 \\ f(x, b) &= 1 \\ f(f(x, b), f(a, y)) &= \end{aligned}$$

Example: Evaluation of Terms

- ▶ Consider a first-order language containing object constants a, b and binary function f
- ▶ Consider universe $\{1, 2\}$ and interpretation I :

$$\begin{aligned} I(a) &= 1 & I(b) &= 2 \\ I(f) &= \{\langle 1, 1 \rangle \mapsto 2, \langle 1, 2 \rangle \mapsto 2, \langle 2, 1 \rangle \mapsto 1, \langle 2, 2 \rangle \mapsto 1\} \end{aligned}$$

- ▶ Consider variable assignment $\sigma : \{x \mapsto 2, y \mapsto 1\}$
- ▶ Under I and σ , what do these terms evaluate to?

$$\begin{aligned} f(a, y) &= 2 \\ f(x, b) &= 1 \\ f(f(x, b), f(a, y)) &= 2 \end{aligned}$$

Evaluation of Formulas, Notation

- ▶ We define evaluation of formula F under structure $S = \langle U, I \rangle$ and variable assignment σ .

Evaluation of Formulas, Notation

- ▶ We define evaluation of formula F under structure $S = \langle U, I \rangle$ and variable assignment σ .
- ▶ If F evaluates to true under U, I, σ , we write $U, I, \sigma \models F$

Evaluation of Formulas, Notation

- ▶ We define evaluation of formula F under structure $S = \langle U, I \rangle$ and variable assignment σ .
- ▶ If F evaluates to true under U, I, σ , we write $U, I, \sigma \models F$
- ▶ If F evaluates to false under U, I, σ , we write $U, I, \sigma \not\models F$

Evaluation of Formulas, Notation

- ▶ We define evaluation of formula F under structure $S = \langle U, I \rangle$ and variable assignment σ .
- ▶ If F evaluates to true under U, I, σ , we write $U, I, \sigma \models F$
- ▶ If F evaluates to false under U, I, σ , we write $U, I, \sigma \not\models F$
- ▶ We define the semantics of \models inductively.

Evaluation of Formulas, Bases Cases

- Base case I:

$$U, I, \sigma \models \top \quad U, I, \sigma \not\models \perp$$

Evaluation of Formulas, Bases Cases

- ▶ Base case I:

$$U, I, \sigma \models \top \quad U, I, \sigma \not\models \perp$$

- ▶ Base case II:

$$U, I, \sigma \models p(t_1, \dots, t_k) \text{ iff } \langle \langle I, \sigma \rangle(t_1), \dots, \langle I, \sigma \rangle(t_k) \rangle \in I(p)$$

Example I: Evaluation of Formulas

- Consider a first-order language containing object constants a, b and unary function f , and binary relation constant p

Example I: Evaluation of Formulas

- ▶ Consider a first-order language containing object constants a, b and unary function f , and binary relation constant p
- ▶ Consider universe $\{\star, \bullet\}$ and interpretation I :

$$\begin{aligned}I(a) &= \star & I(b) &= \bullet \\I(f) &= \{\star \mapsto \bullet, \bullet \mapsto \star\} \\I(p) &= \{\langle \bullet, \star \rangle, \langle \bullet, \bullet \rangle\}\end{aligned}$$

Example I: Evaluation of Formulas

- ▶ Consider a first-order language containing object constants a, b and unary function f , and binary relation constant p
- ▶ Consider universe $\{\star, \bullet\}$ and interpretation I :

$$\begin{aligned}I(a) &= \star & I(b) &= \bullet \\I(f) &= \{\star \mapsto \bullet, \bullet \mapsto \star\} \\I(p) &= \{\langle \bullet, \star \rangle, \langle \bullet, \bullet \rangle\}\end{aligned}$$

- ▶ Consider variable assignment $\sigma : \{x \mapsto \star\}$

Example I: Evaluation of Formulas

- ▶ Consider a first-order language containing object constants a, b and unary function f , and binary relation constant p
- ▶ Consider universe $\{\star, \bullet\}$ and interpretation I :

$$\begin{aligned}I(a) &= \star & I(b) &= \bullet \\I(f) &= \{\star \mapsto \bullet, \bullet \mapsto \star\} \\I(p) &= \{\langle \bullet, \star \rangle, \langle \bullet, \bullet \rangle\}\end{aligned}$$

- ▶ Consider variable assignment $\sigma : \{x \mapsto \star\}$
- ▶ Under U, I and σ , what do these formulas evaluate to?

$$p(f(b), f(x)) \quad =$$

Example I: Evaluation of Formulas

- ▶ Consider a first-order language containing object constants a, b and unary function f , and binary relation constant p
- ▶ Consider universe $\{\star, \bullet\}$ and interpretation I :

$$\begin{aligned}I(a) &= \star & I(b) &= \bullet \\I(f) &= \{\star \mapsto \bullet, \bullet \mapsto \star\} \\I(p) &= \{\langle \bullet, \star \rangle, \langle \bullet, \bullet \rangle\}\end{aligned}$$

- ▶ Consider variable assignment $\sigma : \{x \mapsto \star\}$
- ▶ Under U, I and σ , what do these formulas evaluate to?

$$p(f(b), f(x)) \quad = \quad \textit{false}$$

Example I: Evaluation of Formulas

- ▶ Consider a first-order language containing object constants a, b and unary function f , and binary relation constant p
- ▶ Consider universe $\{\star, \bullet\}$ and interpretation I :

$$\begin{aligned}I(a) &= \star & I(b) &= \bullet \\I(f) &= \{\star \mapsto \bullet, \bullet \mapsto \star\} \\I(p) &= \{\langle \bullet, \star \rangle, \langle \bullet, \bullet \rangle\}\end{aligned}$$

- ▶ Consider variable assignment $\sigma : \{x \mapsto \star\}$
- ▶ Under U, I and σ , what do these formulas evaluate to?

$$\begin{aligned}p(f(b), f(x)) &= \text{false} \\p(f(x), f(b)) &= \end{aligned}$$

Example I: Evaluation of Formulas

- ▶ Consider a first-order language containing object constants a, b and unary function f , and binary relation constant p
- ▶ Consider universe $\{\star, \bullet\}$ and interpretation I :

$$\begin{aligned}I(a) &= \star & I(b) &= \bullet \\I(f) &= \{\star \mapsto \bullet, \bullet \mapsto \star\} \\I(p) &= \{\langle \bullet, \star \rangle, \langle \bullet, \bullet \rangle\}\end{aligned}$$

- ▶ Consider variable assignment $\sigma : \{x \mapsto \star\}$
- ▶ Under U, I and σ , what do these formulas evaluate to?

$$\begin{aligned}p(f(b), f(x)) &= \text{false} \\p(f(x), f(b)) &= \text{true}\end{aligned}$$

Example I: Evaluation of Formulas

- ▶ Consider a first-order language containing object constants a, b and unary function f , and binary relation constant p
- ▶ Consider universe $\{\star, \bullet\}$ and interpretation I :

$$\begin{aligned}I(a) &= \star & I(b) &= \bullet \\I(f) &= \{\star \mapsto \bullet, \bullet \mapsto \star\} \\I(p) &= \{\langle \bullet, \star \rangle, \langle \bullet, \bullet \rangle\}\end{aligned}$$

- ▶ Consider variable assignment $\sigma : \{x \mapsto \star\}$
- ▶ Under U, I and σ , what do these formulas evaluate to?

$$\begin{aligned}p(f(b), f(x)) &= \text{false} \\p(f(x), f(b)) &= \text{true} \\p(a, f(x)) &= \end{aligned}$$

Example I: Evaluation of Formulas

- ▶ Consider a first-order language containing object constants a, b and unary function f , and binary relation constant p
- ▶ Consider universe $\{\star, \bullet\}$ and interpretation I :

$$\begin{aligned}I(a) &= \star & I(b) &= \bullet \\I(f) &= \{\star \mapsto \bullet, \bullet \mapsto \star\} \\I(p) &= \{\langle \bullet, \star \rangle, \langle \bullet, \bullet \rangle\}\end{aligned}$$

- ▶ Consider variable assignment $\sigma : \{x \mapsto \star\}$
- ▶ Under U, I and σ , what do these formulas evaluate to?

$$\begin{aligned}p(f(b), f(x)) &= \text{false} \\p(f(x), f(b)) &= \text{true} \\p(a, f(x)) &= \text{false}\end{aligned}$$

Evaluation of Formulas II

- ▶ Inductive semantics for boolean connectives:

Evaluation of Formulas II

- ▶ Inductive semantics for boolean connectives:

$$U, I, \sigma \models \neg F \quad \text{iff}$$

Evaluation of Formulas II

- ▶ Inductive semantics for boolean connectives:

$$U, I, \sigma \models \neg F \quad \text{iff } U, I, \sigma \not\models F$$

Evaluation of Formulas II

- ▶ Inductive semantics for boolean connectives:

$$\begin{array}{ll} U, I, \sigma \models \neg F & \text{iff } U, I, \sigma \not\models F \\ U, I, \sigma \models F_1 \wedge F_2 & \text{iff} \end{array}$$

Evaluation of Formulas II

- ▶ Inductive semantics for boolean connectives:

$$\begin{array}{ll} U, I, \sigma \models \neg F & \text{iff } U, I, \sigma \not\models F \\ U, I, \sigma \models F_1 \wedge F_2 & \text{iff } U, I, \sigma \models F_1 \text{ and } U, I, \sigma \models F_2 \end{array}$$

Evaluation of Formulas II

- ▶ Inductive semantics for boolean connectives:

$$\begin{array}{ll} U, I, \sigma \models \neg F & \text{iff } U, I, \sigma \not\models F \\ U, I, \sigma \models F_1 \wedge F_2 & \text{iff } U, I, \sigma \models F_1 \text{ and } U, I, \sigma \models F_2 \\ U, I, \sigma \models F_1 \vee F_2 & \text{iff} \end{array}$$

Evaluation of Formulas II

- ▶ Inductive semantics for boolean connectives:

$$\begin{array}{ll} U, I, \sigma \models \neg F & \text{iff } U, I, \sigma \not\models F \\ U, I, \sigma \models F_1 \wedge F_2 & \text{iff } U, I, \sigma \models F_1 \text{ and } U, I, \sigma \models F_2 \\ U, I, \sigma \models F_1 \vee F_2 & \text{iff } U, I, \sigma \models F_1 \text{ or } U, I, \sigma \models F_2 \end{array}$$

Evaluation of Formulas II

- ▶ Inductive semantics for boolean connectives:

$$\begin{array}{ll} U, I, \sigma \models \neg F & \text{iff } U, I, \sigma \not\models F \\ U, I, \sigma \models F_1 \wedge F_2 & \text{iff } U, I, \sigma \models F_1 \text{ and } U, I, \sigma \models F_2 \\ U, I, \sigma \models F_1 \vee F_2 & \text{iff } U, I, \sigma \models F_1 \text{ or } U, I, \sigma \models F_2 \\ U, I, \sigma \models F_1 \rightarrow F_2 & \text{iff,} \end{array}$$

Evaluation of Formulas II

- ▶ Inductive semantics for boolean connectives:

$$\begin{array}{ll} U, I, \sigma \models \neg F & \text{iff } U, I, \sigma \not\models F \\ U, I, \sigma \models F_1 \wedge F_2 & \text{iff } U, I, \sigma \models F_1 \text{ and } U, I, \sigma \models F_2 \\ U, I, \sigma \models F_1 \vee F_2 & \text{iff } U, I, \sigma \models F_1 \text{ or } U, I, \sigma \models F_2 \\ U, I, \sigma \models F_1 \rightarrow F_2 & \text{iff, } U, I, \sigma \not\models F_1 \text{ or } U, I, \sigma \models F_2 \end{array}$$

Evaluation of Formulas II

- ▶ Inductive semantics for boolean connectives:

$$\begin{array}{ll} U, I, \sigma \models \neg F & \text{iff } U, I, \sigma \not\models F \\ U, I, \sigma \models F_1 \wedge F_2 & \text{iff } U, I, \sigma \models F_1 \text{ and } U, I, \sigma \models F_2 \\ U, I, \sigma \models F_1 \vee F_2 & \text{iff } U, I, \sigma \models F_1 \text{ or } U, I, \sigma \models F_2 \\ U, I, \sigma \models F_1 \rightarrow F_2 & \text{iff, } U, I, \sigma \not\models F_1 \text{ or } U, I, \sigma \models F_2 \\ U, I, \sigma \models F_1 \leftrightarrow F_2 & \text{iff,} \end{array}$$

Evaluation of Formulas II

- ▶ Inductive semantics for boolean connectives:

$$\begin{array}{ll} U, I, \sigma \models \neg F & \text{iff } U, I, \sigma \not\models F \\ U, I, \sigma \models F_1 \wedge F_2 & \text{iff } U, I, \sigma \models F_1 \text{ and } U, I, \sigma \models F_2 \\ U, I, \sigma \models F_1 \vee F_2 & \text{iff } U, I, \sigma \models F_1 \text{ or } U, I, \sigma \models F_2 \\ U, I, \sigma \models F_1 \rightarrow F_2 & \text{iff, } U, I, \sigma \not\models F_1 \text{ or } U, I, \sigma \models F_2 \\ U, I, \sigma \models F_1 \leftrightarrow F_2 & \text{iff, } U, I, \sigma \models F_1 \text{ and } U, I, \sigma \models F_2, \\ & \text{or } U, I, \sigma \not\models F_1 \text{ and } U, I, \sigma \not\models F_2 \end{array}$$

Example II: Evaluation of Formulas

- Consider universe $\{\star, \bullet\}$, variable assignment $\sigma : \{x \mapsto \star\}$, and interpretation I :

$$\begin{aligned}I(a) &= \star & I(b) &= \bullet \\I(f) &= \{\star \mapsto \bullet, \bullet \mapsto \star\} \\I(p) &= \{\langle \bullet, \star \rangle, \langle \bullet, \bullet \rangle\}\end{aligned}$$

Example II: Evaluation of Formulas

- Consider universe $\{\star, \bullet\}$, variable assignment $\sigma : \{x \mapsto \star\}$, and interpretation I :

$$\begin{aligned}I(a) &= \star & I(b) &= \bullet \\I(f) &= \{\star \mapsto \bullet, \bullet \mapsto \star\} \\I(p) &= \{\langle \bullet, \star \rangle, \langle \bullet, \bullet \rangle\}\end{aligned}$$

- Under U, I and σ , what do these formulas evaluate to?

$$p(f(b), f(x)) \rightarrow p(f(x), f(b)) \quad =$$

Example II: Evaluation of Formulas

- Consider universe $\{\star, \bullet\}$, variable assignment $\sigma : \{x \mapsto \star\}$, and interpretation I :

$$\begin{aligned}I(a) &= \star & I(b) &= \bullet \\I(f) &= \{\star \mapsto \bullet, \bullet \mapsto \star\} \\I(p) &= \{\langle \bullet, \star \rangle, \langle \bullet, \bullet \rangle\}\end{aligned}$$

- Under U, I and σ , what do these formulas evaluate to?

$$p(f(b), f(x)) \rightarrow p(f(x), f(b)) \quad = \quad \text{true}$$

Example II: Evaluation of Formulas

- Consider universe $\{\star, \bullet\}$, variable assignment $\sigma : \{x \mapsto \star\}$, and interpretation I :

$$\begin{aligned}I(a) &= \star & I(b) &= \bullet \\I(f) &= \{\star \mapsto \bullet, \bullet \mapsto \star\} \\I(p) &= \{\langle \bullet, \star \rangle, \langle \bullet, \bullet \rangle\}\end{aligned}$$

- Under U, I and σ , what do these formulas evaluate to?

$$\begin{aligned}p(f(b), f(x)) \rightarrow p(f(x), f(b)) &= \text{true} \\p(f(x), f(b)) \rightarrow p(f(b), f(x)) &= \end{aligned}$$

Example II: Evaluation of Formulas

- Consider universe $\{\star, \bullet\}$, variable assignment $\sigma : \{x \mapsto \star\}$, and interpretation I :

$$\begin{aligned}I(a) &= \star & I(b) &= \bullet \\I(f) &= \{\star \mapsto \bullet, \bullet \mapsto \star\} \\I(p) &= \{\langle \bullet, \star \rangle, \langle \bullet, \bullet \rangle\}\end{aligned}$$

- Under U, I and σ , what do these formulas evaluate to?

$$\begin{aligned}p(f(b), f(x)) \rightarrow p(f(x), f(b)) &= \text{true} \\p(f(x), f(b)) \rightarrow p(f(b), f(x)) &= \text{false}\end{aligned}$$

Variant of Variable Assignment

- ▶ We still need to evaluate formulas containing quantifiers!

Variant of Variable Assignment

- ▶ We still need to evaluate formulas containing quantifiers!
- ▶ But to do that, we first define an x -variant of a variable assignment.

Variant of Variable Assignment

- ▶ We still need to evaluate formulas containing quantifiers!
- ▶ But to do that, we first define an x -variant of a variable assignment.
- ▶ An x -variant of assignment σ , written $\sigma[x \mapsto c]$, is the assignment that agrees with σ for assignments to all variables except x and assigns x to c .

Variant of Variable Assignment

- ▶ We still need to evaluate formulas containing quantifiers!
- ▶ But to do that, we first define an x -variant of a variable assignment.
- ▶ An x -variant of assignment σ , written $\sigma[x \mapsto c]$, is the assignment that agrees with σ for assignments to all variables except x and assigns x to c .
- ▶ **Example:** If $\sigma : \{x \mapsto 1, y \mapsto 2\}$, what is $\sigma[x \mapsto 3]$?

Variant of Variable Assignment

- ▶ We still need to evaluate formulas containing quantifiers!
- ▶ But to do that, we first define an x -variant of a variable assignment.
- ▶ An x -variant of assignment σ , written $\sigma[x \mapsto c]$, is the assignment that agrees with σ for assignments to all variables except x and assigns x to c .
- ▶ **Example:** If $\sigma : \{x \mapsto 1, y \mapsto 2\}$, what is $\sigma[x \mapsto 3]$? $\sigma : \{x \mapsto 3, y \mapsto 2\}$

Evaluation of Formulas II

- ▶ We can now give semantics to quantifiers:

Evaluation of Formulas II

- ▶ We can now give semantics to quantifiers:
- ▶ Universal quantifier:

$$U, I, \sigma \models \forall x.F \quad \text{iff} \quad \text{for all } o \in U, U, I, \sigma[x \mapsto o] \models F$$

Evaluation of Formulas II

- ▶ We can now give semantics to quantifiers:

- ▶ Universal quantifier:

$$U, I, \sigma \models \forall x.F \quad \text{iff} \quad \text{for all } o \in U, U, I, \sigma[x \mapsto o] \models F$$

- ▶ Existential quantifier:

$$U, I, \sigma \models \exists x.F \quad \text{iff} \quad \text{there exists } o \in U \text{ s.t. } U, I, \sigma[x \mapsto o] \models F$$

Example III: Evaluation of Formulas

- Consider universe $\{\star, \bullet\}$, variable assignment $\sigma : \{x \mapsto \star\}$, and interpretation I :

$$\begin{aligned}I(a) &= \star & I(b) &= \bullet \\I(f) &= \{\star \mapsto \bullet, \bullet \mapsto \star\} \\I(p) &= \{\langle \bullet, \star \rangle, \langle \bullet, \bullet \rangle\}\end{aligned}$$

Example III: Evaluation of Formulas

- Consider universe $\{\star, \bullet\}$, variable assignment $\sigma : \{x \mapsto \star\}$, and interpretation I :

$$\begin{aligned}I(a) &= \star & I(b) &= \bullet \\I(f) &= \{\star \mapsto \bullet, \bullet \mapsto \star\} \\I(p) &= \{\langle \bullet, \star \rangle, \langle \bullet, \bullet \rangle\}\end{aligned}$$

- Under U, I and σ , what do these formulas evaluate to?

$$\forall x.p(x, a) \quad =$$

Example III: Evaluation of Formulas

- Consider universe $\{\star, \bullet\}$, variable assignment $\sigma : \{x \mapsto \star\}$, and interpretation I :

$$\begin{aligned}I(a) &= \star & I(b) &= \bullet \\I(f) &= \{\star \mapsto \bullet, \bullet \mapsto \star\} \\I(p) &= \{\langle \bullet, \star \rangle, \langle \bullet, \bullet \rangle\}\end{aligned}$$

- Under U, I and σ , what do these formulas evaluate to?

$$\forall x.p(x, a) \quad = \quad \text{false}$$

Example III: Evaluation of Formulas

- Consider universe $\{\star, \bullet\}$, variable assignment $\sigma : \{x \mapsto \star\}$, and interpretation I :

$$\begin{aligned}I(a) &= \star & I(b) &= \bullet \\I(f) &= \{\star \mapsto \bullet, \bullet \mapsto \star\} \\I(p) &= \{\langle \bullet, \star \rangle, \langle \bullet, \bullet \rangle\}\end{aligned}$$

- Under U, I and σ , what do these formulas evaluate to?

$$\begin{aligned}\forall x.p(x, a) &= \text{false} \\ \forall x.p(b, x) &= \end{aligned}$$

Example III: Evaluation of Formulas

- Consider universe $\{\star, \bullet\}$, variable assignment $\sigma : \{x \mapsto \star\}$, and interpretation I :

$$\begin{aligned}I(a) &= \star & I(b) &= \bullet \\I(f) &= \{\star \mapsto \bullet, \bullet \mapsto \star\} \\I(p) &= \{\langle \bullet, \star \rangle, \langle \bullet, \bullet \rangle\}\end{aligned}$$

- Under U, I and σ , what do these formulas evaluate to?

$$\begin{aligned}\forall x.p(x, a) &= \textit{false} \\ \forall x.p(b, x) &= \textit{true}\end{aligned}$$

Example III: Evaluation of Formulas

- ▶ Consider universe $\{\star, \bullet\}$, variable assignment $\sigma : \{x \mapsto \star\}$, and interpretation I :

$$\begin{aligned}I(a) &= \star & I(b) &= \bullet \\I(f) &= \{\star \mapsto \bullet, \bullet \mapsto \star\} \\I(p) &= \{\langle \bullet, \star \rangle, \langle \bullet, \bullet \rangle\}\end{aligned}$$

- ▶ Under U, I and σ , what do these formulas evaluate to?

$$\begin{aligned}\forall x.p(x, a) &= \textit{false} \\ \forall x.p(b, x) &= \textit{true} \\ \exists x.p(a, x) &= \end{aligned}$$

Example III: Evaluation of Formulas

- ▶ Consider universe $\{\star, \bullet\}$, variable assignment $\sigma : \{x \mapsto \star\}$, and interpretation I :

$$\begin{aligned}I(a) &= \star & I(b) &= \bullet \\I(f) &= \{\star \mapsto \bullet, \bullet \mapsto \star\} \\I(p) &= \{\langle \bullet, \star \rangle, \langle \bullet, \bullet \rangle\}\end{aligned}$$

- ▶ Under U, I and σ , what do these formulas evaluate to?

$$\begin{aligned}\forall x.p(x, a) &= \textit{false} \\ \forall x.p(b, x) &= \textit{true} \\ \exists x.p(a, x) &= \textit{false}\end{aligned}$$

Example III: Evaluation of Formulas

- Consider universe $\{\star, \bullet\}$, variable assignment $\sigma : \{x \mapsto \star\}$, and interpretation I :

$$\begin{aligned}I(a) &= \star & I(b) &= \bullet \\I(f) &= \{\star \mapsto \bullet, \bullet \mapsto \star\} \\I(p) &= \{\langle \bullet, \star \rangle, \langle \bullet, \bullet \rangle\}\end{aligned}$$

- Under U, I and σ , what do these formulas evaluate to?

$$\begin{aligned}\forall x. p(x, a) &= \text{false} \\ \forall x. p(b, x) &= \text{true} \\ \exists x. p(a, x) &= \text{false} \\ \forall x. (p(a, x) \rightarrow p(b, x)) &= \end{aligned}$$

Example III: Evaluation of Formulas

- Consider universe $\{\star, \bullet\}$, variable assignment $\sigma : \{x \mapsto \star\}$, and interpretation I :

$$\begin{aligned}I(a) &= \star & I(b) &= \bullet \\I(f) &= \{\star \mapsto \bullet, \bullet \mapsto \star\} \\I(p) &= \{\langle \bullet, \star \rangle, \langle \bullet, \bullet \rangle\}\end{aligned}$$

- Under U, I and σ , what do these formulas evaluate to?

$$\begin{aligned}\forall x. p(x, a) &= \textit{false} \\ \forall x. p(b, x) &= \textit{true} \\ \exists x. p(a, x) &= \textit{false} \\ \forall x. (p(a, x) \rightarrow p(b, x)) &= \textit{true}\end{aligned}$$

Example III: Evaluation of Formulas

- Consider universe $\{\star, \bullet\}$, variable assignment $\sigma : \{x \mapsto \star\}$, and interpretation I :

$$\begin{aligned}I(a) &= \star & I(b) &= \bullet \\I(f) &= \{\star \mapsto \bullet, \bullet \mapsto \star\} \\I(p) &= \{\langle \bullet, \star \rangle, \langle \bullet, \bullet \rangle\}\end{aligned}$$

- Under U, I and σ , what do these formulas evaluate to?

$$\begin{aligned}\forall x. p(x, a) &= \text{false} \\ \forall x. p(b, x) &= \text{true} \\ \exists x. p(a, x) &= \text{false} \\ \forall x. (p(a, x) \rightarrow p(b, x)) &= \text{true} \\ \exists x. (p(f(x), f(x)) \rightarrow p(x, x)) &= \end{aligned}$$

Example III: Evaluation of Formulas

- Consider universe $\{\star, \bullet\}$, variable assignment $\sigma : \{x \mapsto \star\}$, and interpretation I :

$$\begin{aligned}I(a) &= \star & I(b) &= \bullet \\I(f) &= \{\star \mapsto \bullet, \bullet \mapsto \star\} \\I(p) &= \{\langle \bullet, \star \rangle, \langle \bullet, \bullet \rangle\}\end{aligned}$$

- Under U, I and σ , what do these formulas evaluate to?

$$\begin{aligned}\forall x. p(x, a) &= \textit{false} \\ \forall x. p(b, x) &= \textit{true} \\ \exists x. p(a, x) &= \textit{false} \\ \forall x. (p(a, x) \rightarrow p(b, x)) &= \textit{true} \\ \exists x. (p(f(x), f(x)) \rightarrow p(x, x)) &= \textit{true}\end{aligned}$$

Satisfiability and Validity of First-Order Formulas

- ▶ A first-order formula F is **satisfiable** iff there exists a structure S and variable assignment σ such that

$$S, \sigma \models F$$

Satisfiability and Validity of First-Order Formulas

- ▶ A first-order formula F is **satisfiable** iff there exists a structure S and variable assignment σ such that

$$S, \sigma \models F$$

- ▶ Otherwise, F is **unsatisfiable**.

Satisfiability and Validity of First-Order Formulas

- ▶ A first-order formula F is **satisfiable** iff there exists a structure S and variable assignment σ such that

$$S, \sigma \models F$$

- ▶ Otherwise, F is **unsatisfiable**.
- ▶ A structure S is a **model** of F , written $S \models F$, if for all variable assignments $\sigma \in X \rightarrow U$, $S, \sigma \models F$.

Satisfiability and Validity of First-Order Formulas

- ▶ A first-order formula F is **satisfiable** iff there exists a structure S and variable assignment σ such that

$$S, \sigma \models F$$

- ▶ Otherwise, F is **unsatisfiable**.
- ▶ A structure S is a **model** of F , written $S \models F$, if for all variable assignments $\sigma \in X \rightarrow U$, $S, \sigma \models F$.
- ▶ A first-order formula F is **valid**, written $\models F$ if for all structures S , and all assignments σ $S, \sigma \models F$

Satisfiability and Validity Examples

- Is the formula $\forall x.\exists y.p(x, y)$ satisfiable?

Satisfiability and Validity Examples

- ▶ Is the formula $\forall x.\exists y.p(x, y)$ satisfiable? **yes**

Satisfiability and Validity Examples

- ▶ Is the formula $\forall x.\exists y.p(x, y)$ satisfiable? **yes**
- ▶ Satisfying interpretation:

Satisfiability and Validity Examples

- ▶ Is the formula $\forall x.\exists y.p(x, y)$ satisfiable? **yes**
- ▶ Satisfying interpretation: $U = \{\star\}$, $I(p) = \{\langle \star, \star \rangle\}$

Satisfiability and Validity Examples

- ▶ Is the formula $\forall x.\exists y.p(x, y)$ satisfiable? **yes**
- ▶ Satisfying interpretation: $U = \{\star\}$, $I(p) = \{\langle \star, \star \rangle\}$
- ▶ Is this formula valid?

Satisfiability and Validity Examples

- ▶ Is the formula $\forall x.\exists y.p(x, y)$ satisfiable? **yes**
- ▶ Satisfying interpretation: $U = \{\star\}$, $I(p) = \{\langle \star, \star \rangle\}$
- ▶ Is this formula valid? **no**

Satisfiability and Validity Examples

- ▶ Is the formula $\forall x.\exists y.p(x, y)$ satisfiable? **yes**
- ▶ Satisfying interpretation: $U = \{\star\}$, $I(p) = \{\langle \star, \star \rangle\}$
- ▶ Is this formula valid? **no**
- ▶ Falsifying interpretation:

Satisfiability and Validity Examples

- ▶ Is the formula $\forall x.\exists y.p(x, y)$ satisfiable? **yes**
- ▶ Satisfying interpretation: $U = \{\star\}$, $I(p) = \{\langle\star, \star\rangle\}$
- ▶ Is this formula valid? **no**
- ▶ Falsifying interpretation: $U = \{\star\}$, $I(p) = \{\}$

Satisfiability and Validity Examples

- ▶ Is the formula $\forall x.\exists y.p(x, y)$ satisfiable? **yes**
- ▶ Satisfying interpretation: $U = \{\star\}$, $I(p) = \{\langle \star, \star \rangle\}$
- ▶ Is this formula valid? **no**
- ▶ Falsifying interpretation: $U = \{\star\}$, $I(p) = \{\}$
- ▶ Is the formula $\forall x.(p(x, x) \rightarrow \exists y.p(x, y))$ valid?

Satisfiability and Validity Examples

- ▶ Is the formula $\forall x.\exists y.p(x, y)$ satisfiable? **yes**
- ▶ Satisfying interpretation: $U = \{\star\}$, $I(p) = \{\langle \star, \star \rangle\}$
- ▶ Is this formula valid? **no**
- ▶ Falsifying interpretation: $U = \{\star\}$, $I(p) = \{\}$
- ▶ Is the formula $\forall x.(p(x, x) \rightarrow \exists y.p(x, y))$ valid? **yes**

Satisfiability and Validity Examples

- ▶ Is the formula $\forall x.\exists y.p(x, y)$ satisfiable? **yes**
- ▶ Satisfying interpretation: $U = \{\star\}$, $I(p) = \{\langle\star, \star\rangle\}$
- ▶ Is this formula valid? **no**
- ▶ Falsifying interpretation: $U = \{\star\}$, $I(p) = \{\}$
- ▶ Is the formula $\forall x.(p(x, x) \rightarrow \exists y.p(x, y))$ valid? **yes**
- ▶ **Intuition:** Consider any object o . If $p(o, o)$ is false, then implication satisfied. If $p(o, o)$ is true, there exists a y (namely o) s.t $p(x, y)$ is also true.

More Satisfiability and Validity Examples

- Is the formula $(\exists x.p(x)) \rightarrow p(x)$ sat, unsat, or valid?

More Satisfiability and Validity Examples

- Is the formula $(\exists x.p(x)) \rightarrow p(x)$ sat, unsat, or valid? **sat**

More Satisfiability and Validity Examples

- ▶ Is the formula $(\exists x.p(x)) \rightarrow p(x)$ sat, unsat, or valid? **sat**
- ▶ Satisfying U, I, σ :

More Satisfiability and Validity Examples

- ▶ Is the formula $(\exists x.p(x)) \rightarrow p(x)$ sat, unsat, or valid? **sat**
- ▶ Satisfying U, I, σ : $U = \{\star, \circ\}, I(p) = \{\}, \sigma(x) = \circ$

More Satisfiability and Validity Examples

- ▶ Is the formula $(\exists x.p(x)) \rightarrow p(x)$ sat, unsat, or valid? **sat**
- ▶ Satisfying U, I, σ : $U = \{\star, \circ\}, I(p) = \{\}, \sigma(x) = \circ$
- ▶ Falsifying interpretation:

More Satisfiability and Validity Examples

- ▶ Is the formula $(\exists x.p(x)) \rightarrow p(x)$ sat, unsat, or valid? **sat**
- ▶ Satisfying U, I, σ : $U = \{\star, \circ\}, I(p) = \{\}, \sigma(x) = \circ$
- ▶ Falsifying interpretation: $U = \{\star, \circ\}, I(p) = \{\langle \star \rangle\}, \sigma(x) = \circ$

More Satisfiability and Validity Examples

- ▶ Is the formula $(\exists x.p(x)) \rightarrow p(x)$ sat, unsat, or valid? **sat**
- ▶ Satisfying U, I, σ : $U = \{\star, \circ\}, I(p) = \{\}, \sigma(x) = \circ$
- ▶ Falsifying interpretation: $U = \{\star, \circ\}, I(p) = \{\langle \star \rangle\}, \sigma(x) = \circ$
- ▶ Is the formula $(\forall x.p(x)) \rightarrow p(x)$ sat, unsat, or valid?

More Satisfiability and Validity Examples

- ▶ Is the formula $(\exists x.p(x)) \rightarrow p(x)$ sat, unsat, or valid? **sat**
- ▶ Satisfying U, I, σ : $U = \{\star, \circ\}, I(p) = \{\}, \sigma(x) = \circ$
- ▶ Falsifying interpretation: $U = \{\star, \circ\}, I(p) = \{\langle \star \rangle\}, \sigma(x) = \circ$
- ▶ Is the formula $(\forall x.p(x)) \rightarrow p(x)$ sat, unsat, or valid? **valid**

More Satisfiability and Validity Examples

- ▶ Is the formula $(\exists x.p(x)) \rightarrow p(x)$ sat, unsat, or valid? **sat**
- ▶ Satisfying U, I, σ : $U = \{\star, \circ\}, I(p) = \{\}, \sigma(x) = \circ$
- ▶ Falsifying interpretation: $U = \{\star, \circ\}, I(p) = \{\langle \star \rangle\}, \sigma(x) = \circ$
- ▶ Is the formula $(\forall x.p(x)) \rightarrow p(x)$ sat, unsat, or valid? **valid**
- ▶ What about $(\forall x.p(x) \rightarrow q(x)) \rightarrow (\exists x.(p(x) \wedge q(x)))$?

More Satisfiability and Validity Examples

- ▶ Is the formula $(\exists x.p(x)) \rightarrow p(x)$ sat, unsat, or valid? **sat**
- ▶ Satisfying U, I, σ : $U = \{\star, \circ\}, I(p) = \{\}, \sigma(x) = \circ$
- ▶ Falsifying interpretation: $U = \{\star, \circ\}, I(p) = \{\langle \star \rangle\}, \sigma(x) = \circ$
- ▶ Is the formula $(\forall x.p(x)) \rightarrow p(x)$ sat, unsat, or valid? **valid**
- ▶ What about $(\forall x.p(x) \rightarrow q(x)) \rightarrow (\exists x.(p(x) \wedge q(x)))$? **sat**

More Satisfiability and Validity Examples

- ▶ Is the formula $(\exists x.p(x)) \rightarrow p(x)$ sat, unsat, or valid? **sat**
- ▶ Satisfying U, I, σ : $U = \{\star, \circ\}, I(p) = \{\}, \sigma(x) = \circ$
- ▶ Falsifying interpretation: $U = \{\star, \circ\}, I(p) = \{\langle \star \rangle\}, \sigma(x) = \circ$
- ▶ Is the formula $(\forall x.p(x)) \rightarrow p(x)$ sat, unsat, or valid? **valid**
- ▶ What about $(\forall x.p(x) \rightarrow q(x)) \rightarrow (\exists x.(p(x) \wedge q(x)))$? **sat**
- ▶ Satisfying interpretation:

More Satisfiability and Validity Examples

- ▶ Is the formula $(\exists x.p(x)) \rightarrow p(x)$ sat, unsat, or valid? **sat**
- ▶ Satisfying U, I, σ : $U = \{\star, \circ\}, I(p) = \{\}, \sigma(x) = \circ$
- ▶ Falsifying interpretation: $U = \{\star, \circ\}, I(p) = \{\langle\star\rangle\}, \sigma(x) = \circ$
- ▶ Is the formula $(\forall x.p(x)) \rightarrow p(x)$ sat, unsat, or valid? **valid**
- ▶ What about $(\forall x.p(x) \rightarrow q(x)) \rightarrow (\exists x.(p(x) \wedge q(x)))$? **sat**
- ▶ Satisfying interpretation: $U = \{\star\}, I(p) = \{\langle\star\rangle\}, I(q) = \{\langle\star\rangle\}$

More Satisfiability and Validity Examples

- ▶ Is the formula $(\exists x.p(x)) \rightarrow p(x)$ sat, unsat, or valid? **sat**
- ▶ Satisfying U, I, σ : $U = \{\star, \circ\}, I(p) = \{\}, \sigma(x) = \circ$
- ▶ Falsifying interpretation: $U = \{\star, \circ\}, I(p) = \{\langle\star\rangle\}, \sigma(x) = \circ$
- ▶ Is the formula $(\forall x.p(x)) \rightarrow p(x)$ sat, unsat, or valid? **valid**
- ▶ What about $(\forall x.p(x) \rightarrow q(x)) \rightarrow (\exists x.(p(x) \wedge q(x)))$? **sat**
- ▶ Satisfying interpretation: $U = \{\star\}, I(p) = \{\langle\star\rangle\}, I(q) = \{\langle\star\rangle\}$
- ▶ Falsifying interpretation:

More Satisfiability and Validity Examples

- ▶ Is the formula $(\exists x.p(x)) \rightarrow p(x)$ sat, unsat, or valid? **sat**
- ▶ Satisfying U, I, σ : $U = \{\star, \circ\}, I(p) = \{\}, \sigma(x) = \circ$
- ▶ Falsifying interpretation: $U = \{\star, \circ\}, I(p) = \{\langle\star\rangle\}, \sigma(x) = \circ$
- ▶ Is the formula $(\forall x.p(x)) \rightarrow p(x)$ sat, unsat, or valid? **valid**
- ▶ What about $(\forall x.p(x) \rightarrow q(x)) \rightarrow (\exists x.(p(x) \wedge q(x)))$? **sat**
- ▶ Satisfying interpretation: $U = \{\star\}, I(p) = \{\langle\star\rangle\}, I(q) = \{\langle\star\rangle\}$
- ▶ Falsifying interpretation: $U = \{\star\}, I(p) = \{\}, I(q) = \{\langle\star\rangle\}$

Understanding Models

- **Recall:** A structure S is a model of a formula if for all σ , $S, \sigma \models F$

Understanding Models

- ▶ **Recall:** A structure S is a model of a formula if for all σ , $S, \sigma \models F$
- ▶ Consider a formula F such that $S, \sigma \models F$. Is S a model F ?

Understanding Models

- ▶ **Recall:** A structure S is a model of a formula if for all σ , $S, \sigma \models F$
- ▶ Consider a formula F such that $S, \sigma \models F$. Is S a model F ? **not necessarily**

Understanding Models

- ▶ **Recall:** A structure S is a model of a formula if for all σ , $S, \sigma \models F$
- ▶ Consider a formula F such that $S, \sigma \models F$. Is S a model F ? **not necessarily**
- ▶ Consider a sentence F such that $S, \sigma \models F$. Is S a model F ?

Understanding Models

- ▶ **Recall:** A structure S is a model of a formula if for all σ , $S, \sigma \models F$
- ▶ Consider a formula F such that $S, \sigma \models F$. Is S a model F ? **not necessarily**
- ▶ Consider a sentence F such that $S, \sigma \models F$. Is S a model F ? **yes**

Understanding Models

- ▶ **Recall:** A structure S is a model of a formula if for all σ , $S, \sigma \models F$
- ▶ Consider a formula F such that $S, \sigma \models F$. Is S a model F ? **not necessarily**
- ▶ Consider a sentence F such that $S, \sigma \models F$. Is S a model F ? **yes**
- ▶ Consider a ground formula F such that $S, \sigma \models F$. Is S a model of F ?

Understanding Models

- ▶ **Recall:** A structure S is a model of a formula if for all σ , $S, \sigma \models F$
- ▶ Consider a formula F such that $S, \sigma \models F$. Is S a model F ? **not necessarily**
- ▶ Consider a sentence F such that $S, \sigma \models F$. Is S a model F ? **yes**
- ▶ Consider a ground formula F such that $S, \sigma \models F$. Is S a model of F ? **yes**

Summary

- ▶ Today: Syntax and formal semantics of FOL
- ▶ Next lecture:
 - ▶ Semantic argument method for FOL
 - ▶ Properties of first-order logic: decidability results, compactness