

# ECE750T-28: Computer-aided Reasoning for Software Engineering

## Lecture 6: First Order Logic Syntax and Semantics

Vijay Ganesh  
(Original notes from Isil Dillig)

# Overview

- ▶ So far: Automated reasoning in propositional logic.

# Overview

- ▶ So far: Automated reasoning in propositional logic.
- ▶ Propositional logic is simple and easy to automate, but not very expressive

# Overview

- ▶ So far: Automated reasoning in propositional logic.
- ▶ Propositional logic is simple and easy to automate, but not very expressive
- ▶ **Today:** First order logic, also known as relational logic, predicate logic, or first-order predicate calculus

# Overview

- ▶ So far: Automated reasoning in propositional logic.
- ▶ Propositional logic is simple and easy to automate, but not very expressive
- ▶ **Today:** First order logic, also known as relational logic, predicate logic, or first-order predicate calculus
- ▶ Much richer and more expressive, but does not admit completely automated reasoning (more on this later)

# The Plan

- ▶ Syntax and informal semantics (review, today's lecture)

# The Plan

- ▶ Syntax and informal semantics (review, today's lecture)
- ▶ Formal semantics, model theory (today's lecture)

# The Plan

- ▶ Syntax and informal semantics (review, today's lecture)
- ▶ Formal semantics, model theory (today's lecture)
- ▶ Semantic argument method for FOL and properties (next lecture)



# The Plan

- ▶ Syntax and informal semantics (review, today's lecture)
- ▶ Formal semantics, model theory (today's lecture)
- ▶ Semantic argument method for FOL and properties (next lecture)
- ▶ Unification, clausal form (third lecture)

# The Plan

- ▶ Syntax and informal semantics (review, today's lecture)
- ▶ Formal semantics, model theory (today's lecture)
- ▶ Semantic argument method for FOL and properties (next lecture)
- ▶ Unification, clausal form (third lecture)
- ▶ Resolution and first-order theorem proving (fourth lecture)

## Constants in First-Order Logic

- ▶ In propositional logic, we had two constants  $\top$  and  $\perp$

## Constants in First-Order Logic

- ▶ In propositional logic, we had two constants  $\top$  and  $\perp$
- ▶ In first order logic, constants are more involved.

# Constants in First-Order Logic

- ▶ In propositional logic, we had two constants  $\top$  and  $\perp$
- ▶ In first order logic, constants are more involved.
- ▶ Three kinds of constants:
  1. object constants
  2. function constants
  3. relation constants

## Object Constants

- ▶ **Object constants** refer to objects in a universe of discourse.

# Object Constants

- ▶ **Object constants** refer to objects in a universe of discourse.
- ▶ Objects can be anything we want to say something about: integers, people, real numbers, geometric shapes, ...

# Object Constants

- ▶ **Object constants** refer to objects in a universe of discourse.
- ▶ Objects can be anything we want to say something about: integers, people, real numbers, geometric shapes, ...
- ▶ **Example:** If our universe of discourse is people, object constants can be *jack, jane, joe, ...*



# Object Constants

- ▶ **Object constants** refer to objects in a universe of discourse.
- ▶ Objects can be anything we want to say something about: integers, people, real numbers, geometric shapes, ...
- ▶ **Example:** If our universe of discourse is people, object constants can be *jack, jane, joe, ...*
- ▶ As a convention, we will use letters starting with  $a - t$  or digits to denote object constants.

# Object Constants

- ▶ **Object constants** refer to objects in a universe of discourse.
- ▶ Objects can be anything we want to say something about: integers, people, real numbers, geometric shapes, ...
- ▶ **Example:** If our universe of discourse is people, object constants can be *jack, jane, joe, ...*
- ▶ As a convention, we will use letters starting with  $a - t$  or digits to denote object constants.
- ▶ Example: *a, art, beth, 1* etc. refer to object constants.

# Function Constants

- ▶ **Function constants** refer to functions

# Function Constants

- ▶ **Function constants** refer to functions
- ▶ Examples: *mother*, *age*, *plus*, *times*, ...

# Function Constants

- ▶ **Function constants** refer to functions
- ▶ Examples: *mother*, *age*, *plus*, *times*, ...
- ▶ Each function constant has an associated **arity** indicating its number of arguments

# Function Constants

- ▶ **Function constants** refer to functions
- ▶ Examples: *mother*, *age*, *plus*, *times*, ...
- ▶ Each function constant has an associated **arity** indicating its number of arguments
- ▶ Example: *mother* has arity 1 (unary), *times* has arity 2 (binary) etc.

# Function Constants

- ▶ **Function constants** refer to functions
- ▶ Examples: *mother*, *age*, *plus*, *times*, ...
- ▶ Each function constant has an associated **arity** indicating its number of arguments
- ▶ Example: *mother* has arity 1 (unary), *times* has arity 2 (binary) etc.
- ▶ An object constant is really a special case of a function constant with arity 0

# Relation Constants

- ▶ **Relation constants** refer to relations between or properties of objects



# Relation Constants

- ▶ **Relation constants** refer to relations between or properties of objects
- ▶ Example: *loves*, *betterthan*, *ishappy*, ...

# Relation Constants

- ▶ **Relation constants** refer to relations between or properties of objects
- ▶ Example: *loves*, *betterthan*, *ishappy*, ...
- ▶ Each relation constant also has an associated arity

# Relation Constants

- ▶ **Relation constants** refer to relations between or properties of objects
- ▶ Example: *loves*, *betterthan*, *ishappy*, ...
- ▶ Each relation constant also has an associated arity
- ▶ Example: *loves* has arity 2, *ishappy* has arity 1 etc.

# Terms

- ▶ A set of object, function, and relation constants  $C$ ,  $F$ ,  $R$  specifies a **first-order language**, written  $L(C, F, R)$

# Terms

- ▶ A set of object, function, and relation constants  $C, F, R$  specifies a **first-order language**, written  $L(C, F, R)$
- ▶  $C, F, R$  form the **signature** of the language.

# Terms

- ▶ A set of object, function, and relation constants  $C, F, R$  specifies a **first-order language**, written  $L(C, F, R)$
- ▶  $C, F, R$  form the **signature** of the language.
- ▶ Terms  $t$  for a first order language are generated using  $C, F$

# Terms

- ▶ A set of object, function, and relation constants  $C, F, R$  specifies a **first-order language**, written  $L(C, F, R)$
- ▶  $C, F, R$  form the **signature** of the language.
- ▶ Terms  $t$  for a first order language are generated using  $C, F$
- ▶ **Basic terms**: Any object constant in  $C$  and **variables**, denoted  $x, y, z, \dots$

# Terms

- ▶ A set of object, function, and relation constants  $C, F, R$  specifies a **first-order language**, written  $L(C, F, R)$
- ▶  $C, F, R$  form the **signature** of the language.
- ▶ Terms  $t$  for a first order language are generated using  $C, F$
- ▶ **Basic terms**: Any object constant in  $C$  and **variables**, denoted  $x, y, z, \dots$
- ▶ **Composite terms**:  $f(t_1, \dots, t_k)$  where  $f \in F$  is function of arity  $k$ , and  $t_1, \dots, t_k$  are terms



# Terms

- ▶ A set of object, function, and relation constants  $C, F, R$  specifies a **first-order language**, written  $L(C, F, R)$
- ▶  $C, F, R$  form the **signature** of the language.
- ▶ Terms  $t$  for a first order language are generated using  $C, F$
- ▶ **Basic terms**: Any object constant in  $C$  and **variables**, denoted  $x, y, z, \dots$
- ▶ **Composite terms**:  $f(t_1, \dots, t_k)$  where  $f \in F$  is function of arity  $k$ , and  $t_1, \dots, t_k$  are terms
- ▶ Examples:  $mary, x, sister(mary), price(x, macys), age(mother(y)), \dots$

# Formulas

- **Formulas** of  $L(C, F, R)$  are formed using the terms of this language, relation constants  $R$ , logical connectives  $\neg, \wedge, \vee, \dots$ , and quantifiers  $\forall, \exists$ .

# Formulas

- ▶ **Formulas** of  $L(C, F, R)$  are formed using the terms of this language, relation constants  $R$ , logical connectives  $\neg, \wedge, \vee, \dots$ , and quantifiers  $\forall, \exists$ .
- ▶ **Atomic formula (predicate)**: Expression  $p(t_1, \dots, t_n)$  where  $p \in R$  (of arity  $n$ ), and  $t_1, \dots, t_n$  are terms of  $L(C, F, R)$

# Formulas

- ▶ **Formulas** of  $L(C, F, R)$  are formed using the terms of this language, relation constants  $R$ , logical connectives  $\neg, \wedge, \vee, \dots$ , and quantifiers  $\forall, \exists$ .
- ▶ **Atomic formula (predicate)**: Expression  $p(t_1, \dots, t_n)$  where  $p \in R$  (of arity  $n$ ), and  $t_1, \dots, t_n$  are terms of  $L(C, F, R)$
- ▶ If  $F_1$  and  $F_2$  are formulas, then so is  $F_1 \star F_2$  where  $\star$  is any binary connective

# Formulas

- ▶ **Formulas** of  $L(C, F, R)$  are formed using the terms of this language, relation constants  $R$ , logical connectives  $\neg, \wedge, \vee, \dots$ , and quantifiers  $\forall, \exists$ .
- ▶ **Atomic formula (predicate)**: Expression  $p(t_1, \dots, t_n)$  where  $p \in R$  (of arity  $n$ ), and  $t_1, \dots, t_n$  are terms of  $L(C, F, R)$
- ▶ If  $F_1$  and  $F_2$  are formulas, then so is  $F_1 \star F_2$  where  $\star$  is any binary connective
- ▶ If  $F$  is a formula, then so is  $\neg F$

# Formulas

- ▶ **Formulas** of  $L(C, F, R)$  are formed using the terms of this language, relation constants  $R$ , logical connectives  $\neg, \wedge, \vee, \dots$ , and quantifiers  $\forall, \exists$ .
- ▶ **Atomic formula (predicate)**: Expression  $p(t_1, \dots, t_n)$  where  $p \in R$  (of arity  $n$ ), and  $t_1, \dots, t_n$  are terms of  $L(C, F, R)$
- ▶ If  $F_1$  and  $F_2$  are formulas, then so is  $F_1 \star F_2$  where  $\star$  is any binary connective
- ▶ If  $F$  is a formula, then so is  $\neg F$
- ▶ If  $F$  is a formula and  $x$  a variable, so are  $\forall x.F$  (asserts facts about *all* objects) and  $\exists x.F$  (asserts facts about *some* object)

## Important Reminder

- ▶ Predicates (e.g.,  $p(x)$ ) and function terms (e.g.,  $f(x)$ ) look similar, but they are very different!

## Important Reminder

- ▶ Predicates (e.g.,  $p(x)$ ) and function terms (e.g.,  $f(x)$ ) look similar, but they are very different!
- ▶ Function terms can be nested within each other and inside relation constants:  $f(f(x)), p(f(x)), \dots$



## Important Reminder

- ▶ Predicates (e.g.,  $p(x)$ ) and function terms (e.g.,  $f(x)$ ) look similar, but they are very different!
- ▶ Function terms can be nested within each other and inside relation constants:  $f(f(x)), p(f(x)), \dots$
- ▶ Predicates such as  $p(x)$  cannot be nested within function terms or other predicates!

## Important Reminder

- ▶ Predicates (e.g.,  $p(x)$ ) and function terms (e.g.,  $f(x)$ ) look similar, but they are very different!
- ▶ Function terms can be nested within each other and inside relation constants:  $f(f(x)), p(f(x)), \dots$
- ▶ Predicates such as  $p(x)$  cannot be nested within function terms or other predicates!
- ▶  $f(p(x)), p(p(x))$  etc. not valid in FOL!

## Quantifiers and Scoping

- ▶ The subformula embedded inside a quantifier is called the **scope** of that quantifier.

# Quantifiers and Scoping

- ▶ The subformula embedded inside a quantifier is called the **scope** of that quantifier.
- ▶ Example:  $\forall y.((\forall x.p(x)) \rightarrow q(x, y))$

# Quantifiers and Scoping

- ▶ The subformula embedded inside a quantifier is called the **scope** of that quantifier.
- ▶ Example:  $\forall y.((\forall x.p(x)) \rightarrow q(x, y))$
- ▶ An occurrence of a variable is **bound** if it is in the scope of some quantifier.

## Quantifiers and Scoping

- ▶ The subformula embedded inside a quantifier is called the **scope** of that quantifier.
- ▶ Example:  $\forall y.((\forall x.p(x)) \rightarrow q(x, y))$
- ▶ An occurrence of a variable is **bound** if it is in the scope of some quantifier.
- ▶ An occurrence of a variable is **free** if it is not in the scope of any quantifiers.

## Free vs. Bound Variable Example

- Consider the formula:

$$\forall y.((\forall x.p(x)) \rightarrow q(x, y))$$

## Free vs. Bound Variable Example

- ▶ Consider the formula:

$$\forall y.((\forall x.p(x)) \rightarrow q(x, y))$$

- ▶ Is variable  $y$  bound or free?



## Free vs. Bound Variable Example

- ▶ Consider the formula:

$$\forall y.((\forall x.p(x)) \rightarrow q(x, y))$$

- ▶ Is variable  $y$  bound or free? **bound**

## Free vs. Bound Variable Example

- ▶ Consider the formula:

$$\forall y.((\forall x.p(x)) \rightarrow q(x, y))$$

- ▶ Is variable  $y$  bound or free? **bound**
- ▶ Is first occurrence of  $x$  bound or free?

## Free vs. Bound Variable Example

- ▶ Consider the formula:

$$\forall y.((\forall x.p(x)) \rightarrow q(x, y))$$

- ▶ Is variable  $y$  bound or free? **bound**
- ▶ Is first occurrence of  $x$  bound or free? **bound**

## Free vs. Bound Variable Example

- ▶ Consider the formula:

$$\forall y.((\forall x.p(x)) \rightarrow q(x, y))$$

- ▶ Is variable  $y$  bound or free? **bound**
- ▶ Is first occurrence of  $x$  bound or free? **bound**
- ▶ What about second occurrence of  $x$ ?

## Free vs. Bound Variable Example

- ▶ Consider the formula:

$$\forall y.((\forall x.p(x)) \rightarrow q(x, y))$$

- ▶ Is variable  $y$  bound or free? **bound**
- ▶ Is first occurrence of  $x$  bound or free? **bound**
- ▶ What about second occurrence of  $x$ ? **free**

## Closed, Open, and Ground Formulas

- ▶ A formula with no free variables is called a **closed** formula.

## Closed, Open, and Ground Formulas

- ▶ A formula with no free variables is called a **closed** formula.
- ▶ A closed formula is also called a **sentence**.

## Closed, Open, and Ground Formulas

- ▶ A formula with no free variables is called a **closed** formula.
- ▶ A closed formula is also called a **sentence**.
- ▶ A formula containing free variables is said to be **open**.



## Closed, Open, and Ground Formulas

- ▶ A formula with no free variables is called a **closed** formula.
- ▶ A closed formula is also called a **sentence**.
- ▶ A formula containing free variables is said to be **open**.
- ▶ Example: Is the formula  $\forall y.((\forall x.p(x)) \rightarrow q(x, y))$  closed or open?

## Closed, Open, and Ground Formulas

- ▶ A formula with no free variables is called a **closed** formula.
- ▶ A closed formula is also called a **sentence**.
- ▶ A formula containing free variables is said to be **open**.
- ▶ Example: Is the formula  $\forall y.((\forall x.p(x)) \rightarrow q(x, y))$  closed or open? **open**

## Closed, Open, and Ground Formulas

- ▶ A formula with no free variables is called a **closed** formula.
- ▶ A closed formula is also called a **sentence**.
- ▶ A formula containing free variables is said to be **open**.
- ▶ Example: Is the formula  $\forall y.((\forall x.p(x)) \rightarrow q(x, y))$  closed or open? **open**
- ▶ Is the formula  $\forall y.((\forall x.p(x)) \rightarrow (\exists x.q(x, y)))$  a sentence?

## Closed, Open, and Ground Formulas

- ▶ A formula with no free variables is called a **closed** formula.
- ▶ A closed formula is also called a **sentence**.
- ▶ A formula containing free variables is said to be **open**.
- ▶ Example: Is the formula  $\forall y.((\forall x.p(x)) \rightarrow q(x, y))$  closed or open? **open**
- ▶ Is the formula  $\forall y.((\forall x.p(x)) \rightarrow (\exists x.q(x, y)))$  a sentence? **yes**

## Closed, Open, and Ground Formulas

- ▶ A formula with no free variables is called a **closed** formula.
- ▶ A closed formula is also called a **sentence**.
- ▶ A formula containing free variables is said to be **open**.
- ▶ Example: Is the formula  $\forall y.((\forall x.p(x)) \rightarrow q(x, y))$  closed or open? **open**
- ▶ Is the formula  $\forall y.((\forall x.p(x)) \rightarrow (\exists x.q(x, y)))$  a sentence? **yes**
- ▶ A formula is said to be **ground** if it contains no variables.

## Closed, Open, and Ground Formulas

- ▶ A formula with no free variables is called a **closed** formula.
- ▶ A closed formula is also called a **sentence**.
- ▶ A formula containing free variables is said to be **open**.
- ▶ Example: Is the formula  $\forall y.((\forall x.p(x)) \rightarrow q(x, y))$  closed or open? **open**
- ▶ Is the formula  $\forall y.((\forall x.p(x)) \rightarrow (\exists x.q(x, y)))$  a sentence? **yes**
- ▶ A formula is said to be **ground** if it contains no variables.
- ▶ Example:  $p(a, f(b)) \rightarrow q(c)$  is ground.

## Closed, Open, and Ground Formulas

- ▶ A formula with no free variables is called a **closed** formula.
- ▶ A closed formula is also called a **sentence**.
- ▶ A formula containing free variables is said to be **open**.
- ▶ Example: Is the formula  $\forall y.((\forall x.p(x)) \rightarrow q(x, y))$  closed or open? **open**
- ▶ Is the formula  $\forall y.((\forall x.p(x)) \rightarrow (\exists x.q(x, y)))$  a sentence? **yes**
- ▶ A formula is said to be **ground** if it contains no variables.
- ▶ Example:  $p(a, f(b)) \rightarrow q(c)$  is ground.
- ▶ Is  $\forall x.p(x)$  ground?

## Closed, Open, and Ground Formulas

- ▶ A formula with no free variables is called a **closed** formula.
- ▶ A closed formula is also called a **sentence**.
- ▶ A formula containing free variables is said to be **open**.
- ▶ Example: Is the formula  $\forall y.((\forall x.p(x)) \rightarrow q(x, y))$  closed or open? **open**
- ▶ Is the formula  $\forall y.((\forall x.p(x)) \rightarrow (\exists x.q(x, y)))$  a sentence? **yes**
- ▶ A formula is said to be **ground** if it contains no variables.
- ▶ Example:  $p(a, f(b)) \rightarrow q(c)$  is ground.
- ▶ Is  $\forall x.p(x)$  ground? **no**



## A Few Examples

- ▶ To give some intuition about meaning of formula in FOL, some examples:

## A Few Examples

- ▶ To give some intuition about meaning of formula in FOL, some examples:
- ▶ Given a relation constant **friend** of arity 2, how do we express the following in FOL?

## A Few Examples

- ▶ To give some intuition about meaning of formula in FOL, some examples:
- ▶ Given a relation constant **friend** of arity 2, how do we express the following in FOL?
- ▶ *“Everyone has a friend”*

## A Few Examples

- ▶ To give some intuition about meaning of formula in FOL, some examples:
- ▶ Given a relation constant **friend** of arity 2, how do we express the following in FOL?
- ▶ *"Everyone has a friend"*
- ▶  $\forall x. \exists y. \text{friend}(x, y)$

## A Few Examples

- ▶ To give some intuition about meaning of formula in FOL, some examples:
- ▶ Given a relation constant **friend** of arity 2, how do we express the following in FOL?
- ▶ *"Everyone has a friend"*
- ▶  $\forall x. \exists y. \text{friend}(x, y)$
- ▶ *"There is someone who is friends with everyone"*

## A Few Examples

- ▶ To give some intuition about meaning of formula in FOL, some examples:
- ▶ Given a relation constant **friend** of arity 2, how do we express the following in FOL?
- ▶ *"Everyone has a friend"*
- ▶  $\forall x. \exists y. \text{friend}(x, y)$
- ▶ *"There is someone who is friends with everyone"*
- ▶  $\exists x. \forall y. \text{friend}(x, y)$

## A Few Examples

- ▶ To give some intuition about meaning of formula in FOL, some examples:
- ▶ Given a relation constant **friend** of arity 2, how do we express the following in FOL?
- ▶ *"Everyone has a friend"*
- ▶  $\forall x. \exists y. \text{friend}(x, y)$
- ▶ *"There is someone who is friends with everyone"*
- ▶  $\exists x. \forall y. \text{friend}(x, y)$
- ▶ If you flip the quantifiers, completely different meaning!

## More Friendship Examples

- ▶ Given unary relation constants `student` and `atWM` and binary relation constant `friend`, how do we say the following in FOL?



## More Friendship Examples

- ▶ Given unary relation constants `student` and `atWM` and binary relation constant `friend`, how do we say the following in FOL?
- ▶ *"Every student has a friend."*

## More Friendship Examples

- ▶ Given unary relation constants `student` and `atWM` and binary relation constant `friend`, how do we say the following in FOL?
- ▶ *"Every student has a friend."*
- ▶  $\forall x. \text{student}(x) \rightarrow \exists y. \text{friend}(x, y)$

## More Friendship Examples

- ▶ Given unary relation constants `student` and `atWM` and binary relation constant `friend`, how do we say the following in FOL?
- ▶ *"Every student has a friend."*
- ▶  $\forall x. \text{student}(x) \rightarrow \exists y. \text{friend}(x, y)$
- ▶ *"No student has a friend unless he/she is at William & Mary."*

## More Friendship Examples

- ▶ Given unary relation constants `student` and `atWM` and binary relation constant `friend`, how do we say the following in FOL?
- ▶ *"Every student has a friend."*
- ▶  $\forall x. \text{student}(x) \rightarrow \exists y. \text{friend}(x, y)$
- ▶ *"No student has a friend unless he/she is at William & Mary."*
- ▶  $\forall x. ((\text{student}(x) \wedge \neg \text{atWM}(x)) \rightarrow \neg \exists y. \text{friend}(x, y))$

## Even More Friendship Examples

- ▶ “William & Mary students only have friends at William & Mary.”

## Even More Friendship Examples

- ▶ “William & Mary students only have friends at William & Mary.”
- ▶  $\forall x.\forall y.(student(x) \wedge atWM(x) \wedge friends(x, y)) \rightarrow atWM(y)$

## Even More Friendship Examples

- ▶ “William & Mary students only have friends at William & Mary.”

- ▶  $\forall x. \forall y. (student(x) \wedge atWM(x) \wedge friends(x, y)) \rightarrow atWM(y)$

- ▶ Another way of saying this:

$$\forall x. (student(x) \wedge atWM(x)) \rightarrow (\forall y. (friends(x, y) \rightarrow atWM(y)))$$

## Even More Friendship Examples

- ▶ “William & Mary students only have friends at William & Mary.”

- ▶  $\forall x. \forall y. (student(x) \wedge atWM(x) \wedge friends(x, y)) \rightarrow atWM(y)$

- ▶ Another way of saying this:

$$\forall x. (student(x) \wedge atWM(x)) \rightarrow (\forall y. (friends(x, y) \rightarrow atWM(y)))$$

- ▶ These two formulas are actually semantically equivalent



## More Friendship Examples

- ▶ “William & Mary students are all friends with each other.”

## More Friendship Examples

- ▶ “William & Mary students are all friends with each other.”

$$\forall x.\forall y. \quad ((student(x) \wedge atWM(x) \wedge student(y) \wedge atWM(y)) \\ \rightarrow friends(x, y))$$

## More Friendship Examples

- ▶ “William & Mary students are all friends with each other.”

$$\forall x. \forall y. \quad ((student(x) \wedge atWM(x) \wedge student(y) \wedge atWM(y)) \\ \rightarrow friends(x, y))$$

- ▶ “Every William & Mary student has friends outside of their own college”

## More Friendship Examples

- ▶ “William & Mary students are all friends with each other.”

$$\forall x. \forall y. \quad ((student(x) \wedge atWM(x) \wedge student(y) \wedge atWM(y)) \\ \rightarrow friends(x, y))$$

- ▶ “Every William & Mary student has friends outside of their own college”

$$\forall x. \quad ((atWM(x) \wedge student(x)) \rightarrow \\ \exists y. (friends(x, y) \wedge \neg atWM(y)))$$

# Mathematical Theorems in FOL

- **Fermat's Last Theorem:** No three positive integers  $a, b, c$  satisfy the equation  $a^n + b^n = c^n$  for any integer  $n$  greater than 2.

# Mathematical Theorems in FOL

- ▶ **Fermat's Last Theorem:** No three positive integers  $a, b, c$  satisfy the equation  $a^n + b^n = c^n$  for any integer  $n$  greater than 2.
- ▶ **Aside:** French (amateur) mathematician Pierre de Fermat wrote this in 1637 in the margin of Arithmetica, and claimed his proof is too large to fit in the margin.

## Mathematical Theorems in FOL

- ▶ **Fermat's Last Theorem:** No three positive integers  $a, b, c$  satisfy the equation  $a^n + b^n = c^n$  for any integer  $n$  greater than 2.
- ▶ **Aside:** French (amateur) mathematician Pierre de Fermat wrote this in 1637 in the margin of Arithmetica, and claimed his proof is too large to fit in the margin.
- ▶ Could not be proved until British mathematician Andrew Wiles proved in 1995 using properties of elliptical curves.

# Mathematical Theorems in FOL

- ▶ **Fermat's Last Theorem:** No three positive integers  $a, b, c$  satisfy the equation  $a^n + b^n = c^n$  for any integer  $n$  greater than 2.
- ▶ **Aside:** French (amateur) mathematician Pierre de Fermat wrote this in 1637 in the margin of Arithmetica, and claimed his proof is too large to fit in the margin.
- ▶ Could not be proved until British mathematician Andrew Wiles proved in 1995 using properties of elliptical curves.
- ▶ How do we express Fermat's last theorem in FOL (given a function constant  $\wedge$ )?



# Mathematical Theorems in FOL

- ▶ **Fermat's Last Theorem:** No three positive integers  $a, b, c$  satisfy the equation  $a^n + b^n = c^n$  for any integer  $n$  greater than 2.
- ▶ **Aside:** French (amateur) mathematician Pierre de Fermat wrote this in 1637 in the margin of Arithmetica, and claimed his proof is too large to fit in the margin.
- ▶ Could not be proved until British mathematician Andrew Wiles proved in 1995 using properties of elliptical curves.
- ▶ How do we express Fermat's last theorem in FOL (given a function constant  $\wedge$ )?

$$\forall n. n > 2 \rightarrow \neg \exists a, b, c. a > 0 \wedge b > 0 \wedge c > 0 \wedge a^n + b^n = c^n$$

## One Last Example

- ▶ Given binary relation constant `friend`, how do we say this in FOL?

## One Last Example

- ▶ Given binary relation constant `friend`, how do we say this in FOL?
- ▶ *“Every pair of friends has something in common”*

## One Last Example

- ▶ Given binary relation constant **friend**, how do we say this in FOL?
- ▶ *“Every pair of friends has something in common”*
- ▶ This cannot be expressed in FOL because it requires quantification over relation constants!

## One Last Example

- ▶ Given binary relation constant **friend**, how do we say this in FOL?
- ▶ *“Every pair of friends has something in common”*
- ▶ This cannot be expressed in FOL because it requires quantification over relation constants!
- ▶ But it can, however, be expressed in **second-order logic**:

$$\forall x, y. \text{friend}(x, y) \rightarrow \exists p. p(x) \wedge p(y)$$

# Semantics of First Order Logic

- ▶ In propositional logic, the concepts of interpretation, satisfiability, validity were all straightforward.

# Semantics of First Order Logic

- ▶ In propositional logic, the concepts of interpretation, satisfiability, validity were all straightforward.
- ▶ In FOL, these concepts are a bit more involved . . .

# Semantics of First Order Logic

- ▶ In propositional logic, the concepts of interpretation, satisfiability, validity were all straightforward.
- ▶ In FOL, these concepts are a bit more involved . . .
- ▶ To give semantics to FOL, we need to talk about a **universe of discourse** (also sometimes called just “universe” or “domain”)



# Universe of Discourse

- ▶ A **universe of discourse** is a *non-empty* set of objects about which we want to say something

# Universe of Discourse

- ▶ A **universe of discourse** is a *non-empty* set of objects about which we want to say something
- ▶ Universe of discourse can be finite, countably infinite, or uncountably infinite, but not empty

# Universe of Discourse

- ▶ A **universe of discourse** is a *non-empty* set of objects about which we want to say something
- ▶ Universe of discourse can be finite, countably infinite, or uncountably infinite, but not empty
- ▶ Example universes:

# Universe of Discourse

- ▶ A **universe of discourse** is a *non-empty* set of objects about which we want to say something
- ▶ Universe of discourse can be finite, countably infinite, or uncountably infinite, but not empty
- ▶ Example universes:
  - ▶ Set of non-negative integers:

# Universe of Discourse

- ▶ A **universe of discourse** is a *non-empty* set of objects about which we want to say something
- ▶ Universe of discourse can be finite, countably infinite, or uncountably infinite, but not empty
- ▶ Example universes:
  - ▶ Set of non-negative integers: countably infinite

# Universe of Discourse

- ▶ A **universe of discourse** is a *non-empty* set of objects about which we want to say something
- ▶ Universe of discourse can be finite, countably infinite, or uncountably infinite, but not empty
- ▶ Example universes:
  - ▶ Set of non-negative integers: countably infinite
  - ▶ Set of real numbers:

# Universe of Discourse

- ▶ A **universe of discourse** is a *non-empty* set of objects about which we want to say something
- ▶ Universe of discourse can be finite, countably infinite, or uncountably infinite, but not empty
- ▶ Example universes:
  - ▶ Set of non-negative integers: countably infinite
  - ▶ Set of real numbers: uncountably infinite

# Universe of Discourse

- ▶ A **universe of discourse** is a *non-empty* set of objects about which we want to say something
- ▶ Universe of discourse can be finite, countably infinite, or uncountably infinite, but not empty
- ▶ Example universes:
  - ▶ Set of non-negative integers: countably infinite
  - ▶ Set of real numbers: uncountably infinite
  - ▶ The set of suits in playing cards  $\{\clubsuit, \diamond, \heartsuit, \spadesuit\}$ : finite



# Universe of Discourse

- ▶ A **universe of discourse** is a *non-empty* set of objects about which we want to say something
- ▶ Universe of discourse can be finite, countably infinite, or uncountably infinite, but not empty
- ▶ Example universes:
  - ▶ Set of non-negative integers: countably infinite
  - ▶ Set of real numbers: uncountably infinite
  - ▶ The set of suits in playing cards  $\{\clubsuit, \diamond, \heartsuit, \spadesuit\}$ : finite
  - ▶ Students in this class: finite

## First-Order Interpretations

- ▶ An **interpretation** for a first order language  $L(C, F, R)$  is a mapping  $I$  from  $C, F, R$  to objects universe of discourse  $U$

## First-Order Interpretations

- ▶ An **interpretation** for a first order language  $L(C, F, R)$  is a mapping  $I$  from  $C, F, R$  to objects universe of discourse  $U$
- ▶  $I$  maps every  $c \in C$  to some member of  $U$ :  $I(c) \in U$

## First-Order Interpretations

- ▶ An **interpretation** for a first order language  $L(C, F, R)$  is a mapping  $I$  from  $C, F, R$  to objects universe of discourse  $U$
- ▶  $I$  maps every  $c \in C$  to some member of  $U$ :  $I(c) \in U$
- ▶  $I$  maps every  $n$ -ary function constant  $f \in F$  to an  $n$ -ary function  $f^I : U^n \rightarrow U$

# First-Order Interpretations

- ▶ An **interpretation** for a first order language  $L(C, F, R)$  is a mapping  $I$  from  $C, F, R$  to objects universe of discourse  $U$
- ▶  $I$  maps every  $c \in C$  to some member of  $U$ :  $I(c) \in U$
- ▶  $I$  maps every  $n$ -ary function constant  $f \in F$  to an  $n$ -ary function  $f^I : U^n \rightarrow U$
- ▶  $I$  maps every  $n$ -ary relation constant  $p \in R$  to an  $n$ -ary relation  $p^I$  such that  $p^I \subseteq U^n$

# First-Order Interpretations

- ▶ An **interpretation** for a first order language  $L(C, F, R)$  is a mapping  $I$  from  $C, F, R$  to objects universe of discourse  $U$
- ▶  $I$  maps every  $c \in C$  to some member of  $U$ :  $I(c) \in U$
- ▶  $I$  maps every  $n$ -ary function constant  $f \in F$  to an  $n$ -ary function  $f^I : U^n \rightarrow U$
- ▶  $I$  maps every  $n$ -ary relation constant  $p \in R$  to an  $n$ -ary relation  $p^I$  such that  $p^I \subseteq U^n$
- ▶ **Observe:** A first-order interpretation does not talk about variables (only constants)

## An Example

- ▶ Consider the first order language containing object constants  $\{a, b, c\}$ , binary function constant  $f$ , and ternary relation constant  $r$ .

## An Example

- ▶ Consider the first order language containing object constants  $\{a, b, c\}$ , binary function constant  $f$ , and ternary relation constant  $r$ .
- ▶ Universe of discourse:  $U = \{1, 2, 3\}$



## An Example

- ▶ Consider the first order language containing object constants  $\{a, b, c\}$ , binary function constant  $f$ , and ternary relation constant  $r$ .
- ▶ Universe of discourse:  $U = \{1, 2, 3\}$
- ▶ Possible interpretation  $I$ :

$$I(a) = 1, I(b) = 2, I(c) = 2$$

## An Example

- ▶ Consider the first order language containing object constants  $\{a, b, c\}$ , binary function constant  $f$ , and ternary relation constant  $r$ .
- ▶ Universe of discourse:  $U = \{1, 2, 3\}$
- ▶ Possible interpretation  $I$ :

$$I(a) = 1, I(b) = 2, I(c) = 2$$

$$I(f) = \{1 \rightarrow 2, 2 \rightarrow 3, 3 \rightarrow 3\}$$

## An Example

- ▶ Consider the first order language containing object constants  $\{a, b, c\}$ , binary function constant  $f$ , and ternary relation constant  $r$ .
- ▶ Universe of discourse:  $U = \{1, 2, 3\}$
- ▶ Possible interpretation  $I$ :

$$I(a) = 1, I(b) = 2, I(c) = 2$$

$$I(f) = \{1 \rightarrow 2, 2 \rightarrow 3, 3 \rightarrow 3\}$$

$$I(r) = \{\langle 1, 2, 1 \rangle, \langle 2, 2, 1 \rangle\}$$

## An Example

- ▶ Consider the first order language containing object constants  $\{a, b, c\}$ , binary function constant  $f$ , and ternary relation constant  $r$ .
- ▶ Universe of discourse:  $U = \{1, 2, 3\}$
- ▶ Possible interpretation  $I$ :

$$I(a) = 1, I(b) = 2, I(c) = 2$$

$$I(f) = \{1 \rightarrow 2, 2 \rightarrow 3, 3 \rightarrow 3\}$$

$$I(r) = \{\langle 1, 2, 1 \rangle, \langle 2, 2, 1 \rangle\}$$

- ▶ **Observe:** Different object constants do not have to map to distinct objects in  $U$ !

## Another Example

- ▶ Consider the universe of discourse  $U = \{\square, \triangle\}$

## Another Example

- ▶ Consider the universe of discourse  $U = \{\square, \triangle\}$
- ▶ For an object constant  $a$ , what are the possible interpretations?

## Another Example

- ▶ Consider the universe of discourse  $U = \{\square, \triangle\}$
- ▶ For an object constant  $a$ , what are the possible interpretations?  $I(a) = \square$   
or  $I(a) = \triangle$

## Another Example

- ▶ Consider the universe of discourse  $U = \{\square, \triangle\}$
- ▶ For an object constant  $a$ , what are the possible interpretations?  $I(a) = \square$   
or  $I(a) = \triangle$
- ▶ For a unary function constant  $f$ , what are the possible interpretations?



## Another Example

- ▶ Consider the universe of discourse  $U = \{\square, \triangle\}$
- ▶ For an object constant  $a$ , what are the possible interpretations?  $I(a) = \square$   
or  $I(a) = \triangle$
- ▶ For a unary function constant  $f$ , what are the possible interpretations?
- ▶  $I(f) = \{\square \mapsto \square, \triangle \mapsto \square\}$ , or  $I(f) = \{\square \mapsto \square, \triangle \mapsto \triangle\}$ , or  
 $I(f) = \{\square \mapsto \triangle, \triangle \mapsto \square\}$  or  $I(f) = \{\square \mapsto \triangle, \triangle \mapsto \triangle\}$

## Another Example

- ▶ Consider the universe of discourse  $U = \{\square, \triangle\}$
- ▶ For an object constant  $a$ , what are the possible interpretations?  $I(a) = \square$   
or  $I(a) = \triangle$
- ▶ For a unary function constant  $f$ , what are the possible interpretations?
- ▶  $I(f) = \{\square \mapsto \square, \triangle \mapsto \square\}$ , or  $I(f) = \{\square \mapsto \square, \triangle \mapsto \triangle\}$ , or  
 $I(f) = \{\square \mapsto \triangle, \triangle \mapsto \square\}$  or  $I(f) = \{\square \mapsto \triangle, \triangle \mapsto \triangle\}$
- ▶ For a unary relation constant  $r$ , what are all the possible interpretations?

## Another Example

- ▶ Consider the universe of discourse  $U = \{\square, \triangle\}$
- ▶ For an object constant  $a$ , what are the possible interpretations?  $I(a) = \square$   
or  $I(a) = \triangle$
- ▶ For a unary function constant  $f$ , what are the possible interpretations?
- ▶  $I(f) = \{\square \mapsto \square, \triangle \mapsto \square\}$ , or  $I(f) = \{\square \mapsto \square, \triangle \mapsto \triangle\}$ , or  
 $I(f) = \{\square \mapsto \triangle, \triangle \mapsto \square\}$  or  $I(f) = \{\square \mapsto \triangle, \triangle \mapsto \triangle\}$
- ▶ For a unary relation constant  $r$ , what are all the possible interpretations?
- ▶  $I(r) = \{\}$ ,  $I(r) = \{\langle \square \rangle\}$ , or  $I(r) = \{\langle \triangle \rangle\}$ , or  $I(r) = \{\langle \square \rangle, \langle \triangle \rangle\}$

## First-Order Structures and Variable Assignments

- ▶ A **structure**  $S = \langle U, I \rangle$  for a first order language consists of a universe of discourse of  $U$  and an interpretation  $I$ .

# First-Order Structures and Variable Assignments

- ▶ A **structure**  $S = \langle U, I \rangle$  for a first order language consists of a universe of discourse of  $U$  and an interpretation  $I$ .
- ▶ This is sometimes also called an **algebra**.

# First-Order Structures and Variable Assignments

- ▶ A **structure**  $S = \langle U, I \rangle$  for a first order language consists of a universe of discourse of  $U$  and an interpretation  $I$ .
- ▶ This is sometimes also called an **algebra**.
- ▶ A **variable assignment**  $\sigma$  (or assignment) to a FOL formula  $\phi$  in a structure  $S = \langle U, I \rangle$  is a mapping from variables in  $\phi$  to an element of  $U$ .

# First-Order Structures and Variable Assignments

- ▶ A **structure**  $S = \langle U, I \rangle$  for a first order language consists of a universe of discourse of  $U$  and an interpretation  $I$ .
- ▶ This is sometimes also called an **algebra**.
- ▶ A **variable assignment**  $\sigma$  (or assignment) to a FOL formula  $\phi$  in a structure  $S = \langle U, I \rangle$  is a mapping from variables in  $\phi$  to an element of  $U$ .
- ▶ **Example:** Given  $U = \{\square, \triangle\}$ , a possible variable assignment for  $x$ :  
 $\sigma(x) = \triangle$

# First-Order Structures and Variable Assignments

- ▶ A **structure**  $S = \langle U, I \rangle$  for a first order language consists of a universe of discourse of  $U$  and an interpretation  $I$ .
- ▶ This is sometimes also called an **algebra**.
- ▶ A **variable assignment**  $\sigma$  (or assignment) to a FOL formula  $\phi$  in a structure  $S = \langle U, I \rangle$  is a mapping from variables in  $\phi$  to an element of  $U$ .
- ▶ **Example:** Given  $U = \{\square, \triangle\}$ , a possible variable assignment for  $x$ :  
 $\sigma(x) = \triangle$
- ▶ **Observe:**  $\sigma$  does not map variables to object constants but to objects in  $U$ !



## Evaluation of Terms

- ▶ We define how to evaluate a term  $t$  under interpretation  $I$  and assignment  $\sigma$ , written  $\langle I, \sigma \rangle(t)$

## Evaluation of Terms

- ▶ We define how to evaluate a term  $t$  under interpretation  $I$  and assignment  $\sigma$ , written  $\langle I, \sigma \rangle(t)$
- ▶ **Object constants:**  $\langle I, \sigma \rangle(a) = I(a)$

# Evaluation of Terms

- ▶ We define how to evaluate a term  $t$  under interpretation  $I$  and assignment  $\sigma$ , written  $\langle I, \sigma \rangle(t)$
- ▶ Object constants:  $\langle I, \sigma \rangle(a) = I(a)$
- ▶ Variable terms:  $\langle I, \sigma \rangle(v) = \sigma(v)$

# Evaluation of Terms

- ▶ We define how to evaluate a term  $t$  under interpretation  $I$  and assignment  $\sigma$ , written  $\langle I, \sigma \rangle(t)$
- ▶ Object constants:  $\langle I, \sigma \rangle(a) = I(a)$
- ▶ Variable terms:  $\langle I, \sigma \rangle(v) = \sigma(v)$
- ▶ Function terms:

$$\langle I, \sigma \rangle(f(t_1, \dots, t_k)) = I(f)(\langle I, \sigma \rangle(t_1), \dots, \langle I, \sigma \rangle(t_k))$$

## Example: Evaluation of Terms

- ▶ Consider a first-order language containing object constants  $a, b$  and binary function  $f$

## Example: Evaluation of Terms

- ▶ Consider a first-order language containing object constants  $a, b$  and binary function  $f$
- ▶ Consider universe  $\{1, 2\}$  and interpretation  $I$ :

$$\begin{aligned} I(a) &= 1 \quad I(b) = 2 \\ I(f) &= \{\langle 1, 1 \rangle \mapsto 2, \langle 1, 2 \rangle \mapsto 2, \langle 2, 1 \rangle \mapsto 1, \langle 2, 2 \rangle \mapsto 1\} \end{aligned}$$

## Example: Evaluation of Terms

- ▶ Consider a first-order language containing object constants  $a, b$  and binary function  $f$
- ▶ Consider universe  $\{1, 2\}$  and interpretation  $I$ :

$$\begin{aligned} I(a) &= 1 \quad I(b) = 2 \\ I(f) &= \{\langle 1, 1 \rangle \mapsto 2, \langle 1, 2 \rangle \mapsto 2, \langle 2, 1 \rangle \mapsto 1, \langle 2, 2 \rangle \mapsto 1\} \end{aligned}$$

- ▶ Consider variable assignment  $\sigma : \{x \mapsto 2, y \mapsto 1\}$

## Example: Evaluation of Terms

- ▶ Consider a first-order language containing object constants  $a, b$  and binary function  $f$
- ▶ Consider universe  $\{1, 2\}$  and interpretation  $I$ :

$$\begin{aligned} I(a) &= 1 \quad I(b) = 2 \\ I(f) &= \{\langle 1, 1 \rangle \mapsto 2, \langle 1, 2 \rangle \mapsto 2, \langle 2, 1 \rangle \mapsto 1, \langle 2, 2 \rangle \mapsto 1\} \end{aligned}$$

- ▶ Consider variable assignment  $\sigma : \{x \mapsto 2, y \mapsto 1\}$
- ▶ Under  $I$  and  $\sigma$ , what do these terms evaluate to?

$$f(a, y) \quad =$$



## Example: Evaluation of Terms

- ▶ Consider a first-order language containing object constants  $a, b$  and binary function  $f$
- ▶ Consider universe  $\{1, 2\}$  and interpretation  $I$ :

$$\begin{aligned} I(a) &= 1 & I(b) &= 2 \\ I(f) &= \{\langle 1, 1 \rangle \mapsto 2, \langle 1, 2 \rangle \mapsto 2, \langle 2, 1 \rangle \mapsto 1, \langle 2, 2 \rangle \mapsto 1\} \end{aligned}$$

- ▶ Consider variable assignment  $\sigma : \{x \mapsto 2, y \mapsto 1\}$
- ▶ Under  $I$  and  $\sigma$ , what do these terms evaluate to?

$$f(a, y) \quad = \quad 2$$

## Example: Evaluation of Terms

- ▶ Consider a first-order language containing object constants  $a, b$  and binary function  $f$
- ▶ Consider universe  $\{1, 2\}$  and interpretation  $I$ :

$$\begin{aligned} I(a) &= 1 & I(b) &= 2 \\ I(f) &= \{\langle 1, 1 \rangle \mapsto 2, \langle 1, 2 \rangle \mapsto 2, \langle 2, 1 \rangle \mapsto 1, \langle 2, 2 \rangle \mapsto 1\} \end{aligned}$$

- ▶ Consider variable assignment  $\sigma : \{x \mapsto 2, y \mapsto 1\}$
- ▶ Under  $I$  and  $\sigma$ , what do these terms evaluate to?

$$\begin{array}{rcl} f(a, y) & = & 2 \\ f(x, b) & = & \end{array}$$

## Example: Evaluation of Terms

- ▶ Consider a first-order language containing object constants  $a, b$  and binary function  $f$
- ▶ Consider universe  $\{1, 2\}$  and interpretation  $I$ :

$$\begin{aligned} I(a) &= 1 & I(b) &= 2 \\ I(f) &= \{\langle 1, 1 \rangle \mapsto 2, \langle 1, 2 \rangle \mapsto 2, \langle 2, 1 \rangle \mapsto 1, \langle 2, 2 \rangle \mapsto 1\} \end{aligned}$$

- ▶ Consider variable assignment  $\sigma : \{x \mapsto 2, y \mapsto 1\}$
- ▶ Under  $I$  and  $\sigma$ , what do these terms evaluate to?

$$\begin{aligned} f(a, y) &= 2 \\ f(x, b) &= 1 \end{aligned}$$

## Example: Evaluation of Terms

- ▶ Consider a first-order language containing object constants  $a, b$  and binary function  $f$
- ▶ Consider universe  $\{1, 2\}$  and interpretation  $I$ :

$$\begin{aligned} I(a) &= 1 & I(b) &= 2 \\ I(f) &= \{\langle 1, 1 \rangle \mapsto 2, \langle 1, 2 \rangle \mapsto 2, \langle 2, 1 \rangle \mapsto 1, \langle 2, 2 \rangle \mapsto 1\} \end{aligned}$$

- ▶ Consider variable assignment  $\sigma : \{x \mapsto 2, y \mapsto 1\}$
- ▶ Under  $I$  and  $\sigma$ , what do these terms evaluate to?

$$\begin{aligned} f(a, y) &= 2 \\ f(x, b) &= 1 \\ f(f(x, b), f(a, y)) &= \end{aligned}$$

## Example: Evaluation of Terms

- ▶ Consider a first-order language containing object constants  $a, b$  and binary function  $f$
- ▶ Consider universe  $\{1, 2\}$  and interpretation  $I$ :

$$\begin{aligned} I(a) &= 1 & I(b) &= 2 \\ I(f) &= \{\langle 1, 1 \rangle \mapsto 2, \langle 1, 2 \rangle \mapsto 2, \langle 2, 1 \rangle \mapsto 1, \langle 2, 2 \rangle \mapsto 1\} \end{aligned}$$

- ▶ Consider variable assignment  $\sigma : \{x \mapsto 2, y \mapsto 1\}$
- ▶ Under  $I$  and  $\sigma$ , what do these terms evaluate to?

$$\begin{aligned} f(a, y) &= 2 \\ f(x, b) &= 1 \\ f(f(x, b), f(a, y)) &= 2 \end{aligned}$$

## Evaluation of Formulas, Notation

- ▶ We define evaluation of formula  $F$  under structure  $S = \langle U, I \rangle$  and variable assignment  $\sigma$ .

## Evaluation of Formulas, Notation

- ▶ We define evaluation of formula  $F$  under structure  $S = \langle U, I \rangle$  and variable assignment  $\sigma$ .
- ▶ If  $F$  evaluates to true under  $U, I, \sigma$ , we write  $U, I, \sigma \models F$

## Evaluation of Formulas, Notation

- ▶ We define evaluation of formula  $F$  under structure  $S = \langle U, I \rangle$  and variable assignment  $\sigma$ .
- ▶ If  $F$  evaluates to true under  $U, I, \sigma$ , we write  $U, I, \sigma \models F$
- ▶ If  $F$  evaluates to false under  $U, I, \sigma$ , we write  $U, I, \sigma \not\models F$



## Evaluation of Formulas, Notation

- ▶ We define evaluation of formula  $F$  under structure  $S = \langle U, I \rangle$  and variable assignment  $\sigma$ .
- ▶ If  $F$  evaluates to true under  $U, I, \sigma$ , we write  $U, I, \sigma \models F$
- ▶ If  $F$  evaluates to false under  $U, I, \sigma$ , we write  $U, I, \sigma \not\models F$
- ▶ We define the semantics of  $\models$  inductively.

## Evaluation of Formulas, Bases Cases

- Base case I:

$$U, I, \sigma \models \top \quad U, I, \sigma \not\models \perp$$

## Evaluation of Formulas, Bases Cases

- ▶ Base case I:

$$U, I, \sigma \models \top \quad U, I, \sigma \not\models \perp$$

- ▶ Base case II:

$$U, I, \sigma \models p(t_1, \dots, t_k) \text{ iff } \langle \langle I, \sigma \rangle(t_1), \dots, \langle I, \sigma \rangle(t_k) \rangle \in I(p)$$

## Example I: Evaluation of Formulas

- Consider a first-order language containing object constants  $a, b$  and unary function  $f$ , and binary relation constant  $p$

## Example I: Evaluation of Formulas

- ▶ Consider a first-order language containing object constants  $a, b$  and unary function  $f$ , and binary relation constant  $p$
- ▶ Consider universe  $\{\star, \bullet\}$  and interpretation  $I$ :

$$\begin{aligned}I(a) &= \star & I(b) &= \bullet \\I(f) &= \{\star \mapsto \bullet, \bullet \mapsto \star\} \\I(p) &= \{\langle \bullet, \star \rangle, \langle \bullet, \bullet \rangle\}\end{aligned}$$

## Example I: Evaluation of Formulas

- ▶ Consider a first-order language containing object constants  $a, b$  and unary function  $f$ , and binary relation constant  $p$
- ▶ Consider universe  $\{\star, \bullet\}$  and interpretation  $I$ :

$$\begin{aligned}I(a) &= \star & I(b) &= \bullet \\I(f) &= \{\star \mapsto \bullet, \bullet \mapsto \star\} \\I(p) &= \{\langle \bullet, \star \rangle, \langle \bullet, \bullet \rangle\}\end{aligned}$$

- ▶ Consider variable assignment  $\sigma : \{x \mapsto \star\}$

## Example I: Evaluation of Formulas

- ▶ Consider a first-order language containing object constants  $a, b$  and unary function  $f$ , and binary relation constant  $p$
- ▶ Consider universe  $\{\star, \bullet\}$  and interpretation  $I$ :

$$\begin{aligned}I(a) &= \star & I(b) &= \bullet \\I(f) &= \{\star \mapsto \bullet, \bullet \mapsto \star\} \\I(p) &= \{\langle \bullet, \star \rangle, \langle \bullet, \bullet \rangle\}\end{aligned}$$

- ▶ Consider variable assignment  $\sigma : \{x \mapsto \star\}$
- ▶ Under  $U, I$  and  $\sigma$ , what do these formulas evaluate to?

$$p(f(b), f(x)) \quad =$$

## Example I: Evaluation of Formulas

- ▶ Consider a first-order language containing object constants  $a, b$  and unary function  $f$ , and binary relation constant  $p$
- ▶ Consider universe  $\{\star, \bullet\}$  and interpretation  $I$ :

$$\begin{aligned}I(a) &= \star & I(b) &= \bullet \\I(f) &= \{\star \mapsto \bullet, \bullet \mapsto \star\} \\I(p) &= \{\langle \bullet, \star \rangle, \langle \bullet, \bullet \rangle\}\end{aligned}$$

- ▶ Consider variable assignment  $\sigma : \{x \mapsto \star\}$
- ▶ Under  $U, I$  and  $\sigma$ , what do these formulas evaluate to?

$$p(f(b), f(x)) \quad = \quad \textit{false}$$



## Example I: Evaluation of Formulas

- ▶ Consider a first-order language containing object constants  $a, b$  and unary function  $f$ , and binary relation constant  $p$
- ▶ Consider universe  $\{\star, \bullet\}$  and interpretation  $I$ :

$$\begin{aligned}I(a) &= \star & I(b) &= \bullet \\I(f) &= \{\star \mapsto \bullet, \bullet \mapsto \star\} \\I(p) &= \{\langle \bullet, \star \rangle, \langle \bullet, \bullet \rangle\}\end{aligned}$$

- ▶ Consider variable assignment  $\sigma : \{x \mapsto \star\}$
- ▶ Under  $U, I$  and  $\sigma$ , what do these formulas evaluate to?

$$\begin{aligned}p(f(b), f(x)) &= \text{false} \\p(f(x), f(b)) &= \end{aligned}$$

## Example I: Evaluation of Formulas

- ▶ Consider a first-order language containing object constants  $a, b$  and unary function  $f$ , and binary relation constant  $p$
- ▶ Consider universe  $\{\star, \bullet\}$  and interpretation  $I$ :

$$\begin{aligned}I(a) &= \star & I(b) &= \bullet \\I(f) &= \{\star \mapsto \bullet, \bullet \mapsto \star\} \\I(p) &= \{\langle \bullet, \star \rangle, \langle \bullet, \bullet \rangle\}\end{aligned}$$

- ▶ Consider variable assignment  $\sigma : \{x \mapsto \star\}$
- ▶ Under  $U, I$  and  $\sigma$ , what do these formulas evaluate to?

$$\begin{aligned}p(f(b), f(x)) &= \text{false} \\p(f(x), f(b)) &= \text{true}\end{aligned}$$

## Example I: Evaluation of Formulas

- ▶ Consider a first-order language containing object constants  $a, b$  and unary function  $f$ , and binary relation constant  $p$
- ▶ Consider universe  $\{\star, \bullet\}$  and interpretation  $I$ :

$$\begin{aligned}I(a) &= \star & I(b) &= \bullet \\I(f) &= \{\star \mapsto \bullet, \bullet \mapsto \star\} \\I(p) &= \{\langle \bullet, \star \rangle, \langle \bullet, \bullet \rangle\}\end{aligned}$$

- ▶ Consider variable assignment  $\sigma : \{x \mapsto \star\}$
- ▶ Under  $U, I$  and  $\sigma$ , what do these formulas evaluate to?

$$\begin{aligned}p(f(b), f(x)) &= \text{false} \\p(f(x), f(b)) &= \text{true} \\p(a, f(x)) &= \end{aligned}$$

## Example I: Evaluation of Formulas

- ▶ Consider a first-order language containing object constants  $a, b$  and unary function  $f$ , and binary relation constant  $p$
- ▶ Consider universe  $\{\star, \bullet\}$  and interpretation  $I$ :

$$\begin{aligned}I(a) &= \star & I(b) &= \bullet \\I(f) &= \{\star \mapsto \bullet, \bullet \mapsto \star\} \\I(p) &= \{\langle \bullet, \star \rangle, \langle \bullet, \bullet \rangle\}\end{aligned}$$

- ▶ Consider variable assignment  $\sigma : \{x \mapsto \star\}$
- ▶ Under  $U, I$  and  $\sigma$ , what do these formulas evaluate to?

$$\begin{aligned}p(f(b), f(x)) &= \text{false} \\p(f(x), f(b)) &= \text{true} \\p(a, f(x)) &= \text{false}\end{aligned}$$

## Evaluation of Formulas II

- ▶ Inductive semantics for boolean connectives:

## Evaluation of Formulas II

- ▶ Inductive semantics for boolean connectives:

$$U, I, \sigma \models \neg F \quad \text{iff}$$

## Evaluation of Formulas II

- ▶ Inductive semantics for boolean connectives:

$$U, I, \sigma \models \neg F \quad \text{iff } U, I, \sigma \not\models F$$

## Evaluation of Formulas II

- ▶ Inductive semantics for boolean connectives:

$$\begin{array}{ll} U, I, \sigma \models \neg F & \text{iff } U, I, \sigma \not\models F \\ U, I, \sigma \models F_1 \wedge F_2 & \text{iff} \end{array}$$



## Evaluation of Formulas II

- ▶ Inductive semantics for boolean connectives:

$$\begin{array}{ll} U, I, \sigma \models \neg F & \text{iff } U, I, \sigma \not\models F \\ U, I, \sigma \models F_1 \wedge F_2 & \text{iff } U, I, \sigma \models F_1 \text{ and } U, I, \sigma \models F_2 \end{array}$$

## Evaluation of Formulas II

- ▶ Inductive semantics for boolean connectives:

$$\begin{array}{ll} U, I, \sigma \models \neg F & \text{iff } U, I, \sigma \not\models F \\ U, I, \sigma \models F_1 \wedge F_2 & \text{iff } U, I, \sigma \models F_1 \text{ and } U, I, \sigma \models F_2 \\ U, I, \sigma \models F_1 \vee F_2 & \text{iff} \end{array}$$

## Evaluation of Formulas II

- ▶ Inductive semantics for boolean connectives:

$$\begin{array}{ll} U, I, \sigma \models \neg F & \text{iff } U, I, \sigma \not\models F \\ U, I, \sigma \models F_1 \wedge F_2 & \text{iff } U, I, \sigma \models F_1 \text{ and } U, I, \sigma \models F_2 \\ U, I, \sigma \models F_1 \vee F_2 & \text{iff } U, I, \sigma \models F_1 \text{ or } U, I, \sigma \models F_2 \end{array}$$

## Evaluation of Formulas II

- ▶ Inductive semantics for boolean connectives:

$$\begin{array}{ll} U, I, \sigma \models \neg F & \text{iff } U, I, \sigma \not\models F \\ U, I, \sigma \models F_1 \wedge F_2 & \text{iff } U, I, \sigma \models F_1 \text{ and } U, I, \sigma \models F_2 \\ U, I, \sigma \models F_1 \vee F_2 & \text{iff } U, I, \sigma \models F_1 \text{ or } U, I, \sigma \models F_2 \\ U, I, \sigma \models F_1 \rightarrow F_2 & \text{iff,} \end{array}$$

## Evaluation of Formulas II

- ▶ Inductive semantics for boolean connectives:

$$\begin{array}{ll} U, I, \sigma \models \neg F & \text{iff } U, I, \sigma \not\models F \\ U, I, \sigma \models F_1 \wedge F_2 & \text{iff } U, I, \sigma \models F_1 \text{ and } U, I, \sigma \models F_2 \\ U, I, \sigma \models F_1 \vee F_2 & \text{iff } U, I, \sigma \models F_1 \text{ or } U, I, \sigma \models F_2 \\ U, I, \sigma \models F_1 \rightarrow F_2 & \text{iff, } U, I, \sigma \not\models F_1 \text{ or } U, I, \sigma \models F_2 \end{array}$$

## Evaluation of Formulas II

- ▶ Inductive semantics for boolean connectives:

$$\begin{array}{ll} U, I, \sigma \models \neg F & \text{iff } U, I, \sigma \not\models F \\ U, I, \sigma \models F_1 \wedge F_2 & \text{iff } U, I, \sigma \models F_1 \text{ and } U, I, \sigma \models F_2 \\ U, I, \sigma \models F_1 \vee F_2 & \text{iff } U, I, \sigma \models F_1 \text{ or } U, I, \sigma \models F_2 \\ U, I, \sigma \models F_1 \rightarrow F_2 & \text{iff, } U, I, \sigma \not\models F_1 \text{ or } U, I, \sigma \models F_2 \\ U, I, \sigma \models F_1 \leftrightarrow F_2 & \text{iff,} \end{array}$$

## Evaluation of Formulas II

- ▶ Inductive semantics for boolean connectives:

$$\begin{array}{ll} U, I, \sigma \models \neg F & \text{iff } U, I, \sigma \not\models F \\ U, I, \sigma \models F_1 \wedge F_2 & \text{iff } U, I, \sigma \models F_1 \text{ and } U, I, \sigma \models F_2 \\ U, I, \sigma \models F_1 \vee F_2 & \text{iff } U, I, \sigma \models F_1 \text{ or } U, I, \sigma \models F_2 \\ U, I, \sigma \models F_1 \rightarrow F_2 & \text{iff, } U, I, \sigma \not\models F_1 \text{ or } U, I, \sigma \models F_2 \\ U, I, \sigma \models F_1 \leftrightarrow F_2 & \text{iff, } U, I, \sigma \models F_1 \text{ and } U, I, \sigma \models F_2, \\ & \text{or } U, I, \sigma \not\models F_1 \text{ and } U, I, \sigma \not\models F_2 \end{array}$$

## Example II: Evaluation of Formulas

- Consider universe  $\{\star, \bullet\}$ , variable assignment  $\sigma : \{x \mapsto \star\}$ , and interpretation  $I$ :

$$\begin{aligned}I(a) &= \star & I(b) &= \bullet \\I(f) &= \{\star \mapsto \bullet, \bullet \mapsto \star\} \\I(p) &= \{\langle \bullet, \star \rangle, \langle \bullet, \bullet \rangle\}\end{aligned}$$



## Example II: Evaluation of Formulas

- Consider universe  $\{\star, \bullet\}$ , variable assignment  $\sigma : \{x \mapsto \star\}$ , and interpretation  $I$ :

$$\begin{aligned}I(a) &= \star & I(b) &= \bullet \\I(f) &= \{\star \mapsto \bullet, \bullet \mapsto \star\} \\I(p) &= \{\langle \bullet, \star \rangle, \langle \bullet, \bullet \rangle\}\end{aligned}$$

- Under  $U, I$  and  $\sigma$ , what do these formulas evaluate to?

$$p(f(b), f(x)) \rightarrow p(f(x), f(b)) \quad =$$

## Example II: Evaluation of Formulas

- Consider universe  $\{\star, \bullet\}$ , variable assignment  $\sigma : \{x \mapsto \star\}$ , and interpretation  $I$ :

$$\begin{aligned}I(a) &= \star & I(b) &= \bullet \\I(f) &= \{\star \mapsto \bullet, \bullet \mapsto \star\} \\I(p) &= \{\langle \bullet, \star \rangle, \langle \bullet, \bullet \rangle\}\end{aligned}$$

- Under  $U, I$  and  $\sigma$ , what do these formulas evaluate to?

$$p(f(b), f(x)) \rightarrow p(f(x), f(b)) \quad = \quad \text{true}$$

## Example II: Evaluation of Formulas

- Consider universe  $\{\star, \bullet\}$ , variable assignment  $\sigma : \{x \mapsto \star\}$ , and interpretation  $I$ :

$$\begin{aligned}I(a) &= \star & I(b) &= \bullet \\I(f) &= \{\star \mapsto \bullet, \bullet \mapsto \star\} \\I(p) &= \{\langle \bullet, \star \rangle, \langle \bullet, \bullet \rangle\}\end{aligned}$$

- Under  $U, I$  and  $\sigma$ , what do these formulas evaluate to?

$$\begin{aligned}p(f(b), f(x)) \rightarrow p(f(x), f(b)) &= \text{true} \\p(f(x), f(b)) \rightarrow p(f(b), f(x)) &= \end{aligned}$$

## Example II: Evaluation of Formulas

- Consider universe  $\{\star, \bullet\}$ , variable assignment  $\sigma : \{x \mapsto \star\}$ , and interpretation  $I$ :

$$\begin{aligned}I(a) &= \star & I(b) &= \bullet \\I(f) &= \{\star \mapsto \bullet, \bullet \mapsto \star\} \\I(p) &= \{\langle \bullet, \star \rangle, \langle \bullet, \bullet \rangle\}\end{aligned}$$

- Under  $U, I$  and  $\sigma$ , what do these formulas evaluate to?

$$\begin{aligned}p(f(b), f(x)) \rightarrow p(f(x), f(b)) &= \text{true} \\p(f(x), f(b)) \rightarrow p(f(b), f(x)) &= \text{false}\end{aligned}$$

## Variant of Variable Assignment

- ▶ We still need to evaluate formulas containing quantifiers!

## Variant of Variable Assignment

- ▶ We still need to evaluate formulas containing quantifiers!
- ▶ But to do that, we first define an  $x$ -variant of a variable assignment.

## Variant of Variable Assignment

- ▶ We still need to evaluate formulas containing quantifiers!
- ▶ But to do that, we first define an  $x$ -variant of a variable assignment.
- ▶ An  $x$ -variant of assignment  $\sigma$ , written  $\sigma[x \mapsto c]$ , is the assignment that agrees with  $\sigma$  for assignments to all variables except  $x$  and assigns  $x$  to  $c$ .

## Variant of Variable Assignment

- ▶ We still need to evaluate formulas containing quantifiers!
- ▶ But to do that, we first define an  $x$ -variant of a variable assignment.
- ▶ An  $x$ -variant of assignment  $\sigma$ , written  $\sigma[x \mapsto c]$ , is the assignment that agrees with  $\sigma$  for assignments to all variables except  $x$  and assigns  $x$  to  $c$ .
- ▶ **Example:** If  $\sigma : \{x \mapsto 1, y \mapsto 2\}$ , what is  $\sigma[x \mapsto 3]$ ?



## Variant of Variable Assignment

- ▶ We still need to evaluate formulas containing quantifiers!
- ▶ But to do that, we first define an  $x$ -variant of a variable assignment.
- ▶ An  $x$ -variant of assignment  $\sigma$ , written  $\sigma[x \mapsto c]$ , is the assignment that agrees with  $\sigma$  for assignments to all variables except  $x$  and assigns  $x$  to  $c$ .
- ▶ **Example:** If  $\sigma : \{x \mapsto 1, y \mapsto 2\}$ , what is  $\sigma[x \mapsto 3]$ ?  $\sigma : \{x \mapsto 3, y \mapsto 2\}$

## Evaluation of Formulas II

- ▶ We can now give semantics to quantifiers:

## Evaluation of Formulas II

- ▶ We can now give semantics to quantifiers:
- ▶ Universal quantifier:

$$U, I, \sigma \models \forall x.F \quad \text{iff} \quad \text{for all } o \in U, U, I, \sigma[x \mapsto o] \models F$$

## Evaluation of Formulas II

- ▶ We can now give semantics to quantifiers:

- ▶ Universal quantifier:

$$U, I, \sigma \models \forall x.F \quad \text{iff} \quad \text{for all } o \in U, U, I, \sigma[x \mapsto o] \models F$$

- ▶ Existential quantifier:

$$U, I, \sigma \models \exists x.F \quad \text{iff} \quad \text{there exists } o \in U \text{ s.t. } U, I, \sigma[x \mapsto o] \models F$$

## Example III: Evaluation of Formulas

- Consider universe  $\{\star, \bullet\}$ , variable assignment  $\sigma : \{x \mapsto \star\}$ , and interpretation  $I$ :

$$\begin{aligned}I(a) &= \star & I(b) &= \bullet \\I(f) &= \{\star \mapsto \bullet, \bullet \mapsto \star\} \\I(p) &= \{\langle \bullet, \star \rangle, \langle \bullet, \bullet \rangle\}\end{aligned}$$

## Example III: Evaluation of Formulas

- Consider universe  $\{\star, \bullet\}$ , variable assignment  $\sigma : \{x \mapsto \star\}$ , and interpretation  $I$ :

$$\begin{aligned}I(a) &= \star & I(b) &= \bullet \\I(f) &= \{\star \mapsto \bullet, \bullet \mapsto \star\} \\I(p) &= \{\langle \bullet, \star \rangle, \langle \bullet, \bullet \rangle\}\end{aligned}$$

- Under  $U, I$  and  $\sigma$ , what do these formulas evaluate to?

$$\forall x.p(x, a) \quad =$$

## Example III: Evaluation of Formulas

- Consider universe  $\{\star, \bullet\}$ , variable assignment  $\sigma : \{x \mapsto \star\}$ , and interpretation  $I$ :

$$\begin{aligned}I(a) &= \star & I(b) &= \bullet \\I(f) &= \{\star \mapsto \bullet, \bullet \mapsto \star\} \\I(p) &= \{\langle \bullet, \star \rangle, \langle \bullet, \bullet \rangle\}\end{aligned}$$

- Under  $U, I$  and  $\sigma$ , what do these formulas evaluate to?

$$\forall x.p(x, a) \quad = \quad \text{false}$$

## Example III: Evaluation of Formulas

- Consider universe  $\{\star, \bullet\}$ , variable assignment  $\sigma : \{x \mapsto \star\}$ , and interpretation  $I$ :

$$\begin{aligned}I(a) &= \star & I(b) &= \bullet \\I(f) &= \{\star \mapsto \bullet, \bullet \mapsto \star\} \\I(p) &= \{\langle \bullet, \star \rangle, \langle \bullet, \bullet \rangle\}\end{aligned}$$

- Under  $U, I$  and  $\sigma$ , what do these formulas evaluate to?

$$\begin{aligned}\forall x.p(x, a) &= \text{false} \\ \forall x.p(b, x) &= \end{aligned}$$



## Example III: Evaluation of Formulas

- Consider universe  $\{\star, \bullet\}$ , variable assignment  $\sigma : \{x \mapsto \star\}$ , and interpretation  $I$ :

$$\begin{aligned}I(a) &= \star & I(b) &= \bullet \\I(f) &= \{\star \mapsto \bullet, \bullet \mapsto \star\} \\I(p) &= \{\langle \bullet, \star \rangle, \langle \bullet, \bullet \rangle\}\end{aligned}$$

- Under  $U, I$  and  $\sigma$ , what do these formulas evaluate to?

$$\begin{aligned}\forall x.p(x, a) &= \textit{false} \\ \forall x.p(b, x) &= \textit{true}\end{aligned}$$

## Example III: Evaluation of Formulas

- ▶ Consider universe  $\{\star, \bullet\}$ , variable assignment  $\sigma : \{x \mapsto \star\}$ , and interpretation  $I$ :

$$\begin{aligned}I(a) &= \star & I(b) &= \bullet \\I(f) &= \{\star \mapsto \bullet, \bullet \mapsto \star\} \\I(p) &= \{\langle \bullet, \star \rangle, \langle \bullet, \bullet \rangle\}\end{aligned}$$

- ▶ Under  $U, I$  and  $\sigma$ , what do these formulas evaluate to?

$$\begin{aligned}\forall x.p(x, a) &= \textit{false} \\ \forall x.p(b, x) &= \textit{true} \\ \exists x.p(a, x) &= \end{aligned}$$

## Example III: Evaluation of Formulas

- ▶ Consider universe  $\{\star, \bullet\}$ , variable assignment  $\sigma : \{x \mapsto \star\}$ , and interpretation  $I$ :

$$\begin{aligned}I(a) &= \star & I(b) &= \bullet \\I(f) &= \{\star \mapsto \bullet, \bullet \mapsto \star\} \\I(p) &= \{\langle \bullet, \star \rangle, \langle \bullet, \bullet \rangle\}\end{aligned}$$

- ▶ Under  $U, I$  and  $\sigma$ , what do these formulas evaluate to?

$$\begin{aligned}\forall x.p(x, a) &= \textit{false} \\ \forall x.p(b, x) &= \textit{true} \\ \exists x.p(a, x) &= \textit{false}\end{aligned}$$

## Example III: Evaluation of Formulas

- Consider universe  $\{\star, \bullet\}$ , variable assignment  $\sigma : \{x \mapsto \star\}$ , and interpretation  $I$ :

$$\begin{aligned}I(a) &= \star & I(b) &= \bullet \\I(f) &= \{\star \mapsto \bullet, \bullet \mapsto \star\} \\I(p) &= \{\langle \bullet, \star \rangle, \langle \bullet, \bullet \rangle\}\end{aligned}$$

- Under  $U, I$  and  $\sigma$ , what do these formulas evaluate to?

$$\begin{aligned}\forall x. p(x, a) &= \text{false} \\ \forall x. p(b, x) &= \text{true} \\ \exists x. p(a, x) &= \text{false} \\ \forall x. (p(a, x) \rightarrow p(b, x)) &= \end{aligned}$$

## Example III: Evaluation of Formulas

- Consider universe  $\{\star, \bullet\}$ , variable assignment  $\sigma : \{x \mapsto \star\}$ , and interpretation  $I$ :

$$\begin{aligned}I(a) &= \star & I(b) &= \bullet \\I(f) &= \{\star \mapsto \bullet, \bullet \mapsto \star\} \\I(p) &= \{\langle \bullet, \star \rangle, \langle \bullet, \bullet \rangle\}\end{aligned}$$

- Under  $U, I$  and  $\sigma$ , what do these formulas evaluate to?

$$\begin{aligned}\forall x. p(x, a) &= \textit{false} \\ \forall x. p(b, x) &= \textit{true} \\ \exists x. p(a, x) &= \textit{false} \\ \forall x. (p(a, x) \rightarrow p(b, x)) &= \textit{true}\end{aligned}$$

## Example III: Evaluation of Formulas

- Consider universe  $\{\star, \bullet\}$ , variable assignment  $\sigma : \{x \mapsto \star\}$ , and interpretation  $I$ :

$$\begin{aligned}I(a) &= \star & I(b) &= \bullet \\I(f) &= \{\star \mapsto \bullet, \bullet \mapsto \star\} \\I(p) &= \{\langle \bullet, \star \rangle, \langle \bullet, \bullet \rangle\}\end{aligned}$$

- Under  $U, I$  and  $\sigma$ , what do these formulas evaluate to?

$$\begin{aligned}\forall x. p(x, a) &= \textit{false} \\ \forall x. p(b, x) &= \textit{true} \\ \exists x. p(a, x) &= \textit{false} \\ \forall x. (p(a, x) \rightarrow p(b, x)) &= \textit{true} \\ \exists x. (p(f(x), f(x)) \rightarrow p(x, x)) &= \end{aligned}$$

## Example III: Evaluation of Formulas

- Consider universe  $\{\star, \bullet\}$ , variable assignment  $\sigma : \{x \mapsto \star\}$ , and interpretation  $I$ :

$$\begin{aligned}I(a) &= \star & I(b) &= \bullet \\I(f) &= \{\star \mapsto \bullet, \bullet \mapsto \star\} \\I(p) &= \{\langle \bullet, \star \rangle, \langle \bullet, \bullet \rangle\}\end{aligned}$$

- Under  $U, I$  and  $\sigma$ , what do these formulas evaluate to?

$$\begin{aligned}\forall x. p(x, a) &= \textit{false} \\ \forall x. p(b, x) &= \textit{true} \\ \exists x. p(a, x) &= \textit{false} \\ \forall x. (p(a, x) \rightarrow p(b, x)) &= \textit{true} \\ \exists x. (p(f(x), f(x)) \rightarrow p(x, x)) &= \textit{true}\end{aligned}$$

## Satisfiability and Validity of First-Order Formulas

- ▶ A first-order formula  $F$  is **satisfiable** iff there exists a structure  $S$  and variable assignment  $\sigma$  such that

$$S, \sigma \models F$$



# Satisfiability and Validity of First-Order Formulas

- ▶ A first-order formula  $F$  is **satisfiable** iff there exists a structure  $S$  and variable assignment  $\sigma$  such that

$$S, \sigma \models F$$

- ▶ Otherwise,  $F$  is **unsatisfiable**.

# Satisfiability and Validity of First-Order Formulas

- ▶ A first-order formula  $F$  is **satisfiable** iff there exists a structure  $S$  and variable assignment  $\sigma$  such that

$$S, \sigma \models F$$

- ▶ Otherwise,  $F$  is **unsatisfiable**.
- ▶ A structure  $S$  is a **model** of  $F$ , written  $S \models F$ , if for all variable assignments  $\sigma \in X \rightarrow U$ ,  $S, \sigma \models F$ .

# Satisfiability and Validity of First-Order Formulas

- ▶ A first-order formula  $F$  is **satisfiable** iff there exists a structure  $S$  and variable assignment  $\sigma$  such that

$$S, \sigma \models F$$

- ▶ Otherwise,  $F$  is **unsatisfiable**.
- ▶ A structure  $S$  is a **model** of  $F$ , written  $S \models F$ , if for all variable assignments  $\sigma \in X \rightarrow U$ ,  $S, \sigma \models F$ .
- ▶ A first-order formula  $F$  is **valid**, written  $\models F$  if for all structures  $S$ ,  $S, \sigma \models F$

## Satisfiability and Validity Examples

- Is the formula  $\forall x.\exists y.p(x, y)$  satisfiable?

## Satisfiability and Validity Examples

- ▶ Is the formula  $\forall x.\exists y.p(x, y)$  satisfiable? **yes**

## Satisfiability and Validity Examples

- ▶ Is the formula  $\forall x.\exists y.p(x, y)$  satisfiable? **yes**
- ▶ Satisfying interpretation:

## Satisfiability and Validity Examples

- ▶ Is the formula  $\forall x.\exists y.p(x, y)$  satisfiable? **yes**
- ▶ Satisfying interpretation:  $U = \{\star\}$ ,  $I(p) = \{\langle \star, \star \rangle\}$

## Satisfiability and Validity Examples

- ▶ Is the formula  $\forall x.\exists y.p(x, y)$  satisfiable? **yes**
- ▶ Satisfying interpretation:  $U = \{\star\}$ ,  $I(p) = \{\langle \star, \star \rangle\}$
- ▶ Is this formula valid?



## Satisfiability and Validity Examples

- ▶ Is the formula  $\forall x.\exists y.p(x, y)$  satisfiable? **yes**
- ▶ Satisfying interpretation:  $U = \{\star\}$ ,  $I(p) = \{\langle \star, \star \rangle\}$
- ▶ Is this formula valid? **no**

## Satisfiability and Validity Examples

- ▶ Is the formula  $\forall x.\exists y.p(x, y)$  satisfiable? **yes**
- ▶ Satisfying interpretation:  $U = \{\star\}$ ,  $I(p) = \{\langle \star, \star \rangle\}$
- ▶ Is this formula valid? **no**
- ▶ Falsifying interpretation:

## Satisfiability and Validity Examples

- ▶ Is the formula  $\forall x.\exists y.p(x, y)$  satisfiable? **yes**
- ▶ Satisfying interpretation:  $U = \{\star\}$ ,  $I(p) = \{\langle \star, \star \rangle\}$
- ▶ Is this formula valid? **no**
- ▶ Falsifying interpretation:  $U = \{\star\}$ ,  $I(p) = \{\}$

## Satisfiability and Validity Examples

- ▶ Is the formula  $\forall x.\exists y.p(x, y)$  satisfiable? **yes**
- ▶ Satisfying interpretation:  $U = \{\star\}$ ,  $I(p) = \{\langle \star, \star \rangle\}$
- ▶ Is this formula valid? **no**
- ▶ Falsifying interpretation:  $U = \{\star\}$ ,  $I(p) = \{\}$
- ▶ Is the formula  $\forall x.(p(x, x) \rightarrow \exists y.p(x, y))$  valid?

## Satisfiability and Validity Examples

- ▶ Is the formula  $\forall x.\exists y.p(x, y)$  satisfiable? **yes**
- ▶ Satisfying interpretation:  $U = \{\star\}$ ,  $I(p) = \{\langle \star, \star \rangle\}$
- ▶ Is this formula valid? **no**
- ▶ Falsifying interpretation:  $U = \{\star\}$ ,  $I(p) = \{\}$
- ▶ Is the formula  $\forall x.(p(x, x) \rightarrow \exists y.p(x, y))$  valid? **yes**

## Satisfiability and Validity Examples

- ▶ Is the formula  $\forall x.\exists y.p(x, y)$  satisfiable? **yes**
- ▶ Satisfying interpretation:  $U = \{\star\}$ ,  $I(p) = \{\langle\star, \star\rangle\}$
- ▶ Is this formula valid? **no**
- ▶ Falsifying interpretation:  $U = \{\star\}$ ,  $I(p) = \{\}$
- ▶ Is the formula  $\forall x.(p(x, x) \rightarrow \exists y.p(x, y))$  valid? **yes**
- ▶ **Intuition:** Consider any object  $o$ . If  $p(o, o)$  is false, then implication satisfied. If  $p(o, o)$  is true, there exists a  $y$  (namely  $o$ ) s.t  $p(x, y)$  is also true.

## More Satisfiability and Validity Examples

- Is the formula  $(\exists x.p(x)) \rightarrow p(x)$  sat, unsat, or valid?

## More Satisfiability and Validity Examples

- Is the formula  $(\exists x.p(x)) \rightarrow p(x)$  sat, unsat, or valid? **sat**



## More Satisfiability and Validity Examples

- ▶ Is the formula  $(\exists x.p(x)) \rightarrow p(x)$  sat, unsat, or valid? **sat**
- ▶ Satisfying  $U, I, \sigma$ :

## More Satisfiability and Validity Examples

- ▶ Is the formula  $(\exists x.p(x)) \rightarrow p(x)$  sat, unsat, or valid? **sat**
- ▶ Satisfying  $U, I, \sigma$ :  $U = \{\star, \circ\}, I(p) = \{\}, \sigma(x) = \circ$

## More Satisfiability and Validity Examples

- ▶ Is the formula  $(\exists x.p(x)) \rightarrow p(x)$  sat, unsat, or valid? **sat**
- ▶ Satisfying  $U, I, \sigma$ :  $U = \{\star, \circ\}, I(p) = \{\}, \sigma(x) = \circ$
- ▶ Falsifying interpretation:

## More Satisfiability and Validity Examples

- ▶ Is the formula  $(\exists x.p(x)) \rightarrow p(x)$  sat, unsat, or valid? **sat**
- ▶ Satisfying  $U, I, \sigma$ :  $U = \{\star, \circ\}, I(p) = \{\}, \sigma(x) = \circ$
- ▶ Falsifying interpretation:  $U = \{\star, \circ\}, I(p) = \{\langle \star \rangle\}, \sigma(x) = \circ$

## More Satisfiability and Validity Examples

- ▶ Is the formula  $(\exists x.p(x)) \rightarrow p(x)$  sat, unsat, or valid? **sat**
- ▶ Satisfying  $U, I, \sigma$ :  $U = \{\star, \circ\}, I(p) = \{\}, \sigma(x) = \circ$
- ▶ Falsifying interpretation:  $U = \{\star, \circ\}, I(p) = \{\langle \star \rangle\}, \sigma(x) = \circ$
- ▶ Is the formula  $(\forall x.p(x)) \rightarrow p(x)$  sat, unsat, or valid?

## More Satisfiability and Validity Examples

- ▶ Is the formula  $(\exists x.p(x)) \rightarrow p(x)$  sat, unsat, or valid? **sat**
- ▶ Satisfying  $U, I, \sigma$ :  $U = \{\star, \circ\}, I(p) = \{\}, \sigma(x) = \circ$
- ▶ Falsifying interpretation:  $U = \{\star, \circ\}, I(p) = \{\langle \star \rangle\}, \sigma(x) = \circ$
- ▶ Is the formula  $(\forall x.p(x)) \rightarrow p(x)$  sat, unsat, or valid? **valid**

## More Satisfiability and Validity Examples

- ▶ Is the formula  $(\exists x.p(x)) \rightarrow p(x)$  sat, unsat, or valid? **sat**
- ▶ Satisfying  $U, I, \sigma$ :  $U = \{\star, \circ\}, I(p) = \{\}, \sigma(x) = \circ$
- ▶ Falsifying interpretation:  $U = \{\star, \circ\}, I(p) = \{\langle \star \rangle\}, \sigma(x) = \circ$
- ▶ Is the formula  $(\forall x.p(x)) \rightarrow p(x)$  sat, unsat, or valid? **valid**
- ▶ What about  $(\forall x.p(x) \rightarrow q(x)) \rightarrow (\exists x.(p(x) \wedge q(x)))$ ?

## More Satisfiability and Validity Examples

- ▶ Is the formula  $(\exists x.p(x)) \rightarrow p(x)$  sat, unsat, or valid? **sat**
- ▶ Satisfying  $U, I, \sigma$ :  $U = \{\star, \circ\}, I(p) = \{\}, \sigma(x) = \circ$
- ▶ Falsifying interpretation:  $U = \{\star, \circ\}, I(p) = \{\langle \star \rangle\}, \sigma(x) = \circ$
- ▶ Is the formula  $(\forall x.p(x)) \rightarrow p(x)$  sat, unsat, or valid? **valid**
- ▶ What about  $(\forall x.p(x) \rightarrow q(x)) \rightarrow (\exists x.(p(x) \wedge q(x)))$ ? **sat**



## More Satisfiability and Validity Examples

- ▶ Is the formula  $(\exists x.p(x)) \rightarrow p(x)$  sat, unsat, or valid? **sat**
- ▶ Satisfying  $U, I, \sigma$ :  $U = \{\star, \circ\}, I(p) = \{\}, \sigma(x) = \circ$
- ▶ Falsifying interpretation:  $U = \{\star, \circ\}, I(p) = \{\langle \star \rangle\}, \sigma(x) = \circ$
- ▶ Is the formula  $(\forall x.p(x)) \rightarrow p(x)$  sat, unsat, or valid? **valid**
- ▶ What about  $(\forall x.p(x) \rightarrow q(x)) \rightarrow (\exists x.(p(x) \wedge q(x)))$ ? **sat**
- ▶ Satisfying interpretation:

## More Satisfiability and Validity Examples

- ▶ Is the formula  $(\exists x.p(x)) \rightarrow p(x)$  sat, unsat, or valid? **sat**
- ▶ Satisfying  $U, I, \sigma$ :  $U = \{\star, \circ\}, I(p) = \{\}, \sigma(x) = \circ$
- ▶ Falsifying interpretation:  $U = \{\star, \circ\}, I(p) = \{\langle\star\rangle\}, \sigma(x) = \circ$
- ▶ Is the formula  $(\forall x.p(x)) \rightarrow p(x)$  sat, unsat, or valid? **valid**
- ▶ What about  $(\forall x.p(x) \rightarrow q(x)) \rightarrow (\exists x.(p(x) \wedge q(x)))$ ? **sat**
- ▶ Satisfying interpretation:  $U = \{\star\}, I(p) = \{\langle\star\rangle\}, I(q) = \{\langle\star\rangle\}$

## More Satisfiability and Validity Examples

- ▶ Is the formula  $(\exists x.p(x)) \rightarrow p(x)$  sat, unsat, or valid? **sat**
- ▶ Satisfying  $U, I, \sigma$ :  $U = \{\star, \circ\}, I(p) = \{\}, \sigma(x) = \circ$
- ▶ Falsifying interpretation:  $U = \{\star, \circ\}, I(p) = \{\langle\star\rangle\}, \sigma(x) = \circ$
- ▶ Is the formula  $(\forall x.p(x)) \rightarrow p(x)$  sat, unsat, or valid? **valid**
- ▶ What about  $(\forall x.p(x) \rightarrow q(x)) \rightarrow (\exists x.(p(x) \wedge q(x)))$ ? **sat**
- ▶ Satisfying interpretation:  $U = \{\star\}, I(p) = \{\langle\star\rangle\}, I(q) = \{\langle\star\rangle\}$
- ▶ Falsifying interpretation:

## More Satisfiability and Validity Examples

- ▶ Is the formula  $(\exists x.p(x)) \rightarrow p(x)$  sat, unsat, or valid? **sat**
- ▶ Satisfying  $U, I, \sigma$ :  $U = \{\star, \circ\}, I(p) = \{\}, \sigma(x) = \circ$
- ▶ Falsifying interpretation:  $U = \{\star, \circ\}, I(p) = \{\langle\star\rangle\}, \sigma(x) = \circ$
- ▶ Is the formula  $(\forall x.p(x)) \rightarrow p(x)$  sat, unsat, or valid? **valid**
- ▶ What about  $(\forall x.p(x) \rightarrow q(x)) \rightarrow (\exists x.(p(x) \wedge q(x)))$ ? **sat**
- ▶ Satisfying interpretation:  $U = \{\star\}, I(p) = \{\langle\star\rangle\}, I(q) = \{\langle\star\rangle\}$
- ▶ Falsifying interpretation:  $U = \{\star\}, I(p) = \{\}, I(q) = \{\langle\star\rangle\}$

# Understanding Models

- **Recall:** A structure  $S$  is a model of a formula if for all  $\sigma$ ,  $S, \sigma \models F$

# Understanding Models

- ▶ **Recall:** A structure  $S$  is a model of a formula if for all  $\sigma$ ,  $S, \sigma \models F$
- ▶ Consider a formula  $F$  such that  $S, \sigma \models F$ . Is  $S$  a model  $F$ ?

# Understanding Models

- ▶ **Recall:** A structure  $S$  is a model of a formula if for all  $\sigma$ ,  $S, \sigma \models F$
- ▶ Consider a formula  $F$  such that  $S, \sigma \models F$ . Is  $S$  a model  $F$ ? **not necessarily**

# Understanding Models

- ▶ **Recall:** A structure  $S$  is a model of a formula if for all  $\sigma$ ,  $S, \sigma \models F$
- ▶ Consider a formula  $F$  such that  $S, \sigma \models F$ . Is  $S$  a model  $F$ ? **not necessarily**
- ▶ Consider a sentence  $F$  such that  $S, \sigma \models F$ . Is  $S$  a model  $F$ ?



# Understanding Models

- ▶ **Recall:** A structure  $S$  is a model of a formula if for all  $\sigma$ ,  $S, \sigma \models F$
- ▶ Consider a formula  $F$  such that  $S, \sigma \models F$ . Is  $S$  a model  $F$ ? **not necessarily**
- ▶ Consider a sentence  $F$  such that  $S, \sigma \models F$ . Is  $S$  a model  $F$ ? **yes**

# Understanding Models

- ▶ **Recall:** A structure  $S$  is a model of a formula if for all  $\sigma$ ,  $S, \sigma \models F$
- ▶ Consider a formula  $F$  such that  $S, \sigma \models F$ . Is  $S$  a model  $F$ ? **not necessarily**
- ▶ Consider a sentence  $F$  such that  $S, \sigma \models F$ . Is  $S$  a model  $F$ ? **yes**
- ▶ Consider a ground formula  $F$  such that  $S, \sigma \models F$ . Is  $S$  a model of  $F$ ?

# Understanding Models

- ▶ **Recall:** A structure  $S$  is a model of a formula if for all  $\sigma$ ,  $S, \sigma \models F$
- ▶ Consider a formula  $F$  such that  $S, \sigma \models F$ . Is  $S$  a model  $F$ ? **not necessarily**
- ▶ Consider a sentence  $F$  such that  $S, \sigma \models F$ . Is  $S$  a model  $F$ ? **yes**
- ▶ Consider a ground formula  $F$  such that  $S, \sigma \models F$ . Is  $S$  a model of  $F$ ? **yes**

# Summary

- ▶ Today: Syntax and formal semantics of FOL
- ▶ Next lecture:
  - ▶ Semantic argument method for FOL
  - ▶ Properties of first-order logic: decidability results, compactness