
A
SIR profile
Date−level PERMANOVA R² = 0.17, p = 0.112; dispersion p = 0.037; stress = 0.09
NMDS2
NMDS1
Sep 23
Jul 23
Nov 23
Jun 23
Oct 23
Aug 23
Dec 23
Mar 24
Feb 24
May 24
Jan 24
Apr 24
Jun 24
Plant−active
Plant−dormant
B
Class trajectories
log2 SIR relative to annual mean
Readily available
Complex
Other
C
Seasonal difference by substrate
Oxalic acid
D−Xylose
Lactic acid
D−Glucose
D−Fructose
D−Trehalose
D−Cellobiose
Succinic acid
Citric acid
Malic acid (steady)
Methyl oleate
Chitin
Lignin
Cellulose
BSA
Chitosan
Hemicellulose (steady)
Salicylic acid
Glycine
L−Glutamine
Phenylethylamine
Diethylamine
L−Leucine
Vanillic acid
Tween40
L−Alanine
N−acetyl−D−glucosamine
L−Arginine
Glutamic acid (steady)
GABA
Glucosamine−HCl
Dormant − active (log2 fold change, 95% CI)
FDR = 0.05
D
Seasonal difference by class
All substrates: 0.42 (95% CI 0.01 to 0.83), p = 0.045
Complex − readily available = 0.16 (−0.13 to 0.45), p = 0.276; exact permutation p = 0.24
Dormant − active (log2 fold change)
Readily available
Complex
Other