
A
BRITE C transcript composition (325 categories)
Date−level PERMANOVA R² = 0.52, exact p = 0.048 (smallest possible 0.048);
sample−level p = 0.048; dispersion p = 1.000; stress = 0.07
NMDS1
NMDS2
Jun 23
Jul 23
Sep 23
Oct 23
May 24
Mar 24
Jan 24
Plant−active
Plant−dormant
B
Class share of total transcript expression
% of total BRITE C expression (mean, 95% CI)
Lipid−like
Aromatic
Amino sugar
Amino acid
C
Categories mapped uniquely to one compound class
Row label: mean share of total expression; dormant dates in bold
Fatty acid degradation (0.1900%)
Glycerophospholipid metabolism (0.0930%)
Glycerolipid metabolism (0.0840%)
Cutin, suberine and wax biosynthesis (0.0006%)
Phenylalanine metabolism (0.1600%)
Tyrosine metabolism (0.1400%)
Benzoate degradation (0.1300%)
Aminobenzoate degradation (0.0650%)
Naphthalene degradation (0.0320%)
Phenylpropanoid biosynthesis (0.0110%)
Amino sugar and nucleotide sugar metabolism (0.2600%)
Butanoate metabolism (0.6800%)
Alanine, aspartate and glutamate metabolism (0.4800%)
Cysteine and methionine metabolism (0.4700%)
Arginine biosynthesis (0.3900%)
Arginine and proline metabolism (0.2900%)
One carbon pool by folate (0.2600%)
Valine, leucine and isoleucine degradation (0.2600%)
Glycine, serine and threonine metabolism (0.2500%)
beta−Alanine metabolism (0.0970%)
z (within category)