

A
Overall level: mean z-score across all 31 substrates
Mean z-score
Harvest
Harvest
Emergence
B
Season comparison
Dormant − active = 0.48
95% CI 0.13 to 0.83
p = 0.013
Date mean of overall level (z)
Active (8 dates)
Dormant (5 dates)
C
Relative response by substrate and date (date means)
Plant-active
Plant-dormant
Plant-active
Ecological association
Plant structural polymers
Cellulose
Hemicellulose
Lignin
Simple sugars and oligosaccharides
Cellobiose
Xylose
Glucose
Fructose
Trehalose
Aromatic / phenolic acids
Vanillic acid
Salicylic acid
Chitinous structural polymers
Chitin
Chitosan
Amino sugars
N-acetylglucosamine
Glucosamine
Organic acids
Citric acid
Malic acid
Oxalic acid
Succinic acid
Lactic acid
Amino acids and related compounds
Glutamic acid
Glutamine
Alanine
Glycine
Leucine
Arginine
GABA
Amines
Phenylethylamine
Diethylamine
Protein
BSA
Fatty acid esters / surfactants
Methyl oleate
Tween 40
plant structural
root-exudate-associated; plant structural
root-exudate-associated
fungal / microbial structural
protein proxy
synthetic lipid proxy
Relative response (z − sample mean z)
D
Class trajectories of relative response (mean over substrates in class; date means)
Relative response
Plant structural polymers
D − A = 0.07 (-0.27 to 0.41)
p = 0.649; Holm = 1.000
Simple sugars and oligosaccharides
D − A = 0.25 (-0.13 to 0.62)
p = 0.162; Holm = 0.650
Aromatic / phenolic acids
Chitinous structural polymers
Amino sugars
Organic acids
D − A = -0.13 (-1.00 to 0.73)
p = 0.723; Holm = 1.000
Amino acids and related compounds
D − A = -0.03 (-0.34 to 0.28)
p = 0.830; Holm = 1.000
Amines
Protein
Fatty acid esters / surfactants
descriptive
E
Polymer minus breakdown product
Season shift (D − A) of polymer z − product z, 95% CI
Pooled: 0.27 (-0.12 to 0.67), p = 0.150
cellulose-cellobiose
hemicellulose-xylose
lignin-vanillic acid
chitin-N-acetylglucosamine
chitosan-glucosamine
Pooled (5 pairs)
Dormant − active (z units)