

PREDICTIVE INFERENCE TOOLS FOR RESEARCHERS

by

Voyze G. Harris III

Copyright © Voyze G. Harris III 2021

A Thesis Submitted to the Faculty of the

STATISTICS AND DATA SCIENCE
GRADUATE INTERDISCIPLINARY PROGRAM

In Partial Fulfillment of the Requirements
For the Degree of

MASTER OF SCIENCE

In the Graduate College

THE UNIVERSITY OF ARIZONA

2021

THE UNIVERSITY OF ARIZONA
GRADUATE COLLEGE

As members of the Master's Committee, we certify that we have read the thesis prepared by Voyze Gabriel Harris III, titled *[Enter Thesis Title]* and recommend that it be accepted as fulfilling the dissertation requirement for the Master's Degree.

Dr. Dean Billheimer

Date: _____

Dr. Edward Bedrick

Date: _____

Dr. Walter Piegorsch

Date: _____

Final approval and acceptance of this thesis is contingent upon the candidate's submission of the final copies of the thesis to the Graduate College.

I hereby certify that I have read this thesis prepared under my direction and recommend that it be accepted as fulfilling the Master's requirement.

Dr. Dean Billheimer
Master's Thesis Committee Chair
Biostatistics

Date: _____



ARIZONA

Contents

1	Thesis Abstract	4
2	Introduction: Predictive Inference	5
2.1	Why is predictive inference important?	5
2.2	Difference between parametric inference and predictive inference	5
2.2.1	When is predictive inference more useful?	5
2.2.2	When is parametric inference more useful?	5
2.3	The Bayesian Parametric Prediction Format	5
2.4	[Maybe] Example of Difference between results from Plug-in estimator and results using Predictive Inference	5
3	Chapter 1: Predictive Problems with Conjugate Priors	6
3.1	Prediction of Future Successes: Beta-Binomial (Geisser p. 73)	6
3.1.1	Derivation	6
3.1.2	R Implementation	7
3.1.3	Example	7
3.2	Survival Time: Exponential-Gamma (Geisser p. 74)	7
3.2.1	Derivation	7
3.2.2	R Implementation	9
3.2.3	Example	9
3.3	Poisson-Gamma Model (Hoff p. 43ff)	11
3.3.1	Derivation	11
3.3.2	R Implementation	13
3.3.3	Example	13
3.4	Normal Observation with Normal-Inverse Gamma Prior	13
3.4.1	One sample	13
3.4.1.1	Derivation	13
3.4.1.2	R Implementation	13
3.4.1.3	Example	13
3.4.2	Two samples	13
3.4.2.1	Derivation	13
3.4.2.2	R Implementation	14
3.4.2.3	Example	14
3.4.3	k samples	14
3.4.3.1	Derivation	14
3.4.3.2	R Implementation	14
3.4.3.3	Example	14
3.4.3.4	Ranking Treatments	14
4	Chapter 2: Normal Regression with Zellner's g-prior	15
4.0.0.1	Derivation	15
4.0.0.2	R Implementation	15
4.0.0.3	Example	15
5	Conclusion	16

1 Thesis Abstract

- (paragraph) Statement of the thesis topic and objectives
- (paragraph) Explanation of R package

2 Introduction: Predictive Inference

2.1 Why is predictive inference important?

2.2 Difference between parametric inference and predictive inference

2.2.1 When is predictive inference more useful?

2.2.2 When is parametric inference more useful?

[examples, comparisons]

2.3 The Bayesian Parametric Prediction Format

[Geisser p. 49]

Let

$$f(x^{(N)}, x_{(M)} | \theta) = f(x_{(M)} | x^{(N)}, \theta) f(x^{(N)} | \theta).$$

Here $x^{(N)}$ represents observed events and $x_{(M)}$ are future events. We calculate

$$f(x_{(M)}, x^{(N)}) = \int f(x^{(N)}, x_{(M)} | \theta) p(\theta) d\theta$$

where $p(\theta)$ is the prior density and

$$f(x_{(M)} | x^{(N)}) = \frac{f(x_{(M)}, x^{(N)})}{f(x^{(N)})} = \int f(x_{(M)} | \theta) p(\theta | x^{(N)}) d\theta$$

where

$$p(\theta | x^{(N)}) \propto f(x^{(N)} | \theta) p(\theta).$$

2.4 [Maybe] Example of Difference between results from Plug-in estimator and results using Predictive Inference

3 Chapter 1: Predictive Problems with Conjugate Priors

[Problems with closed-form solutions. These problems will be what the R package is designed for. Use problems from Geisser, Casella & Berger (Bayesian chapter), other sources. Regression problem—predictive distributions of models that include and exclude some predictor]

3.1 Prediction of Future Successes: Beta-Binomial (Geisser p. 73)

3.1.1 Derivation

Let X_i be independent binary variables with $\Pr(X_i = 1) = \theta$, and let $T = \sum X_i$. Then T has probability

$$\binom{N}{t} \theta^t (1 - \theta)^{N-t}.$$

Assume $\theta \sim \text{Beta}(\alpha, \beta)$, so

$$p(\theta) = \frac{\Gamma(\alpha + \beta) \theta^{\alpha-1} (1 - \theta)^{\beta-1}}{\Gamma(\alpha) \Gamma(\beta)}.$$

Then

$$p(\theta | X^{(N)}) = \frac{\Gamma(N + \alpha + \beta) \theta^{t+\alpha-1} (1 - \theta)^{N-t+\beta-1}}{\Gamma(t + \alpha) \Gamma(N - t + \beta)}$$

So for $R = \sum_{i=1}^M X_{N+i}$ we have Beta-Binomial predictive distribution


$$\begin{aligned} \Pr[R = r | t] &= \int \binom{M}{r} \theta^r (1 - \theta)^{M-r} p(\theta | X^{(N)}) d\theta \\ &= \binom{M}{r} \int \theta^r (1 - \theta)^{M-r} \frac{\Gamma(N + \alpha + \beta)}{\Gamma(t + \alpha) \Gamma(N - t + \beta)} \theta^{t+\alpha-1} (1 - \theta)^{N-t+\beta-1} d\theta \\ &= \frac{M!}{r!(M-r)!} \frac{\Gamma(N + \alpha + \beta)}{\Gamma(t + \alpha) \Gamma(N - t + \beta)} \int \theta^{r+t+\alpha-1} (1 - \theta)^{M-r+N-t+\beta-1} d\theta \\ &= \frac{\Gamma(M+1) \Gamma(N + \alpha + \beta) \Gamma(r+t+\alpha) \Gamma(M-r+N-t+\beta)}{\Gamma(r+1) \Gamma(M-r+1) \Gamma(t+\alpha) \Gamma(N-t+\beta) \Gamma(M+N+\alpha+\beta)} \end{aligned}$$

3.1.2 R Implementation

This result has been used to create “standard” R functions `dpredBB()`, `ppredBB()`, and `rpredBB()` for the Beta-Binomial distribution for density, cumulative probability, and random sampling, respectively (see appendix). These functions are exercised in the following example.

3.1.3 Example

Suppose $t = 5$ successes have been observed out of $N = 10$ binary events, $\alpha = 2$ and $\beta = 8$. For $M = 1000$ future observations, the figures below show the predictive distribution from `dpredBB()`, the cumulative distribution from `ppredBB()`, and a histogram of random draws from `rpredBB()`.



3.2 Survival Time: Exponential-Gamma (Geisser p. 74)

3.2.1 Derivation

Suppose $X^{(N)} = (X^{(d)}, X^{(N-d)})$ where $X^{(d)}$ represents copies fully observed from an exponential survival time density

$$f(x|\theta) = \theta e^{-\theta x}$$

and $X^{(N-d)}$ represents copies censored at x_{d+1}, \dots, x_N , respectively. Hence

$$L(\theta) \propto \theta^d e^{-\theta N\bar{x}}$$

when $N\bar{x} = \sum_{i=1}^N x_i$, as shown below.

The usual exponential likelihood is used for the fully observed copies, whereas for the censored copies we need $\Pr(x > \theta) = 1 - \Pr(x \leq \theta) = 1 - F(x|\theta) = 1 - (1 - e^{-\theta x}) = e^{-\theta x}$. Thus the overall likelihood is

$$L(\theta|x) = \prod_{i=1}^d \theta e^{-\theta x_i} \prod_{i=d+1}^N e^{-\theta x_i} = \theta^d e^{-\theta N\bar{x}}$$

Assuming a Gamma(δ, γ) prior for θ ,

$$p(\theta) = \frac{\gamma^\delta \theta^{\delta-1} e^{-\gamma\theta}}{\Gamma(\delta)}$$

we obtain the posterior

$$\begin{aligned} p(\theta|X^{(N)}) &= \frac{p(x^{(N)}|\theta) p(\theta)}{\int p(X^{(N)}|\theta) p(\theta) d\theta} \\ &= \frac{\theta^d e^{-\theta N\bar{x}} \cdot \frac{\gamma^\delta \theta^{\delta-1} e^{-\gamma\theta}}{\Gamma(\delta)}}{\int \left(\theta^d e^{-\theta N\bar{x}} \cdot \frac{\gamma^\delta \theta^{\delta-1} e^{-\gamma\theta}}{\Gamma(\delta)} \right) d\theta} \\ &= \frac{\cancel{\frac{\gamma^\delta}{\Gamma(\delta)}} (\theta^{d+\delta-1} e^{-\theta(\gamma+N\bar{x})})}{\cancel{\frac{\gamma^\delta}{\Gamma(\delta)}} \int (\theta^{d+\delta-1} e^{-\theta(\gamma+N\bar{x})}) d\theta} \\ &= \frac{\frac{(\gamma+N\bar{x})^{d+\delta}}{\Gamma(d+\delta)} (\theta^{d+\delta-1} e^{-\theta(\gamma+N\bar{x})})}{\cancel{\frac{(\gamma+N\bar{x})^{d+\delta}}{\Gamma(d+\delta)}} \int \cancel{(\theta^{d+\delta-1} e^{-\theta(\gamma+N\bar{x})})} d\theta} \\ &= \frac{(\gamma+N\bar{x})^{d+\delta} \theta^{d+\delta-1} e^{-\theta(\gamma+N\bar{x})}}{\Gamma(d+\delta)} \end{aligned}$$

with the Gamma($d+\delta, \gamma+N\bar{x}$) density in the next to last step integrating to 1.

Thus the survival time predictive probability is

$$\begin{aligned}
P(X = x|\theta, X^{(N)}) &= \int p(\theta|X^{(N)}) p(x|\theta) d\theta \\
&= \int \frac{(\gamma + N\bar{x})^{d+\delta} \theta^{d+\delta-1} e^{-\theta(\gamma+N\bar{x})}}{\Gamma(d+\delta)} \cdot \theta e^{-\theta x} d\theta \\
&= (d+\delta)(\gamma + N\bar{x})^{d+\delta} \int \frac{\theta^{(d+\delta+1)-1} e^{-\theta(\gamma+N\bar{x}+x)}}{(d+\delta)\Gamma(d+\delta)} d\theta \\
&= \frac{(d+\delta)(\gamma + N\bar{x})^{d+\delta}}{(\gamma + N\bar{x} + x)^{d+\delta+1}} \int \frac{(\gamma + N\bar{x} + x)^{d+\delta+1} \theta^{(d+\delta+1)-1} e^{-\theta(\gamma+N\bar{x}+x)}}{\Gamma(d+\delta+1)} d\theta \\
&= \frac{(d+\delta)(\gamma + N\bar{x})^{d+\delta}}{(\gamma + N\bar{x} + x)^{d+\delta+1}}
\end{aligned}$$

(simplifying by constructing a $\text{Gamma}(d + \delta + 1, \gamma + N\bar{x} + x)$ density in the final integrand.)


3.2.2 R Implementation

This result has been used to create standard format R functions `dpredEG()`, `ppredEG()`, and `rpredEG()` for the Gamma-Exponential distribution for density, cumulative probability, and random sampling, respectively (see appendix). These functions are exercised in the following example.


3.2.3 Example

Suppose $d = 800$ out of $N = 1000$ copies have been observed, and the remaining 200 censored. Say $\delta = 20$, $\gamma = 5$, and we are interested in the number of survivors out of $M = 1000$ future observations. The figures below illustrate the predictive probability using `dpredEG()` and `rpredEG()`, along with a histogram of a random sample taken using `rpredEG()`.


Exponential-Gamma Predictive Density



Exponential-Gamma Cumulative Predictive Probability



Histogram of Sample with Density Curve Overlay



3.3 Poisson-Gamma Model (Hoff p. 43ff)

3.3.1 Derivation

[using Hoff's notation and variable names below. Should I convert this to Geisser's $x^{(N)}, x_{(M)}$ convention for uniformity throughout my thesis?]

Suppose $Y_1, \dots, Y_n | \theta \stackrel{i.i.d.}{\sim} \text{Poisson}(\theta)$ with Gamma prior $\theta \sim \text{Gamma}(\alpha, \beta)$. That is,

$$\begin{aligned} P(Y_1 = y_1, \dots, Y_n = y_n | \theta) &= \prod_{i=1}^n p(y_i | \theta) \\ &= \prod_{i=1}^n \frac{1}{y_i!} \theta^{y_i} e^{-\theta} \\ &= \left(\prod_{i=1}^n \frac{1}{y_i!} \right) \theta^{\sum y_i} e^{-n\theta} \\ &= c(y_1, \dots, y_n) \theta^{\sum y_i} e^{-n\theta} \end{aligned}$$

and

$$p(\theta) = \frac{\beta^\alpha}{\Gamma(\alpha)} \theta^{\alpha-1} e^{-\beta\theta}, \theta, \alpha, \beta > 0.$$

Then we have posterior distribution

$$\begin{aligned} p(\theta | y_1, \dots, y_n) &= \frac{p(y_1, \dots, y_n | \theta) p(\theta)}{\sum_{\theta} p(y_1, \dots, y_n | \theta) p(\theta)} \\ &= \frac{p(y_1, \dots, y_n | \theta) p(\theta)}{p(y_1, \dots, y_n)} \\ &= \frac{1}{p(y_1, \dots, y_n)} \theta^{\sum y_i} e^{-n\theta} \frac{\beta^\alpha}{\Gamma(\alpha)} \theta^{\alpha-1} e^{-\beta\theta} \\ &= C(y_1, \dots, y_n, \alpha, \beta) \theta^{\alpha + \sum y_i - 1} e^{-(\beta + n)\theta} \\ &\sim \text{Gamma}\left(\alpha + \sum y_i, \beta + n\right). \end{aligned}$$

Here

$$\begin{aligned}
C(y_1, \dots, y_n, \alpha, \beta) &= \frac{1}{p(y_1, \dots, y_n)} \cdot \frac{\beta^\alpha}{\Gamma(\alpha)} \\
&= \frac{1}{\sum_{\theta} p(y_1, \dots, y_n | \theta) p(\theta)} \cdot \frac{\beta^\alpha}{\Gamma(\alpha)} \\
&= \frac{1}{\sum_{\theta} \left(\prod \frac{1}{y_i!} \right) \theta^{\sum y_i} e^{-n\theta} \left(\frac{\beta^\alpha}{\Gamma(\alpha)} \right) \theta^{\alpha-1} e^{-\beta\theta}} \cdot \left(\frac{\beta^\alpha}{\Gamma(\alpha)} \right) \\
&= \frac{1}{\left(\prod \frac{1}{y_i!} \right) \frac{\Gamma(\alpha + \sum y_i)}{(\beta + n)^{\alpha + \sum y_i}} \sum_{\theta} \frac{(\beta + n)^{\alpha + \sum y_i}}{\Gamma(\alpha + \sum y_i)} \theta^{\sum y_i + \alpha - 1} e^{-(\beta + n)\theta}} \\
&= \frac{\prod_{i=1}^n y_i! (\beta + n)^{\alpha + \sum y_i}}{\Gamma(\alpha + \sum y_i)}
\end{aligned}$$

For future observation \tilde{y} , then, we compute predictive distribution

$$\begin{aligned}
p(\tilde{y}|y_1, \dots, y_n) &= \int_0^\infty p(\tilde{y}|\theta, y_1, \dots, y_n) p(\theta|y_1, \dots, y_n) d\theta \\
&= \int p(\tilde{y}|\theta) p(\theta|y_1, \dots, y_n) d\theta \\
&= C \int \left(\frac{1}{\tilde{y}!} \theta^{\tilde{y}} e^{-\theta} \right) \theta^{\alpha + \sum y_i - 1} e^{-(\beta + n)\theta} d\theta \\
&= \frac{C}{\tilde{y}!} \int \theta^{\tilde{y} + \alpha + \sum y_i - 1} e^{-(\beta + n + 1)\theta} d\theta \\
&= \frac{C \Gamma(\tilde{y} + \alpha + \sum y_i)}{\Gamma(\tilde{y} + 1) (\beta + n + 1)^{\tilde{y} + \alpha + \sum y_i}} \int \frac{(\beta + n + 1)^{\tilde{y} + \alpha + \sum y_i}}{\Gamma(\tilde{y} + \alpha + \sum y_i)} \theta^{\tilde{y} + \alpha + \sum y_i - 1} e^{-(\beta + n + 1)\theta} d\theta \\
&= C \cdot \frac{\Gamma(\tilde{y} + \alpha + \sum y_i)}{\Gamma(\tilde{y} + 1) (\beta + n + 1)^{\tilde{y} + \alpha + \sum y_i}} \\
&= \frac{\prod_{i=1}^n y_i! (\beta + n)^{\alpha + \sum y_i}}{\Gamma(\alpha + \sum y_i)} \cdot \frac{\Gamma(\tilde{y} + \alpha + \sum y_i)}{\Gamma(\tilde{y} + 1) (\beta + n + 1)^{\tilde{y} + \alpha + \sum y_i}} \\
&= \prod_{i=1}^n y_i! \cdot \frac{\Gamma(\tilde{y} + \alpha + \sum y_i)}{\Gamma(\tilde{y} + 1) \Gamma(\alpha + \sum y_i)} \cdot \left(\frac{\beta + n}{\beta + n + 1} \right)^{\alpha + \sum y_i} \cdot \left(\frac{1}{\beta + n + 1} \right)^{\tilde{y}}
\end{aligned}$$

3.3.2 R Implementation

3.3.3 Example

3.4 Normal Observation with Normal-Inverse Gamma Prior

3.4.1 One sample

3.4.1.1 Derivation

3.4.1.2 R Implementation

3.4.1.3 Example

3.4.2 Two samples

3.4.2.1 Derivation

3.4.2.2 R Implementation

3.4.2.3 Example

3.4.3 k samples

3.4.3.1 Derivation

3.4.3.2 R Implementation

3.4.3.3 Example

3.4.3.4 Ranking Treatments

4 Chapter 2: Normal Regression with Zellner's g -prior

4.0.0.1 Derivation

4.0.0.2 R Implementation

4.0.0.3 Example

5 Conclusion