

**CONFIDENTIAL**

Session time: 01 Mar 2026 – 15 Mar 2026(15 days)

User information

CONFIDENTIAL**CONFIDENTIAL**

Height: 170 cm

Weight: 68.1 kg

Monitor information

Monitor usage: 100%

SN: 449FDABC531E

Monitor model: M8

Summary

Average

5.9 mmol/L

(Goal: <8.5 mmol/L)

CV

16.9 %

(Goal: <33%)

eHbA1c

5.3 %

(Goal: <7.0%)

TITR

93.3 %

(3.9 – 7.8 mmol/L)

Time in range

TAR:

4.0 %

(Goal: <25%)

● Very high (>13.9 mmol/L)

--

● High(7.9 – 13.9 mmol/L)

13 hrs 20 mins

TIR:

93.2

(Goal: >70%)

● Target range(3.9 – 7.8 mmol/L)

13 days 1 hour 50 mins

TBR:

2.8


(Goal: <4%)

● Low(3 – 3.8 mmol/L)

3 hrs 55 mins

● Very low(<3 mmol/L)

5 hrs 15 mins



Average glucose - Average glucose - Gives average glucose for the date range selected.

Hyper/Hypoglycemic events - Shows the distribution of low (red) and high (yellow) glucose events across different time periods, including the number of events and their average duration as represented in the charts.

CV - The coefficient of variation - The ratio of the standard deviation of glucose to the mean, the smaller the value, the smaller the fluctuation.

eHbA1c - Estimated HbA1c- An estimate of your average blood sugar level over time, calculated from CGM data. Recommended level: below 7%.

TITR - Time in tight target range - is the time of day when blood glucose is within the range of 3.9 to 7.8 mmol/L. It is a more tightly constricted time range than the TIR, and better reflects the physiology of blood glucose in a near-standard or healthy population. It can be used as a control group parameter for TIR to optimize glycemic control strategies, such as prediction of future complication risk indicators and assessment of the effectiveness of glucose-lowering medications.

I Meal-time glucose summary

	Pre-Meal	Post-meal peak	Post-meal excursion Goal:<2.2	Post-Meal		
				1hours/hour	2hours/hour	3hours/hour
Breakfast	--	--	--	--	--	--
Lunch	5.9	8.4	1.2	6.5	7.1	6.9
Dinner	5.7	8	0.9	5.9	6.6	6.7
Snack	5.9	7.2	0.3	6.1	6.2	6.1

Pre-Meal - Shows the glucose readings when a pre-meal log has been made.

Post-meal peak - Shows the peak glucose readings three hours post-meal.

Post-meal excursion - The difference between the 2-hour post-meal glucose level and the pre-meal glucose level.

I Daily meal-time glucose

	Breakfast		Lunch		Dinner		Snack	
	 (mmol/L)	 (mmol/L)	 (mmol/L)	 (mmol/L)	 (mmol/L)	 (mmol/L)	 (mmol/L)	 (mmol/L)
Average	--	--	5.9	8.4	5.7	8.0	5.9	7.2
01 Mar	--	--	--	--	5.5	5.7	5.4	6.0
02 Mar	--	--	5.6	9.3	5.7	7.8	--	--
03 Mar	--	--	6.2	9.2	5.9	9.1	--	--
04 Mar	--	--	6.0	7.7	5.6	8.6	--	--
05 Mar	--	--	6.3	7.1	5.9	6.2	--	--
06 Mar	--	--	6.0	6.4	5.7	12.3	--	--
07 Mar	--	--	5.8	8.1	5.2	7.2	--	--
08 Mar	--	--	5.2	8.1	--	--	--	--
09 Mar	--	--	6.1	11.2	5.7	8.1	--	--
10 Mar	--	--	6.3	7.4	--	--	--	--
11 Mar	--	--	6.4	8.5	--	--	6.4	7.6
12 Mar	--	--	6.2	10.3	--	--	5.8	7.9
13 Mar	--	--	5.7	9.0	5.7	7.4	--	--
14 Mar	--	--	5.7	6.5	--	--	--	--
15 Mar	--	--	5.5	8.4	--	--	--	--

Average - Shows the average meal and insulin logs for each time block on a "typical" day.

 **Pre-Meal** - Shows the glucose readings when a pre-meal log has been made.

 **Post-meal peak** - Shows the peak glucose readings three hours post-meal.

I Daily Log

The daily log report shows logs of mealtime, insulin and other data, and detailed glucose readings for each day in the date range selected.

