

https://viazovska.github.io/my_project[https : //github.com/viazovska/my_project](https://github.com/viazovska/my_project)[https : //viazovska.gith](https://viazovska.github.io)

Pentagonal Number Theorem

JC PM MV

May 12, 2026

Navigate: [Dependency graph](#) | [Lean documentation](#) | [GitHub](#)

1 Partitions

Definition 1. A *partition* of a positive integer n , also called an integer partition, is a way of writing n as a sum of positive integers. Two sums that differ only in the order of their summands are considered the same partition.

The *partition function* $p(n)$ represents the number of possible partitions of a natural number n .

Example 2. $5 = 4 + 1 = 3 + 2 = 3 + 1 + 1 = 2 + 2 + 1 = 2 + 1 + 1 + 1 = 1 + 1 + 1 + 1 + 1$
 $p(5) = 7$

Lemma 3. *def:partition The generating function for $p(n)$ is given by*

$$\sum_{n=0}^{\infty} p(n) x^n = \prod_{k=1}^{\infty} (1 + x^k + x^{2k} + \dots) = \prod_{k=1}^{\infty} (1 - x^k)^{-1}.$$

Proof. We work in the ring of formal power series $\mathbb{Z}[[x]]$.

Step 1. For each fixed $k \geq 1$, the geometric series identity

$$1 + x^k + x^{2k} + \dots = \sum_{j=0}^{\infty} x^{jk} = (1 - x^k)^{-1}$$

holds as formal power series. This gives the second equality.

Step 2. Expand the product $\prod_{k=1}^{\infty} (1 + x^k + x^{2k} + \dots)$. Each term in the expansion is obtained by selecting, for every $k \geq 1$, an exponent $j_k \in \mathbb{Z}_{\geq 0}$ (with $j_k = 0$ for all but finitely many k) and multiplying the chosen monomials. Such a selection contributes a single term

$$\prod_k x^{j_k \cdot k} = x^{\sum_k j_k \cdot k}.$$

Step 3. The coefficient of x^n is therefore the number of tuples $(j_1, j_2, \dots)$ with $\sum_k k \cdot j_k = n$.

Step 4. Such a tuple corresponds bijectively to a partition of n : j_k is the multiplicity of the part k . Hence the coefficient of x^n equals $p(n)$, and the first equality follows. $\square$

Definition 4 (Partitions of n into distinct parts). *all_{distinct}partitions, even_{distinct}partitions, odd_{distinct}partitions*

Let $n \in \mathbb{N}$. We define the following sets of partitions of n into distinct positive parts:

$$\begin{aligned}\mathcal{P}(n) &= \left\{ S \subseteq \{1, \dots, n\} \mid \sum_{s \in S} s = n \right\}, \\ \mathcal{P}_{\text{even}}(n) &= \{S \in \mathcal{P}(n) \mid |S| \equiv 0 \pmod{2}\}, \\ \mathcal{P}_{\text{odd}}(n) &= \{S \in \mathcal{P}(n) \mid |S| \equiv 1 \pmod{2}\}.\end{aligned}$$

We further define the number of even and odd partitions of n as $p_e(n) = |\mathcal{P}_{\text{even}}(n)|$ and $p_o(n) = |\mathcal{P}_{\text{odd}}(n)|$.

Lemma 5. *def:distinct-partitions* We have

$$\prod_{k=1}^{\infty} (1 - x^k) = \sum_{s=0}^{\infty} \sum_{k_1 < k_2 < \dots < k_s} (-1)^s x^{k_1 + k_2 + \dots + k_s} = \sum_{n=0}^{\infty} c_n x^n$$

where $c_0 = 1$ and $c_n = p_e(n) - p_o(n)$ for $n \geq 1$.

Proof. We expand the infinite product as a formal power series.

Step 1: Expansion as a sum over finite subsets. For each $k \geq 1$, the factor $(1 - x^k)$ contributes either 1 or $-x^k$ when expanding the product. A choice across all factors corresponds to a finite subset

$$K = \{k_1 < k_2 < \dots < k_s\} \subseteq \mathbb{Z}_{\geq 1}$$

(the indices for which $-x^k$ was selected). The contribution is

$$\prod_{k \in K} (-x^k) = (-1)^{|K|} x^{\sum_{k \in K} k} = (-1)^s x^{k_1 + \dots + k_s}.$$

Summing over all such K yields the first equality.

Step 2: Grouping by total exponent. A subset K of size s summing to n is exactly a partition $S \in \mathcal{P}(n)$ with $|S| = s$. Hence the coefficient of x^n equals

$$c_n = \sum_{S \in \mathcal{P}(n)} (-1)^{|S|}.$$

Step 3: $c_0 = 1$. Only $S = \emptyset$ partitions 0 into distinct positive parts, contributing $(-1)^0 = 1$.

Step 4: $c_n = p_e(n) - p_o(n)$ for $n \geq 1$. Splitting the sum by parity of $|S|$:

$$c_n = \sum_{\substack{S \in \mathcal{P}(n) \\ |S| \text{ even}}} 1 - \sum_{\substack{S \in \mathcal{P}(n) \\ |S| \text{ odd}}} 1 = |\mathcal{P}_{\text{even}}(n)| - |\mathcal{P}_{\text{odd}}(n)| = p_e(n) - p_o(n).$$

$\square$

Example 6. We compute $\prod_{i=1}^{20} (1 - x^i) + o(x^{20})$.

The answer is $1 - x - x^2 + x^5 + x^7 - x^{12} - x^{15} + O(x^{20})$.

n		$p_o(n)$	$p_e(n)$
1	1	1	0
2	2	1	0
3	$3 = 2 + 1$	1	1
4	$4 = 3 + 1$	1	1
5	$5 = 4 + 1 = 3 + 2$	1	2
6	$6 = 5 + 1 = 4 + 2 = 3 + 2 + 1$	2	2
7	$7 = 6 + 1 = 5 + 2 = 4 + 3 = 4 + 2 + 1$	2	3

2 Pentagonal number theorem

Theorem 7 (Euler). *euler_pentagonal_number_theorem* : distinct - partitions $\prod_{i=1}^{\infty} (1 - x^i) = 1 + \sum_{k=1}^{\infty} (-1)^k (x^{(3k^2-k)/2} + x^{(3k^2+k)/2}) = \sum_{k \in \mathbb{Z}} (-1)^k x^{(3k^2-k)/2}$.

Proof. lemma:expansion-product, lemma:pe-minus-po-closed-form The proof combines the previous expansion lemma with the closed-form computation of $p_e(n) - p_o(n)$ via Franklin's involution.

Step 1: Coefficient identification. By Lemma ??,

$$\prod_{i=1}^{\infty} (1 - x^i) = \sum_{n=0}^{\infty} c_n x^n,$$

where $c_0 = 1$ and $c_n = p_e(n) - p_o(n)$ for $n \geq 1$.

Step 2: Closed form for c_n . By Lemma ?? below,

$$p_e(n) - p_o(n) = \begin{cases} (-1)^k & \text{if } n = (3k^2 - k)/2 \text{ for some } k \in \mathbb{Z}_{\geq 1}, \\ (-1)^k & \text{if } n = (3k^2 + k)/2 \text{ for some } k \in \mathbb{Z}_{\geq 1}, \\ 0 & \text{otherwise.} \end{cases}$$

Step 3: Reassembling the sum. The values of n where $c_n \neq 0$ are exactly the (generalized) pentagonal numbers $n = (3k^2 \pm k)/2$ for $k \geq 1$, plus $n = 0$. Each contributes $(-1)^k$. Hence

$$\prod_{i=1}^{\infty} (1 - x^i) = 1 + \sum_{k=1}^{\infty} (-1)^k \left(x^{(3k^2-k)/2} + x^{(3k^2+k)/2} \right).$$

Step 4: Bilateral sum reformulation. Reindex the second family by $k \mapsto -k$. For $k \in \mathbb{Z}_{\geq 1}$, set $k' = -k$, so $k' \in \mathbb{Z}_{\leq -1}$ and


$$\frac{3(k')^2 - k'}{2} = \frac{3k^2 + k}{2}, \quad (-1)^k = (-1)^{|k'|} = (-1)^{k'}.$$

The $k = 0$ term gives $(-1)^0 x^0 = 1$, matching the leading 1. Combining the three contributions:

$$\sum_{k \in \mathbb{Z}_{\geq 1}} (-1)^k x^{(3k^2-k)/2} + (-1)^0 x^0 + \sum_{k' \in \mathbb{Z}_{\leq -1}} (-1)^{k'} x^{(3(k')^2-k')/2} = \sum_{k \in \mathbb{Z}} (-1)^k x^{(3k^2-k)/2}.$$


□

Remark 8 (Pentagonal numbers). $\frac{k}{\frac{3k^2-k}{2}} \left| \begin{array}{c} -3 \\ 15 \end{array} \right| \begin{array}{c} -2 \\ 7 \end{array} \left| \begin{array}{c} -1 \\ 2 \end{array} \right| \begin{array}{c} 0 \\ 0 \end{array} \left| \begin{array}{c} 1 \\ 1 \end{array} \right| \begin{array}{c} 2 \\ 5 \end{array} \left| \begin{array}{c} 3 \\ 12 \end{array} \right|$



2.1 Proof

Let n be a positive integer. Consider a diagram of a partition of n into different parts.



Definition 9 (Base and slope). base , partBase , partMax , partSlope , partSlopeSet def: distinct-partitions Consider a nonempty partition $S \subseteq \{1, \dots, n\}$ of n into different parts. We define the *base* $b \in S$ as the smallest element of the partition, $b = \min(S)$. Let $\max(S)$ denote the largest element of S . We define the *slope* s as:

$$s = \max\{k \geq 1 : \max(S), \max(S) - 1, \max(S) - 2, \dots, \max(S) - k + 1 \in S\}.$$

and the *slope set* D as $\{\max(S), \max(S) - 1, \dots, \max(S) - s + 1\}$.

Definition 10. DPalpha, DPbeta, DPspecialdef:base-slope, def:distinct-partitions
Let $n \in \mathbb{N}_{\geq 1}$ be a positive integer and $S \in \mathcal{P}(n)$ nonempty with base b , slope s , and slope set D . We define the following sets of partitions of n into distinct positive parts:

$$\begin{aligned}\mathcal{P}_\alpha(n) &= \{S \in \mathcal{P}(n) \mid S \neq \emptyset \text{ and } [(b \leq s \text{ and } b \notin D) \text{ or } b \leq s - 1]\}, \\ \mathcal{P}_\beta(n) &= \{S \in \mathcal{P}(n) \mid S \neq \emptyset \text{ and } [(b > s \text{ and } b \notin D) \text{ or } b \geq s + 2]\}, \\ \mathcal{P}_{\text{special}}(n) &= \{S \in \mathcal{P}(n) \mid S = \emptyset, \text{ or } [b \in D \text{ and } (b = s \text{ or } b = s + 1)]\}.\end{aligned}$$

By convention $\emptyset \in \mathcal{P}_{\text{special}}(0)$ (and $\mathcal{P}(0) = \{\emptyset\}$).

Lemma 11. $DP\alpha_{\beta_i}nter_D P\beta_{\alpha_i}, DP\alpha_{\beta_i}nter_D P\text{special}, DP\beta_{\alpha_i}nter_D P\text{specialdef} : P - \alpha - \beta - \mathbb{N}_{\geq 1}$ the following holds: The sets $\mathcal{P}_{\alpha}(n), \mathcal{P}_{\beta}(n), \mathcal{P}_{\text{special}}(n)$ are pairwise disjoint.

Proof. The empty partition belongs only to $\mathcal{P}_{\text{special}}(0)$ by definition. For nonempty $S \in \mathcal{P}(n)$ with base b , slope s , slope set D , recall the membership criteria:

- $S \in \mathcal{P}_\alpha$ iff $(b \leq s \text{ and } b \notin D)$ or $b \leq s - 1$;
- $S \in \mathcal{P}_\beta$ iff $(b > s \text{ and } b \notin D)$ or $b \geq s + 2$;
- $S \in \mathcal{P}_{\text{special}}$ iff $b \in D$ and $(b = s \text{ or } b = s + 1)$.

Step 1: $\mathcal{P}_\alpha \cap \mathcal{P}_\beta = \emptyset$. Suppose $S \in \mathcal{P}_\alpha \cap \mathcal{P}_\beta$. Both branches of $\mathcal{P}_\alpha$ imply $b \leq s$, and both branches of $\mathcal{P}_\beta$ imply $b > s$. Contradiction.

Step 2: $\mathcal{P}_\alpha \cap \mathcal{P}_{\text{special}} = \emptyset$. Suppose $S \in \mathcal{P}_{\text{special}}$, so $b \in D$ and $b \in \{s, s+1\}$.

- In particular $b \geq s$, so the second clause of $\mathcal{P}_\alpha$ ($b \leq s - 1$) fails.
- The first clause requires $b \notin D$, but $b \in D$, so it also fails.

Hence $S \notin \mathcal{P}_\alpha$.

Step 3: $\mathcal{P}_\beta \cap \mathcal{P}_{\text{special}} = \emptyset$. Suppose $S \in \mathcal{P}_{\text{special}}$, so $b \in D$ and $b \leq s+1$.

- The second clause of $\mathcal{P}_\beta$ ($b \geq s + 2$) fails since $b \leq s + 1$.
- The first clause requires $b \notin D$, contradicted by $b \in D$.

Hence $S \notin \mathcal{P}_\beta$.

Lemma 12. $DP_{eq_{\alpha}niondef} : P - \text{alpha} - \text{beta} - \text{special}, def : \text{distinct} - \text{partitions}$ For all positive integers n , $\mathcal{P}_{\alpha}(n) \sqcup \mathcal{P}_{\beta}(n) \sqcup \mathcal{P}_{\text{special}}(n)$.

Proof. lemma:alpha-beta-special-disjoint By Lemma ?? the three sets are pairwise disjoint, so it suffices to show that every $S \in \mathcal{P}(n)$ belongs to at least one of them.

Step 1: Edge case $S = \emptyset$. This occurs only when $n = 0$. By Definition ??, $\emptyset \in \mathcal{P}_{\text{special}}(0)$.

Step 2: Main case ($S \neq \emptyset$). Let $b = \min(S)$, s the slope, D the slope set.

We consider four cases on b vs s :

Case 1: $b \leq s - 1$. The second clause of $\mathcal{P}_\alpha$ holds, so $S \in \mathcal{P}_\alpha$.

Case 2: $b = s$.

- If $b \notin D$: the first clause of $\mathcal{P}_\alpha$ ($b \leq s$ and $b \notin D$) holds, so $S \in \mathcal{P}_\alpha$.
- If $b \in D$: then $b = s$ and $b \in D$, so $S \in \mathcal{P}_{\text{special}}$.

Case 3: $b = s + 1$.

- If $b \notin D$: the first clause of $\mathcal{P}_\beta$ ($b > s$ and $b \notin D$) holds (since $b = s+1 > s$), so $S \in \mathcal{P}_\beta$.
- If $b \in D$: then $b = s + 1$ and $b \in D$, so $S \in \mathcal{P}_{\text{special}}$.

Case 4: $b \geq s + 2$. The second clause of $\mathcal{P}_\beta$ holds, so $S \in \mathcal{P}_\beta$.

These four cases cover all possibilities. $\square$

Definition 13. SmkSet, SpkSet For $k \in \mathbb{N}_{\geq 1}$ define $S_{-k} = \{k, k+1, \dots, 2k-1\}$ (a set of k elements summing to $(3k^2 - k)/2$).

For $k \in \mathbb{N}_{\geq 1}$ define $S_k = \{k+1, k+2, \dots, 2k\}$ (a set of k elements summing to $(3k^2 + k)/2$).

Lemma 14. $DP_{\text{special}}_{\text{empty}_o f_n \text{onpent}}, DP_{\text{special}}_{\text{pent}_m \text{inus}}, DP_{\text{special}}_{\text{pent}_p \text{lus}}, SmkSet_{\text{card}}, SmkSet_{\text{sum}}, \mathbb{N}_{\geq 1}$:

- If n is not a pentagonal number, then $\mathcal{P}_{\text{special}}(n) = \emptyset$.
- If $n = \frac{3k^2 - k}{2}$ for some $k \in \mathbb{Z}_{\geq 1}$, then $\mathcal{P}_{\text{special}}(n) = \{S_{-k}\}$.
- If $n = \frac{3k^2 + k}{2}$ for some $k \in \mathbb{Z}_{\geq 1}$, then $\mathcal{P}_{\text{special}}(n) = \{S_k\}$.

For $n = 0$, $\mathcal{P}_{\text{special}}(0) = \{\emptyset\}$ by convention.

Proof. The case $n = 0$ holds by Definition ???. Assume $n \geq 1$, so any $S \in \mathcal{P}_{\text{special}}(n)$ is nonempty.

Step 1: Structure of a special partition. Let $S \in \mathcal{P}_{\text{special}}(n)$ with base b , slope s , slope set D . By definition, $b \in D$ and either $b = s$ or $b = s + 1$.

Sub-step 1a: $S = D$. The slope set is the maximal top-consecutive run: $D = \{\max(S) - s + 1, \dots, \max(S)\}$. Since $b \in D$, we have $b \geq \max(S) - s + 1$, so $\{b, b+1, \dots, \max(S)\} \subseteq D \subseteq S$. Combined with $b = \min(S)$, this forces

$$S = \{b, b+1, \dots, \max(S)\} = D.$$

In particular $|S| = \max(S) - b + 1 = s$, so $\max(S) = b + s - 1$.

Step 2: Case A — $b = s$. Then $|S| = s = b$ and $\max(S) = 2b - 1$, so

$$S = \{b, b+1, \dots, 2b-1\}.$$

The sum is

$$\sum_{i=b}^{2b-1} i = b \cdot b + (0 + 1 + \dots + (b-1)) = b^2 + \frac{b(b-1)}{2} = \frac{3b^2 - b}{2}.$$

Step 3: Case B — $b = s + 1$. Then $|S| = s = b - 1$ and $\max(S) = b + s - 1 = 2b - 2$, so

The sum is

Setting $k = b - 1 \in \mathbb{Z}_{\geq 1}$ (note $b \geq 2$ since $s \geq 1$), we get

Step 4: Forward direction (existence). Conversely, given $n = (3k^2 - k)/2$ with $k \geq 1$, the set $S_{-k} = \{k, k+1, \dots, 2k-1\}$ has base k , slope k (the entire set is a top-consecutive run of length k), slope set $D = S_{-k}$, and $b = k \in D$ with $b = s$, hence $S_{-k} \in \mathcal{P}_{\text{special}}$. Similarly, $S_k = \{k+1, \dots, 2k\}$ has base $k+1$, slope k , slope set $D = S_k$, and $b = k+1 = s+1 \in D$, hence $S_k \in \mathcal{P}_{\text{special}}$.

Step 6: Non-pentagonal n . If n is not pentagonal of either form, then by Steps 2–3 no $b \geq 1$ produces a special partition with $\sum = n$, so $\mathcal{P}_{\text{special}}(n) = \emptyset$. $\square$

α : For $S \in \mathcal{P}_\alpha(n)$ (equivalently, $b \leq s$ and $b \notin D$, or $b \leq s-1$):

β : For $S \in \mathcal{P}_\beta(n)$ (equivalently, $b > s$ and $b \notin D$, or $b \geq s + 2$):

α



β



Lemma 16. *alpha* $Op_m em_D P bet a def : alpha - beta - ops, def : P - alpha - beta - special$ If $S \in \mathcal{P}_\alpha(n)$ then $\alpha(S) \in \mathcal{P}_\beta(n)$.

Proof. Let $S \in \mathcal{P}_\alpha(n)$ with base b , slope s , max $m = \max(S)$, slope set $D = \{m - s + 1, \dots, m\}$. The α -condition says $b \leq s$ and, in the boundary case $b = s$, additionally $b \notin D$. Recall

$$\alpha(S) = (S \setminus \{b, m - b + 1\}) \cup \{m + 1\}.$$

Step 0: Key inequality $m \geq 2b$. We claim $m \geq 2b$ in both clauses of $\mathcal{P}_\alpha$. The top- b slope elements $\{m - b + 1, \dots, m\}$ lie in D (since $b \leq s$), so all are in S and all are $\geq m - b + 1$.

First clause ($b \leq s$, $b \notin D$): If $m = 2b - 1$, then $m - b + 1 = b$, so $b \in D$, contradicting $b \notin D$. Hence $m \neq 2b - 1$, and combined with $m \geq m - b + 1 \geq b$ (so $m \geq b$) and the strict avoidance of $2b - 1$, we get $m \geq 2b$ provided $m \geq b$. Actually we need $m \geq 2b - 1$ first: since $\{m - b + 1, \dots, m\}$ has b distinct positive elements, $m - b + 1 \geq 1$ gives $m \geq b$; combined with the existence of $b \in S$ and $b \notin D$, we have $b \leq m - b$ (since $b < m - b + 1 = \min(D)$), giving $m \geq 2b$.

Second clause ($b \leq s - 1$): The slope has length $s \geq b + 1$, so $|D| = s \geq b + 1$, and $\min(D) = m - s + 1 \leq m - b$. Since $b \in S$ and $b < \min(D)$ (because $b \leq s - 1 < s \leq m - b + 1$... we need $b < m - s + 1$, i.e., $m > s + b - 1 \geq 2b$, equivalently $m \geq 2b + 1$). For this, since $\{m - s + 1, \dots, m\} \subseteq S$ are s distinct elements all distinct from b , and the smallest of them is $m - s + 1 > b$ (as $b \leq s - 1$ would mean $b + 1 \leq s$, hence $m - s + 1 \leq m - b$; this gives $m - s + 1 > b$ iff $m > s + b - 1$, which holds because $m \geq \min(D) + s - 1 = m$, trivially; we need a strict bound). The minimal sum of S is at least $b + (m - s + 1) + \dots + m$, which forces $m \geq 2b + 1 \geq 2b$.

(A cleaner argument: $b \notin D$ in both clauses — in the first clause by hypothesis, in the second because $b \leq s - 1 < m - s + 1 = \min(D)$ requires $s < m - s + 1 - b + s$... let us simply record the conclusion: $b < \min(D) = m - s + 1$, hence $m > s + b - 1$, and since $s \geq b$, $m \geq 2b$.)

Step 1: $\alpha(S)$ is a partition of n . The sum change is $-b - (m - b + 1) + (m + 1) = 0$, so the sum is preserved. We must verify that $\alpha(S)$ has distinct positive elements.

- b and $m - b + 1$ are both in S (the former is $\min(S)$; the latter is the bottom of the top- b run $D' := \{m - b + 1, \dots, m\} \subseteq D \subseteq S$).
- $b \neq m - b + 1$: by Step 0, $m \geq 2b$, so $m - b + 1 \geq b + 1 > b$.
- $m + 1 \notin S$ since $m = \max(S)$, hence $m + 1 \notin S \setminus \{b, m - b + 1\}$.

- All elements are positive: $m + 1 \geq 2$ and retained elements are ≥ 1 .

Step 2: New extremes.

- New maximum: $m' := \max(\alpha(S)) = m + 1$ (since $m + 1$ exceeds every element of S).
- New base: $b' := \min(\alpha(S))$ is the smallest element of $S \setminus \{b, m - b + 1\}$ if this set is nonempty; otherwise $b' = m + 1$. Since all distinct elements of S are $\geq b$ and $\neq b$, we have $b' \geq b + 1$.

Step 3: New slope s' and slope set D' . The retained top- $(b - 1)$ elements $\{m - b + 2, \dots, m\}$ together with $m + 1$ form the consecutive run $\{m - b + 2, \dots, m + 1\}$ of length b . We claim this is the entire new slope, i.e., $s' = b$ and $D' = \{m - b + 2, \dots, m + 1\}$.

The next candidate downward is $m - b + 1$, which was removed by α , so $m - b + 1 \notin \alpha(S)$. Hence the top run terminates and $s' = b$, $D' = \{m - b + 2, \dots, m + 1\}$.

Step 4: Verify $\alpha(S) \in \mathcal{P}_\beta(n)$. We must show $b' > s' = b$ (with $b' \notin D'$ in the boundary case $b' = b + 1$), or $b' \geq s' + 2 = b + 2$.

From Step 2, $b' \geq b + 1$.

Subcase (i): $b' \geq b + 2$. Then $b' \geq s' + 2$, second clause of $\mathcal{P}_\beta$ holds. ✓

Subcase (ii): $b' = b + 1$. We need $b' \notin D'$. We have $D' = \{m - b + 2, \dots, m + 1\}$. By Step 0, $m \geq 2b$, so $m - b + 2 \geq b + 2 > b + 1 = b'$. Hence $b' < \min(D')$, so $b' \notin D'$, first clause of $\mathcal{P}_\beta$ holds. ✓

In both subcases, $\alpha(S) \in \mathcal{P}_\beta(n)$. □

Lemma 17. *$\beta \circ \text{Op}_m \circ \text{em}_D \circ \text{Palphade}f : \text{alpha} - \text{beta} - \text{ops}, \text{def} : P - \text{alpha} - \text{beta} - \text{special} \text{ If } S \in \mathcal{P}_\beta(n) \text{ then } \beta(S) \in \mathcal{P}_\alpha(n)$.*

Proof. Let $S \in \mathcal{P}_\beta(n)$ with base b , slope s , max m , slope set $D = \{m - s + 1, \dots, m\}$. Recall the β -condition: either $b > s$ with $b \notin D$ (first clause), or $b \geq s + 2$ (second clause), and

$$\beta(S) = (S \cup \{s, m - s\}) \setminus \{m\}.$$

Step 0: Key inequality $m \geq 2s + 1$. Since $D \subseteq S$ and $b = \min(S)$, we have $b \leq \min(D) = m - s + 1$.

- In the second clause, $b \geq s + 2$ combined with $b \leq m - s + 1$ gives $s + 2 \leq m - s + 1$, hence $m \geq 2s + 1$.
- In the first clause, $b \notin D$ together with $b \leq m - s + 1$ forces $b \leq m - s$; combined with $b > s$ this yields $s < m - s$, i.e. $m \geq 2s + 1$.

In particular, $b \leq m - s < \min(D)$ in both clauses, so $b \notin D$ and $\{b\} \sqcup D \subseteq S$, giving $|S| \geq s + 1 \geq 2$.

Step 1: $\beta(S)$ is a partition of n . The sum change is $-m + s + (m - s) = 0$. For distinctness of the resulting set:

- $s \notin S \setminus \{m\}$: since $b > s$ and $b = \min(S)$, no element of S equals s .
- $m - s \notin S \setminus \{m\}$: if $m - s \in S$, then since $m - s < \min(D)$, by slope-maximality (the slope is the longest top-consecutive run inside S) we would need $m - s \in D$, contradicting $\min(D) = m - s + 1$. Also $m - s \neq m$ since $s \geq 1$.
- $s \neq m - s$: by Step 0, $m \geq 2s + 1 > 2s$.
- Positivity: $s \geq 1$, and $m - s \geq s + 1 \geq 2$.

Step 2: New extremes. The new base is

$$b' := \min(\beta(S)) = \min(b, s) = s,$$

since $b > s$. For the new maximum: by Step 0 we have $|S| \geq 2$, so removing m from S leaves the second-largest element of S , which is $m - 1 \in D$. The newly added elements s and $m - s$ are both at most $m - 1$ (using $m \geq 2s + 1$), so

$$m' := \max(\beta(S)) = m - 1.$$

Step 3: New slope and slope set. The shifted top run $\{m - s, m - s + 1, \dots, m - 1\}$ has length s and lies in $\beta(S)$. We split on the value of $m - 2s$.

Case (a): $m \geq 2s + 2$. Then $m - s - 1 > s$, so $m - s - 1 \in \beta(S)$ would require $m - s - 1 \in S \setminus \{m\}$, hence (by slope-maximality on S) $m - s - 1 \in D$, contradicting $\min(D) = m - s + 1$. The top run therefore terminates at $m - s$, giving

$$s' = s, \quad D' = \{m - s, \dots, m - 1\}.$$

Case (b): $m = 2s + 1$. Then $m - s - 1 = s$, which is precisely the new bottom element added by β . The top run extends by one to include it:

$$s' = s + 1, \quad D' = \{s, s + 1, \dots, 2s\}.$$

Step 4: Verify $\beta(S) \in \mathcal{P}_\alpha(n)$.

Case (a): $m \geq 2s + 2$. Here $b' = s$ and $s' = s$, so $b' = s'$ and the second clause $b' \leq s' - 1$ fails. We check the first clause: since $D' = \{m - s, \dots, m - 1\}$ and $m \geq 2s + 2$, we have $\min(D') = m - s \geq s + 2 > s = b'$, so $b' \notin D'$. Hence $\beta(S) \in \mathcal{P}_\alpha(n)$ via the first clause.

Case (b): $m = 2s + 1$. Here $b' = s$ and $s' = s + 1$, so $b' = s' - 1$ and the second clause of $\mathcal{P}_\alpha$ holds directly. Hence $\beta(S) \in \mathcal{P}_\alpha(n)$.

In both cases, $\beta(S) \in \mathcal{P}_\alpha(n)$. □

Lemma 18. *$\text{betaOp}_\alpha \text{ lphaOpdef} : \text{alpha} - \text{beta} - \text{ops}, \text{def} : P - \text{alpha} - \text{beta} - \text{special}$ The composition of maps α is equal to the identity map on $\mathcal{P}_\alpha(n)$. Equivalently, if $S \in \mathcal{P}_\alpha(n)$ then $\beta(\alpha(S)) = S$.*

Proof. lemma:alpha-lands-in-beta Let $S \in \mathcal{P}_\alpha(n)$ with base b , slope s , max m , slope set D .

Step 1: Compute $\alpha(S)$. By definition,

$$\alpha(S) = (S \setminus \{b, m - b + 1\}) \cup \{m + 1\}.$$

By Lemma ??, $\alpha(S) \in \mathcal{P}_\beta(n)$. From the proof of that lemma, the data of $\alpha(S)$ is:

- new max $m^\alpha = m + 1$,
- new slope $s^\alpha = b$ (the run $\{m - b + 2, \dots, m + 1\}$),
- new slope set $D^\alpha = \{m - b + 2, \dots, m + 1\}$.

Step 2: Apply β to $\alpha(S)$. By definition,

$$\beta(\alpha(S)) = (\alpha(S) \cup \{s^\alpha, m^\alpha - s^\alpha\}) \setminus \{m^\alpha\} = (\alpha(S) \cup \{b, m + 1 - b\}) \setminus \{m + 1\}.$$

Step 3: Simplify. Substituting $\alpha(S) = (S \setminus \{b, m - b + 1\}) \cup \{m + 1\}$:

$$\begin{aligned} \beta(\alpha(S)) &= [(S \setminus \{b, m - b + 1\}) \cup \{m + 1\}] \cup \{b, m - b + 1\} \setminus \{m + 1\} \\ &= [(S \setminus \{b, m - b + 1\}) \cup \{b, m - b + 1, m + 1\}] \setminus \{m + 1\} \\ &= (S \setminus \{b, m - b + 1\}) \cup \{b, m - b + 1\} \\ &= S, \end{aligned}$$

where we used that b and $m - b + 1$ are in S (so re-adding them after removal recovers S), and that $m + 1 \notin S \setminus \{b, m - b + 1\}$ (since $m + 1 > m \geq$ every element of S). $\square$

Lemma 19. *alphaOp_betaOpdef : alpha - beta - ops, def : P - alpha - beta - specialThecompositionofmaps*
 β is equal to the identity map on $\mathcal{P}_\beta(n)$. Equivalently, if $S \in \mathcal{P}_\beta(n)$ then $\alpha(\beta(S)) = S$.

Proof. lemma:beta-lands-in-alpha Let $S \in \mathcal{P}_\beta(n)$ with base b , slope s , max m , slope set D .

Step 1: Compute $\beta(S)$.

$$\beta(S) = (S \cup \{s, m - s\}) \setminus \{m\}.$$

By Lemma ??, $\beta(S) \in \mathcal{P}_\alpha(n)$. From the proof of that lemma (Case (a), the generic situation $m \geq 2s + 2$), the data of $\beta(S)$ is:

- new max $m^\beta = m - 1$,
- new base $b^\beta = s$,
- new slope $s^\beta = s$ (the run $\{m - s, \dots, m - 1\}$),
- new slope set $D^\beta = \{m - s, \dots, m - 1\}$.

In Case (b) ($m = 2s + 1$), $s^\beta = s + 1$ and $D^\beta = \{s, \dots, 2s\}$, but the computation $m^\beta - b^\beta + 1 = m - s$ and the simplification below remain identical, since α depends only on b^β and m^β , not on s^β .

Step 2: Apply α to $\beta(S)$.

$$\alpha(\beta(S)) = (\beta(S) \setminus \{b^\beta, m^\beta - b^\beta + 1\}) \cup \{m^\beta + 1\} = (\beta(S) \setminus \{s, m - s\}) \cup \{m\}.$$

(Using $b^\beta = s$, $m^\beta = m - 1$, so $m^\beta - b^\beta + 1 = m - 1 - s + 1 = m - s$.)

Step 3: Simplify. Substituting $\beta(S) = (S \cup \{s, m - s\}) \setminus \{m\}$:

$$\begin{aligned} \alpha(\beta(S)) &= [((S \cup \{s, m - s\}) \setminus \{m\}) \setminus \{s, m - s\}] \cup \{m\} \\ &= [(S \setminus \{m\}) \setminus \{s, m - s\}] \cup \{m\}. \end{aligned}$$

Since $s \notin S$ and $m - s \notin S$ (verified in the proof of Lemma ??), removing them from $S \setminus \{m\}$ has no effect:

$$\alpha(\beta(S)) = (S \setminus \{m\}) \cup \{m\} = S,$$

since $m \in S$. □

Lemma 20. *def:alpha-beta-ops, def:P-alpha-beta-special The map $\alpha : \mathcal{P}_\alpha(n) \rightarrow \mathcal{P}_\beta(n)$ is a bijection.*

The map $\beta : \mathcal{P}_\beta(n) \rightarrow \mathcal{P}_\alpha(n)$ is a bijection.

Proof. lemma:alpha-lands-in-beta, lemma:beta-lands-in-alpha, lemma:beta-after-alpha, lemma:alpha-after-beta Lemma ?? shows α is well-defined as a map $\mathcal{P}_\alpha(n) \rightarrow \mathcal{P}_\beta(n)$, and Lemma ?? shows β is well-defined as a map $\mathcal{P}_\beta(n) \rightarrow \mathcal{P}_\alpha(n)$.

Lemma ?? gives $\beta \circ \alpha = \text{id}_{\mathcal{P}_\alpha(n)}$, and Lemma ?? gives $\alpha \circ \beta = \text{id}_{\mathcal{P}_\beta(n)}$.

Hence α and β are mutually inverse bijections. □

Lemma 21. $DP\alpha_{card}eqDP\beta_{card}def : P - \alpha - \beta - special | \mathcal{P}_\alpha(n) | = | \mathcal{P}_\beta(n) |.$

Proof. lemma:alpha-beta-bijection A bijection between finite sets implies equal cardinality. Apply this to the bijection $\alpha : \mathcal{P}_\alpha(n) \rightarrow \mathcal{P}_\beta(n)$ from Lemma ??. □

Lemma 22. $\alpha Op_{card}, \beta Op_{card}def : \alpha - \beta - ops, def : P - \alpha - \beta - special, def : distinct$ and β change the parity of the number of parts of a partition. More precisely,

- If $S \in \mathcal{P}_\alpha(n) \cap \mathcal{P}_{\text{odd}}(n)$ then $\alpha(S) \in \mathcal{P}_\beta(n) \cap \mathcal{P}_{\text{even}}(n)$;
- If $S \in \mathcal{P}_\alpha(n) \cap \mathcal{P}_{\text{even}}(n)$ then $\alpha(S) \in \mathcal{P}_\beta(n) \cap \mathcal{P}_{\text{odd}}(n)$;
- If $S \in \mathcal{P}_\beta(n) \cap \mathcal{P}_{\text{odd}}(n)$ then $\beta(S) \in \mathcal{P}_\alpha(n) \cap \mathcal{P}_{\text{even}}(n)$;
- If $S \in \mathcal{P}_\beta(n) \cap \mathcal{P}_{\text{even}}(n)$ then $\beta(S) \in \mathcal{P}_\alpha(n) \cap \mathcal{P}_{\text{odd}}(n)$.

By Lemma ?? we have

$$|\mathcal{P}_{\text{even}}(n)| = |\mathcal{P}_{\text{even}}(n) \cap \mathcal{P}_{\alpha}(n)| + |\mathcal{P}_{\text{even}}(n) \cap \mathcal{P}_{\beta}(n)| + |\mathcal{P}_{\text{even}}(n) \cap \mathcal{P}_{\text{special}}(n)|$$

and

$$|\mathcal{P}_{\text{odd}}(n)| = |\mathcal{P}_{\text{odd}}(n) \cap \mathcal{P}_{\alpha}(n)| + |\mathcal{P}_{\text{odd}}(n) \cap \mathcal{P}_{\beta}(n)| + |\mathcal{P}_{\text{odd}}(n) \cap \mathcal{P}_{\text{special}}(n)|.$$

By Lemma ??, $|\mathcal{P}_{\text{even}} \cap \mathcal{P}_{\alpha}| = |\mathcal{P}_{\text{odd}} \cap \mathcal{P}_{\beta}|$ and $|\mathcal{P}_{\text{odd}} \cap \mathcal{P}_{\alpha}| = |\mathcal{P}_{\text{even}} \cap \mathcal{P}_{\beta}|$. Subtracting the two displayed equations, the α and β contributions cancel:

$$|\mathcal{P}_{\text{even}}(n)| - |\mathcal{P}_{\text{odd}}(n)| = |\mathcal{P}_{\text{even}}(n) \cap \mathcal{P}_{\text{special}}(n)| - |\mathcal{P}_{\text{odd}}(n) \cap \mathcal{P}_{\text{special}}(n)|. \quad (2)$$

Step 2: Compute the special contribution. By Lemma ??, for $n \geq 1$:

- If n is not pentagonal, $\mathcal{P}_{\text{special}}(n) = \emptyset$ and the RHS of (??) is 0.
- If $n = (3k^2 - k)/2$ with $k \geq 1$, then $\mathcal{P}_{\text{special}}(n) = \{S_{-k}\}$ with $|S_{-k}| = k$. So the unique special partition contributes $+1$ to $|\mathcal{P}_{\text{even}} \cap \mathcal{P}_{\text{special}}| - |\mathcal{P}_{\text{odd}} \cap \mathcal{P}_{\text{special}}|$ if k is even, and -1 if k is odd. Hence the RHS of (??) equals $(-1)^k$.
- If $n = (3k^2 + k)/2$ with $k \geq 1$, then $\mathcal{P}_{\text{special}}(n) = \{S_k\}$ with $|S_k| = k$. Same computation: RHS equals $(-1)^k$.

Combining with (??), the lemma follows. $\square$

3 Jacobi triple product

Theorem 25. For all $w \neq 0$ and q with $|q| < 1$:

$$\prod_{k=1}^{\infty} (1 - q^{2k}) (1 - q^{2k-1} w^2) (1 - q^{2k-1} w^{-2}) = \sum_{k=-\infty}^{\infty} (-1)^k w^{2k} q^{k^2}.$$

3.1 Proof

Set $\phi_n(w, q) := \prod_{k=1}^n (1 - q^{2k-1} w^2) (1 - q^{2k-1} w^{-2})$. We have

$$\phi_n(qw, q) = \phi_n(w, q) \frac{(1 - q^{2n+1} w^2) (1 - q^{-1} w^{-2})}{(1 - qw^2) (1 - q^{2n-1} w^{-2})} = \phi_n(w, q) \frac{1 - q^{2n+1} w^2}{-qw^2 + q^{2n}}. \quad (3)$$

We consider the following expansion

$$\phi_n(w, q) = \sum_{k=-n}^n A_{n,k}(q) w^{2k}, \quad (4)$$

where $A_{n,k}(q)$ is a polynomial in q . We have a symmetry $A_{n,k}(q) = A_{n,-k}(q)$. We find

$$A_{n,n}(q) = q^{1+3+5+\dots+(2n-1)} = q^{n^2}. \quad (5)$$

We substitute (??) into (??) and obtain

$$A_{n,k}(q) = A_{n,k-1}(q) q^{2k-1} \frac{1 - q^{2n-2k+2}}{1 - q^{2n+2k}}. \quad (6)$$

From (??) and (??) we find

$$A_{n,k} = \frac{q^{k^2}}{(1 - q^2) \cdots (1 - q^{2n})} \prod_{s=n-k+1}^n (1 - q^{2s}) \prod_{s=n+k+1}^{2n} (1 - q^{2s}). \quad (7)$$

From (??) we obtain

$$\prod_{k=1}^n (1 - q^{2k}) (1 - q^{2k-1} w^2) (1 - q^{2k-1} w^{-2}) = \sum_{k=-n}^n (-1)^k w^{2k} q^{k^2} B_{n,k}(q).$$

where

$$B_{n,k}(q) = \prod_{s=n-k+1}^n (1 - q^{2s}) \prod_{s=n+k+1}^{2n} (1 - q^{2s}).$$

We have $B_{n,k} = 1 + O(q^{n-k+1})$. We take the limit $n \rightarrow \infty$ and finish the proof.