



# STATISTICS FOR DATA SCIENCE

## HYPOTHESIS and INFERENCE

---

**Dr. Deepa Nair**  
Department of Science and Humanities

# STATISTICS FOR DATA SCIENCE

---

## UNIT-4      HYPOTHESIS and INFERENCE

### Session-3

### Large-Sample Tests for a Population Mean

**Dr. Deepa Nair**

Department of Science and Humanities

# STATISTICS FOR DATA SCIENCE

## Large-Sample Tests for a Population Mean

---



- Let  $X_1, X_2, \dots, X_n$  be a *large* (e.g.,  $n > 30$ ) sample from a population with mean  $\mu$  and standard deviation  $\sigma$ .
- To test a null hypothesis of the form
- $H_0: \mu \leq \mu_0, H_0: \mu \geq \mu_0, \text{ or } H_0: \mu = \mu_0$

# STATISTICS FOR DATA SCIENCE

## Large-Sample Tests for a Population Mean

---



- Compute the z-score:

$$z = \frac{\bar{X} - \mu_0}{\sigma / \sqrt{n}}$$

If  $\sigma$  is unknown it may be approximated with  $s$ .

# STATISTICS FOR DATA SCIENCE

## Large-Sample Tests for a Population Mean

---



- Compute the  $P$ -value.
- The  $P$ -value is an area under the normal curve, which depends on the alternate hypothesis as in the table:

# STATISTICS FOR DATA SCIENCE

## Large-Sample Tests for a Population Mean

Alternate Hypothesis	<i>P</i> -value
$H_1 : \mu > \mu_0$	Area to the right of $z$
$H_1 : \mu < \mu_0$	Area to the left of $z$
$H_1 : \mu = \mu_0$	Sum of the areas in the tails cut off by $z$ and $-z$

# STATISTICS FOR DATA SCIENCE

## Large-Sample Tests for a Population Mean

---



### Example:

- The mean lifetime of a sample of 100 fluorescent light bulbs produced by a company is computed to be 1570 hours with a standard deviation of 120 hours.
- If  $\mu$  is the life time of all the bulbs produced by the company test the hypothesis  $\mu = 1600 \text{ hours}$  against the alternate hypothesis  $\mu \neq 1600 \text{ hours}$ .

# STATISTICS FOR DATA SCIENCE

## Large-Sample Tests for a Population Mean

---



**Solution:**

$$H_0: \mu = 1600 \text{ hours}$$

$$H_1: \mu \neq 1600 \text{ hours}$$

$$\bar{X} = 1570, \sigma = s = 120, n = 100$$

$$z = \frac{\bar{X} - \mu_0}{\sigma / \sqrt{n}} = \frac{1570 - 1600}{120 / \sqrt{100}} = -2.51$$

P- Value is 0.012.

Which is very low . So we reject  $H_0$



# STATISTICS FOR DATA SCIENCE

## Large-Sample Tests for a Population Mean

---



### Example:

- A trucking firm is suspicious of the claim that the average lifetime of certain tires is at least 28,000 miles.
- To check the claim, the firm puts 40 of these tires on its trucks and gets a mean lifetime of 27,463 with a standard deviation 1,348 miles.

Find the P value for testing  $H_0: \mu \geq 28,000 \text{ miles}$   $H_1: \mu < 28,000 \text{ miles}$ ?

# STATISTICS FOR DATA SCIENCE

## Large-Sample Tests for a Population Mean

---



**Solution:**

$$H_0: \mu \geq 28,000 \text{ miles}$$

$$H_1: \mu < 28,000 \text{ miles}$$

$$z = \frac{\bar{X} - \mu}{s/\sqrt{n}} = \frac{27,463 - 28,000}{1,348/\sqrt{40}} = -2.52 < -2.33$$

**P – Value is 0.0059**

**Since P- Value is a very small probability we need to reject  $H_0$**



**Dr. Deepa Nair**

Department of Science and Humanities

---

**[deepanair@pes.edu](mailto:deepanair@pes.edu)**