

# STATISTICS FOR DATA SCIENCE HYPOTHESIS and INFERENCE

Dr. Deepa Nair

Department of Science and Humanities



**UNIT-4** HYPOTHESIS and INFERENCE

**Session-4** 

**Drawing Conclusions from the Results of Hypothesis Tests** 

Dr. Deepa Nair

Department of Science and Humanities

# **Drawing Conclusions from the Results of Hypothesis Tests**



- The only two conclusions that can be reached in a hypothesis test are that
- $H_0$  is false or that  $H_0$  is plausible.
- One can never conclude that  $H_0$  is true.

# **Drawing Conclusions from the Results of Hypothesis Tests**



- How do we know when to reject  $H_0$ ?
- The smaller the P-value, the less plausible  $H_0$  becomes.
- A common rule of thumb is to draw the line at 5%. According to this rule of thumb, if  $P \leq 0.05$ ,  $H_0$  is rejected; otherwise  $H_0$  is not rejected.

## **Drawing Conclusions from the Results of Hypothesis Tests**



• The smaller the P-value, the more certain we can be that  $H_0$  is false.

• The larger the P-value, the more plausible  $H_0$  becomes, but we can never be certain that  $H_0$  is true.

# **Drawing Conclusions from the Results of Hypothesis Tests**



• There is no sharp dividing line between conclusive evidence against  $\boldsymbol{H}_0$ 

 So while this rule of thumb is convenient, it has no real scientific justification.

**Drawing Conclusions from the Results of Hypothesis Tests** 



• A rule of thumb suggests to reject  $H_0$  whenever  $P \leq 0.05$ .

## **Drawing Conclusions from the Results of Hypothesis Tests**



## **Statistical Significance:**

- Whenever the *P*-value is less than a particular threshold, the result is said to be "statistically significant" at that level.
- So, for example, if  $P \le 0.05$ , the result is statistically significant at the 5% level; if  $P \le 0.01$ , the result is statistically significant at the 1% level, and so on.

# **Drawing Conclusions from the Results of Hypothesis Tests**



## **Statistical Significance:**

• If a result is statistically significant at the  $100\alpha\%$  level, we can also say that the null hypothesis is "rejected at level  $100\alpha\%$ ."

# **Drawing Conclusions from the Results of Hypothesis Tests**



- The null hypothesis is rejected at the  $100\alpha\%$  level.
- When reporting the result of a hypothesis test, report the P -value, rather than just comparing it to 5% or 1%.

# **Drawing Conclusions from the Results of Hypothesis Tests**



- Let  $\alpha$  be any value between 0 and 1. Then, if  $P \leq \alpha$ ,
- The result of the test is said to be statistically significant at the  $100\alpha\%$  level.

## **Drawing Conclusions from the Results of Hypothesis Tests**



# The P-value Is Not the Probability That $H_0$ Is True

Since the P-value is a probability, and since small P-values indicate that  $H_0$  is unlikely to be true, it is tempting to think that the P-value represents the probability that  $H_0$  is true. This is emphatically not the case. The concept of probability discussed here is useful only when applied to outcomes that can turn out in different ways when experiments are repeated. It makes sense to define the P-value as the probability of observing an extreme value of a statistic such as X, since the value of X could come out differently if the experiment were repeated. The null hypothesis, on the other hand, either is true or is not true. The truth or falsehood of  $H_0$  cannot be changed by repeating the experiment. It is therefore not correct to discuss the "probability" that  $H_0$  is true.

## **Drawing Conclusions from the Results of Hypothesis Tests**



## **Example:**

- A hypothesis test is performed of the null hypothesis  $H_0$ :  $\mu=0$ . The P-value turns out to be 0.03.
- Is the result statistically significant at the 10% level? The 5% level? The 1% level?
- Is the null hypothesis rejected at the 10% level? The 5% level? The1% level?

## **Drawing Conclusions from the Results of Hypothesis Tests**



#### **Solution:**

- The result is statistically significant at any level greater than or equal to 3%.
- Thus it is statistically significant at the 10% and 5% levels, but not at the 1% level.
- Similarly, we can reject the null hypothesis at any level greater than or equal to  $3\,\%$
- So  $H_0$  is rejected at the 10% and 5% levels, but not at the 1% level.

# **Drawing Conclusions from the Results of Hypothesis Tests**



## **Example:**

- The length of life *X* of certain computers is approximately normally distributed with mean 800 hours and standard deviation 40 hours.
- If a random sample of 30 computers has an average life of 788 hours, test the null hypothesis that  $\mu=800\ hours$  against the alternate hypothesis that  $\mu\neq800\ hours$ . at
- 0.5% b) 1% c)4% d)5% e) 10% f)15% level of significance.

## **Drawing Conclusions from the Results of Hypothesis Tests**



#### **Solution:**

$$H_0$$
:  $\mu = 800 hours$  ,  $H_1$ :  $\mu \neq 800 hours$ 

$$\overline{X}=788$$
,  $n=30$ ,  $\mu=800$ ,  $\sigma=40$ 

$$z = \frac{\overline{X} - \mu_0}{\sigma / \sqrt{n}} = \frac{788 - 800}{40 / \sqrt{30}} = -1.643$$

# **Drawing Conclusions from the Results of Hypothesis Tests**



#### **Solution:**

Case 1:

$$\alpha = 0.5\% = 0.005$$

P- Value 0.101 > 0.005

So we need to reject the null hypothesis at 0.5% level.

# **Drawing Conclusions from the Results of Hypothesis Tests**



#### **Solution:**

Case 2:

$$\alpha = 1\% = 0.01$$

P- Value 0.101 > 0.01

So we need to reject the null hypothesis at 1% level.

# **Drawing Conclusions from the Results of Hypothesis Tests**



#### **Solution:**

## Case 3:

$$\alpha = 4\% = 0.04$$

P- Value 0.101 > 0.04

So we need to reject the null hypothesis at 4% level.

# **Drawing Conclusions from the Results of Hypothesis Tests**



## **Solution:**

Case 4:

$$\alpha = 5\% = 0.05$$

P- Value 0.101 > 0.05

So we need to reject the null hypothesis at 5% level.

# **Drawing Conclusions from the Results of Hypothesis Tests**



## **Solution:**

## Case 5:

$$\alpha = 10\% = 0.10$$

P- Value 0.101 > 0.10

So we need to reject the null hypothesis at 10% level.

# **Drawing Conclusions from the Results of Hypothesis Tests**



## **Solution:**

Case 6:

$$\alpha = 15\% = 0.15$$

P- Value 0.101 < 0.15

So we accept the null hypothesis at 15% level.

## **Drawing Conclusions from the Results of Hypothesis Tests**



## **Example:**

- Mice with an average life span of 32months will live up to 40 months when fed by a certain nutrious food.
- If 64 mice fed on this diet have an average life span of 38 months and standard deviation of 5.8 months.
- Is there any reason to believe that the average life span is less than 40 months.

## **Drawing Conclusions from the Results of Hypothesis Tests**



#### **Solution:**

Let us take 0.01 as the significance level.

*H*<sub>0</sub>: 
$$\mu \ge 40$$
 months, *H*<sub>1</sub>:  $\mu < 40$  months  $\overline{X} = 38$ ,  $n = 64$ ,  $\sigma \to s = 5.8$  
$$z = \frac{38 - 40}{5.8/\sqrt{64}} = -2.76$$

P- Value is 0.00290.0029 < 0.01

We need to reject  $H_0$ 

We will conclude that there is a reason to believe that the average life span of mice with nutrition food is less than 40 months



# Dr. Deepa Nair

Department of Science and Humanities

deepanair@pes.edu