



# STATISTICS FOR DATA SCIENCE

## HYPOTHESIS and INFERENCE

---

**Dr. Deepa Nair**  
Department of Science and Humanities

# STATISTICS FOR DATA SCIENCE

---

## UNIT-4      HYPOTHESIS and INFERENCE

### Session-11

### Fixed Level Testing

**Dr. Deepa Nair**

Department of Science and Humanities

# STATISTICS FOR DATA SCIENCE

## Fixed Level Testing

---



- A hypothesis test measures the plausibility of the null hypothesis by producing a  $P$ -value.
- The smaller the  $P$  –value, the less plausible the null. We have pointed out that there is no scientifically valid dividing line between plausibility and implausibility, Sometimes, however, a decision has to be made.

# STATISTICS FOR DATA SCIENCE

## Fixed Level Testing

---



- Fixed-level testing is just like the hypothesis testing we have been discussing so far, except that a firm rule is set ahead of time for rejecting the null hypothesis.

# STATISTICS FOR DATA SCIENCE

## Fixed Level Testing

---



### To conduct a fixed-level test:

- Choose a number  $\alpha$ , where  $0 < \alpha < 1$ . This is called the significance level, or the level, of the test.
- Compute the  $P$ -value in the usual way.
- If  $P \leq \alpha$ , reject  $H_0$ . If  $P > \alpha$ , do not reject  $H_0$ .

# STATISTICS FOR DATA SCIENCE

## Fixed Level Testing

---



### Example:

- The mean wear in a sample of 45 steel balls was  $\bar{X} = 673.2\mu m$ , and the standard deviation was  $s = 14.9\mu m$ .
- Let  $\mu$  denote the population mean wear. A test of  $H_1: \mu \geq 675$  versus  $H_0: \mu < 675$  yielded a  $P$ -value of 0.209.
- Can we reject  $H_0$  at the 25% level? Can we reject  $H_0$  at the 5% level?

# STATISTICS FOR DATA SCIENCE

## Fixed Level Testing

---



### Solution:

- The  $P$  –value of 0.209 is less than 0.25
- So if we had chosen a significance level of  $\alpha = 0.25$ , we would reject  $H_0$ .
- Thus we reject  $H_0$  at the 25% level.
- Since  $0.209 > 0.05$ , we do not reject  $H_0$  at the 5% level.

# STATISTICS FOR DATA SCIENCE

## Fixed Level Testing

---



### Example:

A process for a certain type of ore is designed to reduce the concentration of impurities to less than 2%.

- It is known that the standard deviation of impurities for processed ore is 0.6%.
- Let  $\mu$  represent the mean impurity level, in percent, for ore specimens treated by this process.



# STATISTICS FOR DATA SCIENCE

## Fixed Level Testing

---



### Example:

- The impurity of 80 ore specimens is measured, and a test of the hypothesis  $H_0: \mu \geq 2$  versus  $H_1: \mu < 2$  will be performed.
  - a. If the test is made at the 5% level, what is the rejection region?
  - b. If the sample mean impurity level is 1.85, will  $H_0$  be rejected at the 10% level?

# STATISTICS FOR DATA SCIENCE

## Fixed Level Testing

---



### Solution 2 (a)

- $H_0: \mu \geq 2$  versus  $H_1: \mu < 2$
- $\sigma = 0.6, n = 80$
- $\frac{\sigma}{\sqrt{n}} = 0.06708$
- Null distribution of  $\bar{X}$ :  
$$\bar{X} \sim N(2, 0.6^2/80)$$
- $\alpha = 0.05$

# STATISTICS FOR DATA SCIENCE

## Fixed Level Testing

---



### Solution 2 (a)

- $Z = -1.645$  (Critical value for 5%)

The rejection region is

$$\Rightarrow \bar{X} = \frac{Z * S}{\sqrt{n}} + 2$$

$$\Rightarrow \bar{X} = \frac{-1.645 * 0.6}{\sqrt{80}} + 2$$

$$\Rightarrow \bar{X} = 1.89$$

# STATISTICS FOR DATA SCIENCE

## Fixed Level Testing

---



### Solution 2 (a)

- Hence,  $H_0$  will be rejected if  $\bar{X} \leq 1.890$ .
- The rejection region consists of all values of  $\bar{X}$  less than or equal to 1.890.

# STATISTICS FOR DATA SCIENCE

## Fixed Level Testing

---



### Solution Using Rejection Region Approach:

$H_0: \mu \geq 2$  versus  $H_1: \mu < 2$

- Null distribution of  $\bar{X}$ :  
$$\bar{X} \sim N(2, 0.6^2/80)$$
- $\alpha = 0.10$
- $Z = -1.28$  (Critical value)

# STATISTICS FOR DATA SCIENCE

## Fixed Level Testing

---



**Solution Using Rejection Region Approach:**

$$\Rightarrow \bar{X} = \frac{z * s}{\sqrt{n}} + 2$$

$$\Rightarrow \bar{X} = \frac{-1.28 * 0.6}{\sqrt{80}} + 2$$

$$\Rightarrow \bar{X} = 1.9141$$

Since  $1.85 < 1.9141$

$\Rightarrow H_0$  will be rejected at the 10% level

# STATISTICS FOR DATA SCIENCE

## Fixed Level Testing

---



### Solution Using P Value Approach:

$H_0: \mu \geq 2$  versus  $H_1: \mu < 2$

- Null distribution of  $\bar{X}$ :

$$\bar{X} \sim N(2, 0.6^2/80)$$

- $\alpha = 0.10$

$$\bar{X} = 1.85,$$

- Finding z-score for 1.85

- $$Z = \frac{\bar{X} - \mu}{\sigma/\sqrt{n}}$$

# STATISTICS FOR DATA SCIENCE

## Fixed Level Testing

---



**Solution Using P Value Approach:**

$$Z = (1.85 - 2) / (0.6 / \sqrt{80}) = -2.24$$

$$\Rightarrow P = P(Z < -2.24) = 0.0125$$

$\Rightarrow P < \alpha \Rightarrow H_0$  will be rejected at the 10% level



# STATISTICS FOR DATA SCIENCE

## Fixed Level Testing

---



### Example:

#### Small Sample test for population mean:

- Before a substance can be deemed safe for landfilling, its chemical properties must be characterized. The article “Landfilling Ash/Sludge Mixtures” (J. Benoit, T. Eighmy, and B. Crannell, *Journal of Geotechnical and Geoenvironmental Engineering*, 1999: 877–888) reports that in a sample of six replicates of sludge from a New Hampshire wastewater treatment plant, the mean pH was 6.68 with a standard deviation of 0.20.
- Can we conclude that the mean pH is less than 7.0?

# STATISTICS FOR DATA SCIENCE

## Fixed Level Testing

---



**Solution:**

**Small Sample test for population mean:**

$$n = 6$$

$$H_0: \mu \geq 7.0 \text{ versus } H_1: \mu < 7.0$$

Under  $H_0$ , the test statistic

$$t = \frac{\bar{X} - 7.0}{s/\sqrt{n}}$$

has a Student's  $t$  distribution with  $n-1$  degrees of freedom.

# STATISTICS FOR DATA SCIENCE

## Fixed Level Testing

---



### Solution:


#### Small Sample test for population mean:

- Has a Student's  $t$  distribution with five degrees of freedom.  
Substituting  $\bar{X} = 6.68, s = 0.20, \text{ and } n = 6$ , the value of the test statistic is
- $t = \frac{6.68 - 7.0}{0.2/\sqrt{6}} = -3.910$

# STATISTICS FOR DATA SCIENCE

## Fixed Level Testing

TABLE A.3 Upper percentage points for the Student's  $t$  distribution



$\nu$	$\alpha$								
	0.40	0.25	0.10	0.05	0.025	0.01	0.005	0.001	0.0005
1	0.325	1.000	3.078	6.314	12.706	31.821	63.657	318.309	636.619
2	0.289	0.816	1.886	2.920	4.303	6.965	9.925	22.327	31.599
3	0.277	0.765	1.638	2.353	3.182	4.541	5.841	10.215	12.924
4	0.271	0.741	1.533	2.132	2.776	3.747	4.604	7.173	8.610
5	0.267	0.727	1.476	2.015	2.571	3.365	4.032	5.893	6.869
6	0.265	0.718	1.440	1.943	2.447	3.143	3.707	5.208	5.959
7	0.263	0.711	1.415	1.895	2.365	2.998	3.499	4.785	5.408
8	0.262	0.706	1.397	1.860	2.306	2.896	3.355	4.501	5.041
9	0.261	0.703	1.383	1.833	2.262	2.821	3.250	4.297	4.781
10	0.260	0.700	1.372	1.812	2.228	2.764	3.169	4.144	4.587
11	0.260	0.697	1.363	1.796	2.201	2.718	3.106	4.025	4.437
12	0.259	0.695	1.356	1.782	2.179	2.681	3.055	3.930	4.318
13	0.259	0.694	1.350	1.771	2.160	2.650	3.012	3.852	4.221
14	0.258	0.692	1.345	1.761	2.145	2.624	2.977	3.787	4.140
15	0.258	0.691	1.341	1.753	2.131	2.602	2.947	3.733	4.073
16	0.258	0.690	1.337	1.746	2.120	2.583	2.921	3.686	4.015
17	0.257	0.689	1.333	1.740	2.110	2.567	2.898	3.646	3.965
18	0.257	0.688	1.330	1.734	2.101	2.552	2.878	3.610	3.922
19	0.257	0.688	1.328	1.729	2.093	2.539	2.861	3.579	3.883
20	0.257	0.687	1.325	1.725	2.086	2.528	2.845	3.552	3.850
21	0.257	0.686	1.323	1.721	2.080	2.518	2.831	3.527	3.819
22	0.256	0.686	1.321	1.717	2.074	2.508	2.819	3.505	3.792
23	0.256	0.685	1.319	1.714	2.069	2.500	2.807	3.485	3.768
24	0.256	0.685	1.318	1.711	2.064	2.492	2.797	3.467	3.745
25	0.256	0.684	1.316	1.708	2.060	2.485	2.787	3.450	3.725
26	0.256	0.684	1.315	1.706	2.056	2.479	2.779	3.435	3.707
27	0.256	0.684	1.314	1.703	2.052	2.473	2.771	3.421	3.690
28	0.256	0.683	1.313	1.701	2.048	2.467	2.763	3.408	3.674
29	0.256	0.683	1.311	1.699	2.045	2.462	2.756	3.396	3.659
30	0.256	0.683	1.310	1.697	2.042	2.457	2.750	3.385	3.646
35	0.255	0.682	1.306	1.690	2.030	2.438	2.724	3.340	3.591
40	0.255	0.681	1.303	1.684	2.021	2.423	2.704	3.307	3.551
60	0.254	0.679	1.296	1.671	2.000	2.390	2.660	3.232	3.460
120	0.254	0.677	1.289	1.658	1.980	2.358	2.617	3.160	3.373
$\infty$	0.253	0.674	1.282	1.645	1.960	2.326	2.576	3.090	3.291

# STATISTICS FOR DATA SCIENCE

## Fixed Level Testing

**Solution:**

**Small Sample test for population mean:**



- The null distribution is Student's  $t$  with five degrees of freedom. The observed value of  $t$  is  $-3.919$ .
- If  $H_0$  is true, the probability that  $t$  takes on a value as extreme as or more extreme than that observed is between 0.005 and 0.01.

# STATISTICS FOR DATA SCIENCE

## Fixed Level Testing

---



### Solution:

#### Small Sample test for population mean:

- Consulting the  $t$  table, we find that the value  $t = -3.365$  cuts off an area of 0.01 in the left-hand tail, and the value  $t = -4.033$  cuts off an area of 0.005 .
- We conclude that the  $P$ -value is between 0.005 and 0.01. There is strong evidence that the mean pH is less than 7.0.



**Dr. Deepa Nair**

Department of Science and Humanities

---

**[deepanair@pes.edu](mailto:deepanair@pes.edu)**