

MA1140: Lecture 8

Eigenvalues and Eigenvectors

Dipankar Ghosh

Department of Mathematics
Indian Institute of Technology Hyderabad

January 28, 2020

Eigenvalues and eigenvectors (of linear operators)

Let $T : V \rightarrow V$ be a linear map,

Eigenvalues and eigenvectors (of linear operators)

Let $T : V \rightarrow V$ be a linear map, which we call **linear operator** (here domain and codomain are same).

Eigenvalues and eigenvectors (of linear operators)

Let $T : V \rightarrow V$ be a linear map, which we call **linear operator** (here domain and codomain are same).

Definition

- 1 A NON-ZERO vector $v \in V$ is called an **eigenvector** of T if $T(v) = \lambda v$ for SOME scalar λ .

Eigenvalues and eigenvectors (of linear operators)

Let $T : V \rightarrow V$ be a linear map, which we call **linear operator** (here domain and codomain are same).

Definition

- 1 A NON-ZERO vector $v \in V$ is called an **eigenvector** of T if $T(v) = \lambda v$ for SOME scalar λ .
- 2 A scalar λ is called an **eigenvalue** of T if there EXISTS a non-zero vector $v \in V$ such that $T(v) = \lambda v$.

Eigenvalues and eigenvectors (of linear operators)

Let $T : V \rightarrow V$ be a linear map, which we call **linear operator** (here domain and codomain are same).

Definition

- 1 A NON-ZERO vector $v \in V$ is called an **eigenvector** of T if $T(v) = \lambda v$ for SOME scalar λ .
- 2 A scalar λ is called an **eigenvalue** of T if there EXISTS a non-zero vector $v \in V$ such that $T(v) = \lambda v$.
- 3 If $T(v) = \lambda v$, then λ is called an eigenvalue of T corresponding to the eigenvector v .

Eigenvalues and eigenvectors (of linear operators)

Let $T : V \rightarrow V$ be a linear map, which we call **linear operator** (here domain and codomain are same).

Definition

- 1 A NON-ZERO vector $v \in V$ is called an **eigenvector** of T if $T(v) = \lambda v$ for SOME scalar λ .
 - 2 A scalar λ is called an **eigenvalue** of T if there EXISTS a non-zero vector $v \in V$ such that $T(v) = \lambda v$.
 - 3 If $T(v) = \lambda v$, then λ is called an eigenvalue of T corresponding to the eigenvector v .
- **Geometrically**, an eigenvector, corresponding to an eigenvalue,

Eigenvalues and eigenvectors (of linear operators)

Let $T : V \rightarrow V$ be a linear map, which we call **linear operator** (here domain and codomain are same).

Definition

- ❶ A NON-ZERO vector $v \in V$ is called an **eigenvector** of T if $T(v) = \lambda v$ for SOME scalar λ .
 - ❷ A scalar λ is called an **eigenvalue** of T if there EXISTS a non-zero vector $v \in V$ such that $T(v) = \lambda v$.
 - ❸ If $T(v) = \lambda v$, then λ is called an eigenvalue of T corresponding to the eigenvector v .
- **Geometrically**, an eigenvector, corresponding to an eigenvalue, points in a direction that is stretched by the transformation,

Eigenvalues and eigenvectors (of linear operators)

Let $T : V \rightarrow V$ be a linear map, which we call **linear operator** (here domain and codomain are same).

Definition

- 1 A NON-ZERO vector $v \in V$ is called an **eigenvector** of T if $T(v) = \lambda v$ for SOME scalar λ .
 - 2 A scalar λ is called an **eigenvalue** of T if there EXISTS a non-zero vector $v \in V$ such that $T(v) = \lambda v$.
 - 3 If $T(v) = \lambda v$, then λ is called an eigenvalue of T corresponding to the eigenvector v .
- **Geometrically**, an eigenvector, corresponding to an eigenvalue, points in a direction that is stretched by the transformation, and the eigenvalue is the factor by which it is stretched.

Eigenvalues and eigenvectors (of linear operators)

Let $T : V \rightarrow V$ be a linear map, which we call **linear operator** (here domain and codomain are same).

Definition

- 1 A NON-ZERO vector $v \in V$ is called an **eigenvector** of T if $T(v) = \lambda v$ for SOME scalar λ .
 - 2 A scalar λ is called an **eigenvalue** of T if there EXISTS a non-zero vector $v \in V$ such that $T(v) = \lambda v$.
 - 3 If $T(v) = \lambda v$, then λ is called an eigenvalue of T corresponding to the eigenvector v .
-
- **Geometrically**, an eigenvector, corresponding to an eigenvalue, points in a direction that is stretched by the transformation, and the eigenvalue is the factor by which it is stretched. If the eigenvalue is negative, the direction is reversed.

Eigenvalues and eigenvectors (of linear operators)

Let $T : V \rightarrow V$ be a linear map, which we call **linear operator** (here domain and codomain are same).

Definition

- 1 A NON-ZERO vector $v \in V$ is called an **eigenvector** of T if $T(v) = \lambda v$ for SOME scalar λ .
 - 2 A scalar λ is called an **eigenvalue** of T if there EXISTS a non-zero vector $v \in V$ such that $T(v) = \lambda v$.
 - 3 If $T(v) = \lambda v$, then λ is called an eigenvalue of T corresponding to the eigenvector v .
- **Geometrically**, an eigenvector, corresponding to an eigenvalue, points in a direction that is stretched by the transformation, and the eigenvalue is the factor by which it is stretched. If the eigenvalue is negative, the direction is reversed.
 - An eigenvalue can be positive, negative or zero.

Eigenvalues and eigenvectors (of square matrices)

Since there is a one to one correspondence between

Eigenvalues and eigenvectors (of square matrices)

Since there is a one to one correspondence between the set of all linear operators from $V (\cong \mathbb{R}^n)$ to itself

Eigenvalues and eigenvectors (of square matrices)

Since there is a one to one correspondence between the set of all linear operators from $V (\cong \mathbb{R}^n)$ to itself and the collection of all $n \times n$ matrices over \mathbb{R} ,

Eigenvalues and eigenvectors (of square matrices)

Since there is a one to one correspondence between the set of all linear operators from $V (\cong \mathbb{R}^n)$ to itself and the collection of all $n \times n$ matrices over \mathbb{R} , it is equivalent to define eigenvalues and eigenvectors of $n \times n$ matrices.

Eigenvalues and eigenvectors (of square matrices)

Since there is a one to one correspondence between the set of all linear operators from $V (\cong \mathbb{R}^n)$ to itself and the collection of all $n \times n$ matrices over \mathbb{R} , it is equivalent to define eigenvalues and eigenvectors of $n \times n$ matrices.

Definition

Let A be an $n \times n$ matrix over \mathbb{R} .

Eigenvalues and eigenvectors (of square matrices)

Since there is a one to one correspondence between the set of all linear operators from $V (\cong \mathbb{R}^n)$ to itself and the collection of all $n \times n$ matrices over \mathbb{R} , it is equivalent to define eigenvalues and eigenvectors of $n \times n$ matrices.

Definition

Let A be an $n \times n$ matrix over \mathbb{R} .

- 1 A NON-ZERO column vector $v \in \mathbb{R}^n$ is called an **eigenvector** of A

Eigenvalues and eigenvectors (of square matrices)

Since there is a one to one correspondence between the set of all linear operators from $V (\cong \mathbb{R}^n)$ to itself and the collection of all $n \times n$ matrices over \mathbb{R} , it is equivalent to define eigenvalues and eigenvectors of $n \times n$ matrices.

Definition

Let A be an $n \times n$ matrix over \mathbb{R} .

- 1 A NON-ZERO column vector $v \in \mathbb{R}^n$ is called an **eigenvector** of A if $Av = \lambda v$ for **SOME** $\lambda \in \mathbb{R}$.

Eigenvalues and eigenvectors (of square matrices)

Since there is a one to one correspondence between the set of all linear operators from $V (\cong \mathbb{R}^n)$ to itself and the collection of all $n \times n$ matrices over \mathbb{R} , it is equivalent to define eigenvalues and eigenvectors of $n \times n$ matrices.

Definition

Let A be an $n \times n$ matrix over \mathbb{R} .

- 1 A NON-ZERO column vector $v \in \mathbb{R}^n$ is called an **eigenvector** of A if $Av = \lambda v$ for SOME $\lambda \in \mathbb{R}$.
- 2 A scalar λ is called an **eigenvalue** of A

Eigenvalues and eigenvectors (of square matrices)

Since there is a one to one correspondence between the set of all linear operators from $V (\cong \mathbb{R}^n)$ to itself and the collection of all $n \times n$ matrices over \mathbb{R} , it is equivalent to define eigenvalues and eigenvectors of $n \times n$ matrices.

Definition

Let A be an $n \times n$ matrix over \mathbb{R} .

- 1 A NON-ZERO column vector $v \in \mathbb{R}^n$ is called an **eigenvector** of A if $Av = \lambda v$ for SOME $\lambda \in \mathbb{R}$.
- 2 A scalar λ is called an **eigenvalue** of A if there EXISTS a non-zero column vector $v \in \mathbb{R}^n$ such that $Av = \lambda v$.

Eigenvalues and eigenvectors (of square matrices)

Since there is a one to one correspondence between the set of all linear operators from $V (\cong \mathbb{R}^n)$ to itself and the collection of all $n \times n$ matrices over \mathbb{R} , it is equivalent to define eigenvalues and eigenvectors of $n \times n$ matrices.

Definition

Let A be an $n \times n$ matrix over \mathbb{R} .

- 1 A NON-ZERO column vector $v \in \mathbb{R}^n$ is called an **eigenvector** of A if $Av = \lambda v$ for SOME $\lambda \in \mathbb{R}$.
- 2 A scalar λ is called an **eigenvalue** of A if there EXISTS a non-zero column vector $v \in \mathbb{R}^n$ such that $Av = \lambda v$.
- 3 If $Av = \lambda v$, then λ is called an eigenvalue of A corresponding to the eigenvector v .

Example 1: eigenvalues and eigenvectors of stretching

Let $c \in \mathbb{R}$. Define $T : \mathbb{R}^2 \longrightarrow \mathbb{R}^2$ by

$$T \begin{pmatrix} x \\ y \end{pmatrix} = c \begin{pmatrix} x \\ y \end{pmatrix} \text{ for every } \begin{pmatrix} x \\ y \end{pmatrix} \in \mathbb{R}^2.$$

Example 1: eigenvalues and eigenvectors of stretching

Let $c \in \mathbb{R}$. Define $T : \mathbb{R}^2 \longrightarrow \mathbb{R}^2$ by

$$T \begin{pmatrix} x \\ y \end{pmatrix} = c \begin{pmatrix} x \\ y \end{pmatrix} \text{ for every } \begin{pmatrix} x \\ y \end{pmatrix} \in \mathbb{R}^2.$$


The matrix representation: $[T] = \begin{pmatrix} c & 0 \\ 0 & c \end{pmatrix}$

Example 1: eigenvalues and eigenvectors of stretching

Let $c \in \mathbb{R}$. Define $T : \mathbb{R}^2 \longrightarrow \mathbb{R}^2$ by

$$T \begin{pmatrix} x \\ y \end{pmatrix} = c \begin{pmatrix} x \\ y \end{pmatrix} \text{ for every } \begin{pmatrix} x \\ y \end{pmatrix} \in \mathbb{R}^2.$$

The matrix representation: $[T] = \begin{pmatrix} c & 0 \\ 0 & c \end{pmatrix}$




Example 1: eigenvalues and eigenvectors of stretching

Let $c \in \mathbb{R}$. Define $T : \mathbb{R}^2 \rightarrow \mathbb{R}^2$ by

$$T \begin{pmatrix} x \\ y \end{pmatrix} = c \begin{pmatrix} x \\ y \end{pmatrix} \text{ for every } \begin{pmatrix} x \\ y \end{pmatrix} \in \mathbb{R}^2.$$

The matrix representation: $[T] = \begin{pmatrix} c & 0 \\ 0 & c \end{pmatrix}$




Example 1: eigenvalues and eigenvectors of stretching

Let $c \in \mathbb{R}$. Define $T : \mathbb{R}^2 \rightarrow \mathbb{R}^2$ by

$$T \begin{pmatrix} x \\ y \end{pmatrix} = c \begin{pmatrix} x \\ y \end{pmatrix} \text{ for every } \begin{pmatrix} x \\ y \end{pmatrix} \in \mathbb{R}^2.$$

The matrix representation: $[T] = \begin{pmatrix} c & 0 \\ 0 & c \end{pmatrix}$




Every $v (\neq 0) \in \mathbb{R}^2$ is an eigenvector of T with the eigenvalue c .

Example 2: eigenvalues and eigenvectors of reflection


$T : \mathbb{R}^2 \longrightarrow \mathbb{R}^2$ is defined by $T \begin{pmatrix} x \\ y \end{pmatrix} = \begin{pmatrix} x \\ -y \end{pmatrix}$.

Matrix repres. $\begin{pmatrix} 1 & 0 \\ 0 & -1 \end{pmatrix} \begin{pmatrix} x \\ y \end{pmatrix} = \begin{pmatrix} x \\ -y \end{pmatrix}$



Example 2: eigenvalues and eigenvectors of reflection


$T : \mathbb{R}^2 \longrightarrow \mathbb{R}^2$ is defined by $T \begin{pmatrix} x \\ y \end{pmatrix} = \begin{pmatrix} x \\ -y \end{pmatrix}$.



For $x \neq 0$, $\begin{pmatrix} x \\ 0 \end{pmatrix}$ is an eigenvector of T with eigenvalue 1.

Example 2: eigenvalues and eigenvectors of reflection

$T : \mathbb{R}^2 \longrightarrow \mathbb{R}^2$ is defined by $T \begin{pmatrix} x \\ y \end{pmatrix} = \begin{pmatrix} x \\ -y \end{pmatrix}$.




For $x \neq 0$, $\begin{pmatrix} x \\ 0 \end{pmatrix}$ is an eigenvector of T with eigenvalue 1.

For $y \neq 0$, $\begin{pmatrix} 0 \\ y \end{pmatrix}$ is an eigenvector of T with eigenvalue -1 .

Example 2: eigenvalues and eigenvectors of reflection

$T : \mathbb{R}^2 \longrightarrow \mathbb{R}^2$ is defined by $T \begin{pmatrix} x \\ y \end{pmatrix} = \begin{pmatrix} x \\ -y \end{pmatrix}$.



For $x \neq 0$, $\begin{pmatrix} x \\ 0 \end{pmatrix}$ is an eigenvector of T with eigenvalue 1.


For $y \neq 0$, $\begin{pmatrix} 0 \\ y \end{pmatrix}$ is an eigenvector of T with eigenvalue -1 .

These are ALL the eigenvectors of T . (Verify it!)

Example 3: eigenvalues and eigenvectors of projection

Define $T : \mathbb{R}^2 \longrightarrow \mathbb{R}^2$ by $T \begin{pmatrix} x \\ y \end{pmatrix} = \begin{pmatrix} x \\ 0 \end{pmatrix}$


Matrix Repres. $\begin{pmatrix} 1 & 0 \\ 0 & 0 \end{pmatrix} \begin{pmatrix} x \\ y \end{pmatrix} = \begin{pmatrix} x \\ 0 \end{pmatrix}$



Example 3: eigenvalues and eigenvectors of projection

Define $T : \mathbb{R}^2 \longrightarrow \mathbb{R}^2$ by $T \begin{pmatrix} x \\ y \end{pmatrix} = \begin{pmatrix} x \\ 0 \end{pmatrix}$

Matrix Repres. $\begin{pmatrix} 1 & 0 \\ 0 & 0 \end{pmatrix} \begin{pmatrix} x \\ y \end{pmatrix} = \begin{pmatrix} x \\ 0 \end{pmatrix}$




For $x \neq 0$, $\begin{pmatrix} x \\ 0 \end{pmatrix}$ is an eigenvector of T with eigenvalue 1.

Example 3: eigenvalues and eigenvectors of projection

Define $T : \mathbb{R}^2 \rightarrow \mathbb{R}^2$ by $T \begin{pmatrix} x \\ y \end{pmatrix} = \begin{pmatrix} x \\ 0 \end{pmatrix}$

Matrix Repres. $\begin{pmatrix} 1 & 0 \\ 0 & 0 \end{pmatrix} \begin{pmatrix} x \\ y \end{pmatrix} = \begin{pmatrix} x \\ 0 \end{pmatrix}$




For $x \neq 0$, $\begin{pmatrix} x \\ 0 \end{pmatrix}$ is an eigenvector of T with eigenvalue 1.

For $y \neq 0$, $\begin{pmatrix} 0 \\ y \end{pmatrix}$ is an eigenvector of T with eigenvalue 0.

Example 3: eigenvalues and eigenvectors of projection

Define $T : \mathbb{R}^2 \rightarrow \mathbb{R}^2$ by $T \begin{pmatrix} x \\ y \end{pmatrix} = \begin{pmatrix} x \\ 0 \end{pmatrix}$

Matrix Repres. $\begin{pmatrix} 1 & 0 \\ 0 & 0 \end{pmatrix} \begin{pmatrix} x \\ y \end{pmatrix} = \begin{pmatrix} x \\ 0 \end{pmatrix}$



For $x \neq 0$, $\begin{pmatrix} x \\ 0 \end{pmatrix}$ is an eigenvector of T with eigenvalue 1.

For $y \neq 0$, $\begin{pmatrix} 0 \\ y \end{pmatrix}$ is an eigenvector of T with eigenvalue 0.

These are ALL the eigenvectors of T . (Verify it!)

Example 4: A may not have eigenvalues and eigenvectors over a particular field

- Consider the matrix $A = \begin{pmatrix} 0 & 1 \\ -1 & 0 \end{pmatrix}$ over \mathbb{R} .

Example 4: A may not have eigenvalues and eigenvectors over a particular field

- Consider the matrix $A = \begin{pmatrix} 0 & 1 \\ -1 & 0 \end{pmatrix}$ over \mathbb{R} .
- Does A have eigenvalues and eigenvectors over \mathbb{R} ?

Example 4: A may not have eigenvalues and eigenvectors over a particular field

- Consider the matrix $A = \begin{pmatrix} 0 & 1 \\ -1 & 0 \end{pmatrix}$ over \mathbb{R} .
- Does A have eigenvalues and eigenvectors over \mathbb{R} ?
- If yes, then there are $\begin{pmatrix} x \\ y \end{pmatrix} \in \mathbb{R}^2 \setminus \left\{ \begin{pmatrix} 0 \\ 0 \end{pmatrix} \right\}$

Example 4: A may not have eigenvalues and eigenvectors over a particular field

- Consider the matrix $A = \begin{pmatrix} 0 & 1 \\ -1 & 0 \end{pmatrix}$ over \mathbb{R} .
- Does A have eigenvalues and eigenvectors over \mathbb{R} ?
- If yes, then there are $\begin{pmatrix} x \\ y \end{pmatrix} \in \mathbb{R}^2 \setminus \left\{ \begin{pmatrix} 0 \\ 0 \end{pmatrix} \right\}$ and $\lambda \in \mathbb{R}$

Example 4: A may not have eigenvalues and eigenvectors over a particular field

- Consider the matrix $A = \begin{pmatrix} 0 & 1 \\ -1 & 0 \end{pmatrix}$ over \mathbb{R} .
- Does A have eigenvalues and eigenvectors over \mathbb{R} ?
- If yes, then there are $\begin{pmatrix} x \\ y \end{pmatrix} \in \mathbb{R}^2 \setminus \left\{ \begin{pmatrix} 0 \\ 0 \end{pmatrix} \right\}$ and $\lambda \in \mathbb{R}$ such that

$$\begin{pmatrix} 0 & 1 \\ -1 & 0 \end{pmatrix} \begin{pmatrix} x \\ y \end{pmatrix} = \lambda \begin{pmatrix} x \\ y \end{pmatrix}$$

Example 4: A may not have eigenvalues and eigenvectors over a particular field

- Consider the matrix $A = \begin{pmatrix} 0 & 1 \\ -1 & 0 \end{pmatrix}$ over \mathbb{R} .
- Does A have eigenvalues and eigenvectors over \mathbb{R} ?
- If yes, then there are $\begin{pmatrix} x \\ y \end{pmatrix} \in \mathbb{R}^2 \setminus \left\{ \begin{pmatrix} 0 \\ 0 \end{pmatrix} \right\}$ and $\lambda \in \mathbb{R}$ such that

$$\begin{aligned} \begin{pmatrix} 0 & 1 \\ -1 & 0 \end{pmatrix} \begin{pmatrix} x \\ y \end{pmatrix} &= \lambda \begin{pmatrix} x \\ y \end{pmatrix} \\ \Rightarrow \begin{pmatrix} 0 & 1 \\ -1 & 0 \end{pmatrix} \begin{pmatrix} x \\ y \end{pmatrix} &= \begin{pmatrix} \lambda & 0 \\ 0 & \lambda \end{pmatrix} \begin{pmatrix} x \\ y \end{pmatrix} \end{aligned}$$

Example 4: A may not have eigenvalues and eigenvectors over a particular field

- Consider the matrix $A = \begin{pmatrix} 0 & 1 \\ -1 & 0 \end{pmatrix}$ over \mathbb{R} .
- Does A have eigenvalues and eigenvectors over \mathbb{R} ?
- If yes, then there are $\begin{pmatrix} x \\ y \end{pmatrix} \in \mathbb{R}^2 \setminus \left\{ \begin{pmatrix} 0 \\ 0 \end{pmatrix} \right\}$ and $\lambda \in \mathbb{R}$ such that

$$\begin{aligned} & \begin{pmatrix} 0 & 1 \\ -1 & 0 \end{pmatrix} \begin{pmatrix} x \\ y \end{pmatrix} = \lambda \begin{pmatrix} x \\ y \end{pmatrix} \\ \Rightarrow & \begin{pmatrix} 0 & 1 \\ -1 & 0 \end{pmatrix} \begin{pmatrix} x \\ y \end{pmatrix} = \begin{pmatrix} \lambda & 0 \\ 0 & \lambda \end{pmatrix} \begin{pmatrix} x \\ y \end{pmatrix} \\ \Rightarrow & \begin{pmatrix} -\lambda & 1 \\ -1 & -\lambda \end{pmatrix} \begin{pmatrix} x \\ y \end{pmatrix} = \begin{pmatrix} 0 \\ 0 \end{pmatrix} \end{aligned}$$

Example 4: A may not have eigenvalues and eigenvectors over a particular field

- Consider the matrix $A = \begin{pmatrix} 0 & 1 \\ -1 & 0 \end{pmatrix}$ over \mathbb{R} .
- Does A have eigenvalues and eigenvectors over \mathbb{R} ?
- If yes, then there are $\begin{pmatrix} x \\ y \end{pmatrix} \in \mathbb{R}^2 \setminus \left\{ \begin{pmatrix} 0 \\ 0 \end{pmatrix} \right\}$ and $\lambda \in \mathbb{R}$ such that

$$\begin{aligned} \begin{pmatrix} 0 & 1 \\ -1 & 0 \end{pmatrix} \begin{pmatrix} x \\ y \end{pmatrix} &= \lambda \begin{pmatrix} x \\ y \end{pmatrix} \\ \Rightarrow \begin{pmatrix} 0 & 1 \\ -1 & 0 \end{pmatrix} \begin{pmatrix} x \\ y \end{pmatrix} &= \begin{pmatrix} \lambda & 0 \\ 0 & \lambda \end{pmatrix} \begin{pmatrix} x \\ y \end{pmatrix} \\ \Rightarrow \begin{pmatrix} -\lambda & 1 \\ -1 & -\lambda \end{pmatrix} \begin{pmatrix} x \\ y \end{pmatrix} &= \begin{pmatrix} 0 \\ 0 \end{pmatrix} \end{aligned}$$

Since $\begin{pmatrix} x \\ y \end{pmatrix} \neq \begin{pmatrix} 0 \\ 0 \end{pmatrix}$,

Example 4: A may not have eigenvalues and eigenvectors over a particular field

- Consider the matrix $A = \begin{pmatrix} 0 & 1 \\ -1 & 0 \end{pmatrix}$ over \mathbb{R} .
- Does A have eigenvalues and eigenvectors over \mathbb{R} ?
- If yes, then there are $\begin{pmatrix} x \\ y \end{pmatrix} \in \mathbb{R}^2 \setminus \left\{ \begin{pmatrix} 0 \\ 0 \end{pmatrix} \right\}$ and $\lambda \in \mathbb{R}$ such that

$$\begin{aligned} \begin{pmatrix} 0 & 1 \\ -1 & 0 \end{pmatrix} \begin{pmatrix} x \\ y \end{pmatrix} &= \lambda \begin{pmatrix} x \\ y \end{pmatrix} \\ \Rightarrow \begin{pmatrix} 0 & 1 \\ -1 & 0 \end{pmatrix} \begin{pmatrix} x \\ y \end{pmatrix} &= \begin{pmatrix} \lambda & 0 \\ 0 & \lambda \end{pmatrix} \begin{pmatrix} x \\ y \end{pmatrix} \\ \Rightarrow \begin{pmatrix} -\lambda & 1 \\ -1 & -\lambda \end{pmatrix} \begin{pmatrix} x \\ y \end{pmatrix} &= \begin{pmatrix} 0 \\ 0 \end{pmatrix} \end{aligned}$$

Since $\begin{pmatrix} x \\ y \end{pmatrix} \neq \begin{pmatrix} 0 \\ 0 \end{pmatrix}$, $\det \begin{pmatrix} -\lambda & 1 \\ -1 & -\lambda \end{pmatrix} = 0$.

Example 4: A may not have eigenvalues and eigenvectors over a particular field

- Consider the matrix $A = \begin{pmatrix} 0 & 1 \\ -1 & 0 \end{pmatrix}$ over \mathbb{R} .
- Does A have eigenvalues and eigenvectors over \mathbb{R} ?
- If yes, then there are $\begin{pmatrix} x \\ y \end{pmatrix} \in \mathbb{R}^2 \setminus \left\{ \begin{pmatrix} 0 \\ 0 \end{pmatrix} \right\}$ and $\lambda \in \mathbb{R}$ such that

$$\begin{aligned} \begin{pmatrix} 0 & 1 \\ -1 & 0 \end{pmatrix} \begin{pmatrix} x \\ y \end{pmatrix} &= \lambda \begin{pmatrix} x \\ y \end{pmatrix} \\ \Rightarrow \begin{pmatrix} 0 & 1 \\ -1 & 0 \end{pmatrix} \begin{pmatrix} x \\ y \end{pmatrix} &= \begin{pmatrix} \lambda & 0 \\ 0 & \lambda \end{pmatrix} \begin{pmatrix} x \\ y \end{pmatrix} \\ \Rightarrow \begin{pmatrix} -\lambda & 1 \\ -1 & -\lambda \end{pmatrix} \begin{pmatrix} x \\ y \end{pmatrix} &= \begin{pmatrix} 0 \\ 0 \end{pmatrix} \end{aligned}$$

Since $\begin{pmatrix} x \\ y \end{pmatrix} \neq \begin{pmatrix} 0 \\ 0 \end{pmatrix}$, $\det \begin{pmatrix} -\lambda & 1 \\ -1 & -\lambda \end{pmatrix} = 0$. Hence $\lambda^2 + 1 = 0$.

Example 4: A may not have eigenvalues and eigenvectors over a particular field

- Consider the matrix $A = \begin{pmatrix} 0 & 1 \\ -1 & 0 \end{pmatrix}$ over \mathbb{R} .
- Does A have eigenvalues and eigenvectors over \mathbb{R} ?
- If yes, then there are $\begin{pmatrix} x \\ y \end{pmatrix} \in \mathbb{R}^2 \setminus \left\{ \begin{pmatrix} 0 \\ 0 \end{pmatrix} \right\}$ and $\lambda \in \mathbb{R}$ such that

$$\begin{aligned} \begin{pmatrix} 0 & 1 \\ -1 & 0 \end{pmatrix} \begin{pmatrix} x \\ y \end{pmatrix} &= \lambda \begin{pmatrix} x \\ y \end{pmatrix} \\ \Rightarrow \begin{pmatrix} 0 & 1 \\ -1 & 0 \end{pmatrix} \begin{pmatrix} x \\ y \end{pmatrix} &= \begin{pmatrix} \lambda & 0 \\ 0 & \lambda \end{pmatrix} \begin{pmatrix} x \\ y \end{pmatrix} \\ \Rightarrow \begin{pmatrix} -\lambda & 1 \\ -1 & -\lambda \end{pmatrix} \begin{pmatrix} x \\ y \end{pmatrix} &= \begin{pmatrix} 0 \\ 0 \end{pmatrix} \end{aligned}$$

Since $\begin{pmatrix} x \\ y \end{pmatrix} \neq \begin{pmatrix} 0 \\ 0 \end{pmatrix}$, $\det \begin{pmatrix} -\lambda & 1 \\ -1 & -\lambda \end{pmatrix} = 0$. Hence $\lambda^2 + 1 = 0$. But no such λ exists in \mathbb{R} .

Example 4: A may not have eigenvalues and eigenvectors over a particular field

- Consider the matrix $A = \begin{pmatrix} 0 & 1 \\ -1 & 0 \end{pmatrix}$ over \mathbb{R} .
- Does A have eigenvalues and eigenvectors over \mathbb{R} ?
- If yes, then there are $\begin{pmatrix} x \\ y \end{pmatrix} \in \mathbb{R}^2 \setminus \left\{ \begin{pmatrix} 0 \\ 0 \end{pmatrix} \right\}$ and $\lambda \in \mathbb{R}$ such that

$$\begin{aligned} \begin{pmatrix} 0 & 1 \\ -1 & 0 \end{pmatrix} \begin{pmatrix} x \\ y \end{pmatrix} &= \lambda \begin{pmatrix} x \\ y \end{pmatrix} \\ \Rightarrow \begin{pmatrix} 0 & 1 \\ -1 & 0 \end{pmatrix} \begin{pmatrix} x \\ y \end{pmatrix} &= \begin{pmatrix} \lambda & 0 \\ 0 & \lambda \end{pmatrix} \begin{pmatrix} x \\ y \end{pmatrix} \\ \Rightarrow \begin{pmatrix} -\lambda & 1 \\ -1 & -\lambda \end{pmatrix} \begin{pmatrix} x \\ y \end{pmatrix} &= \begin{pmatrix} 0 \\ 0 \end{pmatrix} \end{aligned}$$

Since $\begin{pmatrix} x \\ y \end{pmatrix} \neq \begin{pmatrix} 0 \\ 0 \end{pmatrix}$, $\det \begin{pmatrix} -\lambda & 1 \\ -1 & -\lambda \end{pmatrix} = 0$. Hence $\lambda^2 + 1 = 0$. But no such λ exists in \mathbb{R} .

- So A does not have eigenvalues and eigenvectors over \mathbb{R} .

The existence of eigenvalues and eigenvectors

- Consider the matrix $A = \begin{pmatrix} 0 & 1 \\ -1 & 0 \end{pmatrix}$ over \mathbb{C} ,

The existence of eigenvalues and eigenvectors

- Consider the matrix $A = \begin{pmatrix} 0 & 1 \\ -1 & 0 \end{pmatrix}$ over \mathbb{C} , the set of complex numbers.

The existence of eigenvalues and eigenvectors

- Consider the matrix $A = \begin{pmatrix} 0 & 1 \\ -1 & 0 \end{pmatrix}$ over \mathbb{C} , the set of complex numbers.
- Does A have eigenvalues and eigenvectors over \mathbb{C} ?

The existence of eigenvalues and eigenvectors

- Consider the matrix $A = \begin{pmatrix} 0 & 1 \\ -1 & 0 \end{pmatrix}$ over \mathbb{C} , the set of complex numbers.
- Does A have eigenvalues and eigenvectors over \mathbb{C} ? **Ans. Yes.**

The existence of eigenvalues and eigenvectors

- Consider the matrix $A = \begin{pmatrix} 0 & 1 \\ -1 & 0 \end{pmatrix}$ over \mathbb{C} , the set of complex numbers.
- Does A have eigenvalues and eigenvectors over \mathbb{C} ? **Ans. Yes.**
- Note that $\lambda^2 + 1$ has solutions:

The existence of eigenvalues and eigenvectors

- Consider the matrix $A = \begin{pmatrix} 0 & 1 \\ -1 & 0 \end{pmatrix}$ over \mathbb{C} , the set of complex numbers.
- Does A have eigenvalues and eigenvectors over \mathbb{C} ? **Ans. Yes.**
- Note that $\lambda^2 + 1$ has solutions: $\pm i \in \mathbb{C}$.

The existence of eigenvalues and eigenvectors

- Consider the matrix $A = \begin{pmatrix} 0 & 1 \\ -1 & 0 \end{pmatrix}$ over \mathbb{C} , the set of complex numbers.
- Does A have eigenvalues and eigenvectors over \mathbb{C} ? **Ans. Yes.**
- Note that $\lambda^2 + 1$ has solutions: $\pm i \in \mathbb{C}$.
- Then, for each $\lambda = \pm i$,

The existence of eigenvalues and eigenvectors

- Consider the matrix $A = \begin{pmatrix} 0 & 1 \\ -1 & 0 \end{pmatrix}$ over \mathbb{C} , the set of complex numbers.
- Does A have eigenvalues and eigenvectors over \mathbb{C} ? **Ans. Yes.**
- Note that $\lambda^2 + 1$ has solutions: $\pm i \in \mathbb{C}$.
- Then, for each $\lambda = \pm i$, in view of the previous slide, one should solve the system

$$\begin{pmatrix} -\lambda & 1 \\ -1 & -\lambda \end{pmatrix} \begin{pmatrix} x \\ y \end{pmatrix} = \begin{pmatrix} 0 \\ 0 \end{pmatrix}.$$

The existence of eigenvalues and eigenvectors

- Consider the matrix $A = \begin{pmatrix} 0 & 1 \\ -1 & 0 \end{pmatrix}$ over \mathbb{C} , the set of complex numbers.
- Does A have eigenvalues and eigenvectors over \mathbb{C} ? **Ans. Yes.**
- Note that $\lambda^2 + 1$ has solutions: $\pm i \in \mathbb{C}$.
- Then, for each $\lambda = \pm i$, in view of the previous slide, one should solve the system

$$\begin{pmatrix} -\lambda & 1 \\ -1 & -\lambda \end{pmatrix} \begin{pmatrix} x \\ y \end{pmatrix} = \begin{pmatrix} 0 \\ 0 \end{pmatrix}.$$

to get

$$\begin{pmatrix} 0 & 1 \\ -1 & 0 \end{pmatrix} \begin{pmatrix} i \\ -1 \end{pmatrix} = i \begin{pmatrix} i \\ -1 \end{pmatrix}$$

The existence of eigenvalues and eigenvectors

- Consider the matrix $A = \begin{pmatrix} 0 & 1 \\ -1 & 0 \end{pmatrix}$ over \mathbb{C} , the set of complex numbers.
- Does A have eigenvalues and eigenvectors over \mathbb{C} ? **Ans. Yes.**
- Note that $\lambda^2 + 1$ has solutions: $\pm i \in \mathbb{C}$.
- Then, for each $\lambda = \pm i$, in view of the previous slide, one should solve the system

$$\begin{pmatrix} -\lambda & 1 \\ -1 & -\lambda \end{pmatrix} \begin{pmatrix} x \\ y \end{pmatrix} = \begin{pmatrix} 0 \\ 0 \end{pmatrix}.$$

to get

$$\begin{pmatrix} 0 & 1 \\ -1 & 0 \end{pmatrix} \begin{pmatrix} i \\ -1 \end{pmatrix} = i \begin{pmatrix} i \\ -1 \end{pmatrix} \text{ and } \begin{pmatrix} 0 & 1 \\ -1 & 0 \end{pmatrix} \begin{pmatrix} i \\ 1 \end{pmatrix} = -i \begin{pmatrix} i \\ 1 \end{pmatrix}$$

The existence of eigenvalues and eigenvectors

- Consider the matrix $A = \begin{pmatrix} 0 & 1 \\ -1 & 0 \end{pmatrix}$ over \mathbb{C} , the set of complex numbers.
- Does A have eigenvalues and eigenvectors over \mathbb{C} ? **Ans. Yes.**
- Note that $\lambda^2 + 1$ has solutions: $\pm i \in \mathbb{C}$.
- Then, for each $\lambda = \pm i$, in view of the previous slide, one should solve the system

$$\begin{pmatrix} -\lambda & 1 \\ -1 & -\lambda \end{pmatrix} \begin{pmatrix} x \\ y \end{pmatrix} = \begin{pmatrix} 0 \\ 0 \end{pmatrix}.$$

to get

$$\begin{pmatrix} 0 & 1 \\ -1 & 0 \end{pmatrix} \begin{pmatrix} i \\ -1 \end{pmatrix} = i \begin{pmatrix} i \\ -1 \end{pmatrix} \text{ and } \begin{pmatrix} 0 & 1 \\ -1 & 0 \end{pmatrix} \begin{pmatrix} i \\ 1 \end{pmatrix} = -i \begin{pmatrix} i \\ 1 \end{pmatrix}$$

- **Conclusion:** The matrix A has eigenvalues and eigenvectors over \mathbb{C} , but not over \mathbb{R} .

Characteristic polynomial of a matrix

Let A be an $n \times n$ matrix over \mathbb{C} .

Characteristic polynomial of a matrix

Let A be an $n \times n$ matrix over \mathbb{C} . Denote the identity matrix by I_n .

Characteristic polynomial of a matrix

Let A be an $n \times n$ matrix over \mathbb{C} . Denote the identity matrix by I_n .

Lemma

The following statements are equivalent:

Characteristic polynomial of a matrix

Let A be an $n \times n$ matrix over \mathbb{C} . Denote the identity matrix by I_n .

Lemma

The following statements are equivalent:

- $\lambda \in \mathbb{R}$ is an eigenvalue of A .
- $\det(\lambda I_n - A) = 0$.

Characteristic polynomial of a matrix

Let A be an $n \times n$ matrix over \mathbb{C} . Denote the identity matrix by I_n .

Lemma

The following statements are equivalent:

- $\lambda \in \mathbb{R}$ is an eigenvalue of A .
- $\det(\lambda I_n - A) = 0$.

Proof. Note that λ is an eigenvalue of A

Characteristic polynomial of a matrix

Let A be an $n \times n$ matrix over \mathbb{C} . Denote the identity matrix by I_n .

Lemma

The following statements are equivalent:

- $\lambda \in \mathbb{R}$ is an eigenvalue of A .
- $\det(\lambda I_n - A) = 0$.

Proof. Note that λ is an eigenvalue of $A \Leftrightarrow$ there is $v \neq 0$ in \mathbb{C}^n

Characteristic polynomial of a matrix

Let A be an $n \times n$ matrix over \mathbb{C} . Denote the identity matrix by I_n .

Lemma

The following statements are equivalent:

- $\lambda \in \mathbb{R}$ is an eigenvalue of A .
- $\det(\lambda I_n - A) = 0$.

Proof. Note that λ is an eigenvalue of $A \Leftrightarrow$ there is $v \neq 0$ in \mathbb{C}^n such that $Av = \lambda v$,

Characteristic polynomial of a matrix

Let A be an $n \times n$ matrix over \mathbb{C} . Denote the identity matrix by I_n .

Lemma

The following statements are equivalent:

- $\lambda \in \mathbb{R}$ is an eigenvalue of A .
- $\det(\lambda I_n - A) = 0$.

Proof. Note that λ is an eigenvalue of $A \Leftrightarrow$ there is $v \neq 0$ in \mathbb{C}^n such that $Av = \lambda v$, i.e., $(A - \lambda I_n)v = 0$

Characteristic polynomial of a matrix

Let A be an $n \times n$ matrix over \mathbb{C} . Denote the identity matrix by I_n .

Lemma

The following statements are equivalent:

- $\lambda \in \mathbb{R}$ is an eigenvalue of A .
- $\det(\lambda I_n - A) = 0$.

Proof. Note that λ is an eigenvalue of $A \Leftrightarrow$ there is $v \neq 0$ in \mathbb{C}^n such that $Av = \lambda v$, i.e., $(A - \lambda I_n)v = 0 \Leftrightarrow (A - \lambda I_n)X = 0$ has a non-trivial solution

Characteristic polynomial of a matrix

Let A be an $n \times n$ matrix over \mathbb{C} . Denote the identity matrix by I_n .

Lemma

The following statements are equivalent:

- $\lambda \in \mathbb{R}$ is an eigenvalue of A .
- $\det(\lambda I_n - A) = 0$.

Proof. Note that λ is an eigenvalue of $A \Leftrightarrow$ there is $v \neq 0$ in \mathbb{C}^n such that $Av = \lambda v$, i.e., $(A - \lambda I_n)v = 0 \Leftrightarrow (A - \lambda I_n)X = 0$ has a non-trivial solution $\Leftrightarrow \det(A - \lambda I_n) = 0$

Characteristic polynomial of a matrix

Let A be an $n \times n$ matrix over \mathbb{C} . Denote the identity matrix by I_n .

Lemma

The following statements are equivalent:

- $\lambda \in \mathbb{R}$ is an eigenvalue of A .
- $\det(\lambda I_n - A) = 0$.

Proof. Note that λ is an eigenvalue of $A \Leftrightarrow$ there is $v \neq 0$ in \mathbb{C}^n such that $Av = \lambda v$, i.e., $(A - \lambda I_n)v = 0 \Leftrightarrow (A - \lambda I_n)X = 0$ has a non-trivial solution $\Leftrightarrow \det(A - \lambda I_n) = 0 \Leftrightarrow \det(\lambda I_n - A) = 0$.

Definition

Characteristic polynomial of a matrix

Let A be an $n \times n$ matrix over \mathbb{C} . Denote the identity matrix by I_n .

Lemma

The following statements are equivalent:

- $\lambda \in \mathbb{R}$ is an eigenvalue of A .
- $\det(\lambda I_n - A) = 0$.

Proof. Note that λ is an eigenvalue of $A \Leftrightarrow$ there is $v \neq 0$ in \mathbb{C}^n such that $Av = \lambda v$, i.e., $(A - \lambda I_n)v = 0 \Leftrightarrow (A - \lambda I_n)X = 0$ has a non-trivial solution $\Leftrightarrow \det(A - \lambda I_n) = 0 \Leftrightarrow \det(\lambda I_n - A) = 0$.

Definition

- The **characteristic polynomial** of A , denoted by $p_A(x)$,

Characteristic polynomial of a matrix

Let A be an $n \times n$ matrix over \mathbb{C} . Denote the identity matrix by I_n .

Lemma

The following statements are equivalent:

- $\lambda \in \mathbb{R}$ is an eigenvalue of A .
- $\det(\lambda I_n - A) = 0$.

Proof. Note that λ is an eigenvalue of $A \Leftrightarrow$ there is $v \neq 0$ in \mathbb{C}^n such that $Av = \lambda v$, i.e., $(A - \lambda I_n)v = 0 \Leftrightarrow (A - \lambda I_n)X = 0$ has a non-trivial solution $\Leftrightarrow \det(A - \lambda I_n) = 0 \Leftrightarrow \det(\lambda I_n - A) = 0$.

Definition

- The **characteristic polynomial** of A , denoted by $p_A(x)$, is the polynomial defined by $p_A(x) := \det(xI_n - A)$.

Characteristic polynomial of a matrix

Let A be an $n \times n$ matrix over \mathbb{C} . Denote the identity matrix by I_n .

Lemma

The following statements are equivalent:

- $\lambda \in \mathbb{R}$ is an eigenvalue of A .
- $\det(\lambda I_n - A) = 0$.

Proof. Note that λ is an eigenvalue of $A \Leftrightarrow$ there is $v \neq 0$ in \mathbb{C}^n such that $Av = \lambda v$, i.e., $(A - \lambda I_n)v = 0 \Leftrightarrow (A - \lambda I_n)X = 0$ has a non-trivial solution $\Leftrightarrow \det(A - \lambda I_n) = 0 \Leftrightarrow \det(\lambda I_n - A) = 0$.

Definition

- 1 The **characteristic polynomial** of A , denoted by $p_A(x)$, is the polynomial defined by $p_A(x) := \det(xI_n - A)$.
- 2 Thus the eigenvalues of A are nothing but the roots of $p_A(x)$.

Characteristic polynomial of a matrix

Let A be an $n \times n$ matrix over \mathbb{C} . Denote the identity matrix by I_n .

Lemma

The following statements are equivalent:

- $\lambda \in \mathbb{R}$ is an eigenvalue of A .
- $\det(\lambda I_n - A) = 0$.

Proof. Note that λ is an eigenvalue of $A \Leftrightarrow$ there is $v \neq 0$ in \mathbb{C}^n such that $Av = \lambda v$, i.e., $(A - \lambda I_n)v = 0 \Leftrightarrow (A - \lambda I_n)X = 0$ has a non-trivial solution $\Leftrightarrow \det(A - \lambda I_n) = 0 \Leftrightarrow \det(\lambda I_n - A) = 0$.

Definition

- 1 The **characteristic polynomial** of A , denoted by $p_A(x)$, is the polynomial defined by $p_A(x) := \det(xI_n - A)$.
- 2 Thus the eigenvalues of A are nothing but the roots of $p_A(x)$.
- 3 The **algebraic multiplicity** $\text{AM}_A(\lambda)$ of the eigenvalue λ of A is its multiplicity as a root of the characteristic polynomial $p_A(x)$,

Characteristic polynomial of a matrix

Let A be an $n \times n$ matrix over \mathbb{C} . Denote the identity matrix by I_n .

Lemma

The following statements are equivalent:

- $\lambda \in \mathbb{R}$ is an eigenvalue of A .
- $\det(\lambda I_n - A) = 0$.

Proof. Note that λ is an eigenvalue of $A \Leftrightarrow$ there is $v \neq 0$ in \mathbb{C}^n such that $Av = \lambda v$, i.e., $(A - \lambda I_n)v = 0 \Leftrightarrow (A - \lambda I_n)X = 0$ has a non-trivial solution $\Leftrightarrow \det(A - \lambda I_n) = 0 \Leftrightarrow \det(\lambda I_n - A) = 0$.

Definition

- 1 The **characteristic polynomial** of A , denoted by $p_A(x)$, is the polynomial defined by $p_A(x) := \det(xI_n - A)$.
- 2 Thus the eigenvalues of A are nothing but the roots of $p_A(x)$.
- 3 The **algebraic multiplicity** $\text{AM}_A(\lambda)$ of the eigenvalue λ of A is its multiplicity as a root of the characteristic polynomial $p_A(x)$, that is, the largest integer k such that $(x - \lambda)^k$ is a factor of $p_A(x)$.

Eigenspace associated with an eigenvalue

Let A be an $n \times n$ matrix over \mathbb{C} . Denote the identity matrix by I_n .

Eigenspace associated with an eigenvalue

Let A be an $n \times n$ matrix over \mathbb{C} . Denote the identity matrix by I_n .

Lemma

The following statements are equivalent:

Eigenspace associated with an eigenvalue

Let A be an $n \times n$ matrix over \mathbb{C} . Denote the identity matrix by I_n .

Lemma

The following statements are equivalent:

- 1 $v \in \mathbb{C}^n$ is an eigenvector of A with the corr. eigenvalue λ .

Eigenspace associated with an eigenvalue

Let A be an $n \times n$ matrix over \mathbb{C} . Denote the identity matrix by I_n .

Lemma

The following statements are equivalent:

- 1 $v \in \mathbb{C}^n$ is an eigenvector of A with the corr. eigenvalue λ .
- 2 $v \in \mathbb{C}^n$ is a non-trivial solution of the system $(A - \lambda I_n)X = 0$,

Eigenspace associated with an eigenvalue

Let A be an $n \times n$ matrix over \mathbb{C} . Denote the identity matrix by I_n .

Lemma

The following statements are equivalent:

- 1 $v \in \mathbb{C}^n$ is an eigenvector of A with the corr. eigenvalue λ .
- 2 $v \in \mathbb{C}^n$ is a non-trivial solution of the system $(A - \lambda I_n)X = 0$, i.e., $v \in \mathbb{C}^n \setminus \{0\}$ lies in $\text{Null}(A - \lambda I_n)$.

Eigenspace associated with an eigenvalue

Let A be an $n \times n$ matrix over \mathbb{C} . Denote the identity matrix by I_n .

Lemma

The following statements are equivalent:

- 1 $v \in \mathbb{C}^n$ is an eigenvector of A with the corr. eigenvalue λ .
- 2 $v \in \mathbb{C}^n$ is a non-trivial solution of the system $(A - \lambda I_n)X = 0$, i.e., $v \in \mathbb{C}^n \setminus \{0\}$ lies in $\text{Null}(A - \lambda I_n)$.

Proof. Note that $Av = \lambda v$ if and only if $(A - \lambda I_n)v = 0$.

Eigenspace associated with an eigenvalue

Let A be an $n \times n$ matrix over \mathbb{C} . Denote the identity matrix by I_n .

Lemma

The following statements are equivalent:

- 1 $v \in \mathbb{C}^n$ is an eigenvector of A with the corr. eigenvalue λ .
- 2 $v \in \mathbb{C}^n$ is a non-trivial solution of the system $(A - \lambda I_n)X = 0$, i.e., $v \in \mathbb{C}^n \setminus \{0\}$ lies in $\text{Null}(A - \lambda I_n)$.

Proof. Note that $Av = \lambda v$ if and only if $(A - \lambda I_n)v = 0$.

Definition

- 1 Given a particular eigenvalue λ of A .

Eigenspace associated with an eigenvalue

Let A be an $n \times n$ matrix over \mathbb{C} . Denote the identity matrix by I_n .

Lemma

The following statements are equivalent:

- 1 $v \in \mathbb{C}^n$ is an eigenvector of A with the corr. eigenvalue λ .
- 2 $v \in \mathbb{C}^n$ is a non-trivial solution of the system $(A - \lambda I_n)X = 0$, i.e., $v \in \mathbb{C}^n \setminus \{0\}$ lies in $\text{Null}(A - \lambda I_n)$.

Proof. Note that $Av = \lambda v$ if and only if $(A - \lambda I_n)v = 0$.

Definition

- 1 Given a particular eigenvalue λ of A . The set of all eigenvectors of A corresponding to λ , together with the zero vector,

Eigenspace associated with an eigenvalue

Let A be an $n \times n$ matrix over \mathbb{C} . Denote the identity matrix by I_n .

Lemma

The following statements are equivalent:

- 1 $v \in \mathbb{C}^n$ is an eigenvector of A with the corr. eigenvalue λ .
- 2 $v \in \mathbb{C}^n$ is a non-trivial solution of the system $(A - \lambda I_n)X = 0$, i.e., $v \in \mathbb{C}^n \setminus \{0\}$ lies in $\text{Null}(A - \lambda I_n)$.

Proof. Note that $Av = \lambda v$ if and only if $(A - \lambda I_n)v = 0$.

Definition

- 1 Given a particular eigenvalue λ of A . The set of all eigenvectors of A corresponding to λ , together with the zero vector, is called the **eigenspace** of A associated with λ .

Eigenspace associated with an eigenvalue

Let A be an $n \times n$ matrix over \mathbb{C} . Denote the identity matrix by I_n .

Lemma

The following statements are equivalent:

- 1 $v \in \mathbb{C}^n$ is an eigenvector of A with the corr. eigenvalue λ .
- 2 $v \in \mathbb{C}^n$ is a non-trivial solution of the system $(A - \lambda I_n)X = 0$, i.e., $v \in \mathbb{C}^n \setminus \{0\}$ lies in $\text{Null}(A - \lambda I_n)$.

Proof. Note that $Av = \lambda v$ if and only if $(A - \lambda I_n)v = 0$.

Definition

- 1 Given a particular eigenvalue λ of A . The set of all eigenvectors of A corresponding to λ , together with the zero vector, is called the **eigenspace** of A associated with λ . It is denoted by E_λ .

Eigenspace associated with an eigenvalue

Let A be an $n \times n$ matrix over \mathbb{C} . Denote the identity matrix by I_n .

Lemma

The following statements are equivalent:

- 1 $v \in \mathbb{C}^n$ is an eigenvector of A with the corr. eigenvalue λ .
- 2 $v \in \mathbb{C}^n$ is a non-trivial solution of the system $(A - \lambda I_n)X = 0$, i.e., $v \in \mathbb{C}^n \setminus \{0\}$ lies in $\text{Null}(A - \lambda I_n)$.

Proof. Note that $Av = \lambda v$ if and only if $(A - \lambda I_n)v = 0$.

Definition

- 1 Given a particular eigenvalue λ of A . The set of all eigenvectors of A corresponding to λ , together with the zero vector, is called the **eigenspace** of A associated with λ . It is denoted by E_λ . Note that $E_\lambda = \text{Null}(A - \lambda I_n)$.

Eigenspace associated with an eigenvalue

Let A be an $n \times n$ matrix over \mathbb{C} . Denote the identity matrix by I_n .

Lemma

The following statements are equivalent:

- 1 $v \in \mathbb{C}^n$ is an eigenvector of A with the corr. eigenvalue λ .
- 2 $v \in \mathbb{C}^n$ is a non-trivial solution of the system $(A - \lambda I_n)X = 0$, i.e., $v \in \mathbb{C}^n \setminus \{0\}$ lies in $\text{Null}(A - \lambda I_n)$.

Proof. Note that $Av = \lambda v$ if and only if $(A - \lambda I_n)v = 0$.

Definition

- 1 Given a particular eigenvalue λ of A . The set of all eigenvectors of A corresponding to λ , together with the zero vector, is called the **eigenspace** of A associated with λ . It is denoted by E_λ . Note that $E_\lambda = \text{Null}(A - \lambda I_n)$.
- 2 The dimension of $E_\lambda = \text{Null}(A - \lambda I_n)$ is referred to as the **geometric multiplicity** of λ ,

Eigenspace associated with an eigenvalue

Let A be an $n \times n$ matrix over \mathbb{C} . Denote the identity matrix by I_n .

Lemma

The following statements are equivalent:

- 1 $v \in \mathbb{C}^n$ is an eigenvector of A with the corr. eigenvalue λ .
- 2 $v \in \mathbb{C}^n$ is a non-trivial solution of the system $(A - \lambda I_n)X = 0$, i.e., $v \in \mathbb{C}^n \setminus \{0\}$ lies in $\text{Null}(A - \lambda I_n)$.

Proof. Note that $Av = \lambda v$ if and only if $(A - \lambda I_n)v = 0$.

Definition

- 1 Given a particular eigenvalue λ of A . The set of all eigenvectors of A corresponding to λ , together with the zero vector, is called the **eigenspace** of A associated with λ . It is denoted by E_λ . Note that $E_\lambda = \text{Null}(A - \lambda I_n)$.
- 2 The dimension of $E_\lambda = \text{Null}(A - \lambda I_n)$ is referred to as the **geometric multiplicity of λ** , denoted by $\text{GM}_A(\lambda)$.

Some inequalities on algebraic/geometric multiplicities

Theorem

Let A be an $n \times n$ matrix over \mathbb{C} .

Some inequalities on algebraic/geometric multiplicities

Theorem

Let A be an $n \times n$ matrix over \mathbb{C} . For every eigenvalue λ of A , we have

$$\textcircled{1} \quad 1 \leq \text{AM}_A(\lambda) \leq n$$

Some inequalities on algebraic/geometric multiplicities

Theorem

Let A be an $n \times n$ matrix over \mathbb{C} . For every eigenvalue λ of A , we have

$$\textcircled{1} \quad 1 \leq \text{AM}_A(\lambda) \leq n \quad \text{and} \quad 1 \leq \text{GM}_A(\lambda) \leq n.$$

Some inequalities on algebraic/geometric multiplicities

Theorem

Let A be an $n \times n$ matrix over \mathbb{C} . For every eigenvalue λ of A , we have

① $1 \leq \text{AM}_A(\lambda) \leq n$ and $1 \leq \text{GM}_A(\lambda) \leq n$.

② $\sum_{i=1}^r \text{AM}_A(\lambda_i) = n,$

Some inequalities on algebraic/geometric multiplicities

Theorem

Let A be an $n \times n$ matrix over \mathbb{C} . For every eigenvalue λ of A , we have

- ① $1 \leq \text{AM}_A(\lambda) \leq n$ and $1 \leq \text{GM}_A(\lambda) \leq n$.
- ② $\sum_{i=1}^r \text{AM}_A(\lambda_i) = n$, the sum varies over all the eigenvalues of A .

Some inequalities on algebraic/geometric multiplicities

Theorem

Let A be an $n \times n$ matrix over \mathbb{C} . For every eigenvalue λ of A , we have

- ① $1 \leq \text{AM}_A(\lambda) \leq n$ and $1 \leq \text{GM}_A(\lambda) \leq n$.
- ② $\sum_{i=1}^r \text{AM}_A(\lambda_i) = n$, the sum varies over all the eigenvalues of A .
- ③ $1 \leq \text{GM}_A(\lambda) \leq \text{AM}_A(\lambda) \leq n$.

Some inequalities on algebraic/geometric multiplicities

Theorem

Let A be an $n \times n$ matrix over \mathbb{C} . For every eigenvalue λ of A , we have

- ① $1 \leq \text{AM}_A(\lambda) \leq n$ and $1 \leq \text{GM}_A(\lambda) \leq n$.
- ② $\sum_{i=1}^r \text{AM}_A(\lambda_i) = n$, the sum varies over all the eigenvalues of A .
- ③ $1 \leq \text{GM}_A(\lambda) \leq \text{AM}_A(\lambda) \leq n$.

Proof.

- ① Note that $\deg(p_A(x)) = n$

Some inequalities on algebraic/geometric multiplicities

Theorem

Let A be an $n \times n$ matrix over \mathbb{C} . For every eigenvalue λ of A , we have

- ① $1 \leq \text{AM}_A(\lambda) \leq n$ and $1 \leq \text{GM}_A(\lambda) \leq n$.
- ② $\sum_{i=1}^r \text{AM}_A(\lambda_i) = n$, the sum varies over all the eigenvalues of A .
- ③ $1 \leq \text{GM}_A(\lambda) \leq \text{AM}_A(\lambda) \leq n$.

Proof.

- ① Note that $\deg(p_A(x)) = n$ and $p_A(x) = (x - \lambda)^{\text{AM}_A(\lambda)} f(x)$ for some f .

Some inequalities on algebraic/geometric multiplicities

Theorem

Let A be an $n \times n$ matrix over \mathbb{C} . For every eigenvalue λ of A , we have

- ① $1 \leq \text{AM}_A(\lambda) \leq n$ and $1 \leq \text{GM}_A(\lambda) \leq n$.
- ② $\sum_{i=1}^r \text{AM}_A(\lambda_i) = n$, the sum varies over all the eigenvalues of A .
- ③ $1 \leq \text{GM}_A(\lambda) \leq \text{AM}_A(\lambda) \leq n$.

Proof.

- ① Note that $\deg(p_A(x)) = n$ and $p_A(x) = (x - \lambda)^{\text{AM}_A(\lambda)} f(x)$ for some f .
Since $\text{GM}_A(\lambda) = \dim(\text{Null}(A - \lambda I_n))$,

Some inequalities on algebraic/geometric multiplicities

Theorem

Let A be an $n \times n$ matrix over \mathbb{C} . For every eigenvalue λ of A , we have

- ① $1 \leq \text{AM}_A(\lambda) \leq n$ and $1 \leq \text{GM}_A(\lambda) \leq n$.
- ② $\sum_{i=1}^r \text{AM}_A(\lambda_i) = n$, the sum varies over all the eigenvalues of A .
- ③ $1 \leq \text{GM}_A(\lambda) \leq \text{AM}_A(\lambda) \leq n$.

Proof.

- ① Note that $\deg(p_A(x)) = n$ and $p_A(x) = (x - \lambda)^{\text{AM}_A(\lambda)} f(x)$ for some f . Since $\text{GM}_A(\lambda) = \dim(\text{Null}(A - \lambda I_n))$, we have $1 \leq \text{GM}_A(\lambda) \leq n$.

Some inequalities on algebraic/geometric multiplicities

Theorem

Let A be an $n \times n$ matrix over \mathbb{C} . For every eigenvalue λ of A , we have

- ① $1 \leq \text{AM}_A(\lambda) \leq n$ and $1 \leq \text{GM}_A(\lambda) \leq n$.
- ② $\sum_{i=1}^r \text{AM}_A(\lambda_i) = n$, the sum varies over all the eigenvalues of A .
- ③ $1 \leq \text{GM}_A(\lambda) \leq \text{AM}_A(\lambda) \leq n$.

Proof.

- ① Note that $\deg(p_A(x)) = n$ and $p_A(x) = (x - \lambda)^{\text{AM}_A(\lambda)} f(x)$ for some f . Since $\text{GM}_A(\lambda) = \dim(\text{Null}(A - \lambda I_n))$, we have $1 \leq \text{GM}_A(\lambda) \leq n$.
- ② It follows from $p_A(x) = \prod_{i=1}^r (x - \lambda_i)^{\text{AM}_A(\lambda_i)}$

Some inequalities on algebraic/geometric multiplicities

Theorem

Let A be an $n \times n$ matrix over \mathbb{C} . For every eigenvalue λ of A , we have

- ① $1 \leq \text{AM}_A(\lambda) \leq n$ and $1 \leq \text{GM}_A(\lambda) \leq n$.
- ② $\sum_{i=1}^r \text{AM}_A(\lambda_i) = n$, the sum varies over all the eigenvalues of A .
- ③ $1 \leq \text{GM}_A(\lambda) \leq \text{AM}_A(\lambda) \leq n$.

Proof.

- ① Note that $\deg(p_A(x)) = n$ and $p_A(x) = (x - \lambda)^{\text{AM}_A(\lambda)} f(x)$ for some f . Since $\text{GM}_A(\lambda) = \dim(\text{Null}(A - \lambda I_n))$, we have $1 \leq \text{GM}_A(\lambda) \leq n$.
- ② It follows from $p_A(x) = \prod_{i=1}^r (x - \lambda_i)^{\text{AM}_A(\lambda_i)}$ and $\deg(p_A(x)) = n$.

Some inequalities on algebraic/geometric multiplicities

Theorem

Let A be an $n \times n$ matrix over \mathbb{C} . For every eigenvalue λ of A , we have

- 1 $1 \leq \text{AM}_A(\lambda) \leq n$ and $1 \leq \text{GM}_A(\lambda) \leq n$.
- 2 $\sum_{i=1}^r \text{AM}_A(\lambda_i) = n$, the sum varies over all the eigenvalues of A .
- 3 $1 \leq \text{GM}_A(\lambda) \leq \text{AM}_A(\lambda) \leq n$.

Proof.

- 1 Note that $\deg(p_A(x)) = n$ and $p_A(x) = (x - \lambda)^{\text{AM}_A(\lambda)} f(x)$ for some f . Since $\text{GM}_A(\lambda) = \dim(\text{Null}(A - \lambda I_n))$, we have $1 \leq \text{GM}_A(\lambda) \leq n$.
- 2 It follows from $p_A(x) = \prod_{i=1}^r (x - \lambda_i)^{\text{AM}_A(\lambda_i)}$ and $\deg(p_A(x)) = n$.
- 3 We will skip it.



Example: Characteristic polynomial and eigenvalues

- Consider $A = \begin{pmatrix} 1 & 2 \\ 3 & 4 \end{pmatrix}$.

Example: Characteristic polynomial and eigenvalues

- Consider $A = \begin{pmatrix} 1 & 2 \\ 3 & 4 \end{pmatrix}$.
- The characteristic polynomial of A is given by

Example: Characteristic polynomial and eigenvalues

- Consider $A = \begin{pmatrix} 1 & 2 \\ 3 & 4 \end{pmatrix}$.
- The characteristic polynomial of A is given by

$$p_A(x) = \det(xI_2 - A)$$

Example: Characteristic polynomial and eigenvalues

- Consider $A = \begin{pmatrix} 1 & 2 \\ 3 & 4 \end{pmatrix}$.
- The characteristic polynomial of A is given by

$$p_A(x) = \det(xI_2 - A) = \det \begin{pmatrix} x-1 & -2 \\ -3 & x-4 \end{pmatrix}$$

Example: Characteristic polynomial and eigenvalues

- Consider $A = \begin{pmatrix} 1 & 2 \\ 3 & 4 \end{pmatrix}$.
- The characteristic polynomial of A is given by

$$\begin{aligned} p_A(x) &= \det(xI_2 - A) = \det \begin{pmatrix} x-1 & -2 \\ -3 & x-4 \end{pmatrix} \\ &= (x-1)(x-4) - 6 = x^2 - 5x - 2. \end{aligned}$$

Example: Characteristic polynomial and eigenvalues

- Consider $A = \begin{pmatrix} 1 & 2 \\ 3 & 4 \end{pmatrix}$.
- The characteristic polynomial of A is given by

$$\begin{aligned} p_A(x) &= \det(xI_2 - A) = \det \begin{pmatrix} x-1 & -2 \\ -3 & x-4 \end{pmatrix} \\ &= (x-1)(x-4) - 6 = x^2 - 5x - 2. \end{aligned}$$

- The roots of $p_A(x)$ are $\frac{5 \pm \sqrt{33}}{2}$.

Example: Characteristic polynomial and eigenvalues

- Consider $A = \begin{pmatrix} 1 & 2 \\ 3 & 4 \end{pmatrix}$.

- The characteristic polynomial of A is given by

$$\begin{aligned} p_A(x) &= \det(xI_2 - A) = \det \begin{pmatrix} x-1 & -2 \\ -3 & x-4 \end{pmatrix} \\ &= (x-1)(x-4) - 6 = x^2 - 5x - 2. \end{aligned}$$

- The roots of $p_A(x)$ are $\frac{5 \pm \sqrt{33}}{2}$.
- The eigenvalues of A are $\lambda_1 = \frac{5 - \sqrt{33}}{2}$ and $\lambda_2 = \frac{5 + \sqrt{33}}{2}$.

Example: Characteristic polynomial and eigenvalues

- Consider $A = \begin{pmatrix} 1 & 2 \\ 3 & 4 \end{pmatrix}$.
- The characteristic polynomial of A is given by

$$\begin{aligned} p_A(x) &= \det(xI_2 - A) = \det \begin{pmatrix} x-1 & -2 \\ -3 & x-4 \end{pmatrix} \\ &= (x-1)(x-4) - 6 = x^2 - 5x - 2. \end{aligned}$$

- The roots of $p_A(x)$ are $\frac{5 \pm \sqrt{33}}{2}$.
- The eigenvalues of A are $\lambda_1 = \frac{5 - \sqrt{33}}{2}$ and $\lambda_2 = \frac{5 + \sqrt{33}}{2}$.
- The algebraic multiplicities of both λ_1 and λ_2 are 1.

How to compute eigenvalues and eigenvectors

- First compute the characteristic polynomial $p_A(x) = \det(xI_n - A)$ of A .

How to compute eigenvalues and eigenvectors

- First compute the characteristic polynomial $p_A(x) = \det(xI_n - A)$ of A .
- Next compute the roots of $p_A(x)$ by factorizing it into linear factors.

How to compute eigenvalues and eigenvectors

- First compute the characteristic polynomial $p_A(x) = \det(xI_n - A)$ of A .
- Next compute the roots of $p_A(x)$ by factorizing it into linear factors. Which gives the eigenvalues.

How to compute eigenvalues and eigenvectors

- First compute the characteristic polynomial $p_A(x) = \det(xI_n - A)$ of A .
- Next compute the roots of $p_A(x)$ by factorizing it into linear factors. Which gives the eigenvalues.
- Then, for each eigenvalue λ ,

How to compute eigenvalues and eigenvectors

- First compute the characteristic polynomial $p_A(x) = \det(xI_n - A)$ of A .
- Next compute the roots of $p_A(x)$ by factorizing it into linear factors. Which gives the eigenvalues.
- Then, for each eigenvalue λ , solve the homogeneous system:

$$(A - \lambda I_n)X = 0$$

How to compute eigenvalues and eigenvectors

- First compute the characteristic polynomial $p_A(x) = \det(xI_n - A)$ of A .
- Next compute the roots of $p_A(x)$ by factorizing it into linear factors. Which gives the eigenvalues.
- Then, for each eigenvalue λ , solve the homogeneous system:

$$(A - \lambda I_n)X = 0$$

to get eigenspace of A associated with λ .

How to compute eigenvalues and eigenvectors

- First compute the characteristic polynomial $p_A(x) = \det(xI_n - A)$ of A .
- Next compute the roots of $p_A(x)$ by factorizing it into linear factors. Which gives the eigenvalues.
- Then, for each eigenvalue λ , solve the homogeneous system:

$$(A - \lambda I_n)X = 0$$

to get eigenspace of A associated with λ .

- Recall that in order to solve a linear system, you may apply elementary row operations to make it into a system with row reduced echelon coefficient matrix.

Similarity of matrices

Definition

Two $n \times n$ matrices A and B are called **similar**

Similarity of matrices

Definition

Two $n \times n$ matrices A and B are called **similar** if there exists an invertible $n \times n$ matrix P such that $B = P^{-1}AP$.

Similarity of matrices

Definition

Two $n \times n$ matrices A and B are called **similar** if there exists an invertible $n \times n$ matrix P such that $B = P^{-1}AP$.

Some statements (without proof) about importance of similarity of matrices:

Similarity of matrices

Definition

Two $n \times n$ matrices A and B are called **similar** if there exists an invertible $n \times n$ matrix P such that $B = P^{-1}AP$.

Some statements (without proof) about importance of similarity of matrices:

- Two matrices are similar if and only if they represent the same linear operator with respect to (possibly) different bases. (???)

Similarity of matrices

Definition

Two $n \times n$ matrices A and B are called **similar** if there exists an invertible $n \times n$ matrix P such that $B = P^{-1}AP$.

Some statements (without proof) about importance of similarity of matrices:

- Two matrices are similar if and only if they represent the same linear operator with respect to (possibly) different bases. (???)
- Two similar matrices A and B share many properties:

Similarity of matrices

Definition

Two $n \times n$ matrices A and B are called **similar** if there exists an invertible $n \times n$ matrix P such that $B = P^{-1}AP$.

Some statements (without proof) about importance of similarity of matrices:

- Two matrices are similar if and only if they represent the same linear operator with respect to (possibly) different bases. (???)
- Two similar matrices A and B share many properties:
 - $\text{rank}(A) = \text{rank}(B)$ (as operators from \mathbb{R}^n to itself).

Similarity of matrices

Definition

Two $n \times n$ matrices A and B are called **similar** if there exists an invertible $n \times n$ matrix P such that $B = P^{-1}AP$.

Some statements (without proof) about importance of similarity of matrices:

- Two matrices are similar if and only if they represent the same linear operator with respect to (possibly) different bases. (???)
- Two similar matrices A and B share many properties:
 - $\text{rank}(A) = \text{rank}(B)$ (as operators from \mathbb{R}^n to itself).
 - $\det(A) = \det(B)$;

Similarity of matrices

Definition

Two $n \times n$ matrices A and B are called **similar** if there exists an invertible $n \times n$ matrix P such that $B = P^{-1}AP$.

Some statements (without proof) about importance of similarity of matrices:

- Two matrices are similar if and only if they represent the same linear operator with respect to (possibly) different bases. (???)
- Two similar matrices A and B share many properties:
 - $\text{rank}(A) = \text{rank}(B)$ (as operators from \mathbb{R}^n to itself).
 - $\det(A) = \det(B)$; $\text{tr}(A) = \text{tr}(B)$ (sum of all diagonal entries).

Similarity of matrices

Definition

Two $n \times n$ matrices A and B are called **similar** if there exists an invertible $n \times n$ matrix P such that $B = P^{-1}AP$.

Some statements (without proof) about importance of similarity of matrices:

- Two matrices are similar if and only if they represent the same linear operator with respect to (possibly) different bases. (???)
- Two similar matrices A and B share many properties:
 - $\text{rank}(A) = \text{rank}(B)$ (as operators from \mathbb{R}^n to itself).
 - $\det(A) = \det(B)$; $\text{tr}(A) = \text{tr}(B)$ (sum of all diagonal entries).
 - A and B have same characteristic polynomial, $\det(xI_n - A)$.

Similarity of matrices

Definition

Two $n \times n$ matrices A and B are called **similar** if there exists an invertible $n \times n$ matrix P such that $B = P^{-1}AP$.

Some statements (without proof) about importance of similarity of matrices:

- Two matrices are similar if and only if they represent the same linear operator with respect to (possibly) different bases. (???)
- Two similar matrices A and B share many properties:
 - $\text{rank}(A) = \text{rank}(B)$ (as operators from \mathbb{R}^n to itself).
 - $\det(A) = \det(B)$; $\text{tr}(A) = \text{tr}(B)$ (sum of all diagonal entries).
 - A and B have same characteristic polynomial, $\det(xI_n - A)$.
 - Minimal polynomials of A and B are same.

Similarity of matrices

Definition

Two $n \times n$ matrices A and B are called **similar** if there exists an invertible $n \times n$ matrix P such that $B = P^{-1}AP$.

Some statements (without proof) about importance of similarity of matrices:

- Two matrices are similar if and only if they represent the same linear operator with respect to (possibly) different bases. (???)
- Two similar matrices A and B share many properties:
 - $\text{rank}(A) = \text{rank}(B)$ (as operators from \mathbb{R}^n to itself).
 - $\det(A) = \det(B)$; $\text{tr}(A) = \text{tr}(B)$ (sum of all diagonal entries).
 - A and B have same characteristic polynomial, $\det(xI_n - A)$.
 - Minimal polynomials of A and B are same. A monic polynomial $p(X) \in \mathbb{R}[X]$ is said to be a minimal polynomial of A if $p(A) = 0$ (zero matrix)

Similarity of matrices

Definition

Two $n \times n$ matrices A and B are called **similar** if there exists an invertible $n \times n$ matrix P such that $B = P^{-1}AP$.

Some statements (without proof) about importance of similarity of matrices:

- Two matrices are similar if and only if they represent the same linear operator with respect to (possibly) different bases. (???)
- Two similar matrices A and B share many properties:
 - $\text{rank}(A) = \text{rank}(B)$ (as operators from \mathbb{R}^n to itself).
 - $\det(A) = \det(B)$; $\text{tr}(A) = \text{tr}(B)$ (sum of all diagonal entries).
 - A and B have same characteristic polynomial, $\det(xI_n - A)$.
 - Minimal polynomials of A and B are same. A monic polynomial $p(X) \in \mathbb{R}[X]$ is said to be a minimal polynomial of A if $p(A) = 0$ (zero matrix) and p has minimal possible degree.

Similarity of matrices

Definition

Two $n \times n$ matrices A and B are called **similar** if there exists an invertible $n \times n$ matrix P such that $B = P^{-1}AP$.

Some statements (without proof) about importance of similarity of matrices:

- Two matrices are similar if and only if they represent the same linear operator with respect to (possibly) different bases. (???)
- Two similar matrices A and B share many properties:
 - $\text{rank}(A) = \text{rank}(B)$ (as operators from \mathbb{R}^n to itself).
 - $\det(A) = \det(B)$; $\text{tr}(A) = \text{tr}(B)$ (sum of all diagonal entries).
 - A and B have same characteristic polynomial, $\det(xI_n - A)$.
 - Minimal polynomials of A and B are same. A monic polynomial $p(X) \in \mathbb{R}[X]$ is said to be a minimal polynomial of A if $p(A) = 0$ (zero matrix) and p has minimal possible degree.
 - Jordan canonical forms of A and B are same. (???)

Diagonalizable matrices

Motivation: For a matrix, eigenvalues and eigenvectors can be used to decompose the matrix, for example by diagonalizing it.

Diagonalizable matrices

Motivation: For a matrix, eigenvalues and eigenvectors can be used to decompose the matrix, for example by diagonalizing it.

Definition

A matrix A is said to be **diagonalizable**

Diagonalizable matrices

Motivation: For a matrix, eigenvalues and eigenvectors can be used to decompose the matrix, for example by diagonalizing it.

Definition

A matrix A is said to be **diagonalizable** if A is similar to a diagonal matrix D ,

Diagonalizable matrices

Motivation: For a matrix, eigenvalues and eigenvectors can be used to decompose the matrix, for example by diagonalizing it.

Definition

A matrix A is said to be **diagonalizable** if A is similar to a diagonal matrix D , i.e., if there is an invertible matrix P such that

Diagonalizable matrices

Motivation: For a matrix, eigenvalues and eigenvectors can be used to decompose the matrix, for example by diagonalizing it.

Definition

A matrix A is said to be **diagonalizable** if A is similar to a diagonal matrix D , i.e., if there is an invertible matrix P such that

$$P^{-1}AP = \begin{pmatrix} \lambda_1 & 0 & \cdots & 0 \\ 0 & \lambda_2 & \cdots & 0 \\ \vdots & \vdots & \ddots & \vdots \\ 0 & 0 & \cdots & \lambda_n \end{pmatrix} \quad (\text{a diagonal matrix}).$$

Diagonalizable matrices

Motivation: For a matrix, eigenvalues and eigenvectors can be used to decompose the matrix, for example by diagonalizing it.

Definition

A matrix A is said to be **diagonalizable** if A is similar to a diagonal matrix D , i.e., if there is an invertible matrix P such that

$$P^{-1}AP = \begin{pmatrix} \lambda_1 & 0 & \cdots & 0 \\ 0 & \lambda_2 & \cdots & 0 \\ \vdots & \vdots & \ddots & \vdots \\ 0 & 0 & \cdots & \lambda_n \end{pmatrix} \quad (\text{a diagonal matrix}).$$

The set of eigenvectors helps us to test whether a matrix is diagonalizable or not.

The use of eigenvalues and eigenvectors on diagonalization

Theorem

Let A be an $n \times n$ matrix (over \mathbb{C}).

The use of eigenvalues and eigenvectors on diagonalization

Theorem

Let A be an $n \times n$ matrix (over \mathbb{C}). The following are equivalent:

- 1 *A is diagonalizable.*

The use of eigenvalues and eigenvectors on diagonalization

Theorem

Let A be an $n \times n$ matrix (over \mathbb{C}). The following are equivalent:

- 1 *A is diagonalizable.*
- 2 *The eigenvectors of A form a basis of \mathbb{R}^n ,*

The use of eigenvalues and eigenvectors on diagonalization

Theorem

Let A be an $n \times n$ matrix (over \mathbb{C}). The following are equivalent:

- 1 *A is diagonalizable.*
- 2 *The eigenvectors of A form a basis of \mathbb{R}^n , equivalently, A has n linearly independent eigenvectors v_1, \dots, v_n*

The use of eigenvalues and eigenvectors on diagonalization

Theorem

Let A be an $n \times n$ matrix (over \mathbb{C}). The following are equivalent:

- 1 *A is diagonalizable.*
- 2 *The eigenvectors of A form a basis of \mathbb{R}^n , equivalently, A has n linearly independent eigenvectors v_1, \dots, v_n with associated eigenvalues $\lambda_1, \dots, \lambda_n$*

The use of eigenvalues and eigenvectors on diagonalization

Theorem

Let A be an $n \times n$ matrix (over \mathbb{C}). The following are equivalent:

- 1 *A is diagonalizable.*
- 2 *The eigenvectors of A form a basis of \mathbb{R}^n , equivalently, A has n linearly independent eigenvectors v_1, \dots, v_n with associated eigenvalues $\lambda_1, \dots, \lambda_n$ (which need not be distinct).*
- 3 *$\text{GM}_A(\lambda) = \text{AG}_A(\lambda)$ for every eigenvalue λ of A .*

The use of eigenvalues and eigenvectors on diagonalization

Theorem

Let A be an $n \times n$ matrix (over \mathbb{C}). The following are equivalent:

- ❶ *A is diagonalizable.*
- ❷ *The eigenvectors of A form a basis of \mathbb{R}^n , equivalently, A has n linearly independent eigenvectors v_1, \dots, v_n with associated eigenvalues $\lambda_1, \dots, \lambda_n$ (which need not be distinct).*
- ❸ *$\text{GM}_A(\lambda) = \text{AG}_A(\lambda)$ for every eigenvalue λ of A .*
- ❹ *The minimal polynomial of A has distinct roots*

The use of eigenvalues and eigenvectors on diagonalization

Theorem

Let A be an $n \times n$ matrix (over \mathbb{C}). The following are equivalent:

- ❶ *A is diagonalizable.*
- ❷ *The eigenvectors of A form a basis of \mathbb{R}^n , equivalently, A has n linearly independent eigenvectors v_1, \dots, v_n with associated eigenvalues $\lambda_1, \dots, \lambda_n$ (which need not be distinct).*
- ❸ *$\text{GM}_A(\lambda) = \text{AG}_A(\lambda)$ for every eigenvalue λ of A .*
- ❹ *The minimal polynomial of A has distinct roots (equivalently, A satisfies a polynomial $p(x) \in \mathbb{C}[x]$ having distinct roots).*

The use of eigenvalues and eigenvectors on diagonalization

Theorem

Let A be an $n \times n$ matrix (over \mathbb{C}). The following are equivalent:

- ❶ *A is diagonalizable.*
- ❷ *The eigenvectors of A form a basis of \mathbb{R}^n , equivalently, A has n linearly independent eigenvectors v_1, \dots, v_n with associated eigenvalues $\lambda_1, \dots, \lambda_n$ (which need not be distinct).*
- ❸ *$\text{GM}_A(\lambda) = \text{AG}_A(\lambda)$ for every eigenvalue λ of A .*
- ❹ *The minimal polynomial of A has distinct roots (equivalently, A satisfies a polynomial $p(x) \in \mathbb{C}[x]$ having distinct roots).*

Proof. (1) \Rightarrow (2):

The use of eigenvalues and eigenvectors on diagonalization

Theorem

Let A be an $n \times n$ matrix (over \mathbb{C}). The following are equivalent:

- ① A is diagonalizable.
- ② The eigenvectors of A form a basis of \mathbb{R}^n , equivalently, A has n linearly independent eigenvectors v_1, \dots, v_n with associated eigenvalues $\lambda_1, \dots, \lambda_n$ (which need not be distinct).
- ③ $\text{GM}_A(\lambda) = \text{AG}_A(\lambda)$ for every eigenvalue λ of A .
- ④ The minimal polynomial of A has distinct roots (equivalently, A satisfies a polynomial $p(x) \in \mathbb{C}[x]$ having distinct roots).

Proof. (1) \Rightarrow (2): There is an $n \times n$ invertible matrix P such that

$$P^{-1}AP = \begin{pmatrix} \lambda_1 & 0 & \cdots & 0 \\ 0 & \lambda_2 & \cdots & 0 \\ \vdots & \vdots & \ddots & \vdots \\ 0 & 0 & \cdots & \lambda_n \end{pmatrix} \quad \text{for some } \lambda_1, \dots, \lambda_n \in \mathbb{R}.$$

The use of eigenvalues and eigenvectors on diagonalization

Theorem

Let A be an $n \times n$ matrix (over \mathbb{C}). The following are equivalent:

- 1 A is diagonalizable.
- 2 The eigenvectors of A form a basis of \mathbb{R}^n , equivalently, A has n linearly independent eigenvectors v_1, \dots, v_n with associated eigenvalues $\lambda_1, \dots, \lambda_n$ (which need not be distinct).
- 3 $\text{GM}_A(\lambda) = \text{AG}_A(\lambda)$ for every eigenvalue λ of A .
- 4 The minimal polynomial of A has distinct roots (equivalently, A satisfies a polynomial $p(x) \in \mathbb{C}[x]$ having distinct roots).

Proof. (1) \Rightarrow (2): There is an $n \times n$ invertible matrix P such that

$$P^{-1}AP = \begin{pmatrix} \lambda_1 & 0 & \cdots & 0 \\ 0 & \lambda_2 & \cdots & 0 \\ \vdots & \vdots & \ddots & \vdots \\ 0 & 0 & \cdots & \lambda_n \end{pmatrix} \quad \text{for some } \lambda_1, \dots, \lambda_n \in \mathbb{R}.$$

Hence left multiply by P from the left side.

Proof of the theorem contd...

Proof. (1) \Rightarrow (2): ... Thus $AP = P \begin{pmatrix} \lambda_1 & 0 & \cdots & 0 \\ 0 & \lambda_2 & \cdots & 0 \\ \vdots & \vdots & \ddots & \vdots \\ 0 & 0 & \cdots & \lambda_n \end{pmatrix}.$

Proof of the theorem contd...

Proof. (1) \Rightarrow (2): ... Thus $AP = P \begin{pmatrix} \lambda_1 & 0 & \cdots & 0 \\ 0 & \lambda_2 & \cdots & 0 \\ \vdots & \vdots & \ddots & \vdots \\ 0 & 0 & \cdots & \lambda_n \end{pmatrix}$.

Write $P = [v_1 \quad v_2 \quad \cdots \quad v_n]$ for some $v_1, \dots, v_n \in \mathbb{R}^n$.

Proof of the theorem contd...

Proof. (1) \Rightarrow (2): ... Thus $AP = P \begin{pmatrix} \lambda_1 & 0 & \cdots & 0 \\ 0 & \lambda_2 & \cdots & 0 \\ \vdots & \vdots & \ddots & \vdots \\ 0 & 0 & \cdots & \lambda_n \end{pmatrix}$.

Write $P = [v_1 \ v_2 \ \cdots \ v_n]$ for some $v_1, \dots, v_n \in \mathbb{R}^n$.

Then $AP = [Av_1 \ Av_2 \ \cdots \ Av_n]$

Proof of the theorem contd...

Proof. (1) \Rightarrow (2): ... Thus $AP = P \begin{pmatrix} \lambda_1 & 0 & \cdots & 0 \\ 0 & \lambda_2 & \cdots & 0 \\ \vdots & \vdots & \ddots & \vdots \\ 0 & 0 & \cdots & \lambda_n \end{pmatrix}$.

Write $P = [v_1 \ v_2 \ \cdots \ v_n]$ for some $v_1, \dots, v_n \in \mathbb{R}^n$.

Then $AP = [Av_1 \ Av_2 \ \cdots \ Av_n]$ and

$$P \begin{pmatrix} \lambda_1 & 0 & \cdots & 0 \\ 0 & \lambda_2 & \cdots & 0 \\ \vdots & \vdots & \ddots & \vdots \\ 0 & 0 & \cdots & \lambda_n \end{pmatrix} = \left[P \begin{pmatrix} \lambda_1 \\ 0 \\ \vdots \\ 0 \end{pmatrix} \quad P \begin{pmatrix} 0 \\ \lambda_2 \\ \vdots \\ 0 \end{pmatrix} \quad \cdots \quad P \begin{pmatrix} 0 \\ 0 \\ \vdots \\ \lambda_n \end{pmatrix} \right]$$

Proof of the theorem contd...

Proof. (1) \Rightarrow (2): ... Thus $AP = P \begin{pmatrix} \lambda_1 & 0 & \cdots & 0 \\ 0 & \lambda_2 & \cdots & 0 \\ \vdots & \vdots & \ddots & \vdots \\ 0 & 0 & \cdots & \lambda_n \end{pmatrix}$.

Write $P = [v_1 \ v_2 \ \cdots \ v_n]$ for some $v_1, \dots, v_n \in \mathbb{R}^n$.

Then $AP = [Av_1 \ Av_2 \ \cdots \ Av_n]$ and

$$P \begin{pmatrix} \lambda_1 & 0 & \cdots & 0 \\ 0 & \lambda_2 & \cdots & 0 \\ \vdots & \vdots & \ddots & \vdots \\ 0 & 0 & \cdots & \lambda_n \end{pmatrix} = \left[P \begin{pmatrix} \lambda_1 \\ 0 \\ \vdots \\ 0 \end{pmatrix} \quad P \begin{pmatrix} 0 \\ \lambda_2 \\ \vdots \\ 0 \end{pmatrix} \quad \cdots \quad P \begin{pmatrix} 0 \\ 0 \\ \vdots \\ \lambda_n \end{pmatrix} \right]$$

$$= [\lambda_1 v_1 \quad \lambda_2 v_2 \quad \cdots \quad \lambda_n v_n].$$

Proof of the theorem contd...

Proof. (1) \Rightarrow (2): ... Thus $AP = P \begin{pmatrix} \lambda_1 & 0 & \cdots & 0 \\ 0 & \lambda_2 & \cdots & 0 \\ \vdots & \vdots & \ddots & \vdots \\ 0 & 0 & \cdots & \lambda_n \end{pmatrix}$.

Write $P = [v_1 \ v_2 \ \cdots \ v_n]$ for some $v_1, \dots, v_n \in \mathbb{R}^n$.

Then $AP = [Av_1 \ Av_2 \ \cdots \ Av_n]$ and

$$P \begin{pmatrix} \lambda_1 & 0 & \cdots & 0 \\ 0 & \lambda_2 & \cdots & 0 \\ \vdots & \vdots & \ddots & \vdots \\ 0 & 0 & \cdots & \lambda_n \end{pmatrix} = \left[P \begin{pmatrix} \lambda_1 \\ 0 \\ \vdots \\ 0 \end{pmatrix} \quad P \begin{pmatrix} 0 \\ \lambda_2 \\ \vdots \\ 0 \end{pmatrix} \quad \cdots \quad P \begin{pmatrix} 0 \\ 0 \\ \vdots \\ \lambda_n \end{pmatrix} \right]$$

$$= [\lambda_1 v_1 \ \lambda_2 v_2 \ \cdots \ \lambda_n v_n].$$

Therefore $Av_i = \lambda_i v_i$ for every $1 \leq i \leq n$.

Proof of the theorem contd...

Proof. (1) \Rightarrow (2): ... Thus $AP = P \begin{pmatrix} \lambda_1 & 0 & \cdots & 0 \\ 0 & \lambda_2 & \cdots & 0 \\ \vdots & \vdots & \ddots & \vdots \\ 0 & 0 & \cdots & \lambda_n \end{pmatrix}$.

Write $P = [v_1 \ v_2 \ \cdots \ v_n]$ for some $v_1, \dots, v_n \in \mathbb{R}^n$.

Then $AP = [Av_1 \ Av_2 \ \cdots \ Av_n]$ and

$$P \begin{pmatrix} \lambda_1 & 0 & \cdots & 0 \\ 0 & \lambda_2 & \cdots & 0 \\ \vdots & \vdots & \ddots & \vdots \\ 0 & 0 & \cdots & \lambda_n \end{pmatrix} = \left[P \begin{pmatrix} \lambda_1 \\ 0 \\ \vdots \\ 0 \end{pmatrix} \quad P \begin{pmatrix} 0 \\ \lambda_2 \\ \vdots \\ 0 \end{pmatrix} \quad \cdots \quad P \begin{pmatrix} 0 \\ 0 \\ \vdots \\ \lambda_n \end{pmatrix} \right]$$

$$= [\lambda_1 v_1 \ \lambda_2 v_2 \ \cdots \ \lambda_n v_n].$$

Therefore $Av_i = \lambda_i v_i$ for every $1 \leq i \leq n$.

Note that v_1, \dots, v_n are linearly independent,

Proof of the theorem contd...

Proof. (1) \Rightarrow (2): ... Thus $AP = P \begin{pmatrix} \lambda_1 & 0 & \cdots & 0 \\ 0 & \lambda_2 & \cdots & 0 \\ \vdots & \vdots & \ddots & \vdots \\ 0 & 0 & \cdots & \lambda_n \end{pmatrix}$.

Write $P = [v_1 \ v_2 \ \cdots \ v_n]$ for some $v_1, \dots, v_n \in \mathbb{R}^n$.

Then $AP = [Av_1 \ Av_2 \ \cdots \ Av_n]$ and

$$P \begin{pmatrix} \lambda_1 & 0 & \cdots & 0 \\ 0 & \lambda_2 & \cdots & 0 \\ \vdots & \vdots & \ddots & \vdots \\ 0 & 0 & \cdots & \lambda_n \end{pmatrix} = \left[P \begin{pmatrix} \lambda_1 \\ 0 \\ \vdots \\ 0 \end{pmatrix} \quad P \begin{pmatrix} 0 \\ \lambda_2 \\ \vdots \\ 0 \end{pmatrix} \quad \cdots \quad P \begin{pmatrix} 0 \\ 0 \\ \vdots \\ \lambda_n \end{pmatrix} \right]$$

$$= [\lambda_1 v_1 \ \lambda_2 v_2 \ \cdots \ \lambda_n v_n].$$

Therefore $Av_i = \lambda_i v_i$ for every $1 \leq i \leq n$.

Note that v_1, \dots, v_n are linearly independent, since P is invertible.

Proof of the theorem contd...

Proof. (2) \Rightarrow (1): A has n linearly independent eigenvectors $v_1, \dots, v_n \in \mathbb{R}^n$ with associated eigenvalues $\lambda_1, \dots, \lambda_n \in \mathbb{R}$.

Proof of the theorem contd...

Proof. (2) \Rightarrow (1): A has n linearly independent eigenvectors

$v_1, \dots, v_n \in \mathbb{R}^n$ with associated eigenvalues $\lambda_1, \dots, \lambda_n \in \mathbb{R}$.

Set $P := \begin{bmatrix} v_1 & v_2 & \cdots & v_n \end{bmatrix}$.

Proof of the theorem contd...

Proof. (2) \Rightarrow (1): A has n linearly independent eigenvectors $v_1, \dots, v_n \in \mathbb{R}^n$ with associated eigenvalues $\lambda_1, \dots, \lambda_n \in \mathbb{R}$.
Set $P := \begin{bmatrix} v_1 & v_2 & \cdots & v_n \end{bmatrix}$. Clearly P is an $n \times n$ matrix.

Proof of the theorem contd...

Proof. (2) \Rightarrow (1): A has n linearly independent eigenvectors $v_1, \dots, v_n \in \mathbb{R}^n$ with associated eigenvalues $\lambda_1, \dots, \lambda_n \in \mathbb{R}$.
Set $P := [v_1 \ v_2 \ \cdots \ v_n]$. Clearly P is an $n \times n$ matrix.
Since v_1, \dots, v_n are linearly independent, P is invertible.

Proof of the theorem contd...

Proof. (2) \Rightarrow (1): A has n linearly independent eigenvectors

$v_1, \dots, v_n \in \mathbb{R}^n$ with associated eigenvalues $\lambda_1, \dots, \lambda_n \in \mathbb{R}$.

Set $P := [v_1 \ v_2 \ \cdots \ v_n]$. Clearly P is an $n \times n$ matrix.

Since v_1, \dots, v_n are linearly independent, P is invertible.

Moreover

$$AP = [Av_1 \ Av_2 \ \cdots \ Av_n]$$

Proof of the theorem contd...

Proof. (2) \Rightarrow (1): A has n linearly independent eigenvectors

$v_1, \dots, v_n \in \mathbb{R}^n$ with associated eigenvalues $\lambda_1, \dots, \lambda_n \in \mathbb{R}$.

Set $P := [v_1 \ v_2 \ \cdots \ v_n]$. Clearly P is an $n \times n$ matrix.

Since v_1, \dots, v_n are linearly independent, P is invertible.

Moreover

$$AP = [Av_1 \ Av_2 \ \cdots \ Av_n] = [\lambda_1 v_1 \ \lambda_2 v_2 \ \cdots \ \lambda_n v_n]$$

Proof of the theorem contd...

Proof. (2) \Rightarrow (1): A has n linearly independent eigenvectors

$v_1, \dots, v_n \in \mathbb{R}^n$ with associated eigenvalues $\lambda_1, \dots, \lambda_n \in \mathbb{R}$.

Set $P := [v_1 \ v_2 \ \cdots \ v_n]$. Clearly P is an $n \times n$ matrix.

Since v_1, \dots, v_n are linearly independent, P is invertible.

Moreover

$$\begin{aligned} AP &= [Av_1 \ Av_2 \ \cdots \ Av_n] = [\lambda_1 v_1 \ \lambda_2 v_2 \ \cdots \ \lambda_n v_n] \\ &= P \begin{pmatrix} \lambda_1 & 0 & \cdots & 0 \\ 0 & \lambda_2 & \cdots & 0 \\ \vdots & \vdots & \ddots & \vdots \\ 0 & 0 & \cdots & \lambda_n \end{pmatrix}. \end{aligned}$$

Proof of the theorem contd...

Proof. (2) \Rightarrow (1): A has n linearly independent eigenvectors

$v_1, \dots, v_n \in \mathbb{R}^n$ with associated eigenvalues $\lambda_1, \dots, \lambda_n \in \mathbb{R}$.

Set $P := [v_1 \ v_2 \ \cdots \ v_n]$. Clearly P is an $n \times n$ matrix.

Since v_1, \dots, v_n are linearly independent, P is invertible.

Moreover

$$\begin{aligned} AP &= [Av_1 \ Av_2 \ \cdots \ Av_n] = [\lambda_1 v_1 \ \lambda_2 v_2 \ \cdots \ \lambda_n v_n] \\ &= P \begin{pmatrix} \lambda_1 & 0 & \cdots & 0 \\ 0 & \lambda_2 & \cdots & 0 \\ \vdots & \vdots & \ddots & \vdots \\ 0 & 0 & \cdots & \lambda_n \end{pmatrix}. \end{aligned}$$

Therefore

$$P^{-1}AP = \begin{pmatrix} \lambda_1 & 0 & \cdots & 0 \\ 0 & \lambda_2 & \cdots & 0 \\ \vdots & \vdots & \ddots & \vdots \\ 0 & 0 & \cdots & \lambda_n \end{pmatrix}.$$

Proof of the theorem contd...

Proof. (2) \Rightarrow (1): A has n linearly independent eigenvectors

$v_1, \dots, v_n \in \mathbb{R}^n$ with associated eigenvalues $\lambda_1, \dots, \lambda_n \in \mathbb{R}$.

Set $P := [v_1 \ v_2 \ \cdots \ v_n]$. Clearly P is an $n \times n$ matrix.

Since v_1, \dots, v_n are linearly independent, P is invertible.

Moreover

$$\begin{aligned} AP &= [Av_1 \ Av_2 \ \cdots \ Av_n] = [\lambda_1 v_1 \ \lambda_2 v_2 \ \cdots \ \lambda_n v_n] \\ &= P \begin{pmatrix} \lambda_1 & 0 & \cdots & 0 \\ 0 & \lambda_2 & \cdots & 0 \\ \vdots & \vdots & \ddots & \vdots \\ 0 & 0 & \cdots & \lambda_n \end{pmatrix}. \end{aligned}$$

Therefore

$$P^{-1}AP = \begin{pmatrix} \lambda_1 & 0 & \cdots & 0 \\ 0 & \lambda_2 & \cdots & 0 \\ \vdots & \vdots & \ddots & \vdots \\ 0 & 0 & \cdots & \lambda_n \end{pmatrix}.$$

(1) \Leftrightarrow (3) \Leftrightarrow (4): We will skip it.



Cayley-Hamilton Theorem

- Let A be an $n \times n$ matrix over \mathbb{R} .

Cayley-Hamilton Theorem

- Let A be an $n \times n$ matrix over \mathbb{R} .
- Write A^r for the matrix multiplication of r many copies of A .

Cayley-Hamilton Theorem

- Let A be an $n \times n$ matrix over \mathbb{R} .
- Write A^r for the matrix multiplication of r many copies of A .
- For $c \in \mathbb{R}$, cA is just component wise scalar multiplication.

Cayley-Hamilton Theorem

- Let A be an $n \times n$ matrix over \mathbb{R} .
- Write A^r for the matrix multiplication of r many copies of A .
- For $c \in \mathbb{R}$, cA is just component wise scalar multiplication.
- If $f(x) = a_r x^r + \cdots + a_2 x^2 + a_1 x + a_0 \in \mathbb{R}[x]$,

Cayley-Hamilton Theorem

- Let A be an $n \times n$ matrix over \mathbb{R} .
- Write A^r for the matrix multiplication of r many copies of A .
- For $c \in \mathbb{R}$, cA is just component wise scalar multiplication.
- If $f(x) = a_r x^r + \cdots + a_2 x^2 + a_1 x + a_0 \in \mathbb{R}[x]$, then
 $f(A) = a_r A^r + \cdots + a_2 A^2 + a_1 A + a_0 I_n$ is an $n \times n$ matrix/ \mathbb{R} .

Cayley-Hamilton Theorem

- Let A be an $n \times n$ matrix over \mathbb{R} .
- Write A^r for the matrix multiplication of r many copies of A .
- For $c \in \mathbb{R}$, cA is just component wise scalar multiplication.
- If $f(x) = a_r x^r + \cdots + a_2 x^2 + a_1 x + a_0 \in \mathbb{R}[x]$, then
$$f(A) = a_r A^r + \cdots + a_2 A^2 + a_1 A + a_0 I_n$$
 is an $n \times n$ matrix/ \mathbb{R} .

Theorem (Cayley-Hamilton)

Consider the characteristic polynomial $p_A(x) := \det(xI_n - A)$.

Cayley-Hamilton Theorem

- Let A be an $n \times n$ matrix over \mathbb{R} .
- Write A^r for the matrix multiplication of r many copies of A .
- For $c \in \mathbb{R}$, cA is just component wise scalar multiplication.
- If $f(x) = a_r x^r + \cdots + a_2 x^2 + a_1 x + a_0 \in \mathbb{R}[x]$, then
$$f(A) = a_r A^r + \cdots + a_2 A^2 + a_1 A + a_0 I_n$$
 is an $n \times n$ matrix/ \mathbb{R} .

Theorem (Cayley-Hamilton)

Consider the characteristic polynomial $p_A(x) := \det(xI_n - A)$. Then $p_A(A) = 0$

Cayley-Hamilton Theorem

- Let A be an $n \times n$ matrix over \mathbb{R} .
- Write A^r for the matrix multiplication of r many copies of A .
- For $c \in \mathbb{R}$, cA is just component wise scalar multiplication.
- If $f(x) = a_r x^r + \cdots + a_2 x^2 + a_1 x + a_0 \in \mathbb{R}[x]$, then
$$f(A) = a_r A^r + \cdots + a_2 A^2 + a_1 A + a_0 I_n$$
 is an $n \times n$ matrix/ \mathbb{R} .

Theorem (Cayley-Hamilton)

Consider the characteristic polynomial $p_A(x) := \det(xI_n - A)$. Then $p_A(A) = 0$ (zero matrix of order $n \times n$).

Cayley-Hamilton Theorem

- Let A be an $n \times n$ matrix over \mathbb{R} .
- Write A^r for the matrix multiplication of r many copies of A .
- For $c \in \mathbb{R}$, cA is just component wise scalar multiplication.
- If $f(x) = a_r x^r + \cdots + a_2 x^2 + a_1 x + a_0 \in \mathbb{R}[x]$, then
$$f(A) = a_r A^r + \cdots + a_2 A^2 + a_1 A + a_0 I_n$$
 is an $n \times n$ matrix/ \mathbb{R} .

Theorem (Cayley-Hamilton)

Consider the characteristic polynomial $p_A(x) := \det(xI_n - A)$. Then $p_A(A) = 0$ (zero matrix of order $n \times n$).

Warning: $p_A(A) \neq \det(AI_n - A)$.

Cayley-Hamilton Theorem

- Let A be an $n \times n$ matrix over \mathbb{R} .
- Write A^r for the matrix multiplication of r many copies of A .
- For $c \in \mathbb{R}$, cA is just component wise scalar multiplication.
- If $f(x) = a_r x^r + \cdots + a_2 x^2 + a_1 x + a_0 \in \mathbb{R}[x]$, then
$$f(A) = a_r A^r + \cdots + a_2 A^2 + a_1 A + a_0 I_n$$
 is an $n \times n$ matrix/ \mathbb{R} .

Theorem (Cayley-Hamilton)

Consider the characteristic polynomial $p_A(x) := \det(xI_n - A)$. Then $p_A(A) = 0$ (zero matrix of order $n \times n$).

Warning: $p_A(A) \neq \det(AI_n - A)$. LHS is a matrix;

Cayley-Hamilton Theorem

- Let A be an $n \times n$ matrix over \mathbb{R} .
- Write A^r for the matrix multiplication of r many copies of A .
- For $c \in \mathbb{R}$, cA is just component wise scalar multiplication.
- If $f(x) = a_r x^r + \cdots + a_2 x^2 + a_1 x + a_0 \in \mathbb{R}[x]$, then
$$f(A) = a_r A^r + \cdots + a_2 A^2 + a_1 A + a_0 I_n$$
 is an $n \times n$ matrix/ \mathbb{R} .

Theorem (Cayley-Hamilton)

Consider the characteristic polynomial $p_A(x) := \det(xI_n - A)$. Then $p_A(A) = 0$ (zero matrix of order $n \times n$).

Warning: $p_A(A) \neq \det(AI_n - A)$. LHS is a matrix; RHS is a scalar.

Cayley-Hamilton Theorem

- Let A be an $n \times n$ matrix over \mathbb{R} .
- Write A^r for the matrix multiplication of r many copies of A .
- For $c \in \mathbb{R}$, cA is just component wise scalar multiplication.
- If $f(x) = a_r x^r + \cdots + a_2 x^2 + a_1 x + a_0 \in \mathbb{R}[x]$, then
$$f(A) = a_r A^r + \cdots + a_2 A^2 + a_1 A + a_0 I_n$$
 is an $n \times n$ matrix/ \mathbb{R} .

Theorem (Cayley-Hamilton)

Consider the characteristic polynomial $p_A(x) := \det(xI_n - A)$. Then $p_A(A) = 0$ (zero matrix of order $n \times n$).

Warning: $p_A(A) \neq \det(AI_n - A)$. LHS is a matrix; RHS is a scalar.

Example

$$\text{If } A = \begin{pmatrix} 1 & 2 \\ 3 & 4 \end{pmatrix},$$

Cayley-Hamilton Theorem

- Let A be an $n \times n$ matrix over \mathbb{R} .
- Write A^r for the matrix multiplication of r many copies of A .
- For $c \in \mathbb{R}$, cA is just component wise scalar multiplication.
- If $f(x) = a_r x^r + \cdots + a_2 x^2 + a_1 x + a_0 \in \mathbb{R}[x]$, then
$$f(A) = a_r A^r + \cdots + a_2 A^2 + a_1 A + a_0 I_n \text{ is an } n \times n \text{ matrix/}\mathbb{R}.$$

Theorem (Cayley-Hamilton)

Consider the characteristic polynomial $p_A(x) := \det(xI_n - A)$. Then $p_A(A) = 0$ (zero matrix of order $n \times n$).

Warning: $p_A(A) \neq \det(AI_n - A)$. LHS is a matrix; RHS is a scalar.

Example

If $A = \begin{pmatrix} 1 & 2 \\ 3 & 4 \end{pmatrix}$, then $p_A(x) = x^2 - 5x - 2$.

Cayley-Hamilton Theorem

- Let A be an $n \times n$ matrix over \mathbb{R} .
- Write A^r for the matrix multiplication of r many copies of A .
- For $c \in \mathbb{R}$, cA is just component wise scalar multiplication.
- If $f(x) = a_r x^r + \cdots + a_2 x^2 + a_1 x + a_0 \in \mathbb{R}[x]$, then
$$f(A) = a_r A^r + \cdots + a_2 A^2 + a_1 A + a_0 I_n \text{ is an } n \times n \text{ matrix/}\mathbb{R}.$$

Theorem (Cayley-Hamilton)

Consider the characteristic polynomial $p_A(x) := \det(xI_n - A)$. Then $p_A(A) = 0$ (zero matrix of order $n \times n$).

Warning: $p_A(A) \neq \det(AI_n - A)$. LHS is a matrix; RHS is a scalar.

Example

If $A = \begin{pmatrix} 1 & 2 \\ 3 & 4 \end{pmatrix}$, then $p_A(x) = x^2 - 5x - 2$. The Cayley-Hamilton

Theorem says that $A^2 - 5A - 2I_2 = \begin{pmatrix} 0 & 0 \\ 0 & 0 \end{pmatrix}$.

Similar matrices have same characteristic polynomial

Theorem

Let A and B be similar,

Similar matrices have same characteristic polynomial

Theorem

Let A and B be similar, i.e., $B = P^{-1}AP$ for some invertible matrix P .

Similar matrices have same characteristic polynomial

Theorem

Let A and B be similar, i.e., $B = P^{-1}AP$ for some invertible matrix P . Then $\det(xI_n - A) = \det(xI_n - B)$.

Proof. $\det(xI_n - B)$

Similar matrices have same characteristic polynomial

Theorem

Let A and B be similar, i.e., $B = P^{-1}AP$ for some invertible matrix P . Then $\det(xI_n - A) = \det(xI_n - B)$.

Proof. $\det(xI_n - B) = \det(xP^{-1}I_nP - P^{-1}AP)$

Similar matrices have same characteristic polynomial

Theorem

Let A and B be similar, i.e., $B = P^{-1}AP$ for some invertible matrix P . Then $\det(xI_n - A) = \det(xI_n - B)$.

Proof. $\det(xI_n - B) = \det(xP^{-1}I_nP - P^{-1}AP) = \det(P^{-1}(xI_n - A)P)$

Similar matrices have same characteristic polynomial

Theorem

Let A and B be similar, i.e., $B = P^{-1}AP$ for some invertible matrix P . Then $\det(xI_n - A) = \det(xI_n - B)$.

Proof. $\det(xI_n - B) = \det(xP^{-1}I_nP - P^{-1}AP) = \det(P^{-1}(xI_n - A)P)$
 $= (1/\det(P)) \det(xI_n - A) \det(P)$

Similar matrices have same characteristic polynomial

Theorem

Let A and B be similar, i.e., $B = P^{-1}AP$ for some invertible matrix P . Then $\det(xI_n - A) = \det(xI_n - B)$.

Proof. $\det(xI_n - B) = \det(xP^{-1}I_nP - P^{-1}AP) = \det(P^{-1}(xI_n - A)P)$
 $= (1/\det(P)) \det(xI_n - A) \det(P) = \det(xI_n - A).$ □

Theorem

Similar matrices have same characteristic polynomial

Theorem

Let A and B be similar, i.e., $B = P^{-1}AP$ for some invertible matrix P . Then $\det(xI_n - A) = \det(xI_n - B)$.

Proof. $\det(xI_n - B) = \det(xP^{-1}I_nP - P^{-1}AP) = \det(P^{-1}(xI_n - A)P)$
 $= (1/\det(P)) \det(xI_n - A) \det(P) = \det(xI_n - A).$ □

Theorem

Let A and B be similar,

Similar matrices have same characteristic polynomial

Theorem

Let A and B be similar, i.e., $B = P^{-1}AP$ for some invertible matrix P . Then $\det(xI_n - A) = \det(xI_n - B)$.

Proof. $\det(xI_n - B) = \det(xP^{-1}I_nP - P^{-1}AP) = \det(P^{-1}(xI_n - A)P)$
 $= (1/\det(P)) \det(xI_n - A) \det(P) = \det(xI_n - A).$ □

Theorem

Let A and B be similar, i.e., $B = P^{-1}AP$.

Similar matrices have same characteristic polynomial

Theorem

Let A and B be similar, i.e., $B = P^{-1}AP$ for some invertible matrix P . Then $\det(xI_n - A) = \det(xI_n - B)$.

Proof. $\det(xI_n - B) = \det(xP^{-1}I_nP - P^{-1}AP) = \det(P^{-1}(xI_n - A)P)$
 $= (1/\det(P)) \det(xI_n - A) \det(P) = \det(xI_n - A).$ □

Theorem

Let A and B be similar, i.e., $B = P^{-1}AP$. For a polynomial $f(x) \in \mathbb{R}[x]$,

Similar matrices have same characteristic polynomial

Theorem

Let A and B be similar, i.e., $B = P^{-1}AP$ for some invertible matrix P . Then $\det(xI_n - A) = \det(xI_n - B)$.

Proof. $\det(xI_n - B) = \det(xP^{-1}I_nP - P^{-1}AP) = \det(P^{-1}(xI_n - A)P)$
 $= (1/\det(P)) \det(xI_n - A) \det(P) = \det(xI_n - A).$ □

Theorem

Let A and B be similar, i.e., $B = P^{-1}AP$. For a polynomial $f(x) \in \mathbb{R}[x]$, $f(A) = 0$ if and only if $f(B) = 0$ (zero matrix).

Similar matrices have same characteristic polynomial

Theorem

Let A and B be similar, i.e., $B = P^{-1}AP$ for some invertible matrix P . Then $\det(xI_n - A) = \det(xI_n - B)$.

Proof. $\det(xI_n - B) = \det(xP^{-1}I_nP - P^{-1}AP) = \det(P^{-1}(xI_n - A)P)$
 $= (1/\det(P)) \det(xI_n - A) \det(P) = \det(xI_n - A).$ □

Theorem

Let A and B be similar, i.e., $B = P^{-1}AP$. For a polynomial $f(x) \in \mathbb{R}[x]$, $f(A) = 0$ if and only if $f(B) = 0$ (zero matrix).

Proof. Note: $P^{-1}A^rP = (P^{-1}AP)(P^{-1}AP) \cdots (P^{-1}AP) = B^r,$

Similar matrices have same characteristic polynomial

Theorem

Let A and B be similar, i.e., $B = P^{-1}AP$ for some invertible matrix P . Then $\det(xI_n - A) = \det(xI_n - B)$.

Proof. $\det(xI_n - B) = \det(xP^{-1}I_nP - P^{-1}AP) = \det(P^{-1}(xI_n - A)P)$
 $= (1/\det(P)) \det(xI_n - A) \det(P) = \det(xI_n - A).$ \square

Theorem

Let A and B be similar, i.e., $B = P^{-1}AP$. For a polynomial $f(x) \in \mathbb{R}[x]$, $f(A) = 0$ if and only if $f(B) = 0$ (zero matrix).

Proof. Note: $P^{-1}A^rP = (P^{-1}AP)(P^{-1}AP) \cdots (P^{-1}AP) = B^r$,
and $P^{-1}(c_1D_1 + c_2D_2)P = c_1(P^{-1}D_1P) + c_2(P^{-1}D_2P).$

Similar matrices have same characteristic polynomial

Theorem

Let A and B be similar, i.e., $B = P^{-1}AP$ for some invertible matrix P . Then $\det(xI_n - A) = \det(xI_n - B)$.

Proof. $\det(xI_n - B) = \det(xP^{-1}I_nP - P^{-1}AP) = \det(P^{-1}(xI_n - A)P)$
 $= (1/\det(P)) \det(xI_n - A) \det(P) = \det(xI_n - A).$ \square

Theorem

Let A and B be similar, i.e., $B = P^{-1}AP$. For a polynomial $f(x) \in \mathbb{R}[x]$, $f(A) = 0$ if and only if $f(B) = 0$ (zero matrix).

Proof. Note: $P^{-1}A^rP = (P^{-1}AP)(P^{-1}AP) \cdots (P^{-1}AP) = B^r$,
and $P^{-1}(c_1D_1 + c_2D_2)P = c_1(P^{-1}D_1P) + c_2(P^{-1}D_2P)$.
Verify that $P^{-1}f(A)P = f(B)$

Similar matrices have same characteristic polynomial

Theorem

Let A and B be similar, i.e., $B = P^{-1}AP$ for some invertible matrix P . Then $\det(xI_n - A) = \det(xI_n - B)$.

Proof. $\det(xI_n - B) = \det(xP^{-1}I_nP - P^{-1}AP) = \det(P^{-1}(xI_n - A)P)$
 $= (1/\det(P)) \det(xI_n - A) \det(P) = \det(xI_n - A).$ \square

Theorem

Let A and B be similar, i.e., $B = P^{-1}AP$. For a polynomial $f(x) \in \mathbb{R}[x]$, $f(A) = 0$ if and only if $f(B) = 0$ (zero matrix).

Proof. Note: $P^{-1}A^rP = (P^{-1}AP)(P^{-1}AP) \cdots (P^{-1}AP) = B^r$,
and $P^{-1}(c_1D_1 + c_2D_2)P = c_1(P^{-1}D_1P) + c_2(P^{-1}D_2P)$.
Verify that $P^{-1}f(A)P = f(B)$ and $Pf(B)P^{-1} = f(A)$.

Similar matrices have same characteristic polynomial

Theorem

Let A and B be similar, i.e., $B = P^{-1}AP$ for some invertible matrix P . Then $\det(xI_n - A) = \det(xI_n - B)$.

Proof. $\det(xI_n - B) = \det(xP^{-1}I_nP - P^{-1}AP) = \det(P^{-1}(xI_n - A)P)$
 $= (1/\det(P)) \det(xI_n - A) \det(P) = \det(xI_n - A).$ \square

Theorem

Let A and B be similar, i.e., $B = P^{-1}AP$. For a polynomial $f(x) \in \mathbb{R}[x]$, $f(A) = 0$ if and only if $f(B) = 0$ (zero matrix).

Proof. Note: $P^{-1}A^rP = (P^{-1}AP)(P^{-1}AP) \cdots (P^{-1}AP) = B^r$,
and $P^{-1}(c_1D_1 + c_2D_2)P = c_1(P^{-1}D_1P) + c_2(P^{-1}D_2P)$.
Verify that $P^{-1}f(A)P = f(B)$ and $Pf(B)P^{-1} = f(A)$.
Hence the theorem follows. \square

Proof of the Cayley-Hamilton Theorem for diagonalizable matrix

Let A be a diagonalizable matrix,

Proof of the Cayley-Hamilton Theorem for diagonalizable matrix

Let A be a diagonalizable matrix, i.e., there is P such that

$$P^{-1}AP = \begin{pmatrix} \lambda_1 & 0 & \cdots & 0 \\ 0 & \lambda_2 & \cdots & 0 \\ \vdots & \vdots & \ddots & \vdots \\ 0 & 0 & \cdots & \lambda_n \end{pmatrix} = B \text{ (say).}$$

Proof of the Cayley-Hamilton Theorem for diagonalizable matrix

Let A be a diagonalizable matrix, i.e., there is P such that

$$P^{-1}AP = \begin{pmatrix} \lambda_1 & 0 & \cdots & 0 \\ 0 & \lambda_2 & \cdots & 0 \\ \vdots & \vdots & \ddots & \vdots \\ 0 & 0 & \cdots & \lambda_n \end{pmatrix} = B \text{ (say).}$$

Note that $\det(xI_n - A) = \det(xI_n - B) = (x - \lambda_1) \cdots (x - \lambda_n)$.

Proof of the Cayley-Hamilton Theorem for diagonalizable matrix

Let A be a diagonalizable matrix, i.e., there is P such that

$$P^{-1}AP = \begin{pmatrix} \lambda_1 & 0 & \cdots & 0 \\ 0 & \lambda_2 & \cdots & 0 \\ \vdots & \vdots & \ddots & \vdots \\ 0 & 0 & \cdots & \lambda_n \end{pmatrix} = B \text{ (say).}$$

Note that $\det(xI_n - A) = \det(xI_n - B) = (x - \lambda_1) \cdots (x - \lambda_n)$.

By induction on n , one can verify that

$$(B - \lambda_1 I_n)(B - \lambda_2 I_n) \cdots (B - \lambda_n I_n) = 0.$$

Proof of the Cayley-Hamilton Theorem for diagonalizable matrix

Let A be a diagonalizable matrix, i.e., there is P such that

$$P^{-1}AP = \begin{pmatrix} \lambda_1 & 0 & \cdots & 0 \\ 0 & \lambda_2 & \cdots & 0 \\ \vdots & \vdots & \ddots & \vdots \\ 0 & 0 & \cdots & \lambda_n \end{pmatrix} = B \text{ (say).}$$

Note that $\det(xI_n - A) = \det(xI_n - B) = (x - \lambda_1) \cdots (x - \lambda_n)$.

By induction on n , one can verify that

$$(B - \lambda_1 I_n)(B - \lambda_2 I_n) \cdots (B - \lambda_n I_n) = 0.$$

Hence, multiplying P on left

Proof of the Cayley-Hamilton Theorem for diagonalizable matrix

Let A be a diagonalizable matrix, i.e., there is P such that

$$P^{-1}AP = \begin{pmatrix} \lambda_1 & 0 & \cdots & 0 \\ 0 & \lambda_2 & \cdots & 0 \\ \vdots & \vdots & \ddots & \vdots \\ 0 & 0 & \cdots & \lambda_n \end{pmatrix} = B \text{ (say).}$$

Note that $\det(xI_n - A) = \det(xI_n - B) = (x - \lambda_1) \cdots (x - \lambda_n)$.

By induction on n , one can verify that

$$(B - \lambda_1 I_n)(B - \lambda_2 I_n) \cdots (B - \lambda_n I_n) = 0.$$

Hence, multiplying P on left and P^{-1} on right,

Proof of the Cayley-Hamilton Theorem for diagonalizable matrix

Let A be a diagonalizable matrix, i.e., there is P such that

$$P^{-1}AP = \begin{pmatrix} \lambda_1 & 0 & \cdots & 0 \\ 0 & \lambda_2 & \cdots & 0 \\ \vdots & \vdots & \ddots & \vdots \\ 0 & 0 & \cdots & \lambda_n \end{pmatrix} = B \text{ (say).}$$

Note that $\det(xI_n - A) = \det(xI_n - B) = (x - \lambda_1) \cdots (x - \lambda_n)$.

By induction on n , one can verify that

$$(B - \lambda_1 I_n)(B - \lambda_2 I_n) \cdots (B - \lambda_n I_n) = 0.$$

Hence, multiplying P on left and P^{-1} on right, we get

$$(A - \lambda_1 I_n)(A - \lambda_2 I_n) \cdots (A - \lambda_n I_n) = 0$$

Proof of the Cayley-Hamilton Theorem for diagonalizable matrix

Let A be a diagonalizable matrix, i.e., there is P such that

$$P^{-1}AP = \begin{pmatrix} \lambda_1 & 0 & \cdots & 0 \\ 0 & \lambda_2 & \cdots & 0 \\ \vdots & \vdots & \ddots & \vdots \\ 0 & 0 & \cdots & \lambda_n \end{pmatrix} = B \text{ (say).}$$

Note that $\det(xI_n - A) = \det(xI_n - B) = (x - \lambda_1) \cdots (x - \lambda_n)$.

By induction on n , one can verify that

$$(B - \lambda_1 I_n)(B - \lambda_2 I_n) \cdots (B - \lambda_n I_n) = 0.$$

Hence, multiplying P on left and P^{-1} on right, we get

$$(A - \lambda_1 I_n)(A - \lambda_2 I_n) \cdots (A - \lambda_n I_n) = 0$$

i.e., $p_A(A) = 0$ (zero matrix).



Proof of the Cayley-Hamilton Theorem for diagonalizable matrix

Let A be a diagonalizable matrix, i.e., there is P such that

$$P^{-1}AP = \begin{pmatrix} \lambda_1 & 0 & \cdots & 0 \\ 0 & \lambda_2 & \cdots & 0 \\ \vdots & \vdots & \ddots & \vdots \\ 0 & 0 & \cdots & \lambda_n \end{pmatrix} = B \text{ (say).}$$

Note that $\det(xI_n - A) = \det(xI_n - B) = (x - \lambda_1) \cdots (x - \lambda_n)$.

By induction on n , one can verify that

$$(B - \lambda_1 I_n)(B - \lambda_2 I_n) \cdots (B - \lambda_n I_n) = 0.$$

Hence, multiplying P on left and P^{-1} on right, we get

$$(A - \lambda_1 I_n)(A - \lambda_2 I_n) \cdots (A - \lambda_n I_n) = 0$$

i.e., $p_A(A) = 0$ (zero matrix). □

Remark

It can be observed from the proof that if A is diagonalizable, then A satisfies a polynomial having distinct roots.

Applications of Eigenvalues and Eigenvectors

Some real life applications of the use of eigenvalues and eigenvectors in science, engineering and computer science can be found here:

[https://www.intmath.com/matrices-determinants/
8-applications-eigenvalues-eigenvectors.php](https://www.intmath.com/matrices-determinants/8-applications-eigenvalues-eigenvectors.php)

Thank You!