

# Differential Equations (MA 1150)

D. Sukumar

Lecture 1

April 7, 2020

## Course information

## Introduction

- ODE and its order

- General and particular solution

- IVP

- Geometry

## Solving methods

- Separation of variables

## Section 1

### Course information

## Information about this course

- ▶ Instructor: D Sukumar
- ▶ Office: Academic Block A, 506 [What is the use?](#)
- ▶ Email: [suku@math.iith.ac.in](mailto:suku@math.iith.ac.in)

## Information about this course

- ▶ Instructor: D Sukumar
- ▶ Office: Academic Block A, 506 [What is the use?](#)
- ▶ Email: [suku@math.iith.ac.in](mailto:suku@math.iith.ac.in)
- ▶ [Reference Textbooks](#)
  - ▶ [Elementary Differential Equations by William Trench](#), available at [ramanujan.math.trinity.edu/wtrench/texts/index.shtml](http://ramanujan.math.trinity.edu/wtrench/texts/index.shtml).

## Information about this course

- ▶ Instructor: D Sukumar
- ▶ Office: Academic Block A, 506 [What is the use?](#)
- ▶ Email: [suku@math.iith.ac.in](mailto:suku@math.iith.ac.in)
- ▶ [Reference Textbooks](#)
  - ▶ [Elementary Differential Equations by William Trench](#), available at [ramanujan.math.trinity.edu/wtrench/texts/index.shtml](http://ramanujan.math.trinity.edu/wtrench/texts/index.shtml).

**Marks Distribution.** Check the Course plan file uploaded in Google Classroom

Reference textbook: Elementary Differential Equations by William Trench

**Syllabus.** Ordinary Differential equations: First order linear equations, Bernoulli's equations, Exact equations and integrating factor, second order and higher order linear differential equations with constant coefficients.

- ▶ Lectures 1 – 4: Ordinary Differential equations: First order linear equations, Bernoulli's equations, Exact equations and integrating factor.
- ▶ Lectures 5 – 8: Second order and higher order linear differential equations with constant coefficients.
- ▶ Revision in Lecture 9.

►  $f : \mathbb{N} \rightarrow \mathbb{R}$ , Sequence and properties



- ▶  $f : \mathbb{N} \rightarrow \mathbb{R}$ , Sequence and properties
  - ▶ Convergence, divergence

- ▶  $f : \mathbb{N} \rightarrow \mathbb{R}$ , Sequence and properties
  - ▶ Convergence, divergence
  - ▶ Series, Test for convergence

- ▶  $f : \mathbb{N} \rightarrow \mathbb{R}$ , Sequence and properties
  - ▶ Convergence, divergence
  - ▶ Series, Test for convergence
- ▶  $f : \mathbb{R} \rightarrow \mathbb{R}$ ,  $f : [a, b] \subseteq \mathbb{R} \rightarrow \mathbb{R}$ ,

- ▶  $f : \mathbb{N} \rightarrow \mathbb{R}$ , Sequence and properties
  - ▶ Convergence, divergence
  - ▶ Series, Test for convergence
- ▶  $f : \mathbb{R} \rightarrow \mathbb{R}$ ,  $f : [a, b] \subseteq \mathbb{R} \rightarrow \mathbb{R}$ ,
  - ▶ Continuous, differentiable and integrable,

- ▶  $f : \mathbb{N} \rightarrow \mathbb{R}$ , Sequence and properties
  - ▶ Convergence, divergence
  - ▶ Series, Test for convergence
- ▶  $f : \mathbb{R} \rightarrow \mathbb{R}$ ,  $f : [a, b] \subseteq \mathbb{R} \rightarrow \mathbb{R}$ ,
  - ▶ Continuous, differentiable and integrable,
  - ▶ Maximum & Minimum, *Taylor Series*, *Fourier Series*.

- ▶  $f : \mathbb{N} \rightarrow \mathbb{R}$ , Sequence and properties
  - ▶ Convergence, divergence
  - ▶ Series, Test for convergence
- ▶  $f : \mathbb{R} \rightarrow \mathbb{R}$ ,  $f : [a, b] \subseteq \mathbb{R} \rightarrow \mathbb{R}$ ,
  - ▶ Continuous, differentiable and integrable,
  - ▶ Maximum & Minimum, *Taylor Series*, *Fourier Series*.
- ▶  $f : \mathbb{R}^n \rightarrow \mathbb{R}$ ,  $f : D \subseteq \mathbb{R}^n \rightarrow \mathbb{R}$ ,

- ▶  $f : \mathbb{N} \rightarrow \mathbb{R}$ , Sequence and properties
  - ▶ Convergence, divergence
  - ▶ Series, Test for convergence
- ▶  $f : \mathbb{R} \rightarrow \mathbb{R}$ ,  $f : [a, b] \subseteq \mathbb{R} \rightarrow \mathbb{R}$ ,
  - ▶ Continuous, differentiable and integrable,
  - ▶ Maximum & Minimum, *Taylor Series*, *Fourier Series*.
- ▶  $f : \mathbb{R}^n \rightarrow \mathbb{R}$ ,  $f : D \subseteq \mathbb{R}^n \rightarrow \mathbb{R}$ ,
  - ▶ Continuous, Partial Derivatives,

- ▶  $f : \mathbb{N} \rightarrow \mathbb{R}$ , Sequence and properties
  - ▶ Convergence, divergence
  - ▶ Series, Test for convergence
- ▶  $f : \mathbb{R} \rightarrow \mathbb{R}$ ,  $f : [a, b] \subseteq \mathbb{R} \rightarrow \mathbb{R}$ ,
  - ▶ Continuous, differentiable and integrable,
  - ▶ Maximum & Minimum, *Taylor Series*, *Fourier Series*.
- ▶  $f : \mathbb{R}^n \rightarrow \mathbb{R}$ ,  $f : D \subseteq \mathbb{R}^n \rightarrow \mathbb{R}$ ,
  - ▶ Continuous, Partial Derivatives,
  - ▶ Maximum & Minimum, *Taylor Series*, *Fourier Series*.



- ▶  $f : \mathbb{N} \rightarrow \mathbb{R}$ , Sequence and properties
  - ▶ Convergence, divergence
  - ▶ Series, Test for convergence
- ▶  $f : \mathbb{R} \rightarrow \mathbb{R}$ ,  $f : [a, b] \subseteq \mathbb{R} \rightarrow \mathbb{R}$ ,
  - ▶ Continuous, differentiable and integrable,
  - ▶ Maximum & Minimum, *Taylor Series*, *Fourier Series*.
- ▶  $f : \mathbb{R}^n \rightarrow \mathbb{R}$ ,  $f : D \subseteq \mathbb{R}^n \rightarrow \mathbb{R}$ ,
  - ▶ Continuous, Partial Derivatives,
  - ▶ Maximum & Minimum, *Taylor Series*, *Fourier Series*.
  - ▶ Integration

- ▶  $f : \mathbb{N} \rightarrow \mathbb{R}$ , Sequence and properties
  - ▶ Convergence, divergence
  - ▶ Series, Test for convergence
- ▶  $f : \mathbb{R} \rightarrow \mathbb{R}$ ,  $f : [a, b] \subseteq \mathbb{R} \rightarrow \mathbb{R}$ ,
  - ▶ Continuous, differentiable and integrable,
  - ▶ Maximum & Minimum, *Taylor Series, Fourier Series.*
- ▶  $f : \mathbb{R}^n \rightarrow \mathbb{R}$ ,  $f : D \subseteq \mathbb{R}^n \rightarrow \mathbb{R}$ ,
  - ▶ Continuous, Partial Derivatives,
  - ▶ Maximum & Minimum, *Taylor Series, Fourier Series.*
  - ▶ Integration
- ▶  $f : \mathbb{R} \rightarrow \mathbb{R}^m$ , plane and space curves

- ▶  $f : \mathbb{N} \rightarrow \mathbb{R}$ , Sequence and properties
  - ▶ Convergence, divergence
  - ▶ Series, Test for convergence
- ▶  $f : \mathbb{R} \rightarrow \mathbb{R}$ ,  $f : [a, b] \subseteq \mathbb{R} \rightarrow \mathbb{R}$ ,
  - ▶ Continuous, differentiable and integrable,
  - ▶ Maximum & Minimum, *Taylor Series*, *Fourier Series*.
- ▶  $f : \mathbb{R}^n \rightarrow \mathbb{R}$ ,  $f : D \subseteq \mathbb{R}^n \rightarrow \mathbb{R}$ ,
  - ▶ Continuous, Partial Derivatives,
  - ▶ Maximum & Minimum, *Taylor Series*, *Fourier Series*.
  - ▶ Integration
- ▶  $f : \mathbb{R} \rightarrow \mathbb{R}^m$ , plane and space curves
- ▶  $f : \mathbb{R}^n \rightarrow \mathbb{R}^m$ , Integration and Differentiation

- ▶  $f : \mathbb{N} \rightarrow \mathbb{R}$ , Sequence and properties
  - ▶ Convergence, divergence
  - ▶ Series, Test for convergence
- ▶  $f : \mathbb{R} \rightarrow \mathbb{R}$ ,  $f : [a, b] \subseteq \mathbb{R} \rightarrow \mathbb{R}$ ,
  - ▶ Continuous, differentiable and integrable,
  - ▶ Maximum & Minimum, *Taylor Series, Fourier Series.*
- ▶  $f : \mathbb{R}^n \rightarrow \mathbb{R}$ ,  $f : D \subseteq \mathbb{R}^n \rightarrow \mathbb{R}$ ,
  - ▶ Continuous, Partial Derivatives,
  - ▶ Maximum & Minimum, *Taylor Series, Fourier Series.*
  - ▶ Integration
- ▶  $f : \mathbb{R} \rightarrow \mathbb{R}^m$ , plane and space curves
- ▶  $f : \mathbb{R}^n \rightarrow \mathbb{R}^m$ , Integration and Differentiation
- ▶  $f : \mathbb{R}^n \rightarrow \mathbb{R}^m$ , Linear algebra

- ▶  $f : \mathbb{N} \rightarrow \mathbb{R}$ , Sequence and properties
  - ▶ Convergence, divergence
  - ▶ Series, Test for convergence
- ▶  $f : \mathbb{R} \rightarrow \mathbb{R}$ ,  $f : [a, b] \subseteq \mathbb{R} \rightarrow \mathbb{R}$ ,
  - ▶ Continuous, differentiable and integrable,
  - ▶ Maximum & Minimum, *Taylor Series, Fourier Series.*
- ▶  $f : \mathbb{R}^n \rightarrow \mathbb{R}$ ,  $f : D \subseteq \mathbb{R}^n \rightarrow \mathbb{R}$ ,
  - ▶ Continuous, Partial Derivatives,
  - ▶ Maximum & Minimum, *Taylor Series, Fourier Series.*
  - ▶ Integration
- ▶  $f : \mathbb{R} \rightarrow \mathbb{R}^m$ , plane and space curves
- ▶  $f : \mathbb{R}^n \rightarrow \mathbb{R}^m$ , Integration and Differentiation
- ▶  $f : \mathbb{R}^n \rightarrow \mathbb{R}^m$ , Linear algebra
- ▶ Differential Equations (ordinary and partial)

- ▶  $f : \mathbb{N} \rightarrow \mathbb{R}$ , Sequence and properties
  - ▶ Convergence, divergence
  - ▶ Series, Test for convergence
- ▶  $f : \mathbb{R} \rightarrow \mathbb{R}$ ,  $f : [a, b] \subseteq \mathbb{R} \rightarrow \mathbb{R}$ ,
  - ▶ Continuous, differentiable and integrable,
  - ▶ Maximum & Minimum, *Taylor Series*, *Fourier Series*.
- ▶  $f : \mathbb{R}^n \rightarrow \mathbb{R}$ ,  $f : D \subseteq \mathbb{R}^n \rightarrow \mathbb{R}$ ,
  - ▶ Continuous, Partial Derivatives,
  - ▶ Maximum & Minimum, *Taylor Series*, *Fourier Series*.
  - ▶ Integration
- ▶  $f : \mathbb{R} \rightarrow \mathbb{R}^m$ , plane and space curves
- ▶  $f : \mathbb{R}^n \rightarrow \mathbb{R}^m$ , Integration and Differentiation
- ▶  $f : \mathbb{R}^n \rightarrow \mathbb{R}^m$ , Linear algebra
- ▶ Differential Equations (ordinary and partial)
- ▶ Modeling and solving

## Section 2

### Introduction

## Subsection 1

ODE and its order



## Ordinary Differential Equation

**Definition.** Let  $y = y(x)$  be a function of  $x$ . An Ordinary differential equation is an equation involving **atleast one** derivative of  $y$ .

# Ordinary Differential Equation

**Definition.** Let  $y = y(x)$  be a function of  $x$ . An Ordinary differential equation is an equation involving **atleast one** derivative of  $y$ .

**Notation.**

- ▶ Ordinary differential equation by **ODE**.

## Ordinary Differential Equation

**Definition.** Let  $y = y(x)$  be a function of  $x$ . An Ordinary differential equation is an equation involving **atleast one** derivative of  $y$ .

**Notation.**

- ▶ Ordinary differential equation by **ODE**.
- ▶ Derivative of  $y$  by  $y'$ ,  $\frac{dy}{dx}$  or  $y^{(1)}$ .

The order of an ODE is the highest order of derivative of  $y$  occurring in the ODE.

**Examples.**

- ▶  $y' = x^3y^4 + y$  is a **1st** order ODE.
- ▶  $y'' + x^5y' + y = \cos x$  is a **2nd** order ODE.
- ▶  $y^{(4)} + xy^{(1)}y^{(2)} + 2xy = \sin x$  is a **4th** order ODE.

## Open Intervals

Given any  $a, b \in \mathbb{R}$ , we define the **open interval** from  $a$  to  $b$  to be the set

$$(a, b) = \{x \in \mathbb{R} : a < x < b\},$$

and the **closed interval** from  $a$  to  $b$  to be the set  $[a, b] = \{x \in \mathbb{R} : a \leq x \leq b\}$ .

## Open Intervals

Given any  $a, b \in \mathbb{R}$ , we define the **open interval** from  $a$  to  $b$  to be the set

$$(a, b) = \{x \in \mathbb{R} : a < x < b\},$$

and the **closed interval** from  $a$  to  $b$  to be the set  $[a, b] = \{x \in \mathbb{R} : a \leq x \leq b\}$ .

It is often useful to consider the symbols  $\infty$

## Open Intervals

Given any  $a, b \in \mathbb{R}$ , we define the **open interval** from  $a$  to  $b$  to be the set

$$(a, b) = \{x \in \mathbb{R} : a < x < b\},$$

and the **closed interval** from  $a$  to  $b$  to be the set  $[a, b] = \{x \in \mathbb{R} : a \leq x \leq b\}$ .

It is often useful to consider the symbols  $\infty$  (called infinity) and  $-\infty$  (called minus infinity), which may be thought as the fictional (right and left) endpoints of the number line.

## Open Intervals

Given any  $a, b \in \mathbb{R}$ , we define the **open interval** from  $a$  to  $b$  to be the set

$$(a, b) = \{x \in \mathbb{R} : a < x < b\},$$

and the **closed interval** from  $a$  to  $b$  to be the set  $[a, b] = \{x \in \mathbb{R} : a \leq x \leq b\}$ .

It is often useful to consider the symbols  $\infty$  (called infinity) and  $-\infty$  (called minus infinity), which may be thought as the fictional (right and left) endpoints of the number line.

We use the symbols  $\infty$  and  $-\infty$  to define, for any  $a \in \mathbb{R}$ , the following semi-infinite (open) intervals:

$$(-\infty, b) = \{x \in \mathbb{R} : x < b\} \text{ and } (a, \infty) = \{x \in \mathbb{R} : a < x\}.$$

The set  $\mathbb{R}$  can also be thought of as the doubly infinite (open) interval  $(-\infty, \infty)$ .

## Subsection 2

### General and particular solution



## Solution of an ODE

**Definition.** A solution of an ODE is a function  $y = f(x)$  which satisfies the ODE on some open interval.

## Solution of an ODE

**Definition.** A solution of an ODE is a function  $y = f(x)$  which satisfies the ODE on some open interval.

**Example** Consider the ODE:  $y' = \cos x$ .

## Solution of an ODE

**Definition.** A solution of an ODE is a function  $y = f(x)$  which satisfies the ODE on some open interval.

**Example** Consider the ODE:  $y' = \cos x$ .

Note that  $y(x) = \sin x + c$  is an explicit solution to given ODE on  $\mathbb{R}$ , where  $c$  is arbitrary constant. (Why is this a solution?)

## Solution of an ODE

**Definition.** A solution of an ODE is a function  $y = f(x)$  which satisfies the ODE on some open interval.

**Example** Consider the ODE:  $y' = \cos x$ .

Note that  $y(x) = \sin x + c$  is an explicit solution to given ODE on  $\mathbb{R}$ , where  $c$  is arbitrary constant. (Why is this a solution?)

**Question** Do ODE always have explicit solutions?

## Solution of an ODE

**Definition.** A solution of an ODE is a function  $y = f(x)$  which satisfies the ODE on some open interval.

**Example** Consider the ODE:  $y' = \cos x$ .

Note that  $y(x) = \sin x + c$  is an explicit solution to given ODE on  $\mathbb{R}$ , where  $c$  is arbitrary constant. (Why is this a solution?)

**Question** Do ODE always have explicit solutions? (No!)

## Solution of an ODE

**Definition.** A solution of an ODE is a function  $y = f(x)$  which satisfies the ODE on some open interval.

**Example** Consider the ODE:  $y' = \cos x$ .

Note that  $y(x) = \sin x + c$  is an explicit solution to given ODE on  $\mathbb{R}$ , where  $c$  is arbitrary constant. (Why is this a solution?)

**Question** Do ODE always have explicit solutions? (No!)

ODE can have implicit solutions as well. Try with ODE:  $yy' = -x$ . (Be careful in writing solution!!)

## Solution of an ODE

**Definition.** A solution of an ODE is a function  $y = f(x)$  which satisfies the ODE on some open interval.

**Example** Consider the ODE:  $y' = \cos x$ .

Note that  $y(x) = \sin x + c$  is an explicit solution to given ODE on  $\mathbb{R}$ , where  $c$  is arbitrary constant. (Why is this a solution?)

**Question** Do ODE always have explicit solutions? (No!)

ODE can have implicit solutions as well. Try with ODE:  $yy' = -x$ . (Be careful in writing solution!! (why!))

## Solution of an ODE

**Definition.** A solution of an ODE is a function  $y = f(x)$  which satisfies the ODE on some open interval.

**Example** Consider the ODE:  $y' = \cos x$ .

Note that  $y(x) = \sin x + c$  is an explicit solution to given ODE on  $\mathbb{R}$ , where  $c$  is arbitrary constant. (Why is this a solution?)

**Question** Do ODE always have explicit solutions? (No!)

ODE can have implicit solutions as well. Try with ODE:  $yy' = -x$ . (Be careful in writing solution!! (why!) We want  $y$  to be a function.)



## Solution of an ODE

**Definition.** A solution of an ODE is a function  $y = f(x)$  which satisfies the ODE on some open interval.

**Example** Consider the ODE:  $y' = \cos x$ .

Note that  $y(x) = \sin x + c$  is an explicit solution to given ODE on  $\mathbb{R}$ , where  $c$  is arbitrary constant. (Why is this a solution?)

**Question** Do ODE always have explicit solutions? (No!)

ODE can have implicit solutions as well. Try with ODE:  $yy' = -x$ . (Be careful in writing solution!! (why!) We want  $y$  to be a function.)

**Question** Do ODE always have solutions?

## Solution of an ODE

**Definition.** A solution of an ODE is a function  $y = f(x)$  which satisfies the ODE on some open interval.

**Example** Consider the ODE:  $y' = \cos x$ .

Note that  $y(x) = \sin x + c$  is an explicit solution to given ODE on  $\mathbb{R}$ , where  $c$  is arbitrary constant. (Why is this a solution?)

**Question** Do ODE always have explicit solutions? (No!)

ODE can have implicit solutions as well. Try with ODE:  $yy' = -x$ . (Be careful in writing solution!! (why!) We want  $y$  to be a function.)

**Question** Do ODE always have solutions?

## Solution of an ODE

**Example** Consider the ODE:  $y' = \cos x$ .

## Solution of an ODE

**Example** Consider the ODE:  $y' = \cos x$ .

$y = \sin x + c$  is a solution to given ODE.

## Solution of an ODE

**Example** Consider the ODE:  $y' = \cos x$ .

$y = \sin x + c$  is a solution to given ODE.

Draw the diagram for  $y = \sin x + c$  with  $c = -2, -1, 0, 1, 2$ .

## Solution of an ODE

**Example** Consider the ODE:  $y' = \cos x$ .

$y = \sin x + c$  is a solution to given ODE.

Draw the diagram for  $y = \sin x + c$  with  $c = -2, -1, 0, 1, 2$ .

**General and Particular solution**  $y = \sin x + c$  is a general solution and if we choose specific  $c$ , e.g. say  $c = 1$ , then  $y = \sin x + 1$  is a particular solution.

## Solution of an ODE

**Example** Consider the ODE:  $y' = \cos x$ .

$y = \sin x + c$  is a solution to given ODE.

Draw the diagram for  $y = \sin x + c$  with  $c = -2, -1, 0, 1, 2$ .

**General and Particular solution**  $y = \sin x + c$  is a general solution and if we choose specific  $c$ , e.g. say  $c = 1$ , then  $y = \sin x + 1$  is a particular solution.

**Question** Can we recover all possible solutions of a given ODE from general solution?

## Solution of an ODE

**Example** Consider the ODE:  $y' = \cos x$ .

$y = \sin x + c$  is a solution to given ODE.

Draw the diagram for  $y = \sin x + c$  with  $c = -2, -1, 0, 1, 2$ .

**General and Particular solution**  $y = \sin x + c$  is a general solution and if we choose specific  $c$ , e.g. say  $c = 1$ , then  $y = \sin x + 1$  is a particular solution.

**Question** Can we recover all possible solutions of a given ODE from general solution? (Think! you may refer to Textbook).



## Applications. Modeling. Initial Value Problems

Many scientific and in particular physical laws appear in the form of differential equations.

## Applications. Modeling. Initial Value Problems

Many scientific and in particular physical laws appear in the form of differential equations.

**Modeling.** Converting physical system to a mathematical model. Then we find solution to mathematical model, and give physical interpretation of the result.

## Applications. Modeling. Initial Value Problems

Many scientific and in particular physical laws appear in the form of differential equations.

**Modeling.** Converting physical system to a mathematical model. Then we find solution to mathematical model, and give physical interpretation of the result.

**Examples** Radioactive substance decay, Growth of bacteria, Falling stone from certain height, Pendulum movement, Population growth etc. (For more details, refer to textbook).

## Subsection 3

IVP

## Initial Value Problem (IVP)

**Definition.** An Initial value problem (IVP) for 1st order ODE is

$$y' = f(x, y), \text{ where } y(x_0) = y_0.$$

## Initial Value Problem (IVP)

**Definition.** An Initial value problem (IVP) for 1st order ODE is

$$y' = f(x, y), \text{ where } y(x_0) = y_0.$$

**Solution to IVP.** A function  $y = y(x)$  defined on some open interval  $(a, b)$  containing  $x_0$  is a solution of the IVP if  $y$  satisfies the ODE on open interval  $(a, b)$  and  $y(x_0) = y_0$ .

## Subsection 4

### Geometry

Geogebra Webpage



## Section 3

### Solving methods

## Subsection 1

### Separation of variables

## Separation of Variables to solve ODE: Example

Solve.  $y' + 2xy = 0$ ?

## Separation of Variables to solve ODE: Example

Solve.  $y' + 2xy = 0$ ?

Rewrite this equation as  $\frac{1}{y} \frac{dy}{dx} = -2x$ .

## Separation of Variables to solve ODE: Example

**Solve.**  $y' + 2xy = 0$ ?

Rewrite this equation as  $\frac{1}{y} \frac{dy}{dx} = -2x$ .

We can solve this equation by separating the variables.

$$\begin{aligned}\frac{1}{y} dy &= -2x \, dx \\ \Rightarrow \int \frac{1}{y} dy &= \int -2x \, dx, \\ \Rightarrow \ln |y| &= -x^2 + c, \\ \Rightarrow y &= c_1 e^{-x^2}.\end{aligned}$$

## Separation of Variables to solve ODE: Example

**Solve.**  $y' + 2xy = 0$ ?

Rewrite this equation as  $\frac{1}{y} \frac{dy}{dx} = -2x$ .

We can solve this equation by separating the variables.

$$\begin{aligned}\frac{1}{y} dy &= -2x \, dx \\ \Rightarrow \int \frac{1}{y} dy &= \int -2x \, dx, \\ \Rightarrow \ln |y| &= -x^2 + c, \\ \Rightarrow y &= c_1 e^{-x^2}.\end{aligned}$$

**Question.** Is this  $y$  solution to given ODE?

## Separation of Variables: General Method

Let  $M(x, y) + N(x, y)\frac{dy}{dx} = 0$  be a differential equation.

## Separation of Variables: General Method

Let  $M(x, y) + N(x, y) \frac{dy}{dx} = 0$  be a differential equation.

This equation is said to be **separable** if it is possible to choose  $M$  and  $N$  such that  $M$  is a function only in  $x$  and  $N$  is a function only in  $y$ .



## Separation of Variables: General Method

Let  $M(x, y) + N(x, y) \frac{dy}{dx} = 0$  be a differential equation.

This equation is said to be **separable** if it is possible to choose  $M$  and  $N$  such that  $M$  is a function only in  $x$  and  $N$  is a function only in  $y$ .

Assume that ODE is separable.

## Separation of Variables: General Method

Let  $M(x, y) + N(x, y) \frac{dy}{dx} = 0$  be a differential equation.

This equation is said to be **separable** if it is possible to choose  $M$  and  $N$  such that  $M$  is a function only in  $x$  and  $N$  is a function only in  $y$ .

Assume that ODE is separable.

Let  $H_1$  and  $H_2$  be anti derivatives of  $M$  and  $N$  respectively. This means  $H_1'(x) = M(x)$  and  $H_2'(y) = N(y)$ . Then our ODE is

$$H_1'(x) + H_2'(y) \frac{dy}{dx} = 0.$$

## Separation of Variables: General Method (continued...)

$H_1'(x) + H_2'(y) \frac{dy}{dx} = 0$  can be re-written as

$$\frac{d}{dx} [H_1(x) + H_2(y(x))] = 0.$$

## Separation of Variables: General Method (continued...)

$H_1'(x) + H_2'(y)\frac{dy}{dx} = 0$  can be re-written as

$$\frac{d}{dx} [H_1(x) + H_2(y(x))] = 0.$$

Thus we get that the solution to the ODE will satisfy the equation

$$H_1(x) + H_2(y(x)) = c.$$

## Separation of Variables: General Method (continued...)

$H_1'(x) + H_2'(y) \frac{dy}{dx} = 0$  can be re-written as

$$\frac{d}{dx} [H_1(x) + H_2(y(x))] = 0.$$

Thus we get that the solution to the ODE will satisfy the equation

$$H_1(x) + H_2(y(x)) = c.$$

In general, the separable variables method only gives us the implicit solution to the given ODE.

## Separable ODEs - Example

**Example** Solve  $y' = 2xy^2$ .

## Separable ODEs - Example

**Example** Solve  $y' = 2xy^2$ .

Re-write ODE as

$$\frac{y'}{y^2} = 2x.$$

## Separable ODEs - Example

**Example** Solve  $y' = 2xy^2$ .

Re-write ODE as

$$\frac{y'}{y^2} = 2x.$$

Integrating, we get  $\frac{-1}{y} = x^2 + c$ . Thus

$$y = \frac{-1}{x^2 + c}$$



## Separable ODEs - Example

**Example** Solve  $y' = 2xy^2$ .

Re-write ODE as

$$\frac{y'}{y^2} = 2x.$$

Integrating, we get  $\frac{-1}{y} = x^2 + c$ . Thus

$$y = \frac{-1}{x^2 + c}$$

**Question.** Is this  $y$  solution to given ODE? Under what conditions,  $y$  will be solution to given ODE?

## Separable ODEs - Example

**Example** Solve  $y' = 2xy^2$ .

Re-write ODE as

$$\frac{y'}{y^2} = 2x.$$

Integrating, we get  $\frac{-1}{y} = x^2 + c$ . Thus

$$y = \frac{-1}{x^2 + c}$$

**Question.** Is this  $y$  solution to given ODE? Under what conditions,  $y$  will be solution to given ODE?

**Remark** Note that  $y \equiv 0$  is also a solution. However this solution cannot be obtained for any choice of  $c$ .

## Separable ODEs - Example

**Example** Solve  $y' = 2xy^2$ .

Re-write ODE as

$$\frac{y'}{y^2} = 2x.$$

Integrating, we get  $\frac{-1}{y} = x^2 + c$ . Thus

$$y = \frac{-1}{x^2 + c}$$

**Question.** Is this  $y$  solution to given ODE? Under what conditions,  $y$  will be solution to given ODE?

**Remark** Note that  $y \equiv 0$  is also a solution. However this solution cannot be obtained for any choice of  $c$ . (Does this remind you of something!!!)

## Converting to Separable Equation

Solve  $xy' = x + y$ .

## Converting to Separable Equation

**Solve**  $xy' = x + y$ .

Re-write it as

$$y' = \frac{y}{x} + 1.$$

## Converting to Separable Equation

**Solve**  $xy' = x + y$ .

Re-write it as

$$y' = \frac{y}{x} + 1.$$

We can make this ODE separable by changing our variables.

Let  $v = \frac{y}{x}$  or  $y = vx$ . Then  $y' = v'x + v$ .

Substituting we get  $v'x = 1$ .

## Converting to Separable Equation

**Solve**  $xy' = x + y$ .

Re-write it as

$$y' = \frac{y}{x} + 1.$$

We can make this ODE separable by changing our variables.

Let  $v = \frac{y}{x}$  or  $y = vx$ . Then  $y' = v'x + v$ .

Substituting we get  $v'x = 1$ .

Apply separation of variables to get  $y = (\ln |x| + C)x$ .

## Converting to Separable Equation

**Solve**  $xy' = x + y$ .

Re-write it as

$$y' = \frac{y}{x} + 1.$$

We can make this ODE separable by changing our variables.

Let  $v = \frac{y}{x}$  or  $y = vx$ . Then  $y' = v'x + v$ .

Substituting we get  $v'x = 1$ .

Apply separation of variables to get  $y = (\ln|x| + C)x$ .

**Question.** Is this  $y$  solution to given ODE? Under what conditions,  $y$  will be solution to given ODE?



## Converting to Separable Equation

**Solve**  $xy' = x + y$ .

Re-write it as

$$y' = \frac{y}{x} + 1.$$

We can make this ODE separable by changing our variables.

Let  $v = \frac{y}{x}$  or  $y = vx$ . Then  $y' = v'x + v$ .

Substituting we get  $v'x = 1$ .

Apply separation of variables to get  $y = (\ln|x| + C)x$ .

**Question.** Is this  $y$  solution to given ODE? Under what conditions,  $y$  will be solution to given ODE?

- Any non linear ODE of the form  $y' = q\left(\frac{y}{x}\right)$  can be converted into a separable ODE by substituting  $y = vx$ .

## Example

**Example** Solve  $x^2 y' = y^2 + xy - x^2$ .

## Example

**Example** Solve  $x^2 y' = y^2 + xy - x^2$ .

Re-write it as

$$y' = \frac{y^2 + xy - x^2}{x^2} = \left(\frac{y}{x}\right)^2 + \frac{y}{x} - 1.$$

## Example

**Example** Solve  $x^2 y' = y^2 + xy - x^2$ .

Re-write it as

$$y' = \frac{y^2 + xy - x^2}{x^2} = \left(\frac{y}{x}\right)^2 + \frac{y}{x} - 1.$$

Substituting  $y = vx$ , we get

$$v'x + v = v^2 + v - 1.$$

## Example

**Example** Solve  $x^2 y' = y^2 + xy - x^2$ .

Re-write it as

$$y' = \frac{y^2 + xy - x^2}{x^2} = \left(\frac{y}{x}\right)^2 + \frac{y}{x} - 1.$$

Substituting  $y = vx$ , we get

$$v'x + v = v^2 + v - 1.$$

Re-write it as

$$\begin{aligned}\frac{v'}{v^2 - 1} &= \frac{1}{x} \\ \frac{1}{2} \left( \frac{1}{v-1} - \frac{1}{v+1} \right) v' &= \frac{1}{x}.\end{aligned}$$

## Example

**Example** Solve  $x^2 y' = y^2 + xy - x^2$ .

Re-write it as

$$y' = \frac{y^2 + xy - x^2}{x^2} = \left(\frac{y}{x}\right)^2 + \frac{y}{x} - 1.$$

Substituting  $y = vx$ , we get

$$v'x + v = v^2 + v - 1.$$

Re-write it as

$$\begin{aligned}\frac{v'}{v^2 - 1} &= \frac{1}{x} \\ \frac{1}{2} \left( \frac{1}{v-1} - \frac{1}{v+1} \right) v' &= \frac{1}{x}.\end{aligned}$$

Integrating, we get

$$\frac{1}{2} (\ln|v-1| - \ln|v+1|) = \ln|x| + c_1.$$

## Example (Continued ...)

$$\frac{v-1}{v+1} = cx^2 \implies v = \frac{1+cx^2}{1-cx^2}.$$

## Example (Continued ...)

$$\frac{v-1}{v+1} = cx^2 \implies v = \frac{1+cx^2}{1-cx^2}.$$

Hence

$$y = x \frac{1+cx^2}{1-cx^2}.$$

**Question.** Is this  $y$  solution to given ODE? Under what conditions,  $y$  will be solution to given ODE?



## Example (Continued ...)

$$\frac{v-1}{v+1} = cx^2 \implies v = \frac{1+cx^2}{1-cx^2}.$$

Hence

$$y = x \frac{1+cx^2}{1-cx^2}.$$

**Question.** Is this  $y$  solution to given ODE? Under what conditions,  $y$  will be solution to given ODE?

**Question** Are these all the solutions?

## Example (Continued ...)

$$\frac{v-1}{v+1} = cx^2 \implies v = \frac{1+cx^2}{1-cx^2}.$$

Hence

$$y = x \frac{1+cx^2}{1-cx^2}.$$

**Question.** Is this  $y$  solution to given ODE? Under what conditions,  $y$  will be solution to given ODE?

**Question** Are these all the solutions?

**Ans** No.

## Example (Continued ...)

$$\frac{v-1}{v+1} = cx^2 \implies v = \frac{1+cx^2}{1-cx^2}.$$

Hence

$$y = x \frac{1+cx^2}{1-cx^2}.$$

**Question.** Is this  $y$  solution to given ODE? Under what conditions,  $y$  will be solution to given ODE?

**Question** Are these all the solutions?

**Ans** No.

Both  $y = x$  and  $y = -x$  are also solutions, but only  $y = x$  can be obtained by choosing a particular value of  $c = 0$ .

## Converting to Separable Equation

**Example** The following ODE

$$y' = \frac{2x + y + 1}{x + 2y - 4}$$

can be converted to a separable equation.

## Converting to Separable Equation

**Example** The following ODE

$$y' = \frac{2x + y + 1}{x + 2y - 4}$$

can be converted to a separable equation.

Use substitution  $X = x + 2$ , and  $Y = y - 3$ .

## Example

Consider the ODE

$$y' + ay = 0, \quad a \in \mathbb{R}.$$

## Example

Consider the ODE

$$y' + ay = 0, \quad a \in \mathbb{R}.$$

We can try to solve this differential equation as follows. Rewrite it as

$$\frac{y'}{y} = -a$$

Thus, we get

$$\ln |y| = -ax + c$$

that is,

$$y = e^c e^{-ax}$$

## Example

Consider the ODE

$$y' + ay = 0, \quad a \in \mathbb{R}.$$

We can try to solve this differential equation as follows. Rewrite it as

$$\frac{y'}{y} = -a$$

Thus, we get

$$\ln |y| = -ax + c$$

that is,

$$y = e^c e^{-ax}$$

**Question.** Is this  $y$  an explicit solution of ODE  $y' + ay = 0$ ?