

# Lectures 6 and 7

## Linear Transformation and Rank-Nullity Theorem

Dipankar Ghosh

Department of Mathematics  
Indian Institute of Technology Hyderabad

January 21, 2020

# Linear Transformations

A 'Linear Transformation' is nothing but a map between vector spaces.

# Linear Transformations

A 'Linear Transformation' is nothing but a map between vector spaces. Let us start with some well known maps:

# Linear Transformations

A 'Linear Transformation' is nothing but a map between vector spaces. Let us start with some well known maps:

$$T : \mathbb{R}^2 \longrightarrow \mathbb{R}^2$$

# Linear Transformations

A 'Linear Transformation' is nothing but a map between vector spaces. Let us start with some well known maps:

$$T : \mathbb{R}^2 \longrightarrow \mathbb{R}^2$$

$$\begin{pmatrix} x \\ y \end{pmatrix} \mapsto c \begin{pmatrix} x \\ y \end{pmatrix}$$

# Linear Transformations

A 'Linear Transformation' is nothing but a map between vector spaces. Let us start with some well known maps:

$$T : \mathbb{R}^2 \longrightarrow \mathbb{R}^2$$


$$\begin{pmatrix} x \\ y \end{pmatrix} \mapsto c \begin{pmatrix} x \\ y \end{pmatrix} \quad \text{where } c \in \mathbb{R}$$

# Linear Transformations

A 'Linear Transformation' is nothing but a map between vector spaces. Let us start with some well known maps:

$$T : \mathbb{R}^2 \longrightarrow \mathbb{R}^2$$

$$\begin{pmatrix} x \\ y \end{pmatrix} \mapsto c \begin{pmatrix} x \\ y \end{pmatrix} \quad \text{where } c \in \mathbb{R}$$




# Linear Transformations

A 'Linear Transformation' is nothing but a map between vector spaces. Let us start with some well known maps:

$$T : \mathbb{R}^2 \longrightarrow \mathbb{R}^2$$

$$\begin{pmatrix} x \\ y \end{pmatrix} \mapsto c \begin{pmatrix} x \\ y \end{pmatrix} \quad \text{where } c \in \mathbb{R}$$






# Linear Transformations

A 'Linear Transformation' is nothing but a map between vector spaces. Let us start with some well known maps:

$$T : \mathbb{R}^2 \longrightarrow \mathbb{R}^2$$

$$\begin{pmatrix} x \\ y \end{pmatrix} \mapsto c \begin{pmatrix} x \\ y \end{pmatrix} \quad \text{where } c \in \mathbb{R}$$




# Linear Transformations

A 'Linear Transformation' is nothing but a map between vector spaces. Let us start with some well known maps:

$$T : \mathbb{R}^2 \longrightarrow \mathbb{R}^2$$

$$\begin{pmatrix} x \\ y \end{pmatrix} \mapsto c \begin{pmatrix} x \\ y \end{pmatrix} \quad \text{where } c \in \mathbb{R}$$



# Linear Transformations


A 'Linear Transformation' is nothing but a map between vector spaces. Let us start with some well known maps:

$$T : \mathbb{R}^2 \longrightarrow \mathbb{R}^2$$


$$\begin{pmatrix} x \\ y \end{pmatrix} \mapsto c \begin{pmatrix} x \\ y \end{pmatrix} \quad \text{where } c \in \mathbb{R}$$

$$\begin{pmatrix} c & 0 \\ 0 & c \end{pmatrix} \begin{pmatrix} x \\ y \end{pmatrix} = \begin{pmatrix} cx \\ cy \end{pmatrix}$$


$$c \cdot (x, y) = (cx, cy)$$




# Reflection with $x$ -axis as mirror




# Reflection with $x$ -axis as mirror



# Reflection with $x$ -axis as mirror




# Reflection with $x$ -axis as mirror



# Reflection with $x$ -axis as mirror

$$T : \mathbb{R}^2 \longrightarrow \mathbb{R}^2$$






# Reflection with $x$ -axis as mirror

$$T : \mathbb{R}^2 \longrightarrow \mathbb{R}^2$$

$$\begin{pmatrix} x \\ y \end{pmatrix} \mapsto \begin{pmatrix} x \\ -y \end{pmatrix}$$




# Reflection with $x$ -axis as mirror


$$T : \mathbb{R}^2 \longrightarrow \mathbb{R}^2$$

$$\begin{pmatrix} x \\ y \end{pmatrix} \mapsto \begin{pmatrix} x \\ -y \end{pmatrix}$$


$$\begin{pmatrix} 1 & 0 \\ 0 & -1 \end{pmatrix} \begin{pmatrix} x \\ y \end{pmatrix} = \begin{pmatrix} x \\ -y \end{pmatrix}$$




# Projection on the $x$ -axis



# Projection on the $x$ -axis




# Projection on the $x$ -axis



projection on  $x$ -axis


# Projection on the $x$ -axis



projection on  $x$ -axis

# Projection on the $x$ -axis

$$T : \mathbb{R}^2 \longrightarrow \mathbb{R}^2$$




projection on  $x$ -axis

# Projection on the $x$ -axis

$$T : \mathbb{R}^2 \longrightarrow \mathbb{R}^2$$

$$\begin{pmatrix} x \\ y \end{pmatrix} \mapsto \begin{pmatrix} x \\ 0 \end{pmatrix}$$



projection on  $x$ -axis




# Projection on the $x$ -axis

$$T : \mathbb{R}^2 \longrightarrow \mathbb{R}^2$$

$$\begin{pmatrix} x \\ y \end{pmatrix} \mapsto \begin{pmatrix} x \\ 0 \end{pmatrix}$$

$$\begin{pmatrix} 1 & 0 \\ 0 & 0 \end{pmatrix} \begin{pmatrix} x \\ y \end{pmatrix} = \begin{pmatrix} x \\ 0 \end{pmatrix}$$



projection on  $x$ -axis


# Projection on the $x$ -axis

$$S : \mathbb{R}^2 \longrightarrow \mathbb{R}$$

$$T : \mathbb{R}^2 \longrightarrow \mathbb{R}^2$$

$$\begin{pmatrix} x \\ y \end{pmatrix} \mapsto \begin{pmatrix} x \\ 0 \end{pmatrix}$$

$$\begin{pmatrix} 1 & 0 \\ 0 & 0 \end{pmatrix} \begin{pmatrix} x \\ y \end{pmatrix} = \begin{pmatrix} x \\ 0 \end{pmatrix}$$



projection on  $x$ -axis

# Projection on the $x$ -axis


$$S : \mathbb{R}^2 \longrightarrow \mathbb{R}$$

$$\begin{pmatrix} x \\ y \end{pmatrix} \mapsto x$$

$$T : \mathbb{R}^2 \longrightarrow \mathbb{R}^2$$

$$\begin{pmatrix} x \\ y \end{pmatrix} \mapsto \begin{pmatrix} x \\ 0 \end{pmatrix}$$

$$\begin{pmatrix} 1 & 0 \\ 0 & 0 \end{pmatrix} \begin{pmatrix} x \\ y \end{pmatrix} = \begin{pmatrix} x \\ 0 \end{pmatrix}$$



projection on  $x$ -axis

# Projection on the $x$ -axis

$$S : \mathbb{R}^2 \longrightarrow \mathbb{R}$$


$$\begin{pmatrix} x \\ y \end{pmatrix} \mapsto x$$

$$\begin{bmatrix} 1 & 0 \end{bmatrix} \begin{pmatrix} x \\ y \end{pmatrix} = x$$

$$T : \mathbb{R}^2 \longrightarrow \mathbb{R}^2$$

$$\begin{pmatrix} x \\ y \end{pmatrix} \mapsto \begin{pmatrix} x \\ 0 \end{pmatrix}$$

$$\begin{pmatrix} 1 & 0 \\ 0 & 0 \end{pmatrix} \begin{pmatrix} x \\ y \end{pmatrix} = \begin{pmatrix} x \\ 0 \end{pmatrix}$$



projection on  $x$ -axis

# Rotation in Euclidean plane by an angle $\theta$

Consider the map  $T : \mathbb{R}^2 \rightarrow \mathbb{R}^2$  which performs a rotation in the  $xy$ -plane counterclockwise by an angle  $\theta$  about the origin.

# Rotation in Euclidean plane by an angle $\theta$


Consider the map  $T : \mathbb{R}^2 \rightarrow \mathbb{R}^2$  which performs a rotation in the  $xy$ -plane counterclockwise by an angle  $\theta$  about the origin.

$$T \begin{pmatrix} x \\ y \end{pmatrix} = \begin{pmatrix} \cos \theta & -\sin \theta \\ \sin \theta & \cos \theta \end{pmatrix} \begin{pmatrix} x \\ y \end{pmatrix} = \begin{pmatrix} x \cos \theta - y \sin \theta \\ x \sin \theta + y \cos \theta \end{pmatrix}.$$

# Rotation in Euclidean plane by an angle $\theta$

Consider the map  $T : \mathbb{R}^2 \rightarrow \mathbb{R}^2$  which performs a rotation in the  $xy$ -plane counterclockwise by an angle  $\theta$  about the origin.


$$T \begin{pmatrix} x \\ y \end{pmatrix} = \begin{pmatrix} \cos \theta & -\sin \theta \\ \sin \theta & \cos \theta \end{pmatrix} \begin{pmatrix} x \\ y \end{pmatrix} = \begin{pmatrix} x \cos \theta - y \sin \theta \\ x \sin \theta + y \cos \theta \end{pmatrix}.$$



# Rotation in Euclidean plane by an angle $\theta$

Consider the map  $T : \mathbb{R}^2 \rightarrow \mathbb{R}^2$  which performs a rotation in the  $xy$ -plane counterclockwise by an angle  $\theta$  about the origin.

$$T \begin{pmatrix} x \\ y \end{pmatrix} = \begin{pmatrix} \cos \theta & -\sin \theta \\ \sin \theta & \cos \theta \end{pmatrix} \begin{pmatrix} x \\ y \end{pmatrix} = \begin{pmatrix} x \cos \theta - y \sin \theta \\ x \sin \theta + y \cos \theta \end{pmatrix}.$$






# Rotation in Euclidean plane by an angle $\theta$

Consider the map  $T : \mathbb{R}^2 \rightarrow \mathbb{R}^2$  which performs a rotation in the  $xy$ -plane counterclockwise by an angle  $\theta$  about the origin.


$$T \begin{pmatrix} x \\ y \end{pmatrix} = \begin{pmatrix} \cos \theta & -\sin \theta \\ \sin \theta & \cos \theta \end{pmatrix} \begin{pmatrix} x \\ y \end{pmatrix} = \begin{pmatrix} x \cos \theta - y \sin \theta \\ x \sin \theta + y \cos \theta \end{pmatrix}.$$



# Rotation in Euclidean plane by an angle $\theta$

Consider the map  $T : \mathbb{R}^2 \rightarrow \mathbb{R}^2$  which performs a rotation in the  $xy$ -plane counterclockwise by an angle  $\theta$  about the origin.

$$T \begin{pmatrix} x \\ y \end{pmatrix} = \begin{pmatrix} \cos \theta & -\sin \theta \\ \sin \theta & \cos \theta \end{pmatrix} \begin{pmatrix} x \\ y \end{pmatrix} = \begin{pmatrix} x \cos \theta - y \sin \theta \\ x \sin \theta + y \cos \theta \end{pmatrix}.$$



# Linear transformation, or linear map

## Definition

A map between vector spaces which satisfies the rule of linearity is called linear transformation

# Linear transformation, or linear map

## Definition

A map between vector spaces which satisfies the rule of linearity is called linear transformation (or linear map).

# Linear transformation, or linear map

## Definition

A map between vector spaces which satisfies the rule of linearity is called linear transformation (or linear map).

More precisely, let  $V$  and  $W$  be vector spaces over  $\mathbb{R}$ .

# Linear transformation, or linear map

## Definition

A map between vector spaces which satisfies the rule of linearity is called linear transformation (or linear map).

More precisely, let  $V$  and  $W$  be vector spaces over  $\mathbb{R}$ . A linear transformation  $T : V \rightarrow W$  is a function

# Linear transformation, or linear map

## Definition

A map between vector spaces which satisfies the rule of linearity is called linear transformation (or linear map).

More precisely, let  $V$  and  $W$  be vector spaces over  $\mathbb{R}$ . A linear transformation  $T : V \rightarrow W$  is a function such that

$$T(c_1v_1 + c_2v_2) = c_1T(v_1) + c_2T(v_2)$$

for all  $c_1, c_2 \in \mathbb{R}$  and  $v_1, v_2 \in V$ .

# Linear transformation, or linear map

## Definition

A map between vector spaces which satisfies the rule of linearity is called linear transformation (or linear map).

More precisely, let  $V$  and  $W$  be vector spaces over  $\mathbb{R}$ . A linear transformation  $T : V \rightarrow W$  is a function such that

$$T(c_1v_1 + c_2v_2) = c_1T(v_1) + c_2T(v_2)$$

for all  $c_1, c_2 \in \mathbb{R}$  and  $v_1, v_2 \in V$ . Equivalently, a linear map is a map which respects both vector addition and scalar multiplication.



# Linear transformation, or linear map

## Definition

A map between vector spaces which satisfies the rule of linearity is called linear transformation (or linear map).

More precisely, let  $V$  and  $W$  be vector spaces over  $\mathbb{R}$ . A linear transformation  $T : V \rightarrow W$  is a function such that

$$T(c_1v_1 + c_2v_2) = c_1T(v_1) + c_2T(v_2)$$

for all  $c_1, c_2 \in \mathbb{R}$  and  $v_1, v_2 \in V$ . Equivalently, a linear map is a map which respects both vector addition and scalar multiplication.

## Example (A matrix can be thought of as a linear map)

Starting with an  $m \times n$  matrix  $A$  over  $\mathbb{R}$ ,

# Linear transformation, or linear map

## Definition

A map between vector spaces which satisfies the rule of linearity is called linear transformation (or linear map).

More precisely, let  $V$  and  $W$  be vector spaces over  $\mathbb{R}$ . A linear transformation  $T : V \rightarrow W$  is a function such that

$$T(c_1v_1 + c_2v_2) = c_1T(v_1) + c_2T(v_2)$$

for all  $c_1, c_2 \in \mathbb{R}$  and  $v_1, v_2 \in V$ . Equivalently, a linear map is a map which respects both vector addition and scalar multiplication.

## Example (A matrix can be thought of as a linear map)

Starting with an  $m \times n$  matrix  $A$  over  $\mathbb{R}$ , one can construct a linear map  $T_A : \mathbb{R}^n \rightarrow \mathbb{R}^m$

# Linear transformation, or linear map

## Definition

A map between vector spaces which satisfies the rule of linearity is called linear transformation (or linear map).

More precisely, let  $V$  and  $W$  be vector spaces over  $\mathbb{R}$ . A linear transformation  $T : V \rightarrow W$  is a function such that

$$T(c_1v_1 + c_2v_2) = c_1T(v_1) + c_2T(v_2)$$

for all  $c_1, c_2 \in \mathbb{R}$  and  $v_1, v_2 \in V$ . Equivalently, a linear map is a map which respects both vector addition and scalar multiplication.

## Example (A matrix can be thought of as a linear map)

Starting with an  $m \times n$  matrix  $A$  over  $\mathbb{R}$ , one can construct a linear map  $T_A : \mathbb{R}^n \rightarrow \mathbb{R}^m$  defined by  $T_A(X) := AX$  for all  $X \in \mathbb{R}^n$ .

# Linear transformation, or linear map

## Definition

A map between vector spaces which satisfies the rule of linearity is called linear transformation (or linear map).

More precisely, let  $V$  and  $W$  be vector spaces over  $\mathbb{R}$ . A linear transformation  $T : V \rightarrow W$  is a function such that

$$T(c_1v_1 + c_2v_2) = c_1T(v_1) + c_2T(v_2)$$

for all  $c_1, c_2 \in \mathbb{R}$  and  $v_1, v_2 \in V$ . Equivalently, a linear map is a map which respects both vector addition and scalar multiplication.

## Example (A matrix can be thought of as a linear map)

Starting with an  $m \times n$  matrix  $A$  over  $\mathbb{R}$ , one can construct a linear map  $T_A : \mathbb{R}^n \rightarrow \mathbb{R}^m$  defined by  $T_A(X) := AX$  for all  $X \in \mathbb{R}^n$ .

*Proof.*  $T_A(X + Y)$

# Linear transformation, or linear map

## Definition

A map between vector spaces which satisfies the rule of linearity is called linear transformation (or linear map).

More precisely, let  $V$  and  $W$  be vector spaces over  $\mathbb{R}$ . A linear transformation  $T : V \rightarrow W$  is a function such that

$$T(c_1v_1 + c_2v_2) = c_1T(v_1) + c_2T(v_2)$$

for all  $c_1, c_2 \in \mathbb{R}$  and  $v_1, v_2 \in V$ . Equivalently, a linear map is a map which respects both vector addition and scalar multiplication.

## Example (A matrix can be thought of as a linear map)

Starting with an  $m \times n$  matrix  $A$  over  $\mathbb{R}$ , one can construct a linear map  $T_A : \mathbb{R}^n \rightarrow \mathbb{R}^m$  defined by  $T_A(X) := AX$  for all  $X \in \mathbb{R}^n$ .

*Proof.*  $T_A(X + Y) = A(X + Y)$

# Linear transformation, or linear map

## Definition

A map between vector spaces which satisfies the rule of linearity is called linear transformation (or linear map).

More precisely, let  $V$  and  $W$  be vector spaces over  $\mathbb{R}$ . A linear transformation  $T : V \rightarrow W$  is a function such that

$$T(c_1v_1 + c_2v_2) = c_1T(v_1) + c_2T(v_2)$$

for all  $c_1, c_2 \in \mathbb{R}$  and  $v_1, v_2 \in V$ . Equivalently, a linear map is a map which respects both vector addition and scalar multiplication.

## Example (A matrix can be thought of as a linear map)

Starting with an  $m \times n$  matrix  $A$  over  $\mathbb{R}$ , one can construct a linear map  $T_A : \mathbb{R}^n \rightarrow \mathbb{R}^m$  defined by  $T_A(X) := AX$  for all  $X \in \mathbb{R}^n$ .

*Proof.*  $T_A(X + Y) = A(X + Y) = AX + AY$

# Linear transformation, or linear map

## Definition

A map between vector spaces which satisfies the rule of linearity is called linear transformation (or linear map).

More precisely, let  $V$  and  $W$  be vector spaces over  $\mathbb{R}$ . A linear transformation  $T : V \rightarrow W$  is a function such that

$$T(c_1v_1 + c_2v_2) = c_1T(v_1) + c_2T(v_2)$$

for all  $c_1, c_2 \in \mathbb{R}$  and  $v_1, v_2 \in V$ . Equivalently, a linear map is a map which respects both vector addition and scalar multiplication.

## Example (A matrix can be thought of as a linear map)

Starting with an  $m \times n$  matrix  $A$  over  $\mathbb{R}$ , one can construct a linear map  $T_A : \mathbb{R}^n \rightarrow \mathbb{R}^m$  defined by  $T_A(X) := AX$  for all  $X \in \mathbb{R}^n$ .

*Proof.*  $T_A(X + Y) = A(X + Y) = AX + AY = T_A(X) + T_A(Y)$

# Linear transformation, or linear map

## Definition

A map between vector spaces which satisfies the rule of linearity is called linear transformation (or linear map).

More precisely, let  $V$  and  $W$  be vector spaces over  $\mathbb{R}$ . A linear transformation  $T : V \rightarrow W$  is a function such that

$$T(c_1v_1 + c_2v_2) = c_1T(v_1) + c_2T(v_2)$$

for all  $c_1, c_2 \in \mathbb{R}$  and  $v_1, v_2 \in V$ . Equivalently, a linear map is a map which respects both vector addition and scalar multiplication.

## Example (A matrix can be thought of as a linear map)

Starting with an  $m \times n$  matrix  $A$  over  $\mathbb{R}$ , one can construct a linear map  $T_A : \mathbb{R}^n \rightarrow \mathbb{R}^m$  defined by  $T_A(X) := AX$  for all  $X \in \mathbb{R}^n$ .

*Proof.*  $T_A(X + Y) = A(X + Y) = AX + AY = T_A(X) + T_A(Y)$  and  $T_A(cX)$



# Linear transformation, or linear map

## Definition

A map between vector spaces which satisfies the rule of linearity is called linear transformation (or linear map).

More precisely, let  $V$  and  $W$  be vector spaces over  $\mathbb{R}$ . A linear transformation  $T : V \rightarrow W$  is a function such that

$$T(c_1v_1 + c_2v_2) = c_1T(v_1) + c_2T(v_2)$$

for all  $c_1, c_2 \in \mathbb{R}$  and  $v_1, v_2 \in V$ . Equivalently, a linear map is a map which respects both vector addition and scalar multiplication.

## Example (A matrix can be thought of as a linear map)

Starting with an  $m \times n$  matrix  $A$  over  $\mathbb{R}$ , one can construct a linear map  $T_A : \mathbb{R}^n \rightarrow \mathbb{R}^m$  defined by  $T_A(X) := AX$  for all  $X \in \mathbb{R}^n$ .

*Proof.*  $T_A(X + Y) = A(X + Y) = AX + AY = T_A(X) + T_A(Y)$  and  $T_A(cX) = A(cX)$

# Linear transformation, or linear map

## Definition

A map between vector spaces which satisfies the rule of linearity is called linear transformation (or linear map).

More precisely, let  $V$  and  $W$  be vector spaces over  $\mathbb{R}$ . A linear transformation  $T : V \rightarrow W$  is a function such that

$$T(c_1v_1 + c_2v_2) = c_1T(v_1) + c_2T(v_2)$$

for all  $c_1, c_2 \in \mathbb{R}$  and  $v_1, v_2 \in V$ . Equivalently, a linear map is a map which respects both vector addition and scalar multiplication.

## Example (A matrix can be thought of as a linear map)

Starting with an  $m \times n$  matrix  $A$  over  $\mathbb{R}$ , one can construct a linear map  $T_A : \mathbb{R}^n \rightarrow \mathbb{R}^m$  defined by  $T_A(X) := AX$  for all  $X \in \mathbb{R}^n$ .

*Proof.*  $T_A(X + Y) = A(X + Y) = AX + AY = T_A(X) + T_A(Y)$  and  $T_A(cX) = A(cX) = c(AX) = cT_A(X)$

# Linear transformation, or linear map

## Definition

A map between vector spaces which satisfies the rule of linearity is called linear transformation (or linear map).

More precisely, let  $V$  and  $W$  be vector spaces over  $\mathbb{R}$ . A linear transformation  $T : V \rightarrow W$  is a function such that

$$T(c_1v_1 + c_2v_2) = c_1T(v_1) + c_2T(v_2)$$

for all  $c_1, c_2 \in \mathbb{R}$  and  $v_1, v_2 \in V$ . Equivalently, a linear map is a map which respects both vector addition and scalar multiplication.

## Example (A matrix can be thought of as a linear map)

Starting with an  $m \times n$  matrix  $A$  over  $\mathbb{R}$ , one can construct a linear map  $T_A : \mathbb{R}^n \rightarrow \mathbb{R}^m$  defined by  $T_A(X) := AX$  for all  $X \in \mathbb{R}^n$ .

*Proof.*  $T_A(X + Y) = A(X + Y) = AX + AY = T_A(X) + T_A(Y)$  and  $T_A(cX) = A(cX) = c(AX) = cT_A(X)$ .

# Differentiation and integration transformation

## Example (Differentiation transformation)

Let  $V = \mathbb{R}[x]$ ,

# Differentiation and integration transformation

## Example (Differentiation transformation)

Let  $V = \mathbb{R}[x]$ , the set of all polynomials in  $x$  over  $\mathbb{R}$ .

# Differentiation and integration transformation

## Example (Differentiation transformation)

Let  $V = \mathbb{R}[x]$ , the set of all polynomials in  $x$  over  $\mathbb{R}$ . Define a map  $D : V \rightarrow V$  as follows:

# Differentiation and integration transformation

## Example (Differentiation transformation)

Let  $V = \mathbb{R}[x]$ , the set of all polynomials in  $x$  over  $\mathbb{R}$ . Define a map  $D : V \rightarrow V$  as follows: If  $f = a_0 + a_1x + a_2x^2 + \cdots + a_rx^r$ ,

# Differentiation and integration transformation

## Example (Differentiation transformation)

Let  $V = \mathbb{R}[x]$ , the set of all polynomials in  $x$  over  $\mathbb{R}$ . Define a map  $D : V \rightarrow V$  as follows: If  $f = a_0 + a_1x + a_2x^2 + \cdots + a_rx^r$ , then

$$D(f) := a_1 + 2a_2x + \cdots + ra_rx^{r-1}.$$



# Differentiation and integration transformation

## Example (Differentiation transformation)

Let  $V = \mathbb{R}[x]$ , the set of all polynomials in  $x$  over  $\mathbb{R}$ . Define a map  $D : V \rightarrow V$  as follows: If  $f = a_0 + a_1x + a_2x^2 + \cdots + a_rx^r$ , then

$$D(f) := a_1 + 2a_2x + \cdots + ra_rx^{r-1}.$$

Then  $D$  is a linear transformation.

# Differentiation and integration transformation

## Example (Differentiation transformation)

Let  $V = \mathbb{R}[x]$ , the set of all polynomials in  $x$  over  $\mathbb{R}$ . Define a map  $D : V \rightarrow V$  as follows: If  $f = a_0 + a_1x + a_2x^2 + \cdots + a_rx^r$ , then

$$D(f) := a_1 + 2a_2x + \cdots + ra_rx^{r-1}.$$

Then  $D$  is a linear transformation.

## Example (Integration transformation)

Let  $V$  be the set of all continuous functions from  $\mathbb{R}$  into  $\mathbb{R}$ .

# Differentiation and integration transformation

## Example (Differentiation transformation)

Let  $V = \mathbb{R}[x]$ , the set of all polynomials in  $x$  over  $\mathbb{R}$ . Define a map  $D : V \rightarrow V$  as follows: If  $f = a_0 + a_1x + a_2x^2 + \cdots + a_rx^r$ , then

$$D(f) := a_1 + 2a_2x + \cdots + ra_rx^{r-1}.$$

Then  $D$  is a linear transformation.

## Example (Integration transformation)

Let  $V$  be the set of all continuous functions from  $\mathbb{R}$  into  $\mathbb{R}$ . Define a map  $T : V \rightarrow V$  as follows:

# Differentiation and integration transformation

## Example (Differentiation transformation)

Let  $V = \mathbb{R}[x]$ , the set of all polynomials in  $x$  over  $\mathbb{R}$ . Define a map  $D : V \rightarrow V$  as follows: If  $f = a_0 + a_1x + a_2x^2 + \cdots + a_rx^r$ , then

$$D(f) := a_1 + 2a_2x + \cdots + ra_rx^{r-1}.$$

Then  $D$  is a linear transformation.

## Example (Integration transformation)

Let  $V$  be the set of all continuous functions from  $\mathbb{R}$  into  $\mathbb{R}$ . Define a map  $T : V \rightarrow V$  as follows: If  $f \in V$ ,

# Differentiation and integration transformation

## Example (Differentiation transformation)

Let  $V = \mathbb{R}[x]$ , the set of all polynomials in  $x$  over  $\mathbb{R}$ . Define a map  $D : V \rightarrow V$  as follows: If  $f = a_0 + a_1x + a_2x^2 + \cdots + a_rx^r$ , then

$$D(f) := a_1 + 2a_2x + \cdots + ra_rx^{r-1}.$$

Then  $D$  is a linear transformation.

## Example (Integration transformation)

Let  $V$  be the set of all continuous functions from  $\mathbb{R}$  into  $\mathbb{R}$ . Define a map  $T : V \rightarrow V$  as follows: If  $f \in V$ , then  $T(f)$  is given by

$$T(f)(x) = \int_0^x f(t)dt \quad \text{for all } x \in \mathbb{R}.$$

# Differentiation and integration transformation

## Example (Differentiation transformation)

Let  $V = \mathbb{R}[x]$ , the set of all polynomials in  $x$  over  $\mathbb{R}$ . Define a map  $D : V \rightarrow V$  as follows: If  $f = a_0 + a_1x + a_2x^2 + \cdots + a_rx^r$ , then

$$D(f) := a_1 + 2a_2x + \cdots + ra_rx^{r-1}.$$

Then  $D$  is a linear transformation.

## Example (Integration transformation)

Let  $V$  be the set of all continuous functions from  $\mathbb{R}$  into  $\mathbb{R}$ . Define a map  $T : V \rightarrow V$  as follows: If  $f \in V$ , then  $T(f)$  is given by

$$T(f)(x) = \int_0^x f(t)dt \quad \text{for all } x \in \mathbb{R}.$$

Then  $T$  is a linear transformation.

# What is $T(0)$ ?

- Let  $T : V \rightarrow W$  be a linear transformation.

# What is $T(0)$ ?

- Let  $T : V \rightarrow W$  be a linear transformation.
- What is  $T(0)$ ?



# What is $T(0)$ ?

- Let  $T : V \rightarrow W$  be a linear transformation.
- What is  $T(0)$ ?
- Answer:  $T(0) = 0$ ,

# What is $T(0)$ ?

- Let  $T : V \rightarrow W$  be a linear transformation.
- What is  $T(0)$ ?
- Answer:  $T(0) = 0$ , because  $T(0) = T(0 + 0) = T(0) + T(0)$ .

# An observation on matrix multiplication

- Multiplying a matrix  $A$  with a column vector  $b$  yields a linear combination of the columns of  $A$ .

# An observation on matrix multiplication

- Multiplying a matrix  $A$  with a column vector  $b$  yields a linear combination of the columns of  $A$ .

- $$\begin{pmatrix} 1 & 2 & 3 & 4 \\ 5 & 6 & 7 & 8 \\ 9 & 10 & 11 & 12 \end{pmatrix} \begin{pmatrix} x_1 \\ x_2 \\ x_3 \\ x_4 \end{pmatrix} =$$

$$x_1 \begin{pmatrix} 1 \\ 5 \\ 9 \end{pmatrix} + x_2 \begin{pmatrix} 2 \\ 6 \\ 10 \end{pmatrix} + x_3 \begin{pmatrix} 3 \\ 7 \\ 11 \end{pmatrix} + x_4 \begin{pmatrix} 4 \\ 8 \\ 12 \end{pmatrix}.$$

# An observation on matrix multiplication

- Multiplying a matrix  $A$  with a column vector  $b$  yields a linear combination of the columns of  $A$ .

- $$\begin{pmatrix} 1 & 2 & 3 & 4 \\ 5 & 6 & 7 & 8 \\ 9 & 10 & 11 & 12 \end{pmatrix} \begin{pmatrix} x_1 \\ x_2 \\ x_3 \\ x_4 \end{pmatrix} =$$

$$x_1 \begin{pmatrix} 1 \\ 5 \\ 9 \end{pmatrix} + x_2 \begin{pmatrix} 2 \\ 6 \\ 10 \end{pmatrix} + x_3 \begin{pmatrix} 3 \\ 7 \\ 11 \end{pmatrix} + x_4 \begin{pmatrix} 4 \\ 8 \\ 12 \end{pmatrix}.$$

- $$\begin{bmatrix} C_1 & C_2 & \cdots & C_n \end{bmatrix}_{m \times n} \begin{pmatrix} x_1 \\ x_2 \\ \vdots \\ x_n \end{pmatrix} = x_1(C_1) + x_2(C_2) + \cdots + x_n(C_n),$$

# An observation on matrix multiplication

- Multiplying a matrix  $A$  with a column vector  $b$  yields a linear combination of the columns of  $A$ .

- $$\begin{pmatrix} 1 & 2 & 3 & 4 \\ 5 & 6 & 7 & 8 \\ 9 & 10 & 11 & 12 \end{pmatrix} \begin{pmatrix} x_1 \\ x_2 \\ x_3 \\ x_4 \end{pmatrix} =$$

$$x_1 \begin{pmatrix} 1 \\ 5 \\ 9 \end{pmatrix} + x_2 \begin{pmatrix} 2 \\ 6 \\ 10 \end{pmatrix} + x_3 \begin{pmatrix} 3 \\ 7 \\ 11 \end{pmatrix} + x_4 \begin{pmatrix} 4 \\ 8 \\ 12 \end{pmatrix}.$$

- $$[C_1 \quad C_2 \quad \cdots \quad C_n]_{m \times n} \begin{pmatrix} x_1 \\ x_2 \\ \vdots \\ x_n \end{pmatrix} = x_1(C_1) + x_2(C_2) + \cdots + x_n(C_n),$$

where  $C_1, \dots, C_n \in \mathbb{R}^m$ .

# Matrix representation of a linear map $T : \mathbb{R}^n \rightarrow \mathbb{R}^m$

## Theorem

*Let  $T : \mathbb{R}^n \rightarrow \mathbb{R}^m$  be a linear transformation.*

# Matrix representation of a linear map $T : \mathbb{R}^n \rightarrow \mathbb{R}^m$

## Theorem

*Let  $T : \mathbb{R}^n \rightarrow \mathbb{R}^m$  be a linear transformation. Then there exists an  $m \times n$  matrix  $A$*



# Matrix representation of a linear map $T : \mathbb{R}^n \rightarrow \mathbb{R}^m$

## Theorem

*Let  $T : \mathbb{R}^n \rightarrow \mathbb{R}^m$  be a linear transformation. Then there exists an  $m \times n$  matrix  $A$  such that  $T$  can be represented by  $A$ ,*

# Matrix representation of a linear map $T : \mathbb{R}^n \rightarrow \mathbb{R}^m$

## Theorem

*Let  $T : \mathbb{R}^n \rightarrow \mathbb{R}^m$  be a linear transformation. Then there exists an  $m \times n$  matrix  $A$  such that  $T$  can be represented by  $A$ , i.e.,  $T(X) = AX$  for every  $X \in \mathbb{R}^n$ .*

# Matrix representation of a linear map $T : \mathbb{R}^n \rightarrow \mathbb{R}^m$

## Theorem

*Let  $T : \mathbb{R}^n \rightarrow \mathbb{R}^m$  be a linear transformation. Then there exists an  $m \times n$  matrix  $A$  such that  $T$  can be represented by  $A$ , i.e.,  $T(X) = AX$  for every  $X \in \mathbb{R}^n$ .*

**Proof.** Let  $\{e_1, \dots, e_n\}$  be the standard basis of  $\mathbb{R}^n$ .

# Matrix representation of a linear map $T : \mathbb{R}^n \rightarrow \mathbb{R}^m$

## Theorem

*Let  $T : \mathbb{R}^n \rightarrow \mathbb{R}^m$  be a linear transformation. Then there exists an  $m \times n$  matrix  $A$  such that  $T$  can be represented by  $A$ , i.e.,  $T(X) = AX$  for every  $X \in \mathbb{R}^n$ .*

**Proof.** Let  $\{e_1, \dots, e_n\}$  be the standard basis of  $\mathbb{R}^n$ . Consider

$$X = \begin{pmatrix} x_1 \\ x_2 \\ \vdots \\ x_n \end{pmatrix} \in \mathbb{R}^n.$$

# Matrix representation of a linear map $T : \mathbb{R}^n \rightarrow \mathbb{R}^m$

## Theorem

*Let  $T : \mathbb{R}^n \rightarrow \mathbb{R}^m$  be a linear transformation. Then there exists an  $m \times n$  matrix  $A$  such that  $T$  can be represented by  $A$ , i.e.,  $T(X) = AX$  for every  $X \in \mathbb{R}^n$ .*

**Proof.** Let  $\{e_1, \dots, e_n\}$  be the standard basis of  $\mathbb{R}^n$ . Consider

$$X = \begin{pmatrix} x_1 \\ x_2 \\ \vdots \\ x_n \end{pmatrix} \in \mathbb{R}^n. \text{ Then } X = x_1 \begin{pmatrix} 1 \\ 0 \\ \vdots \\ 0 \end{pmatrix} + \dots + x_n \begin{pmatrix} 0 \\ \vdots \\ 0 \\ 1 \end{pmatrix}$$

# Matrix representation of a linear map $T : \mathbb{R}^n \rightarrow \mathbb{R}^m$

## Theorem

*Let  $T : \mathbb{R}^n \rightarrow \mathbb{R}^m$  be a linear transformation. Then there exists an  $m \times n$  matrix  $A$  such that  $T$  can be represented by  $A$ , i.e.,  $T(X) = AX$  for every  $X \in \mathbb{R}^n$ .*

**Proof.** Let  $\{e_1, \dots, e_n\}$  be the standard basis of  $\mathbb{R}^n$ . Consider

$$X = \begin{pmatrix} x_1 \\ x_2 \\ \vdots \\ x_n \end{pmatrix} \in \mathbb{R}^n. \text{ Then } X = x_1 \begin{pmatrix} 1 \\ 0 \\ \vdots \\ 0 \end{pmatrix} + \dots + x_n \begin{pmatrix} 0 \\ \vdots \\ 0 \\ 1 \end{pmatrix} = \sum_{i=1}^n x_i e_i.$$

# Matrix representation of a linear map $T : \mathbb{R}^n \rightarrow \mathbb{R}^m$

## Theorem

*Let  $T : \mathbb{R}^n \rightarrow \mathbb{R}^m$  be a linear transformation. Then there exists an  $m \times n$  matrix  $A$  such that  $T$  can be represented by  $A$ , i.e.,  $T(X) = AX$  for every  $X \in \mathbb{R}^n$ .*

**Proof.** Let  $\{e_1, \dots, e_n\}$  be the standard basis of  $\mathbb{R}^n$ . Consider

$$X = \begin{pmatrix} x_1 \\ x_2 \\ \vdots \\ x_n \end{pmatrix} \in \mathbb{R}^n. \text{ Then } X = x_1 \begin{pmatrix} 1 \\ 0 \\ \vdots \\ 0 \end{pmatrix} + \dots + x_n \begin{pmatrix} 0 \\ \vdots \\ 0 \\ 1 \end{pmatrix} = \sum_{i=1}^n x_i e_i.$$

Applying  $T$  on the above equalities,

# Matrix representation of a linear map $T : \mathbb{R}^n \rightarrow \mathbb{R}^m$

## Theorem

*Let  $T : \mathbb{R}^n \rightarrow \mathbb{R}^m$  be a linear transformation. Then there exists an  $m \times n$  matrix  $A$  such that  $T$  can be represented by  $A$ , i.e.,  $T(X) = AX$  for every  $X \in \mathbb{R}^n$ .*

**Proof.** Let  $\{e_1, \dots, e_n\}$  be the standard basis of  $\mathbb{R}^n$ . Consider

$$X = \begin{pmatrix} x_1 \\ x_2 \\ \vdots \\ x_n \end{pmatrix} \in \mathbb{R}^n. \text{ Then } X = x_1 \begin{pmatrix} 1 \\ 0 \\ \vdots \\ 0 \end{pmatrix} + \dots + x_n \begin{pmatrix} 0 \\ \vdots \\ 0 \\ 1 \end{pmatrix} = \sum_{i=1}^n x_i e_i.$$

Applying  $T$  on the above equalities, we have that  $T(X)$



# Matrix representation of a linear map $T : \mathbb{R}^n \rightarrow \mathbb{R}^m$

## Theorem

*Let  $T : \mathbb{R}^n \rightarrow \mathbb{R}^m$  be a linear transformation. Then there exists an  $m \times n$  matrix  $A$  such that  $T$  can be represented by  $A$ , i.e.,  $T(X) = AX$  for every  $X \in \mathbb{R}^n$ .*

**Proof.** Let  $\{e_1, \dots, e_n\}$  be the standard basis of  $\mathbb{R}^n$ . Consider

$$X = \begin{pmatrix} x_1 \\ x_2 \\ \vdots \\ x_n \end{pmatrix} \in \mathbb{R}^n. \text{ Then } X = x_1 \begin{pmatrix} 1 \\ 0 \\ \vdots \\ 0 \end{pmatrix} + \dots + x_n \begin{pmatrix} 0 \\ \vdots \\ 0 \\ 1 \end{pmatrix} = \sum_{i=1}^n x_i e_i.$$

Applying  $T$  on the above equalities, we have that

$$T(X) = x_1 T(e_1) + x_2 T(e_2) + \dots + x_n T(e_n)$$

# Matrix representation of a linear map $T : \mathbb{R}^n \rightarrow \mathbb{R}^m$

## Theorem

*Let  $T : \mathbb{R}^n \rightarrow \mathbb{R}^m$  be a linear transformation. Then there exists an  $m \times n$  matrix  $A$  such that  $T$  can be represented by  $A$ , i.e.,  $T(X) = AX$  for every  $X \in \mathbb{R}^n$ .*

**Proof.** Let  $\{e_1, \dots, e_n\}$  be the standard basis of  $\mathbb{R}^n$ . Consider

$$X = \begin{pmatrix} x_1 \\ x_2 \\ \vdots \\ x_n \end{pmatrix} \in \mathbb{R}^n. \text{ Then } X = x_1 \begin{pmatrix} 1 \\ 0 \\ \vdots \\ 0 \end{pmatrix} + \dots + x_n \begin{pmatrix} 0 \\ \vdots \\ 0 \\ 1 \end{pmatrix} = \sum_{i=1}^n x_i e_i.$$

Applying  $T$  on the above equalities, we have that

$$\begin{aligned} T(X) &= x_1 T(e_1) + x_2 T(e_2) + \dots + x_n T(e_n) \\ &= [T(e_1) \quad T(e_2) \quad \dots \quad T(e_n)] X \quad [\text{by the observation}]. \end{aligned}$$

# Matrix representation of a linear map $T : \mathbb{R}^n \rightarrow \mathbb{R}^m$

## Theorem

*Let  $T : \mathbb{R}^n \rightarrow \mathbb{R}^m$  be a linear transformation. Then there exists an  $m \times n$  matrix  $A$  such that  $T$  can be represented by  $A$ , i.e.,  $T(X) = AX$  for every  $X \in \mathbb{R}^n$ .*

**Proof.** Let  $\{e_1, \dots, e_n\}$  be the standard basis of  $\mathbb{R}^n$ . Consider

$$X = \begin{pmatrix} x_1 \\ x_2 \\ \vdots \\ x_n \end{pmatrix} \in \mathbb{R}^n. \text{ Then } X = x_1 \begin{pmatrix} 1 \\ 0 \\ \vdots \\ 0 \end{pmatrix} + \dots + x_n \begin{pmatrix} 0 \\ \vdots \\ 0 \\ 1 \end{pmatrix} = \sum_{i=1}^n x_i e_i.$$

Applying  $T$  on the above equalities, we have that

$$\begin{aligned} T(X) &= x_1 T(e_1) + x_2 T(e_2) + \dots + x_n T(e_n) \\ &= [T(e_1) \quad T(e_2) \quad \dots \quad T(e_n)] X \quad [\text{by the observation}]. \end{aligned}$$

The theorem follows by setting  $A := [T(e_1) \quad T(e_2) \quad \dots \quad T(e_n)]$ .

# Matrix representation of a linear map $T : \mathbb{R}^n \rightarrow \mathbb{R}^m$

## Theorem

*Let  $T : \mathbb{R}^n \rightarrow \mathbb{R}^m$  be a linear transformation. Then there exists an  $m \times n$  matrix  $A$  such that  $T$  can be represented by  $A$ , i.e.,  $T(X) = AX$  for every  $X \in \mathbb{R}^n$ .*

**Proof.** Let  $\{e_1, \dots, e_n\}$  be the standard basis of  $\mathbb{R}^n$ . Consider

$$X = \begin{pmatrix} x_1 \\ x_2 \\ \vdots \\ x_n \end{pmatrix} \in \mathbb{R}^n. \text{ Then } X = x_1 \begin{pmatrix} 1 \\ 0 \\ \vdots \\ 0 \end{pmatrix} + \dots + x_n \begin{pmatrix} 0 \\ \vdots \\ 0 \\ 1 \end{pmatrix} = \sum_{i=1}^n x_i e_i.$$

Applying  $T$  on the above equalities, we have that

$$\begin{aligned} T(X) &= x_1 T(e_1) + x_2 T(e_2) + \dots + x_n T(e_n) \\ &= [T(e_1) \quad T(e_2) \quad \dots \quad T(e_n)] X \quad [\text{by the observation}]. \end{aligned}$$

The theorem follows by setting  $A := [T(e_1) \quad T(e_2) \quad \dots \quad T(e_n)]$ .

**Remark.** A linear map  $T : \mathbb{R}^n \rightarrow \mathbb{R}^m$  is uniquely determined by its action on  $\{e_1, \dots, e_n\}$ , i.e., by  $T(e_i)$  for all  $1 \leq i \leq n$ .

# Correspondence between linear maps and matrices

## Corollary

*There is a one to one correspondence between the set of all linear transformations from  $\mathbb{R}^n$  to  $\mathbb{R}^m$*

# Correspondence between linear maps and matrices

## Corollary

*There is a one to one correspondence between the set of all linear transformations from  $\mathbb{R}^n$  to  $\mathbb{R}^m$  and the collection of all  $m \times n$  matrices over  $\mathbb{R}$ .*

# Correspondence between linear maps and matrices

## Corollary

*There is a one to one correspondence between the set of all linear transformations from  $\mathbb{R}^n$  to  $\mathbb{R}^m$  and the collection of all  $m \times n$  matrices over  $\mathbb{R}$ .*

*Proof.* The correspondences are given by

$$\varphi : T \mapsto [T(e_1) \quad T(e_2) \quad \cdots \quad T(e_n)] \quad \text{and} \quad \psi : A \mapsto T_A.$$

# Correspondence between linear maps and matrices

## Corollary

*There is a one to one correspondence between the set of all linear transformations from  $\mathbb{R}^n$  to  $\mathbb{R}^m$  and the collection of all  $m \times n$  matrices over  $\mathbb{R}$ .*

*Proof.* The correspondences are given by

$$\varphi : T \mapsto [T(e_1) \quad T(e_2) \quad \cdots \quad T(e_n)] \quad \text{and} \quad \psi : A \mapsto T_A.$$

It can be verified that the compositions  $\varphi \circ \psi$



# Correspondence between linear maps and matrices

## Corollary

*There is a one to one correspondence between the set of all linear transformations from  $\mathbb{R}^n$  to  $\mathbb{R}^m$  and the collection of all  $m \times n$  matrices over  $\mathbb{R}$ .*

*Proof.* The correspondences are given by

$$\varphi : T \mapsto [T(e_1) \quad T(e_2) \quad \cdots \quad T(e_n)] \quad \text{and} \quad \psi : A \mapsto T_A.$$

It can be verified that the compositions  $\varphi \circ \psi$  and  $\psi \circ \varphi$  are the identity maps on the respective domains.

# Matrix representation of a linear map $T : V \rightarrow W$

## Theorem

*Let  $T : V \rightarrow W$  be a linear transformation.*

# Matrix representation of a linear map $T : V \rightarrow W$

## Theorem

*Let  $T : V \rightarrow W$  be a linear transformation. Let  $\mathcal{B} = \{v_1, \dots, v_n\}$*

# Matrix representation of a linear map $T : V \rightarrow W$

## Theorem

*Let  $T : V \rightarrow W$  be a linear transformation. Let  $\mathcal{B} = \{v_1, \dots, v_n\}$  and  $\mathcal{B}' = \{w_1, \dots, w_m\}$  be two ordered bases of  $V$  and  $W$  respectively.*

# Matrix representation of a linear map $T : V \rightarrow W$

## Theorem

*Let  $T : V \rightarrow W$  be a linear transformation. Let  $\mathcal{B} = \{v_1, \dots, v_n\}$  and  $\mathcal{B}' = \{w_1, \dots, w_m\}$  be two ordered bases of  $V$  and  $W$  respectively. Then there exists an  $m \times n$  matrix  $A$  such that  $T$  can be represented by  $A$ ,*

# Matrix representation of a linear map $T : V \rightarrow W$

## Theorem

*Let  $T : V \rightarrow W$  be a linear transformation. Let  $\mathcal{B} = \{v_1, \dots, v_n\}$  and  $\mathcal{B}' = \{w_1, \dots, w_m\}$  be two ordered bases of  $V$  and  $W$  respectively. Then there exists an  $m \times n$  matrix  $A$  such that  $T$  can be represented by  $A$ , i.e.,  $A[v]_{\mathcal{B}} = [T(v)]_{\mathcal{B}'}$  for every  $v \in V$ .*

# Matrix representation of a linear map $T : V \rightarrow W$

## Theorem


*Let  $T : V \rightarrow W$  be a linear transformation. Let  $\mathcal{B} = \{v_1, \dots, v_n\}$  and  $\mathcal{B}' = \{w_1, \dots, w_m\}$  be two ordered bases of  $V$  and  $W$  respectively. Then there exists an  $m \times n$  matrix  $A$  such that  $T$  can be represented by  $A$ , i.e.,  $A[v]_{\mathcal{B}} = [T(v)]_{\mathcal{B}'}$  for every  $v \in V$ . The  $i$ th column of  $A$ , which is same as  $Ae_i$ , will be obtained by  $[T(v_i)]_{\mathcal{B}'}$ .*

# Matrix representation of a linear map $T : V \rightarrow W$

## Theorem

*Let  $T : V \rightarrow W$  be a linear transformation. Let  $\mathcal{B} = \{v_1, \dots, v_n\}$  and  $\mathcal{B}' = \{w_1, \dots, w_m\}$  be two ordered bases of  $V$  and  $W$  respectively. Then there exists an  $m \times n$  matrix  $A$  such that  $T$  can be represented by  $A$ , i.e.,  $A[v]_{\mathcal{B}} = [T(v)]_{\mathcal{B}'}$  for every  $v \in V$ . The  $i$ th column of  $A$ , which is same as  $Ae_i$ , will be obtained by  $[T(v_i)]_{\mathcal{B}'}$ .*

*Sketch of the Proof.*






# Matrix representation of a linear map $T : V \rightarrow W$

## Theorem

Let  $T : V \rightarrow W$  be a linear transformation. Let  $\mathcal{B} = \{v_1, \dots, v_n\}$  and  $\mathcal{B}' = \{w_1, \dots, w_m\}$  be two ordered bases of  $V$  and  $W$  respectively. Then there exists an  $m \times n$  matrix  $A$  such that  $T$  can be represented by  $A$ , i.e.,  $A[v]_{\mathcal{B}} = [T(v)]_{\mathcal{B}'}$  for every  $v \in V$ . The  $i$ th column of  $A$ , which is same as  $Ae_i$ , will be obtained by  $[T(v_i)]_{\mathcal{B}'}$ .

*Sketch of the Proof.*



# Matrix representation of a linear map $T : V \rightarrow W$

## Theorem

Let  $T : V \rightarrow W$  be a linear transformation. Let  $\mathcal{B} = \{v_1, \dots, v_n\}$  and  $\mathcal{B}' = \{w_1, \dots, w_m\}$  be two ordered bases of  $V$  and  $W$  respectively. Then there exists an  $m \times n$  matrix  $A$  such that  $T$  can be represented by  $A$ , i.e.,  $A[v]_{\mathcal{B}} = [T(v)]_{\mathcal{B}'}$  for every  $v \in V$ . The  $i$ th column of  $A$ , which is same as  $Ae_i$ , will be obtained by  $[T(v_i)]_{\mathcal{B}'}$ .

*Sketch of the Proof.*

$$\begin{array}{ccccc} V & \xrightarrow{T} & W & & v & \xrightarrow{T} & T(v) & & V & \xrightarrow{T} & W \\ \updownarrow & & \updownarrow & & \updownarrow & & \updownarrow & & \updownarrow & & \updownarrow \\ \mathbb{R}^n & & \mathbb{R}^m & & [v]_{\mathcal{B}} & & [T(v)]_{\mathcal{B}'} & & \mathbb{R}^n & \xrightarrow{???} & \mathbb{R}^m \end{array}$$

# Matrix representation of a linear map $T : V \rightarrow W$

## Theorem

Let  $T : V \rightarrow W$  be a linear transformation. Let  $\mathcal{B} = \{v_1, \dots, v_n\}$  and  $\mathcal{B}' = \{w_1, \dots, w_m\}$  be two ordered bases of  $V$  and  $W$  respectively. Then there exists an  $m \times n$  matrix  $A$  such that  $T$  can be represented by  $A$ , i.e.,  $A[v]_{\mathcal{B}} = [T(v)]_{\mathcal{B}'}$  for every  $v \in V$ . The  $i$ th column of  $A$ , which is same as  $Ae_i$ , will be obtained by  $[T(v_i)]_{\mathcal{B}'}$ .

*Sketch of the Proof.*

$$\begin{array}{ccc} V & \xrightarrow{T} & W \\ \updownarrow & & \updownarrow \\ \mathbb{R}^n & & \mathbb{R}^m \end{array} \quad \begin{array}{ccc} v & \xrightarrow{T} & T(v) \\ \updownarrow & & \updownarrow \\ [v]_{\mathcal{B}} & & [T(v)]_{\mathcal{B}'} \end{array} \quad \begin{array}{ccc} V & \xrightarrow{T} & W \\ \updownarrow & & \updownarrow \\ \mathbb{R}^n & \xrightarrow{???} & \mathbb{R}^m \end{array} .$$

By the last theorem, there exists  $A$  such that the following diagram is commutative.

$$\begin{array}{ccc} V & \xrightarrow{T} & W \\ \updownarrow & & \updownarrow \\ \mathbb{R}^n & \xrightarrow{A} & \mathbb{R}^m \end{array} .$$

# Matrix representation of a linear map $T : V \rightarrow W$

## Theorem

Let  $T : V \rightarrow W$  be a linear transformation. Let  $\mathcal{B} = \{v_1, \dots, v_n\}$  and  $\mathcal{B}' = \{w_1, \dots, w_m\}$  be two ordered bases of  $V$  and  $W$  respectively. Then there exists an  $m \times n$  matrix  $A$  such that  $T$  can be represented by  $A$ , i.e.,  $A[v]_{\mathcal{B}} = [T(v)]_{\mathcal{B}'}$  for every  $v \in V$ . The  $i$ th column of  $A$ , which is same as  $Ae_i$ , will be obtained by  $[T(v_i)]_{\mathcal{B}'}$ .

*Sketch of the Proof.*

$$\begin{array}{ccc} V & \xrightarrow{T} & W \\ \updownarrow & & \updownarrow \\ \mathbb{R}^n & & \mathbb{R}^m \end{array} \quad \begin{array}{ccc} v & \xrightarrow{T} & T(v) \\ \updownarrow & & \updownarrow \\ [v]_{\mathcal{B}} & & [T(v)]_{\mathcal{B}'} \end{array} \quad \begin{array}{ccc} V & \xrightarrow{T} & W \\ \updownarrow & & \updownarrow \\ \mathbb{R}^n & \xrightarrow{???} & \mathbb{R}^m \end{array} .$$

By the last theorem, there exists  $A$  such that the following diagram is commutative.

$$\begin{array}{ccc} V & \xrightarrow{T} & W \\ \updownarrow & & \updownarrow \\ \mathbb{R}^n & \xrightarrow{A} & \mathbb{R}^m \end{array} . \quad \text{Observe that the } i\text{th column of } A \text{ is } [T(v_i)]_{\mathcal{B}'} .$$

A linear map  $T : \mathbb{R}^n \rightarrow \mathbb{R}^m$  is determined by  $T(e_i)$

### Theorem

*Consider the standard basis  $\{e_1, \dots, e_n\}$  of  $\mathbb{R}^n$ .*

# A linear map $T : \mathbb{R}^n \rightarrow \mathbb{R}^m$ is determined by $T(e_i)$

## Theorem

*Consider the standard basis  $\{e_1, \dots, e_n\}$  of  $\mathbb{R}^n$ .*

*Then any linear transformation  $T : \mathbb{R}^n \rightarrow \mathbb{R}^m$  is uniquely determined by  $T(e_i)$  for all  $1 \leq i \leq n$ .*

# A linear map $T : \mathbb{R}^n \rightarrow \mathbb{R}^m$ is determined by $T(e_i)$

## Theorem

*Consider the standard basis  $\{e_1, \dots, e_n\}$  of  $\mathbb{R}^n$ .*

*Then any linear transformation  $T : \mathbb{R}^n \rightarrow \mathbb{R}^m$  is uniquely determined by  $T(e_i)$  for all  $1 \leq i \leq n$ .*

**Proof.** Every vector  $v = \begin{pmatrix} x_1 \\ x_2 \\ \vdots \\ x_n \end{pmatrix} \in \mathbb{R}^n$  has a unique expression:

$$v = x_1 \begin{pmatrix} 1 \\ 0 \\ \vdots \\ 0 \end{pmatrix} + x_2 \begin{pmatrix} 0 \\ 1 \\ \vdots \\ 0 \end{pmatrix} + \cdots + x_n \begin{pmatrix} 0 \\ 0 \\ \vdots \\ 1 \end{pmatrix} = x_1 e_1 + x_2 e_2 + \cdots + x_n e_n.$$

# A linear map $T : \mathbb{R}^n \rightarrow \mathbb{R}^m$ is determined by $T(e_i)$

## Theorem

*Consider the standard basis  $\{e_1, \dots, e_n\}$  of  $\mathbb{R}^n$ .*

*Then any linear transformation  $T : \mathbb{R}^n \rightarrow \mathbb{R}^m$  is uniquely determined by  $T(e_i)$  for all  $1 \leq i \leq n$ .*

**Proof.** Every vector  $v = \begin{pmatrix} x_1 \\ x_2 \\ \vdots \\ x_n \end{pmatrix} \in \mathbb{R}^n$  has a unique expression:

$$v = x_1 \begin{pmatrix} 1 \\ 0 \\ \vdots \\ 0 \end{pmatrix} + x_2 \begin{pmatrix} 0 \\ 1 \\ \vdots \\ 0 \end{pmatrix} + \cdots + x_n \begin{pmatrix} 0 \\ 0 \\ \vdots \\ 1 \end{pmatrix} = x_1 e_1 + x_2 e_2 + \cdots + x_n e_n.$$

Hence, by linearity,  $T(v) = x_1 T(e_1) + \cdots + x_n T(e_n)$ ,



# A linear map $T : \mathbb{R}^n \rightarrow \mathbb{R}^m$ is determined by $T(e_i)$

## Theorem

*Consider the standard basis  $\{e_1, \dots, e_n\}$  of  $\mathbb{R}^n$ .*

*Then any linear transformation  $T : \mathbb{R}^n \rightarrow \mathbb{R}^m$  is uniquely determined by  $T(e_i)$  for all  $1 \leq i \leq n$ .*

**Proof.** Every vector  $v = \begin{pmatrix} x_1 \\ x_2 \\ \vdots \\ x_n \end{pmatrix} \in \mathbb{R}^n$  has a unique expression:

$$v = x_1 \begin{pmatrix} 1 \\ 0 \\ \vdots \\ 0 \end{pmatrix} + x_2 \begin{pmatrix} 0 \\ 1 \\ \vdots \\ 0 \end{pmatrix} + \cdots + x_n \begin{pmatrix} 0 \\ 0 \\ \vdots \\ 1 \end{pmatrix} = x_1 e_1 + x_2 e_2 + \cdots + x_n e_n.$$

Hence, by linearity,  $T(v) = x_1 T(e_1) + \cdots + x_n T(e_n)$ , which has a unique choice

# A linear map $T : \mathbb{R}^n \rightarrow \mathbb{R}^m$ is determined by $T(e_i)$

## Theorem

*Consider the standard basis  $\{e_1, \dots, e_n\}$  of  $\mathbb{R}^n$ .*

*Then any linear transformation  $T : \mathbb{R}^n \rightarrow \mathbb{R}^m$  is uniquely determined by  $T(e_i)$  for all  $1 \leq i \leq n$ .*

**Proof.** Every vector  $v = \begin{pmatrix} x_1 \\ x_2 \\ \vdots \\ x_n \end{pmatrix} \in \mathbb{R}^n$  has a unique expression:

$$v = x_1 \begin{pmatrix} 1 \\ 0 \\ \vdots \\ 0 \end{pmatrix} + x_2 \begin{pmatrix} 0 \\ 1 \\ \vdots \\ 0 \end{pmatrix} + \cdots + x_n \begin{pmatrix} 0 \\ 0 \\ \vdots \\ 1 \end{pmatrix} = x_1 e_1 + x_2 e_2 + \cdots + x_n e_n.$$

Hence, by linearity,  $T(v) = x_1 T(e_1) + \cdots + x_n T(e_n)$ , which has a unique choice once  $T(e_i)$  is given for every  $i$ .

# A linear map is uniquely determined by its action on a basis

## Theorem

*Let  $V$  be finite dimensional,*

# A linear map is uniquely determined by its action on a basis

## Theorem

*Let  $V$  be finite dimensional, and  $\{v_1, \dots, v_n\}$  be a basis of  $V$ .*

# A linear map is uniquely determined by its action on a basis

## Theorem

*Let  $V$  be finite dimensional, and  $\{v_1, \dots, v_n\}$  be a basis of  $V$ . Then any linear transformation  $T : V \rightarrow W$*

# A linear map is uniquely determined by its action on a basis

## Theorem

*Let  $V$  be finite dimensional, and  $\{v_1, \dots, v_n\}$  be a basis of  $V$ . Then any linear transformation  $T : V \rightarrow W$  is uniquely determined by  $T(v_i)$  for all  $1 \leq i \leq n$ .*

# A linear map is uniquely determined by its action on a basis

## Theorem

*Let  $V$  be finite dimensional, and  $\{v_1, \dots, v_n\}$  be a basis of  $V$ . Then any linear transformation  $T : V \rightarrow W$  is uniquely determined by  $T(v_i)$  for all  $1 \leq i \leq n$ .*

*Proof.* Every vector  $v \in V$

# A linear map is uniquely determined by its action on a basis

## Theorem

*Let  $V$  be finite dimensional, and  $\{v_1, \dots, v_n\}$  be a basis of  $V$ . Then any linear transformation  $T : V \rightarrow W$  is uniquely determined by  $T(v_i)$  for all  $1 \leq i \leq n$ .*

*Proof.* Every vector  $v \in V$  has a unique expression:

$$v = c_1 v_1 + c_2 v_2 + \dots + c_n v_n,$$



# A linear map is uniquely determined by its action on a basis

## Theorem

*Let  $V$  be finite dimensional, and  $\{v_1, \dots, v_n\}$  be a basis of  $V$ . Then any linear transformation  $T : V \rightarrow W$  is uniquely determined by  $T(v_i)$  for all  $1 \leq i \leq n$ .*

*Proof.* Every vector  $v \in V$  has a unique expression:

$v = c_1v_1 + c_2v_2 + \dots + c_nv_n$ , because if  $v = d_1v_1 + d_2v_2 + \dots + d_nv_n$  is another expression,

# A linear map is uniquely determined by its action on a basis

## Theorem

*Let  $V$  be finite dimensional, and  $\{v_1, \dots, v_n\}$  be a basis of  $V$ . Then any linear transformation  $T : V \rightarrow W$  is uniquely determined by  $T(v_i)$  for all  $1 \leq i \leq n$ .*

**Proof.** Every vector  $v \in V$  has a unique expression:

$v = c_1v_1 + c_2v_2 + \dots + c_nv_n$ , because if  $v = d_1v_1 + d_2v_2 + \dots + d_nv_n$  is another expression, then

$$(c_1 - d_1)v_1 + \dots + (c_n - d_n)v_n = 0$$

# A linear map is uniquely determined by its action on a basis

## Theorem

*Let  $V$  be finite dimensional, and  $\{v_1, \dots, v_n\}$  be a basis of  $V$ . Then any linear transformation  $T : V \rightarrow W$  is uniquely determined by  $T(v_i)$  for all  $1 \leq i \leq n$ .*

**Proof.** Every vector  $v \in V$  has a unique expression:

$v = c_1v_1 + c_2v_2 + \dots + c_nv_n$ , because if  $v = d_1v_1 + d_2v_2 + \dots + d_nv_n$  is another expression, then

$$(c_1 - d_1)v_1 + \dots + (c_n - d_n)v_n = 0 \implies c_i = d_i \text{ for all } i.$$

# A linear map is uniquely determined by its action on a basis

## Theorem

*Let  $V$  be finite dimensional, and  $\{v_1, \dots, v_n\}$  be a basis of  $V$ . Then any linear transformation  $T : V \rightarrow W$  is uniquely determined by  $T(v_i)$  for all  $1 \leq i \leq n$ .*

*Proof.* Every vector  $v \in V$  has a unique expression:

$v = c_1v_1 + c_2v_2 + \dots + c_nv_n$ , because if  $v = d_1v_1 + d_2v_2 + \dots + d_nv_n$  is another expression, then

$$(c_1 - d_1)v_1 + \dots + (c_n - d_n)v_n = 0 \implies c_i = d_i \text{ for all } i.$$

Hence, by linearity,  $T(v) = c_1T(v_1) + \dots + c_nT(v_n)$ ,

# A linear map is uniquely determined by its action on a basis

## Theorem

*Let  $V$  be finite dimensional, and  $\{v_1, \dots, v_n\}$  be a basis of  $V$ . Then any linear transformation  $T : V \rightarrow W$  is uniquely determined by  $T(v_i)$  for all  $1 \leq i \leq n$ .*

**Proof.** Every vector  $v \in V$  has a unique expression:

$v = c_1v_1 + c_2v_2 + \dots + c_nv_n$ , because if  $v = d_1v_1 + d_2v_2 + \dots + d_nv_n$  is another expression, then

$$(c_1 - d_1)v_1 + \dots + (c_n - d_n)v_n = 0 \implies c_i = d_i \text{ for all } i.$$

Hence, by linearity,  $T(v) = c_1T(v_1) + \dots + c_nT(v_n)$ , which has a unique choice,

# A linear map is uniquely determined by its action on a basis

## Theorem

*Let  $V$  be finite dimensional, and  $\{v_1, \dots, v_n\}$  be a basis of  $V$ . Then any linear transformation  $T : V \rightarrow W$  is uniquely determined by  $T(v_i)$  for all  $1 \leq i \leq n$ .*

**Proof.** Every vector  $v \in V$  has a unique expression:

$v = c_1v_1 + c_2v_2 + \dots + c_nv_n$ , because if  $v = d_1v_1 + d_2v_2 + \dots + d_nv_n$  is another expression, then

$$(c_1 - d_1)v_1 + \dots + (c_n - d_n)v_n = 0 \implies c_i = d_i \text{ for all } i.$$

Hence, by linearity,  $T(v) = c_1T(v_1) + \dots + c_nT(v_n)$ , which has a unique choice, once  $T(v_i)$  is given for every  $i$ .

# A linear map is determined by its action on a basis

## Theorem

*Let  $V$  be finite dimensional,*

# A linear map is determined by its action on a basis

## Theorem

*Let  $V$  be finite dimensional, and  $\{v_1, \dots, v_n\}$  be a basis of  $V$ .*



# A linear map is determined by its action on a basis

## Theorem

*Let  $V$  be finite dimensional, and  $\{v_1, \dots, v_n\}$  be a basis of  $V$ .  
Let  $\{w_1, \dots, w_n\}$  be any collection of  $n$  vectors in  $W$ .*

# A linear map is determined by its action on a basis

## Theorem

*Let  $V$  be finite dimensional, and  $\{v_1, \dots, v_n\}$  be a basis of  $V$ .*

*Let  $\{w_1, \dots, w_n\}$  be any collection of  $n$  vectors in  $W$ .*

*Then there is EXACTLY one linear transformation  $T : V \rightarrow W$*

# A linear map is determined by its action on a basis

## Theorem

*Let  $V$  be finite dimensional, and  $\{v_1, \dots, v_n\}$  be a basis of  $V$ .*

*Let  $\{w_1, \dots, w_n\}$  be any collection of  $n$  vectors in  $W$ .*

*Then there is EXACTLY one linear transformation  $T : V \rightarrow W$  such that  $T(v_i) = w_i$  for all  $1 \leq i \leq n$ .*

# A linear map is determined by its action on a basis

## Theorem

*Let  $V$  be finite dimensional, and  $\{v_1, \dots, v_n\}$  be a basis of  $V$ .*

*Let  $\{w_1, \dots, w_n\}$  be any collection of  $n$  vectors in  $W$ .*

*Then there is EXACTLY one linear transformation  $T : V \rightarrow W$  such that  $T(v_i) = w_i$  for all  $1 \leq i \leq n$ .*

*Proof.* Once we show the existence,

# A linear map is determined by its action on a basis

## Theorem

*Let  $V$  be finite dimensional, and  $\{v_1, \dots, v_n\}$  be a basis of  $V$ .*

*Let  $\{w_1, \dots, w_n\}$  be any collection of  $n$  vectors in  $W$ .*

*Then there is EXACTLY one linear transformation  $T : V \rightarrow W$  such that  $T(v_i) = w_i$  for all  $1 \leq i \leq n$ .*

*Proof.* Once we show the existence, uniqueness follows from the last theorem.

# A linear map is determined by its action on a basis

## Theorem

*Let  $V$  be finite dimensional, and  $\{v_1, \dots, v_n\}$  be a basis of  $V$ .*

*Let  $\{w_1, \dots, w_n\}$  be any collection of  $n$  vectors in  $W$ .*

*Then there is EXACTLY one linear transformation  $T : V \rightarrow W$  such that  $T(v_i) = w_i$  for all  $1 \leq i \leq n$ .*

*Proof.* Once we show the existence, uniqueness follows from the last theorem. We define a map as follows:

# A linear map is determined by its action on a basis

## Theorem

*Let  $V$  be finite dimensional, and  $\{v_1, \dots, v_n\}$  be a basis of  $V$ .*

*Let  $\{w_1, \dots, w_n\}$  be any collection of  $n$  vectors in  $W$ .*

*Then there is EXACTLY one linear transformation  $T : V \rightarrow W$  such that  $T(v_i) = w_i$  for all  $1 \leq i \leq n$ .*

*Proof.* Once we show the existence, uniqueness follows from the last theorem. We define a map as follows: Every vector  $v \in V$  has a UNIQUE expression:

# A linear map is determined by its action on a basis

## Theorem

*Let  $V$  be finite dimensional, and  $\{v_1, \dots, v_n\}$  be a basis of  $V$ .*

*Let  $\{w_1, \dots, w_n\}$  be any collection of  $n$  vectors in  $W$ .*

*Then there is EXACTLY one linear transformation  $T : V \rightarrow W$  such that  $T(v_i) = w_i$  for all  $1 \leq i \leq n$ .*

*Proof.* Once we show the existence, uniqueness follows from the last theorem. We define a map as follows: Every vector  $v \in V$  has a UNIQUE expression:  $v = c_1 v_1 + \dots + c_n v_n$  as before.



# A linear map is determined by its action on a basis

## Theorem

*Let  $V$  be finite dimensional, and  $\{v_1, \dots, v_n\}$  be a basis of  $V$ .*

*Let  $\{w_1, \dots, w_n\}$  be any collection of  $n$  vectors in  $W$ .*

*Then there is EXACTLY one linear transformation  $T : V \rightarrow W$  such that  $T(v_i) = w_i$  for all  $1 \leq i \leq n$ .*

*Proof.* Once we show the existence, uniqueness follows from the last theorem. We define a map as follows: Every vector  $v \in V$  has a UNIQUE expression:  $v = c_1 v_1 + \dots + c_n v_n$  as before.

Define  $T(v)$

# A linear map is determined by its action on a basis

## Theorem

*Let  $V$  be finite dimensional, and  $\{v_1, \dots, v_n\}$  be a basis of  $V$ .*

*Let  $\{w_1, \dots, w_n\}$  be any collection of  $n$  vectors in  $W$ .*

*Then there is EXACTLY one linear transformation  $T : V \rightarrow W$  such that  $T(v_i) = w_i$  for all  $1 \leq i \leq n$ .*

*Proof.* Once we show the existence, uniqueness follows from the last theorem. We define a map as follows: Every vector  $v \in V$  has a UNIQUE expression:  $v = c_1 v_1 + \dots + c_n v_n$  as before.

Define  $T(v) := c_1 w_1 + \dots + c_n w_n$ .

# A linear map is determined by its action on a basis

## Theorem

*Let  $V$  be finite dimensional, and  $\{v_1, \dots, v_n\}$  be a basis of  $V$ .  
Let  $\{w_1, \dots, w_n\}$  be any collection of  $n$  vectors in  $W$ .  
Then there is EXACTLY one linear transformation  $T : V \rightarrow W$  such that  $T(v_i) = w_i$  for all  $1 \leq i \leq n$ .*

*Proof.* Once we show the existence, uniqueness follows from the last theorem. We define a map as follows: Every vector  $v \in V$  has a UNIQUE expression:  $v = c_1 v_1 + \dots + c_n v_n$  as before.

Define  $T(v) := c_1 w_1 + \dots + c_n w_n$ . Then

- $T : V \rightarrow W$  is a linear map because:

# A linear map is determined by its action on a basis

## Theorem

*Let  $V$  be finite dimensional, and  $\{v_1, \dots, v_n\}$  be a basis of  $V$ .  
Let  $\{w_1, \dots, w_n\}$  be any collection of  $n$  vectors in  $W$ .  
Then there is EXACTLY one linear transformation  $T : V \rightarrow W$  such that  $T(v_i) = w_i$  for all  $1 \leq i \leq n$ .*

*Proof.* Once we show the existence, uniqueness follows from the last theorem. We define a map as follows: Every vector  $v \in V$  has a UNIQUE expression:  $v = c_1 v_1 + \dots + c_n v_n$  as before.

Define  $T(v) := c_1 w_1 + \dots + c_n w_n$ . Then

- $T : V \rightarrow W$  is a linear map because:
- If  $v = c_1 v_1 + \dots + c_n v_n$  and  $u = d_1 v_1 + \dots + d_n v_n$ ,

# A linear map is determined by its action on a basis

## Theorem

*Let  $V$  be finite dimensional, and  $\{v_1, \dots, v_n\}$  be a basis of  $V$ .  
Let  $\{w_1, \dots, w_n\}$  be any collection of  $n$  vectors in  $W$ .  
Then there is EXACTLY one linear transformation  $T : V \rightarrow W$  such that  $T(v_i) = w_i$  for all  $1 \leq i \leq n$ .*

*Proof.* Once we show the existence, uniqueness follows from the last theorem. We define a map as follows: Every vector  $v \in V$  has a UNIQUE expression:  $v = c_1v_1 + \dots + c_nv_n$  as before.

Define  $T(v) := c_1w_1 + \dots + c_nw_n$ . Then

- $T : V \rightarrow W$  is a linear map because:
- If  $v = c_1v_1 + \dots + c_nv_n$  and  $u = d_1v_1 + \dots + d_nv_n$ , then  
 $v + u = (c_1 + d_1)v_1 + \dots + (c_n + d_n)v_n$ .

# A linear map is determined by its action on a basis

## Theorem

*Let  $V$  be finite dimensional, and  $\{v_1, \dots, v_n\}$  be a basis of  $V$ .  
Let  $\{w_1, \dots, w_n\}$  be any collection of  $n$  vectors in  $W$ .  
Then there is EXACTLY one linear transformation  $T : V \rightarrow W$  such that  $T(v_i) = w_i$  for all  $1 \leq i \leq n$ .*

*Proof.* Once we show the existence, uniqueness follows from the last theorem. We define a map as follows: Every vector  $v \in V$  has a UNIQUE expression:  $v = c_1 v_1 + \dots + c_n v_n$  as before.

Define  $T(v) := c_1 w_1 + \dots + c_n w_n$ . Then

- $T : V \rightarrow W$  is a linear map because:
- If  $v = c_1 v_1 + \dots + c_n v_n$  and  $u = d_1 v_1 + \dots + d_n v_n$ , then  $v + u = (c_1 + d_1)v_1 + \dots + (c_n + d_n)v_n$ . Hence  $T(v + u)$

# A linear map is determined by its action on a basis

## Theorem

*Let  $V$  be finite dimensional, and  $\{v_1, \dots, v_n\}$  be a basis of  $V$ .  
Let  $\{w_1, \dots, w_n\}$  be any collection of  $n$  vectors in  $W$ .  
Then there is EXACTLY one linear transformation  $T : V \rightarrow W$  such that  $T(v_i) = w_i$  for all  $1 \leq i \leq n$ .*

*Proof.* Once we show the existence, uniqueness follows from the last theorem. We define a map as follows: Every vector  $v \in V$  has a UNIQUE expression:  $v = c_1 v_1 + \dots + c_n v_n$  as before.

Define  $T(v) := c_1 w_1 + \dots + c_n w_n$ . Then

- $T : V \rightarrow W$  is a linear map because:
- If  $v = c_1 v_1 + \dots + c_n v_n$  and  $u = d_1 v_1 + \dots + d_n v_n$ , then  $v + u = (c_1 + d_1)v_1 + \dots + (c_n + d_n)v_n$ . Hence  $T(v + u) = (c_1 + d_1)w_1 + \dots + (c_n + d_n)w_n$

# A linear map is determined by its action on a basis

## Theorem

*Let  $V$  be finite dimensional, and  $\{v_1, \dots, v_n\}$  be a basis of  $V$ .  
Let  $\{w_1, \dots, w_n\}$  be any collection of  $n$  vectors in  $W$ .  
Then there is EXACTLY one linear transformation  $T : V \rightarrow W$  such that  $T(v_i) = w_i$  for all  $1 \leq i \leq n$ .*

*Proof.* Once we show the existence, uniqueness follows from the last theorem. We define a map as follows: Every vector  $v \in V$  has a UNIQUE expression:  $v = c_1 v_1 + \dots + c_n v_n$  as before.

Define  $T(v) := c_1 w_1 + \dots + c_n w_n$ . Then

- $T : V \rightarrow W$  is a linear map because:
- If  $v = c_1 v_1 + \dots + c_n v_n$  and  $u = d_1 v_1 + \dots + d_n v_n$ , then  $v + u = (c_1 + d_1)v_1 + \dots + (c_n + d_n)v_n$ . Hence  $T(v + u) = (c_1 + d_1)w_1 + \dots + (c_n + d_n)w_n = T(v) + T(u)$ .



# A linear map is determined by its action on a basis

## Theorem

*Let  $V$  be finite dimensional, and  $\{v_1, \dots, v_n\}$  be a basis of  $V$ .  
Let  $\{w_1, \dots, w_n\}$  be any collection of  $n$  vectors in  $W$ .  
Then there is EXACTLY one linear transformation  $T : V \rightarrow W$  such that  $T(v_i) = w_i$  for all  $1 \leq i \leq n$ .*

*Proof.* Once we show the existence, uniqueness follows from the last theorem. We define a map as follows: Every vector  $v \in V$  has a UNIQUE expression:  $v = c_1v_1 + \dots + c_nv_n$  as before.

Define  $T(v) := c_1w_1 + \dots + c_nw_n$ . Then

- $T : V \rightarrow W$  is a linear map because:
- If  $v = c_1v_1 + \dots + c_nv_n$  and  $u = d_1v_1 + \dots + d_nv_n$ , then  $v + u = (c_1 + d_1)v_1 + \dots + (c_n + d_n)v_n$ . Hence  $T(v + u) = (c_1 + d_1)w_1 + \dots + (c_n + d_n)w_n = T(v) + T(u)$ .
- If  $v = c_1v_1 + \dots + c_nv_n$ ,

# A linear map is determined by its action on a basis

## Theorem

*Let  $V$  be finite dimensional, and  $\{v_1, \dots, v_n\}$  be a basis of  $V$ .  
Let  $\{w_1, \dots, w_n\}$  be any collection of  $n$  vectors in  $W$ .  
Then there is EXACTLY one linear transformation  $T : V \rightarrow W$  such that  $T(v_i) = w_i$  for all  $1 \leq i \leq n$ .*

*Proof.* Once we show the existence, uniqueness follows from the last theorem. We define a map as follows: Every vector  $v \in V$  has a UNIQUE expression:  $v = c_1v_1 + \dots + c_nv_n$  as before.

Define  $T(v) := c_1w_1 + \dots + c_nw_n$ . Then

- $T : V \rightarrow W$  is a linear map because:
- If  $v = c_1v_1 + \dots + c_nv_n$  and  $u = d_1v_1 + \dots + d_nv_n$ , then  $v + u = (c_1 + d_1)v_1 + \dots + (c_n + d_n)v_n$ . Hence  $T(v + u) = (c_1 + d_1)w_1 + \dots + (c_n + d_n)w_n = T(v) + T(u)$ .
- If  $v = c_1v_1 + \dots + c_nv_n$ , then  $cv = (cc_1)v_1 + \dots + (cc_n)v_n$ .

# A linear map is determined by its action on a basis

## Theorem

*Let  $V$  be finite dimensional, and  $\{v_1, \dots, v_n\}$  be a basis of  $V$ .  
Let  $\{w_1, \dots, w_n\}$  be any collection of  $n$  vectors in  $W$ .  
Then there is EXACTLY one linear transformation  $T : V \rightarrow W$  such that  $T(v_i) = w_i$  for all  $1 \leq i \leq n$ .*

*Proof.* Once we show the existence, uniqueness follows from the last theorem. We define a map as follows: Every vector  $v \in V$  has a UNIQUE expression:  $v = c_1v_1 + \dots + c_nv_n$  as before.

Define  $T(v) := c_1w_1 + \dots + c_nw_n$ . Then

- $T : V \rightarrow W$  is a linear map because:
- If  $v = c_1v_1 + \dots + c_nv_n$  and  $u = d_1v_1 + \dots + d_nv_n$ , then  $v + u = (c_1 + d_1)v_1 + \dots + (c_n + d_n)v_n$ . Hence  $T(v + u) = (c_1 + d_1)w_1 + \dots + (c_n + d_n)w_n = T(v) + T(u)$ .
- If  $v = c_1v_1 + \dots + c_nv_n$ , then  $cv = (cc_1)v_1 + \dots + (cc_n)v_n$ . Hence  $T(cv) = cT(v)$ .

# A linear map is determined by its action on a basis

## Theorem

*Let  $V$  be finite dimensional, and  $\{v_1, \dots, v_n\}$  be a basis of  $V$ .  
Let  $\{w_1, \dots, w_n\}$  be any collection of  $n$  vectors in  $W$ .  
Then there is EXACTLY one linear transformation  $T : V \rightarrow W$  such that  $T(v_i) = w_i$  for all  $1 \leq i \leq n$ .*

*Proof.* Once we show the existence, uniqueness follows from the last theorem. We define a map as follows: Every vector  $v \in V$  has a UNIQUE expression:  $v = c_1v_1 + \dots + c_nv_n$  as before.

Define  $T(v) := c_1w_1 + \dots + c_nw_n$ . Then

- $T : V \rightarrow W$  is a linear map because:
- If  $v = c_1v_1 + \dots + c_nv_n$  and  $u = d_1v_1 + \dots + d_nv_n$ , then  $v + u = (c_1 + d_1)v_1 + \dots + (c_n + d_n)v_n$ . Hence  $T(v + u) = (c_1 + d_1)w_1 + \dots + (c_n + d_n)w_n = T(v) + T(u)$ .
- If  $v = c_1v_1 + \dots + c_nv_n$ , then  $cv = (cc_1)v_1 + \dots + (cc_n)v_n$ . Hence  $T(cv) = (cc_1)w_1 + \dots + (cc_n)w_n$

# A linear map is determined by its action on a basis

## Theorem

*Let  $V$  be finite dimensional, and  $\{v_1, \dots, v_n\}$  be a basis of  $V$ .  
Let  $\{w_1, \dots, w_n\}$  be any collection of  $n$  vectors in  $W$ .  
Then there is EXACTLY one linear transformation  $T : V \rightarrow W$  such that  $T(v_i) = w_i$  for all  $1 \leq i \leq n$ .*

*Proof.* Once we show the existence, uniqueness follows from the last theorem. We define a map as follows: Every vector  $v \in V$  has a UNIQUE expression:  $v = c_1v_1 + \dots + c_nv_n$  as before.

Define  $T(v) := c_1w_1 + \dots + c_nw_n$ . Then

- $T : V \rightarrow W$  is a linear map because:
- If  $v = c_1v_1 + \dots + c_nv_n$  and  $u = d_1v_1 + \dots + d_nv_n$ , then  $v + u = (c_1 + d_1)v_1 + \dots + (c_n + d_n)v_n$ . Hence  $T(v + u) = (c_1 + d_1)w_1 + \dots + (c_n + d_n)w_n = T(v) + T(u)$ .
- If  $v = c_1v_1 + \dots + c_nv_n$ , then  $cv = (cc_1)v_1 + \dots + (cc_n)v_n$ . Hence  $T(cv) = (cc_1)w_1 + \dots + (cc_n)w_n = cT(v)$ .

# Null space and nullity of a linear transformation

- Let  $T : V \rightarrow W$  be a linear transformation. Then

# Null space and nullity of a linear transformation

- Let  $T : V \rightarrow W$  be a linear transformation. Then
- $\text{Null}(T) := \{v \in V : T(v) = 0\}$  is a subspace of  $V$ ,

# Null space and nullity of a linear transformation

- Let  $T : V \rightarrow W$  be a linear transformation. Then
- $\text{Null}(T) := \{v \in V : T(v) = 0\}$  is a subspace of  $V$ , because:
- It is non-empty as  $0 \in \text{Null}(T)$ .



# Null space and nullity of a linear transformation

- Let  $T : V \rightarrow W$  be a linear transformation. Then
- $\text{Null}(T) := \{v \in V : T(v) = 0\}$  is a subspace of  $V$ , because:
- It is non-empty as  $0 \in \text{Null}(T)$ .
- If  $u, v \in \text{Null}(T)$  and  $c, d \in \mathbb{R}$ ,

# Null space and nullity of a linear transformation

- Let  $T : V \rightarrow W$  be a linear transformation. Then
- $\text{Null}(T) := \{v \in V : T(v) = 0\}$  is a subspace of  $V$ , because:
- It is non-empty as  $0 \in \text{Null}(T)$ .
- If  $u, v \in \text{Null}(T)$  and  $c, d \in \mathbb{R}$ ,  
then  $T(cu + dv) = cT(u) + dT(v) = 0$ ,

# Null space and nullity of a linear transformation

- Let  $T : V \rightarrow W$  be a linear transformation. Then
- $\text{Null}(T) := \{v \in V : T(v) = 0\}$  is a subspace of  $V$ , because:
- It is non-empty as  $0 \in \text{Null}(T)$ .
- If  $u, v \in \text{Null}(T)$  and  $c, d \in \mathbb{R}$ ,  
then  $T(cu + dv) = cT(u) + dT(v) = 0$ ,  
hence  $cu + dv \in \text{Null}(T)$ .

# Null space and nullity of a linear transformation

- Let  $T : V \rightarrow W$  be a linear transformation. Then
- $\text{Null}(T) := \{v \in V : T(v) = 0\}$  is a subspace of  $V$ , because:
- It is non-empty as  $0 \in \text{Null}(T)$ .
- If  $u, v \in \text{Null}(T)$  and  $c, d \in \mathbb{R}$ ,  
then  $T(cu + dv) = cT(u) + dT(v) = 0$ ,  
hence  $cu + dv \in \text{Null}(T)$ .

## Definition (Null space and nullity)

- $\text{Null}(T) := \{v \in V : T(v) = 0\}$  is called the **null space** of  $T$ .

# Null space and nullity of a linear transformation

- Let  $T : V \rightarrow W$  be a linear transformation. Then
- $\text{Null}(T) := \{v \in V : T(v) = 0\}$  is a subspace of  $V$ , because:
- It is non-empty as  $0 \in \text{Null}(T)$ .
- If  $u, v \in \text{Null}(T)$  and  $c, d \in \mathbb{R}$ ,  
then  $T(cu + dv) = cT(u) + dT(v) = 0$ ,  
hence  $cu + dv \in \text{Null}(T)$ .

## Definition (Null space and nullity)

- $\text{Null}(T) := \{v \in V : T(v) = 0\}$  is called the **null space** of  $T$ .
- The **nullity** of  $T$  is the dimension of the null space of  $T$ .

# Range (or Image) of a linear transformation, and rank

- Let  $T : V \rightarrow W$  be a linear transformation.

# Range (or Image) of a linear transformation, and rank

- Let  $T : V \rightarrow W$  be a linear transformation. Then
- $\text{Image}(T) := \{w \in W : w = T(v) \text{ for some } v \in V\}$

# Range (or Image) of a linear transformation, and rank

- Let  $T : V \rightarrow W$  be a linear transformation. Then
- $\text{Image}(T) := \{w \in W : w = T(v) \text{ for some } v \in V\}$  is a subspace of  $W$ ,



# Range (or Image) of a linear transformation, and rank

- Let  $T : V \rightarrow W$  be a linear transformation. Then
- $\text{Image}(T) := \{w \in W : w = T(v) \text{ for some } v \in V\}$  is a subspace of  $W$ , because:
- It is non-empty as  $0 \in \text{Image}(T)$ .

# Range (or Image) of a linear transformation, and rank

- Let  $T : V \rightarrow W$  be a linear transformation. Then
- $\text{Image}(T) := \{w \in W : w = T(v) \text{ for some } v \in V\}$  is a subspace of  $W$ , because:
- It is non-empty as  $0 \in \text{Image}(T)$ .
- If  $w_1, w_2 \in \text{Image}(T)$  and  $c_1, c_2 \in \mathbb{R}$ ,

# Range (or Image) of a linear transformation, and rank

- Let  $T : V \rightarrow W$  be a linear transformation. Then
- $\text{Image}(T) := \{w \in W : w = T(v) \text{ for some } v \in V\}$  is a subspace of  $W$ , because:
- It is non-empty as  $0 \in \text{Image}(T)$ .
- If  $w_1, w_2 \in \text{Image}(T)$  and  $c_1, c_2 \in \mathbb{R}$ , then  $w_1 = T(v_1)$  and  $w_2 = T(v_2)$  for some  $v_1, v_2 \in V$ ,

# Range (or Image) of a linear transformation, and rank

- Let  $T : V \rightarrow W$  be a linear transformation. Then
- $\text{Image}(T) := \{w \in W : w = T(v) \text{ for some } v \in V\}$  is a subspace of  $W$ , because:
  - It is non-empty as  $0 \in \text{Image}(T)$ .
  - If  $w_1, w_2 \in \text{Image}(T)$  and  $c_1, c_2 \in \mathbb{R}$ , then  $w_1 = T(v_1)$  and  $w_2 = T(v_2)$  for some  $v_1, v_2 \in V$ , hence  $c_1 w_1 + c_2 w_2 = T(c_1 v_1 + c_2 v_2) \in \text{Image}(T)$ .

# Range (or Image) of a linear transformation, and rank

- Let  $T : V \rightarrow W$  be a linear transformation. Then
- $\text{Image}(T) := \{w \in W : w = T(v) \text{ for some } v \in V\}$  is a subspace of  $W$ , because:
  - It is non-empty as  $0 \in \text{Image}(T)$ .
  - If  $w_1, w_2 \in \text{Image}(T)$  and  $c_1, c_2 \in \mathbb{R}$ , then  $w_1 = T(v_1)$  and  $w_2 = T(v_2)$  for some  $v_1, v_2 \in V$ , hence  $c_1 w_1 + c_2 w_2 = T(c_1 v_1 + c_2 v_2) \in \text{Image}(T)$ .

## Definition (Range space and rank)

- $\text{Image}(T) := \{w \in W : w = T(v) \text{ for some } v \in V\}$  is called the **range space** of  $T$ .

# Range (or Image) of a linear transformation, and rank

- Let  $T : V \rightarrow W$  be a linear transformation. Then
- $\text{Image}(T) := \{w \in W : w = T(v) \text{ for some } v \in V\}$  is a subspace of  $W$ , because:
  - It is non-empty as  $0 \in \text{Image}(T)$ .
  - If  $w_1, w_2 \in \text{Image}(T)$  and  $c_1, c_2 \in \mathbb{R}$ , then  $w_1 = T(v_1)$  and  $w_2 = T(v_2)$  for some  $v_1, v_2 \in V$ , hence  $c_1 w_1 + c_2 w_2 = T(c_1 v_1 + c_2 v_2) \in \text{Image}(T)$ .

## Definition (Range space and rank)

- $\text{Image}(T) := \{w \in W : w = T(v) \text{ for some } v \in V\}$  is called the **range space** of  $T$ .
- The **rank** of  $T$  is the dimension of the range space of  $T$ .

# Rank-Nullity Theorem

## Theorem

*Let  $T : V \rightarrow W$  be a linear transformation, where  $\dim(V)$  is finite.*

# Rank-Nullity Theorem

## Theorem

*Let  $T : V \rightarrow W$  be a linear transformation, where  $\dim(V)$  is finite.  
Then  $\text{rank}(T) + \text{nullity}(T) = \dim(V)$ .*



# Rank-Nullity Theorem

## Theorem

*Let  $T : V \rightarrow W$  be a linear transformation, where  $\dim(V)$  is finite.  
Then  $\text{rank}(T) + \text{nullity}(T) = \dim(V)$ .*

**Proof.** Start with a basis  $\{u_1, \dots, u_n\}$  of  $\text{Null}(T)$ .

# Rank-Nullity Theorem

## Theorem

*Let  $T : V \rightarrow W$  be a linear transformation, where  $\dim(V)$  is finite. Then  $\text{rank}(T) + \text{nullity}(T) = \dim(V)$ .*

**Proof.** Start with a basis  $\{u_1, \dots, u_n\}$  of  $\text{Null}(T)$ . Extend this to a basis  $\{u_1, \dots, u_n, v_1, \dots, v_r\}$  of  $V$ .

# Rank-Nullity Theorem

## Theorem

*Let  $T : V \rightarrow W$  be a linear transformation, where  $\dim(V)$  is finite. Then  $\text{rank}(T) + \text{nullity}(T) = \dim(V)$ .*

**Proof.** Start with a basis  $\{u_1, \dots, u_n\}$  of  $\text{Null}(T)$ . Extend this to a basis  $\{u_1, \dots, u_n, v_1, \dots, v_r\}$  of  $V$ . It is enough to prove that

$\{T(v_1), \dots, T(v_r)\}$  is a basis of  $\text{Image}(T)$ .

# Rank-Nullity Theorem

## Theorem

*Let  $T : V \rightarrow W$  be a linear transformation, where  $\dim(V)$  is finite. Then  $\text{rank}(T) + \text{nullity}(T) = \dim(V)$ .*

**Proof.** Start with a basis  $\{u_1, \dots, u_n\}$  of  $\text{Null}(T)$ . Extend this to a basis  $\{u_1, \dots, u_n, v_1, \dots, v_r\}$  of  $V$ . It is enough to prove that

$$\{T(v_1), \dots, T(v_r)\} \text{ is a basis of } \text{Image}(T).$$

**Spanning:** Any vector of  $\text{Image}(T)$  looks like  $T(v)$  for some  $v \in V$ .

# Rank-Nullity Theorem

## Theorem

*Let  $T : V \rightarrow W$  be a linear transformation, where  $\dim(V)$  is finite. Then  $\text{rank}(T) + \text{nullity}(T) = \dim(V)$ .*

**Proof.** Start with a basis  $\{u_1, \dots, u_n\}$  of  $\text{Null}(T)$ . Extend this to a basis  $\{u_1, \dots, u_n, v_1, \dots, v_r\}$  of  $V$ . It is enough to prove that

$$\{T(v_1), \dots, T(v_r)\} \text{ is a basis of } \text{Image}(T).$$

**Spanning:** Any vector of  $\text{Image}(T)$  looks like  $T(v)$  for some  $v \in V$ . Write  $v = c_1 u_1 + \dots + c_n u_n + d_1 v_1 + \dots + d_r v_r$ .

# Rank-Nullity Theorem

## Theorem

*Let  $T : V \rightarrow W$  be a linear transformation, where  $\dim(V)$  is finite. Then  $\text{rank}(T) + \text{nullity}(T) = \dim(V)$ .*

**Proof.** Start with a basis  $\{u_1, \dots, u_n\}$  of  $\text{Null}(T)$ . Extend this to a basis  $\{u_1, \dots, u_n, v_1, \dots, v_r\}$  of  $V$ . It is enough to prove that

$\{T(v_1), \dots, T(v_r)\}$  is a basis of  $\text{Image}(T)$ .

**Spanning:** Any vector of  $\text{Image}(T)$  looks like  $T(v)$  for some  $v \in V$ .

Write  $v = c_1 u_1 + \dots + c_n u_n + d_1 v_1 + \dots + d_r v_r$ . Then

$$T(v) = c_1 T(u_1) + \dots + c_n T(u_n) + d_1 T(v_1) + \dots + d_r T(v_r)$$

# Rank-Nullity Theorem

## Theorem

*Let  $T : V \rightarrow W$  be a linear transformation, where  $\dim(V)$  is finite. Then  $\text{rank}(T) + \text{nullity}(T) = \dim(V)$ .*

**Proof.** Start with a basis  $\{u_1, \dots, u_n\}$  of  $\text{Null}(T)$ . Extend this to a basis  $\{u_1, \dots, u_n, v_1, \dots, v_r\}$  of  $V$ . It is enough to prove that

$$\{T(v_1), \dots, T(v_r)\} \text{ is a basis of } \text{Image}(T).$$

**Spanning:** Any vector of  $\text{Image}(T)$  looks like  $T(v)$  for some  $v \in V$ .

Write  $v = c_1 u_1 + \dots + c_n u_n + d_1 v_1 + \dots + d_r v_r$ . Then

$$\begin{aligned} T(v) &= c_1 T(u_1) + \dots + c_n T(u_n) + d_1 T(v_1) + \dots + d_r T(v_r) \\ &= d_1 T(v_1) + \dots + d_r T(v_r). \end{aligned}$$

# Rank-Nullity Theorem

## Theorem

*Let  $T : V \rightarrow W$  be a linear transformation, where  $\dim(V)$  is finite. Then  $\text{rank}(T) + \text{nullity}(T) = \dim(V)$ .*

**Proof.** Start with a basis  $\{u_1, \dots, u_n\}$  of  $\text{Null}(T)$ . Extend this to a basis  $\{u_1, \dots, u_n, v_1, \dots, v_r\}$  of  $V$ . It is enough to prove that

$$\{T(v_1), \dots, T(v_r)\} \text{ is a basis of } \text{Image}(T).$$

**Spanning:** Any vector of  $\text{Image}(T)$  looks like  $T(v)$  for some  $v \in V$ .

Write  $v = c_1 u_1 + \dots + c_n u_n + d_1 v_1 + \dots + d_r v_r$ . Then

$$\begin{aligned} T(v) &= c_1 T(u_1) + \dots + c_n T(u_n) + d_1 T(v_1) + \dots + d_r T(v_r) \\ &= d_1 T(v_1) + \dots + d_r T(v_r). \end{aligned}$$

**Lin. Independence:** Let  $b_1 T(v_1) + \dots + b_r T(v_r) = 0$ .



# Rank-Nullity Theorem

## Theorem

*Let  $T : V \rightarrow W$  be a linear transformation, where  $\dim(V)$  is finite. Then  $\text{rank}(T) + \text{nullity}(T) = \dim(V)$ .*

**Proof.** Start with a basis  $\{u_1, \dots, u_n\}$  of  $\text{Null}(T)$ . Extend this to a basis  $\{u_1, \dots, u_n, v_1, \dots, v_r\}$  of  $V$ . It is enough to prove that

$$\{T(v_1), \dots, T(v_r)\} \text{ is a basis of } \text{Image}(T).$$

**Spanning:** Any vector of  $\text{Image}(T)$  looks like  $T(v)$  for some  $v \in V$ .

Write  $v = c_1 u_1 + \dots + c_n u_n + d_1 v_1 + \dots + d_r v_r$ . Then

$$\begin{aligned} T(v) &= c_1 T(u_1) + \dots + c_n T(u_n) + d_1 T(v_1) + \dots + d_r T(v_r) \\ &= d_1 T(v_1) + \dots + d_r T(v_r). \end{aligned}$$

**Lin. Independence:** Let  $b_1 T(v_1) + \dots + b_r T(v_r) = 0$ .

This implies that  $b_1 v_1 + \dots + b_r v_r \in \text{Null}(T)$ .

# Rank-Nullity Theorem

## Theorem

Let  $T : V \rightarrow W$  be a linear transformation, where  $\dim(V)$  is finite. Then  $\text{rank}(T) + \text{nullity}(T) = \dim(V)$ .

**Proof.** Start with a basis  $\{u_1, \dots, u_n\}$  of  $\text{Null}(T)$ . Extend this to a basis  $\{u_1, \dots, u_n, v_1, \dots, v_r\}$  of  $V$ . It is enough to prove that

$$\{T(v_1), \dots, T(v_r)\} \text{ is a basis of } \text{Image}(T).$$

**Spanning:** Any vector of  $\text{Image}(T)$  looks like  $T(v)$  for some  $v \in V$ .

Write  $v = c_1 u_1 + \dots + c_n u_n + d_1 v_1 + \dots + d_r v_r$ . Then

$$\begin{aligned} T(v) &= c_1 T(u_1) + \dots + c_n T(u_n) + d_1 T(v_1) + \dots + d_r T(v_r) \\ &= d_1 T(v_1) + \dots + d_r T(v_r). \end{aligned}$$

**Lin. Independence:** Let  $b_1 T(v_1) + \dots + b_r T(v_r) = 0$ .

This implies that  $b_1 v_1 + \dots + b_r v_r \in \text{Null}(T)$ .

So  $b_1 v_1 + \dots + b_r v_r = a_1 u_1 + \dots + a_n u_n$  for some  $a_i \in \mathbb{R}$ .

# Rank-Nullity Theorem

## Theorem

*Let  $T : V \rightarrow W$  be a linear transformation, where  $\dim(V)$  is finite. Then  $\text{rank}(T) + \text{nullity}(T) = \dim(V)$ .*

**Proof.** Start with a basis  $\{u_1, \dots, u_n\}$  of  $\text{Null}(T)$ . Extend this to a basis  $\{u_1, \dots, u_n, v_1, \dots, v_r\}$  of  $V$ . It is enough to prove that

$$\{T(v_1), \dots, T(v_r)\} \text{ is a basis of } \text{Image}(T).$$

**Spanning:** Any vector of  $\text{Image}(T)$  looks like  $T(v)$  for some  $v \in V$ .

Write  $v = c_1 u_1 + \dots + c_n u_n + d_1 v_1 + \dots + d_r v_r$ . Then

$$\begin{aligned} T(v) &= c_1 T(u_1) + \dots + c_n T(u_n) + d_1 T(v_1) + \dots + d_r T(v_r) \\ &= d_1 T(v_1) + \dots + d_r T(v_r). \end{aligned}$$

**Lin. Independence:** Let  $b_1 T(v_1) + \dots + b_r T(v_r) = 0$ .

This implies that  $b_1 v_1 + \dots + b_r v_r \in \text{Null}(T)$ .

So  $b_1 v_1 + \dots + b_r v_r = a_1 u_1 + \dots + a_n u_n$  for some  $a_i \in \mathbb{R}$ .

Thus  $b_1 v_1 + \dots + b_r v_r - a_1 u_1 - \dots - a_n u_n = 0$ .

# Rank-Nullity Theorem

## Theorem

Let  $T : V \rightarrow W$  be a linear transformation, where  $\dim(V)$  is finite. Then  $\text{rank}(T) + \text{nullity}(T) = \dim(V)$ .

**Proof.** Start with a basis  $\{u_1, \dots, u_n\}$  of  $\text{Null}(T)$ . Extend this to a basis  $\{u_1, \dots, u_n, v_1, \dots, v_r\}$  of  $V$ . It is enough to prove that

$$\{T(v_1), \dots, T(v_r)\} \text{ is a basis of } \text{Image}(T).$$

**Spanning:** Any vector of  $\text{Image}(T)$  looks like  $T(v)$  for some  $v \in V$ .

Write  $v = c_1 u_1 + \dots + c_n u_n + d_1 v_1 + \dots + d_r v_r$ . Then

$$\begin{aligned} T(v) &= c_1 T(u_1) + \dots + c_n T(u_n) + d_1 T(v_1) + \dots + d_r T(v_r) \\ &= d_1 T(v_1) + \dots + d_r T(v_r). \end{aligned}$$

**Lin. Independence:** Let  $b_1 T(v_1) + \dots + b_r T(v_r) = 0$ .

This implies that  $b_1 v_1 + \dots + b_r v_r \in \text{Null}(T)$ .

So  $b_1 v_1 + \dots + b_r v_r = a_1 u_1 + \dots + a_n u_n$  for some  $a_i \in \mathbb{R}$ .

Thus  $b_1 v_1 + \dots + b_r v_r - a_1 u_1 - \dots - a_n u_n = 0$ .

Therefore  $b_1 = \dots = b_r = 0$ .

# Row and column spaces

## Definition

- Let  $A$  be an  $m \times n$  matrix over  $\mathbb{R}$ .

# Row and column spaces

## Definition

- Let  $A$  be an  $m \times n$  matrix over  $\mathbb{R}$ .
- The subspace of  $\mathbb{R}^m$  generated by all columns (column vectors) of  $A$  is called the **column space** of  $A$ .

# Row and column spaces

## Definition

- Let  $A$  be an  $m \times n$  matrix over  $\mathbb{R}$ .
- The subspace of  $\mathbb{R}^m$  generated by all columns (column vectors) of  $A$  is called the **column space** of  $A$ .
- The subspace of  $\mathbb{R}^n$  generated by all rows (row vectors) of  $A$  is called the **row space** of  $A$ .

# Row and column spaces

## Definition

- Let  $A$  be an  $m \times n$  matrix over  $\mathbb{R}$ .
- The subspace of  $\mathbb{R}^m$  generated by all columns (column vectors) of  $A$  is called the **column space** of  $A$ .
- The subspace of  $\mathbb{R}^n$  generated by all rows (row vectors) of  $A$  is called the **row space** of  $A$ .

## Example

- Let  $A = \begin{pmatrix} 1 & 2 & 3 & 4 \\ 5 & 6 & 7 & 8 \\ 9 & 10 & 11 & 12 \end{pmatrix}$ .



# Row and column spaces

## Definition

- Let  $A$  be an  $m \times n$  matrix over  $\mathbb{R}$ .
- The subspace of  $\mathbb{R}^m$  generated by all columns (column vectors) of  $A$  is called the **column space** of  $A$ .
- The subspace of  $\mathbb{R}^n$  generated by all rows (row vectors) of  $A$  is called the **row space** of  $A$ .

## Example

- Let  $A = \begin{pmatrix} 1 & 2 & 3 & 4 \\ 5 & 6 & 7 & 8 \\ 9 & 10 & 11 & 12 \end{pmatrix}$ .
- Column space of  $A$  is **Span**  $\left\{ \begin{pmatrix} 1 \\ 5 \\ 9 \end{pmatrix}, \begin{pmatrix} 2 \\ 6 \\ 10 \end{pmatrix}, \begin{pmatrix} 3 \\ 7 \\ 11 \end{pmatrix}, \begin{pmatrix} 4 \\ 8 \\ 12 \end{pmatrix} \right\}$ .

# Row and column spaces

## Definition

- Let  $A$  be an  $m \times n$  matrix over  $\mathbb{R}$ .
- The subspace of  $\mathbb{R}^m$  generated by all columns (column vectors) of  $A$  is called the **column space** of  $A$ .
- The subspace of  $\mathbb{R}^n$  generated by all rows (row vectors) of  $A$  is called the **row space** of  $A$ .

## Example

- Let  $A = \begin{pmatrix} 1 & 2 & 3 & 4 \\ 5 & 6 & 7 & 8 \\ 9 & 10 & 11 & 12 \end{pmatrix}$ .
- Column space of  $A$  is **Span**  $\left\{ \begin{pmatrix} 1 \\ 5 \\ 9 \end{pmatrix}, \begin{pmatrix} 2 \\ 6 \\ 10 \end{pmatrix}, \begin{pmatrix} 3 \\ 7 \\ 11 \end{pmatrix}, \begin{pmatrix} 4 \\ 8 \\ 12 \end{pmatrix} \right\}$ .
- Column space of  $A$  is a subspace of  $\mathbb{R}^3$ .

# Examples: Row and column spaces

## Example

- Let  $A = \begin{pmatrix} 1 & 2 & 3 & 4 \\ 5 & 6 & 7 & 8 \\ 9 & 10 & 11 & 12 \end{pmatrix}$ .

# Examples: Row and column spaces

## Example

- Let  $A = \begin{pmatrix} 1 & 2 & 3 & 4 \\ 5 & 6 & 7 & 8 \\ 9 & 10 & 11 & 12 \end{pmatrix}$ .
- Row space of  $A$  is **Span**  $\left\{ \begin{pmatrix} 1 \\ 2 \\ 3 \\ 4 \end{pmatrix}, \begin{pmatrix} 5 \\ 6 \\ 7 \\ 8 \end{pmatrix}, \begin{pmatrix} 9 \\ 10 \\ 11 \\ 12 \end{pmatrix} \right\}$ .

# Examples: Row and column spaces

## Example

- Let  $A = \begin{pmatrix} 1 & 2 & 3 & 4 \\ 5 & 6 & 7 & 8 \\ 9 & 10 & 11 & 12 \end{pmatrix}$ .
- Row space of  $A$  is  $\text{Span} \left\{ \begin{pmatrix} 1 \\ 2 \\ 3 \\ 4 \end{pmatrix}, \begin{pmatrix} 5 \\ 6 \\ 7 \\ 8 \end{pmatrix}, \begin{pmatrix} 9 \\ 10 \\ 11 \\ 12 \end{pmatrix} \right\}$ .
- Row space of  $A$  is a subspace of  $\mathbb{R}^4$ .

# Examples: Row and column spaces

## Example

- Let  $A = \begin{pmatrix} 1 & 2 & 3 & 4 \\ 5 & 6 & 7 & 8 \\ 9 & 10 & 11 & 12 \end{pmatrix}$ .
- Row space of  $A$  is  $\text{Span} \left\{ \begin{pmatrix} 1 \\ 2 \\ 3 \\ 4 \end{pmatrix}, \begin{pmatrix} 5 \\ 6 \\ 7 \\ 8 \end{pmatrix}, \begin{pmatrix} 9 \\ 10 \\ 11 \\ 12 \end{pmatrix} \right\}$ .
- Row space of  $A$  is a subspace of  $\mathbb{R}^4$ .

## Example

$$\text{If } A = \begin{pmatrix} 1 & 0 & 2 & 0 \\ 0 & 1 & 3 & 0 \\ 0 & 0 & 0 & 0 \end{pmatrix},$$

# Examples: Row and column spaces

## Example

- Let  $A = \begin{pmatrix} 1 & 2 & 3 & 4 \\ 5 & 6 & 7 & 8 \\ 9 & 10 & 11 & 12 \end{pmatrix}$ .
- Row space of  $A$  is  $\text{Span} \left\{ \begin{pmatrix} 1 \\ 2 \\ 3 \\ 4 \end{pmatrix}, \begin{pmatrix} 5 \\ 6 \\ 7 \\ 8 \end{pmatrix}, \begin{pmatrix} 9 \\ 10 \\ 11 \\ 12 \end{pmatrix} \right\}$ .
- Row space of  $A$  is a subspace of  $\mathbb{R}^4$ .

## Example

If  $A = \begin{pmatrix} 1 & 0 & 2 & 0 \\ 0 & 1 & 3 & 0 \\ 0 & 0 & 0 & 0 \end{pmatrix}$ , Column Sp. is  $\left\{ \begin{pmatrix} x_1 \\ x_2 \\ x_3 \end{pmatrix} \in \mathbb{R}^3 : x_3 = 0 \right\}$ .

# Row rank and column rank

## Definition

- Let  $A$  be an  $m \times n$  matrix over  $\mathbb{R}$ .



# Row rank and column rank

## Definition

- Let  $A$  be an  $m \times n$  matrix over  $\mathbb{R}$ .
- The dimension of the column space of  $A$  is called the **column rank** of  $A$ .

# Row rank and column rank

## Definition

- Let  $A$  be an  $m \times n$  matrix over  $\mathbb{R}$ .
- The dimension of the column space of  $A$  is called the **column rank** of  $A$ .
- The dimension of the row space of  $A$  is called the **row rank** of  $A$ .

# Row rank and column rank

## Definition

- Let  $A$  be an  $m \times n$  matrix over  $\mathbb{R}$ .
- The dimension of the column space of  $A$  is called the **column rank** of  $A$ .
- The dimension of the row space of  $A$  is called the **row rank** of  $A$ .

## Example

• Let  $A = \begin{pmatrix} 1 & 0 & 2 & 0 \\ 0 & 1 & 3 & 0 \\ 0 & 0 & 0 & 0 \end{pmatrix}$

# Row rank and column rank

## Definition

- Let  $A$  be an  $m \times n$  matrix over  $\mathbb{R}$ .
- The dimension of the column space of  $A$  is called the **column rank** of  $A$ .
- The dimension of the row space of  $A$  is called the **row rank** of  $A$ .

## Example

- Let  $A = \begin{pmatrix} 1 & 0 & 2 & 0 \\ 0 & 1 & 3 & 0 \\ 0 & 0 & 0 & 0 \end{pmatrix}$  and  $B = \begin{pmatrix} 1 & 1 & 1 & 1 \\ 0 & 2 & 1 & 1 \\ 0 & 0 & 3 & 1 \end{pmatrix}$ .

# Row rank and column rank

## Definition

- Let  $A$  be an  $m \times n$  matrix over  $\mathbb{R}$ .
- The dimension of the column space of  $A$  is called the **column rank** of  $A$ .
- The dimension of the row space of  $A$  is called the **row rank** of  $A$ .

## Example

- Let  $A = \begin{pmatrix} 1 & 0 & 2 & 0 \\ 0 & 1 & 3 & 0 \\ 0 & 0 & 0 & 0 \end{pmatrix}$  and  $B = \begin{pmatrix} 1 & 1 & 1 & 1 \\ 0 & 2 & 1 & 1 \\ 0 & 0 & 3 & 1 \end{pmatrix}$ .
- Column rank of  $A$  is 2.

# Row rank and column rank

## Definition

- Let  $A$  be an  $m \times n$  matrix over  $\mathbb{R}$ .
- The dimension of the column space of  $A$  is called the **column rank** of  $A$ .
- The dimension of the row space of  $A$  is called the **row rank** of  $A$ .

## Example

- Let  $A = \begin{pmatrix} 1 & 0 & 2 & 0 \\ 0 & 1 & 3 & 0 \\ 0 & 0 & 0 & 0 \end{pmatrix}$  and  $B = \begin{pmatrix} 1 & 1 & 1 & 1 \\ 0 & 2 & 1 & 1 \\ 0 & 0 & 3 & 1 \end{pmatrix}$ .
- Column rank of  $A$  is 2. Row rank of  $A$  is 2.

# Row rank and column rank

## Definition

- Let  $A$  be an  $m \times n$  matrix over  $\mathbb{R}$ .
- The dimension of the column space of  $A$  is called the **column rank** of  $A$ .
- The dimension of the row space of  $A$  is called the **row rank** of  $A$ .

## Example

- Let  $A = \begin{pmatrix} 1 & 0 & 2 & 0 \\ 0 & 1 & 3 & 0 \\ 0 & 0 & 0 & 0 \end{pmatrix}$  and  $B = \begin{pmatrix} 1 & 1 & 1 & 1 \\ 0 & 2 & 1 & 1 \\ 0 & 0 & 3 & 1 \end{pmatrix}$ .
- Column rank of  $A$  is 2. Row rank of  $A$  is 2.
- Column rank of  $B$  is 3.

# Row rank and column rank

## Definition

- Let  $A$  be an  $m \times n$  matrix over  $\mathbb{R}$ .
- The dimension of the column space of  $A$  is called the **column rank** of  $A$ .
- The dimension of the row space of  $A$  is called the **row rank** of  $A$ .

## Example

- Let  $A = \begin{pmatrix} 1 & 0 & 2 & 0 \\ 0 & 1 & 3 & 0 \\ 0 & 0 & 0 & 0 \end{pmatrix}$  and  $B = \begin{pmatrix} 1 & 1 & 1 & 1 \\ 0 & 2 & 1 & 1 \\ 0 & 0 & 3 & 1 \end{pmatrix}$ .
- Column rank of  $A$  is 2. Row rank of  $A$  is 2.
- Column rank of  $B$  is 3. Row rank of  $B$  is 3.



# Row rank and column rank

## Definition

- Let  $A$  be an  $m \times n$  matrix over  $\mathbb{R}$ .
- The dimension of the column space of  $A$  is called the **column rank** of  $A$ .
- The dimension of the row space of  $A$  is called the **row rank** of  $A$ .

## Example

- Let  $A = \begin{pmatrix} 1 & 0 & 2 & 0 \\ 0 & 1 & 3 & 0 \\ 0 & 0 & 0 & 0 \end{pmatrix}$  and  $B = \begin{pmatrix} 1 & 1 & 1 & 1 \\ 0 & 2 & 1 & 1 \\ 0 & 0 & 3 & 1 \end{pmatrix}$ .
- Column rank of  $A$  is 2. Row rank of  $A$  is 2.
- Column rank of  $B$  is 3. Row rank of  $B$  is 3.

As a consequence of Rank-Nullity Theorem, we will prove that for an arbitrary matrix  $D$ ,  $\text{row rank}(D) = \text{column rank}(D)$ .

For every matrix, row rank = column rank (an application of the Rank-Nullity Theorem)

### Theorem

*For an  $m \times n$  matrix  $A$  over  $\mathbb{R}$ ,*

For every matrix, row rank = column rank (an application of the Rank-Nullity Theorem)

### Theorem

*For an  $m \times n$  matrix  $A$  over  $\mathbb{R}$ ,  $\text{row rank}(A) = \text{column rank}(A)$ .*

# Some observations to prove: row rank = column rank

- 1 Consider  $A$  as a linear map  $A : \mathbb{R}^n \rightarrow \mathbb{R}^m$ .

# Some observations to prove: row rank = column rank

- 1 Consider  $A$  as a linear map  $A : \mathbb{R}^n \rightarrow \mathbb{R}^m$ .
- 2 An element of  $\text{Image}(A)$  looks like  $AX$  for some  $X \in \mathbb{R}^n$ .

# Some observations to prove: row rank = column rank

- 1 Consider  $A$  as a linear map  $A : \mathbb{R}^n \rightarrow \mathbb{R}^m$ .
- 2 An element of  $\text{Image}(A)$  looks like  $AX$  for some  $X \in \mathbb{R}^n$ .
- 3 But  $AX = x_1A_1 + \cdots + x_nA_n$ ,

# Some observations to prove: row rank = column rank

- 1 Consider  $A$  as a linear map  $A : \mathbb{R}^n \rightarrow \mathbb{R}^m$ .
- 2 An element of  $\text{Image}(A)$  looks like  $AX$  for some  $X \in \mathbb{R}^n$ .
- 3 But  $AX = x_1A_1 + \cdots + x_nA_n$ , where  $A_i$  is the  $i$ th column of  $A$ .

# Some observations to prove: row rank = column rank

- 1 Consider  $A$  as a linear map  $A : \mathbb{R}^n \rightarrow \mathbb{R}^m$ .
- 2 An element of  $\text{Image}(A)$  looks like  $AX$  for some  $X \in \mathbb{R}^n$ .
- 3 But  $AX = x_1A_1 + \cdots + x_nA_n$ , where  $A_i$  is the  $i$ th column of  $A$ .
- 4 Therefore  $\text{Image}(A)$



# Some observations to prove: row rank = column rank

- 1 Consider  $A$  as a linear map  $A : \mathbb{R}^n \rightarrow \mathbb{R}^m$ .
- 2 An element of  $\text{Image}(A)$  looks like  $AX$  for some  $X \in \mathbb{R}^n$ .
- 3 But  $AX = x_1A_1 + \cdots + x_nA_n$ , where  $A_i$  is the  $i$ th column of  $A$ .
- 4 Therefore  $\text{Image}(A) = \text{Span}\{A_1, \dots, A_n\}$

# Some observations to prove: row rank = column rank

- 1 Consider  $A$  as a linear map  $A : \mathbb{R}^n \rightarrow \mathbb{R}^m$ .
- 2 An element of  $\text{Image}(A)$  looks like  $AX$  for some  $X \in \mathbb{R}^n$ .
- 3 But  $AX = x_1A_1 + \cdots + x_nA_n$ , where  $A_i$  is the  $i$ th column of  $A$ .
- 4 Therefore  $\text{Image}(A) = \text{Span}\{A_1, \dots, A_n\} = \text{Column Sp.}(A)$ .

# Some observations to prove: row rank = column rank

- 1 Consider  $A$  as a linear map  $A : \mathbb{R}^n \rightarrow \mathbb{R}^m$ .
- 2 An element of  $\text{Image}(A)$  looks like  $AX$  for some  $X \in \mathbb{R}^n$ .
- 3 But  $AX = x_1A_1 + \cdots + x_nA_n$ , where  $A_i$  is the  $i$ th column of  $A$ .
- 4 Therefore  $\text{Image}(A) = \text{Span}\{A_1, \dots, A_n\} = \text{Column Sp.}(A)$ .
- 5 Hence  $\text{rank}(A) := \dim(\text{Image}(A))$

# Some observations to prove: row rank = column rank

- 1 Consider  $A$  as a linear map  $A : \mathbb{R}^n \rightarrow \mathbb{R}^m$ .
- 2 An element of  $\text{Image}(A)$  looks like  $AX$  for some  $X \in \mathbb{R}^n$ .
- 3 But  $AX = x_1A_1 + \cdots + x_nA_n$ , where  $A_i$  is the  $i$ th column of  $A$ .
- 4 Therefore  $\text{Image}(A) = \text{Span}\{A_1, \dots, A_n\} = \text{Column Sp.}(A)$ .
- 5 Hence  $\text{rank}(A) := \dim(\text{Image}(A)) = \text{column rank}(A)$ .

# Some observations to prove: row rank = column rank

- 1 Consider  $A$  as a linear map  $A : \mathbb{R}^n \rightarrow \mathbb{R}^m$ .
- 2 An element of  $\text{Image}(A)$  looks like  $AX$  for some  $X \in \mathbb{R}^n$ .
- 3 But  $AX = x_1 A_1 + \cdots + x_n A_n$ , where  $A_i$  is the  $i$ th column of  $A$ .
- 4 Therefore  $\text{Image}(A) = \text{Span}\{A_1, \dots, A_n\} = \text{Column Sp.}(A)$ .
- 5 Hence  $\text{rank}(A) := \dim(\text{Image}(A)) = \text{column rank}(A)$ .
- 6 **Rank-nullity theorem:**  $\text{rank}(A) + \text{nullity}(A) = \dim(\mathbb{R}^n)$ .

# Some observations to prove: row rank = column rank

- 1 Consider  $A$  as a linear map  $A : \mathbb{R}^n \rightarrow \mathbb{R}^m$ .
- 2 An element of  $\text{Image}(A)$  looks like  $AX$  for some  $X \in \mathbb{R}^n$ .
- 3 But  $AX = x_1 A_1 + \cdots + x_n A_n$ , where  $A_i$  is the  $i$ th column of  $A$ .
- 4 Therefore  $\text{Image}(A) = \text{Span}\{A_1, \dots, A_n\} = \text{Column Sp.}(A)$ .
- 5 Hence  $\text{rank}(A) := \dim(\text{Image}(A)) = \text{column rank}(A)$ .
- 6 **Rank-nullity theorem:**  $\text{rank}(A) + \text{nullity}(A) = \dim(\mathbb{R}^n)$ .
- 7 Therefore  $\text{column rank}(A) = n - \text{nullity}(A)$ .

# Some observations to prove: row rank = column rank

- 1 Consider  $A$  as a linear map  $A : \mathbb{R}^n \rightarrow \mathbb{R}^m$ .
- 2 An element of  $\text{Image}(A)$  looks like  $AX$  for some  $X \in \mathbb{R}^n$ .
- 3 But  $AX = x_1 A_1 + \cdots + x_n A_n$ , where  $A_i$  is the  $i$ th column of  $A$ .
- 4 Therefore  $\text{Image}(A) = \text{Span}\{A_1, \dots, A_n\} = \text{Column Sp.}(A)$ .
- 5 Hence  $\text{rank}(A) := \dim(\text{Image}(A)) = \text{column rank}(A)$ .
- 6 **Rank-nullity theorem:**  $\text{rank}(A) + \text{nullity}(A) = \dim(\mathbb{R}^n)$ .
- 7 Therefore  $\text{column rank}(A) = n - \text{nullity}(A)$ .

So it is enough to show that

$$\text{row rank}(A) = n - \text{nullity}(A).$$

# Elementary row operations preserve row space, hence rank

## Theorem

*Let  $A$  and  $B$  be row equivalent.*



# Elementary row operations preserve row space, hence rank

## Theorem

*Let  $A$  and  $B$  be row equivalent. Then  $A$  and  $B$  have the same row space.*

# Elementary row operations preserve row space, hence rank

## Theorem

*Let  $A$  and  $B$  be row equivalent. Then  $A$  and  $B$  have the same row space. In particular,  $\text{row rank}(A) = \text{row rank}(B)$ .*

**Proof.** Note that  $A$  and  $B$  have the same order (say,  $m \times n$ ).  
Let  $R_1, \dots, R_m \in \mathbb{R}^n$  be the row vectors of  $A$ .

# Elementary row operations preserve row space, hence rank

## Theorem

*Let  $A$  and  $B$  be row equivalent. Then  $A$  and  $B$  have the same row space. In particular,  $\text{row rank}(A) = \text{row rank}(B)$ .*

**Proof.** Note that  $A$  and  $B$  have the same order (say,  $m \times n$ ).

Let  $R_1, \dots, R_m \in \mathbb{R}^n$  be the row vectors of  $A$ . We observe that the elementary row operations preserve the row space:

# Elementary row operations preserve row space, hence rank

## Theorem

*Let  $A$  and  $B$  be row equivalent. Then  $A$  and  $B$  have the same row space. In particular,  $\text{row rank}(A) = \text{row rank}(B)$ .*

**Proof.** Note that  $A$  and  $B$  have the same order (say,  $m \times n$ ).

Let  $R_1, \dots, R_m \in \mathbb{R}^n$  be the row vectors of  $A$ . We observe that the elementary row operations preserve the row space:

- 1 Effect of the **1st type** elementary row operation, e.g.,

# Elementary row operations preserve row space, hence rank

## Theorem

*Let  $A$  and  $B$  be row equivalent. Then  $A$  and  $B$  have the same row space. In particular,  $\text{row rank}(A) = \text{row rank}(B)$ .*

**Proof.** Note that  $A$  and  $B$  have the same order (say,  $m \times n$ ).

Let  $R_1, \dots, R_m \in \mathbb{R}^n$  be the row vectors of  $A$ . We observe that the elementary row operations preserve the row space:

- 1 Effect of the **1st type** elementary row operation, e.g.,  
 $\text{Span}\{\mathbf{R}_1, \mathbf{R}_2, R_3, \dots, R_m\} = \text{Span}\{\mathbf{R}_2, \mathbf{R}_1, R_3, \dots, R_m\}.$

# Elementary row operations preserve row space, hence rank

## Theorem

*Let  $A$  and  $B$  be row equivalent. Then  $A$  and  $B$  have the same row space. In particular,  $\text{row rank}(A) = \text{row rank}(B)$ .*

**Proof.** Note that  $A$  and  $B$  have the same order (say,  $m \times n$ ).

Let  $R_1, \dots, R_m \in \mathbb{R}^n$  be the row vectors of  $A$ . We observe that the elementary row operations preserve the row space:

- 1 Effect of the **1st type** elementary row operation, e.g.,  
 $\text{Span}\{\mathbf{R}_1, \mathbf{R}_2, R_3, \dots, R_m\} = \text{Span}\{\mathbf{R}_2, \mathbf{R}_1, R_3, \dots, R_m\}.$
- 2 Effect of the **2nd type** elementary row operation, e.g.,

# Elementary row operations preserve row space, hence rank

## Theorem

*Let  $A$  and  $B$  be row equivalent. Then  $A$  and  $B$  have the same row space. In particular,  $\text{row rank}(A) = \text{row rank}(B)$ .*

**Proof.** Note that  $A$  and  $B$  have the same order (say,  $m \times n$ ).

Let  $R_1, \dots, R_m \in \mathbb{R}^n$  be the row vectors of  $A$ . We observe that the elementary row operations preserve the row space:

- 1 Effect of the **1st type** elementary row operation, e.g.,  
 $\text{Span}\{\mathbf{R}_1, \mathbf{R}_2, R_3, \dots, R_m\} = \text{Span}\{\mathbf{R}_2, \mathbf{R}_1, R_3, \dots, R_m\}.$
- 2 Effect of the **2nd type** elementary row operation, e.g.,  
 $\text{Span}\{R_1, \mathbf{R}_2, R_3, \dots, R_m\} = \text{Span}\{R_1, c \cdot \mathbf{R}_2, R_3, \dots, R_m\},$  where  $c \neq 0$  (important!).

# Elementary row operations preserve row space, hence rank

## Theorem

*Let  $A$  and  $B$  be row equivalent. Then  $A$  and  $B$  have the same row space. In particular,  $\text{row rank}(A) = \text{row rank}(B)$ .*

**Proof.** Note that  $A$  and  $B$  have the same order (say,  $m \times n$ ).

Let  $R_1, \dots, R_m \in \mathbb{R}^n$  be the row vectors of  $A$ . We observe that the elementary row operations preserve the row space:

- 1 Effect of the **1st type** elementary row operation, e.g.,  
 $\text{Span}\{\mathbf{R}_1, \mathbf{R}_2, R_3, \dots, R_m\} = \text{Span}\{\mathbf{R}_2, \mathbf{R}_1, R_3, \dots, R_m\}.$
- 2 Effect of the **2nd type** elementary row operation, e.g.,  
 $\text{Span}\{R_1, \mathbf{R}_2, R_3, \dots, R_m\} = \text{Span}\{R_1, c \cdot \mathbf{R}_2, R_3, \dots, R_m\},$  where  $c \neq 0$  (important!).
- 3 Effect of the **3rd type** elementary row operation, e.g.,



# Elementary row operations preserve row space, hence rank

## Theorem

*Let  $A$  and  $B$  be row equivalent. Then  $A$  and  $B$  have the same row space. In particular,  $\text{row rank}(A) = \text{row rank}(B)$ .*

**Proof.** Note that  $A$  and  $B$  have the same order (say,  $m \times n$ ).

Let  $R_1, \dots, R_m \in \mathbb{R}^n$  be the row vectors of  $A$ . We observe that the elementary row operations preserve the row space:

- ① Effect of the **1st type** elementary row operation, e.g.,  
 $\text{Span}\{\mathbf{R}_1, \mathbf{R}_2, R_3, \dots, R_m\} = \text{Span}\{\mathbf{R}_2, \mathbf{R}_1, R_3, \dots, R_m\}.$
- ② Effect of the **2nd type** elementary row operation, e.g.,  
 $\text{Span}\{R_1, \mathbf{R}_2, R_3, \dots, R_m\} = \text{Span}\{R_1, c \cdot \mathbf{R}_2, R_3, \dots, R_m\},$  where  $c \neq 0$  (important!).
- ③ Effect of the **3rd type** elementary row operation, e.g.,  
 $\text{Span}\{R_1, \mathbf{R}_2, R_3, \dots, R_m\} = \text{Span}\{R_1, \mathbf{R}_2 - c \cdot \mathbf{R}_1, R_3, \dots, R_m\},$  where  $c \in \mathbb{R}.$

# Elementary row operations preserve the nullity of a matrix

- 1 Let  $A$  and  $B$  be row equivalent matrices over  $\mathbb{R}$ .

# Elementary row operations preserve the nullity of a matrix

- 1 Let  $A$  and  $B$  be row equivalent matrices over  $\mathbb{R}$ .
- 2 Then  $AX = 0$  and  $BX = 0$  have the same solution set,

# Elementary row operations preserve the nullity of a matrix

- 1 Let  $A$  and  $B$  be row equivalent matrices over  $\mathbb{R}$ .
- 2 Then  $AX = 0$  and  $BX = 0$  have the same solution set, i.e.,  $\text{Null}(A) = \text{Null}(B)$ .

# Elementary row operations preserve the nullity of a matrix

- 1 Let  $A$  and  $B$  be row equivalent matrices over  $\mathbb{R}$ .
- 2 Then  $AX = 0$  and  $BX = 0$  have the same solution set, i.e.,  $\text{Null}(A) = \text{Null}(B)$ .
- 3 Therefore  $\text{nullity}(A) = \text{nullity}(B)$ .

# Proof of “row rank( $A$ ) = $n$ - nullity( $A$ )”

- 1 Let  $A$  be an  $m \times n$  matrix over  $\mathbb{R}$ .

# Proof of “ $\text{row rank}(A) = n - \text{nullity}(A)$ ”

- 1 Let  $A$  be an  $m \times n$  matrix over  $\mathbb{R}$ .
- 2  $A$  is row-equivalent to a **row-reduced echelon** matrix  $B$ .

# Proof of “ $\text{row rank}(A) = n - \text{nullity}(A)$ ”

- 1 Let  $A$  be an  $m \times n$  matrix over  $\mathbb{R}$ .
- 2  $A$  is row-equivalent to a **row-reduced echelon** matrix  $B$ .
- 3 Since  $\text{row rank}(A) = \text{row rank}(B)$  and  $\text{nullity}(A) = \text{nullity}(B)$ ,



# Proof of “ $\text{row rank}(A) = n - \text{nullity}(A)$ ”

- 1 Let  $A$  be an  $m \times n$  matrix over  $\mathbb{R}$ .
- 2  $A$  is row-equivalent to a **row-reduced echelon** matrix  $B$ .
- 3 Since  $\text{row rank}(A) = \text{row rank}(B)$  and  $\text{nullity}(A) = \text{nullity}(B)$ , it is enough to prove that

$$\text{row rank}(B) + \text{nullity}(B) = n.$$

# Proof of “ $\text{row rank}(A) = n - \text{nullity}(A)$ ”

- 1 Let  $A$  be an  $m \times n$  matrix over  $\mathbb{R}$ .
- 2  $A$  is row-equivalent to a **row-reduced echelon** matrix  $B$ .
- 3 Since  $\text{row rank}(A) = \text{row rank}(B)$  and  $\text{nullity}(A) = \text{nullity}(B)$ , it is enough to prove that

$$\text{row rank}(B) + \text{nullity}(B) = n.$$

- 4 We will study some examples to observe this inequality.

# Proof of “ $\text{row rank}(A) = n - \text{nullity}(A)$ ”

- 1 Let  $A$  be an  $m \times n$  matrix over  $\mathbb{R}$ .
- 2  $A$  is row-equivalent to a **row-reduced echelon** matrix  $B$ .
- 3 Since  $\text{row rank}(A) = \text{row rank}(B)$  and  $\text{nullity}(A) = \text{nullity}(B)$ , it is enough to prove that

$$\text{row rank}(B) + \text{nullity}(B) = n.$$

- 4 We will study some examples to observe this inequality. But I will leave it as an exercise to verify this inequality in the general situation.

For a row-reduced echelon  $B$ ,  
 $\text{row rank}(B) = n - \text{nullity}(B)$

Consider a row-reduced echelon matrix  $B = \begin{pmatrix} \textcolor{red}{1} & 0 & \frac{3}{5} & \frac{7}{5} \\ 0 & \textcolor{red}{1} & -\frac{1}{5} & \frac{1}{5} \\ 0 & 0 & 0 & 0 \end{pmatrix}$ .

For a row-reduced echelon  $B$ ,  
 $\text{row rank}(B) = n - \text{nullity}(B)$

Consider a row-reduced echelon matrix  $B = \begin{pmatrix} \color{red}{1} & 0 & \frac{3}{5} & \frac{7}{5} \\ 0 & \color{red}{1} & -\frac{1}{5} & \frac{1}{5} \\ 0 & 0 & 0 & 0 \end{pmatrix}$ .

- We shall show that  $\text{row rank}(B) + \text{nullity}(B) = 4$ .

For a row-reduced echelon  $B$ ,  
 $\text{row rank}(B) = n - \text{nullity}(B)$

Consider a row-reduced echelon matrix  $B = \begin{pmatrix} \textcolor{red}{1} & 0 & \frac{3}{5} & \frac{7}{5} \\ 0 & \textcolor{red}{1} & -\frac{1}{5} & \frac{1}{5} \\ 0 & 0 & 0 & 0 \end{pmatrix}$ .

- We shall show that  $\text{row rank}(B) + \text{nullity}(B) = 4$ .
- It can be observed that  $\text{row rank}(B)$  is the number of non-zero rows of  $B$ ,

For a row-reduced echelon  $B$ ,  
 $\text{row rank}(B) = n - \text{nullity}(B)$

Consider a row-reduced echelon matrix  $B = \begin{pmatrix} \textcolor{red}{1} & 0 & \frac{3}{5} & \frac{7}{5} \\ 0 & \textcolor{red}{1} & -\frac{1}{5} & \frac{1}{5} \\ 0 & 0 & 0 & 0 \end{pmatrix}$ .

- We shall show that  $\text{row rank}(B) + \text{nullity}(B) = 4$ .
- It can be observed that  $\text{row rank}(B)$  is the number of non-zero rows of  $B$ , i.e., the number of pivots of  $B$ .

For a row-reduced echelon  $B$ ,  
 $\text{row rank}(B) = n - \text{nullity}(B)$

Consider a row-reduced echelon matrix  $B = \begin{pmatrix} \textcolor{red}{1} & 0 & \frac{3}{5} & \frac{7}{5} \\ 0 & \textcolor{red}{1} & -\frac{1}{5} & \frac{1}{5} \\ 0 & 0 & 0 & 0 \end{pmatrix}$ .

- We shall show that  $\text{row rank}(B) + \text{nullity}(B) = 4$ .
- It can be observed that  $\text{row rank}(B)$  is the number of non-zero rows of  $B$ , i.e., the number of pivots of  $B$ .  
So  $\text{row rank}(B) = 2$ .



For a row-reduced echelon  $B$ ,  
 $\text{row rank}(B) = n - \text{nullity}(B)$

Consider a row-reduced echelon matrix  $B = \begin{pmatrix} \textcolor{red}{1} & 0 & \frac{3}{5} & \frac{7}{5} \\ 0 & \textcolor{red}{1} & -\frac{1}{5} & \frac{1}{5} \\ 0 & 0 & 0 & 0 \end{pmatrix}$ .

- We shall show that  $\text{row rank}(B) + \text{nullity}(B) = 4$ .
- It can be observed that  $\text{row rank}(B)$  is the number of non-zero rows of  $B$ , i.e., the number of pivots of  $B$ .  
So  $\text{row rank}(B) = 2$ .
- Consider the system  $BX = 0$ .

For a row-reduced echelon  $B$ ,  
 $\text{row rank}(B) = n - \text{nullity}(B)$

Consider a row-reduced echelon matrix  $B = \begin{pmatrix} 1 & 0 & \frac{3}{5} & \frac{7}{5} \\ 0 & 1 & -\frac{1}{5} & \frac{1}{5} \\ 0 & 0 & 0 & 0 \end{pmatrix}$ .

- We shall show that  $\text{row rank}(B) + \text{nullity}(B) = 4$ .
- It can be observed that  $\text{row rank}(B)$  is the number of non-zero rows of  $B$ , i.e., the number of pivots of  $B$ .  
So  $\text{row rank}(B) = 2$ .
- Consider the system  $BX = 0$ . The pivot variables are  $x_1, x_2$ .

# For a row-reduced echelon $B$ , $\text{row rank}(B) = n - \text{nullity}(B)$

Consider a row-reduced echelon matrix  $B = \begin{pmatrix} 1 & 0 & \frac{3}{5} & \frac{7}{5} \\ 0 & 1 & -\frac{1}{5} & \frac{1}{5} \\ 0 & 0 & 0 & 0 \end{pmatrix}$ .

- We shall show that  $\text{row rank}(B) + \text{nullity}(B) = 4$ .
- It can be observed that  $\text{row rank}(B)$  is the number of non-zero rows of  $B$ , i.e., the number of pivots of  $B$ .  
So  $\text{row rank}(B) = 2$ .
- Consider the system  $BX = 0$ . The pivot variables are  $x_1, x_2$ . The free variables are  $x_3$  and  $x_4$ .

# For a row-reduced echelon $B$ , $\text{row rank}(B) = n - \text{nullity}(B)$

Consider a row-reduced echelon matrix  $B = \begin{pmatrix} \color{red}{1} & 0 & \frac{3}{5} & \frac{7}{5} \\ 0 & \color{red}{1} & -\frac{1}{5} & \frac{1}{5} \\ 0 & 0 & 0 & 0 \end{pmatrix}$ .

- We shall show that  $\text{row rank}(B) + \text{nullity}(B) = 4$ .
- It can be observed that  $\text{row rank}(B)$  is the number of non-zero rows of  $B$ , i.e., the number of pivots of  $B$ .  
So  $\text{row rank}(B) = 2$ .
- Consider the system  $BX = 0$ . The pivot variables are  $x_1, x_2$ . The free variables are  $x_3$  and  $x_4$ . The system  $BX = 0$  is

$$\begin{aligned} \color{red}{x_1} + \frac{3}{5}x_3 + \frac{7}{5}x_4 &= 0 \\ \color{red}{x_2} - \frac{1}{5}x_3 + \frac{1}{5}x_4 &= 0 \end{aligned}$$

# For a row-reduced echelon $B$ , $\text{row rank}(B) = n - \text{nullity}(B)$

Consider a row-reduced echelon matrix  $B = \begin{pmatrix} 1 & 0 & \frac{3}{5} & \frac{7}{5} \\ 0 & 1 & -\frac{1}{5} & \frac{1}{5} \\ 0 & 0 & 0 & 0 \end{pmatrix}$ .

- We shall show that  $\text{row rank}(B) + \text{nullity}(B) = 4$ .
- It can be observed that  $\text{row rank}(B)$  is the number of non-zero rows of  $B$ , i.e., the number of pivots of  $B$ .  
So  $\text{row rank}(B) = 2$ .
- Consider the system  $BX = 0$ . The pivot variables are  $x_1, x_2$ . The free variables are  $x_3$  and  $x_4$ . The system  $BX = 0$  is

$$\begin{aligned} x_1 + \frac{3}{5}x_3 + \frac{7}{5}x_4 &= 0 \\ x_2 - \frac{1}{5}x_3 + \frac{1}{5}x_4 &= 0 \end{aligned}$$

- We claim that  $\text{nullity}(B)$  is the number of free variables, because  
...

# How to solve $BX = 0$ when $B$ is row-reduced echelon?

- Consider  $B = \begin{pmatrix} 1 & 0 & \frac{3}{5} & \frac{7}{5} \\ 0 & 1 & -\frac{1}{5} & \frac{1}{5} \\ 0 & 0 & 0 & 0 \end{pmatrix}$ , a row-reduced echelon matrix.

# How to solve $BX = 0$ when $B$ is row-reduced echelon?

- Consider  $B = \begin{pmatrix} 1 & 0 & \frac{3}{5} & \frac{7}{5} \\ 0 & 1 & -\frac{1}{5} & \frac{1}{5} \\ 0 & 0 & 0 & 0 \end{pmatrix}$ , a row-reduced echelon matrix.
- The corresponding homogeneous system can be written as

$$x_1 + \frac{3}{5}x_3 + \frac{7}{5}x_4 = 0$$

$$x_2 - \frac{1}{5}x_3 + \frac{1}{5}x_4 = 0$$

# How to solve $BX = 0$ when $B$ is row-reduced echelon?

- Consider  $B = \begin{pmatrix} \textcolor{red}{1} & 0 & \frac{3}{5} & \frac{7}{5} \\ 0 & \textcolor{red}{1} & -\frac{1}{5} & \frac{1}{5} \\ 0 & 0 & 0 & 0 \end{pmatrix}$ , a row-reduced echelon matrix.
- The corresponding homogeneous system can be written as

$$\textcolor{red}{x}_1 + \frac{3}{5}x_3 + \frac{7}{5}x_4 = 0$$

$$\textcolor{red}{x}_2 - \frac{1}{5}x_3 + \frac{1}{5}x_4 = 0$$

- The solutions of the system are given by



# How to solve $BX = 0$ when $B$ is row-reduced echelon?

- Consider  $B = \begin{pmatrix} 1 & 0 & \frac{3}{5} & \frac{7}{5} \\ 0 & 1 & -\frac{1}{5} & \frac{1}{5} \\ 0 & 0 & 0 & 0 \end{pmatrix}$ , a row-reduced echelon matrix.
- The corresponding homogeneous system can be written as

$$x_1 + \frac{3}{5}x_3 + \frac{7}{5}x_4 = 0$$

$$x_2 - \frac{1}{5}x_3 + \frac{1}{5}x_4 = 0$$

- The solutions of the system are given by

$$\begin{pmatrix} x_1 \\ x_2 \\ x_3 \\ x_4 \end{pmatrix}$$

# How to solve $BX = 0$ when $B$ is row-reduced echelon?

- Consider  $B = \begin{pmatrix} 1 & 0 & \frac{3}{5} & \frac{7}{5} \\ 0 & 1 & -\frac{1}{5} & \frac{1}{5} \\ 0 & 0 & 0 & 0 \end{pmatrix}$ , a row-reduced echelon matrix.
- The corresponding homogeneous system can be written as

$$x_1 + \frac{3}{5}x_3 + \frac{7}{5}x_4 = 0$$

$$x_2 - \frac{1}{5}x_3 + \frac{1}{5}x_4 = 0$$

- The solutions of the system are given by

$$\begin{pmatrix} x_1 \\ x_2 \\ x_3 \\ x_4 \end{pmatrix} = \begin{pmatrix} -\frac{3}{5}x_3 - \frac{7}{5}x_4 \\ \frac{1}{5}x_3 - \frac{1}{5}x_4 \\ x_3 \\ x_4 \end{pmatrix}$$

# How to solve $BX = 0$ when $B$ is row-reduced echelon?

- Consider  $B = \begin{pmatrix} 1 & 0 & \frac{3}{5} & \frac{7}{5} \\ 0 & 1 & -\frac{1}{5} & \frac{1}{5} \\ 0 & 0 & 0 & 0 \end{pmatrix}$ , a row-reduced echelon matrix.
- The corresponding homogeneous system can be written as

$$x_1 + \frac{3}{5}x_3 + \frac{7}{5}x_4 = 0$$

$$x_2 - \frac{1}{5}x_3 + \frac{1}{5}x_4 = 0$$

- The solutions of the system are given by

$$\begin{pmatrix} x_1 \\ x_2 \\ x_3 \\ x_4 \end{pmatrix} = \begin{pmatrix} -\frac{3}{5}x_3 - \frac{7}{5}x_4 \\ \frac{1}{5}x_3 - \frac{1}{5}x_4 \\ x_3 \\ x_4 \end{pmatrix} = x_3 \begin{pmatrix} -3/5 \\ 1/5 \\ 1 \\ 0 \end{pmatrix} + x_4 \begin{pmatrix} -7/5 \\ -1/5 \\ 0 \\ 1 \end{pmatrix}$$

# How to solve $BX = 0$ when $B$ is row-reduced echelon?

- Consider  $B = \begin{pmatrix} 1 & 0 & \frac{3}{5} & \frac{7}{5} \\ 0 & 1 & -\frac{1}{5} & \frac{1}{5} \\ 0 & 0 & 0 & 0 \end{pmatrix}$ , a row-reduced echelon matrix.
- The corresponding homogeneous system can be written as

$$x_1 + \frac{3}{5}x_3 + \frac{7}{5}x_4 = 0$$

$$x_2 - \frac{1}{5}x_3 + \frac{1}{5}x_4 = 0$$

- The solutions of the system are given by

$$\begin{pmatrix} x_1 \\ x_2 \\ x_3 \\ x_4 \end{pmatrix} = \begin{pmatrix} -\frac{3}{5}x_3 - \frac{7}{5}x_4 \\ \frac{1}{5}x_3 - \frac{1}{5}x_4 \\ x_3 \\ x_4 \end{pmatrix} = x_3 \begin{pmatrix} -3/5 \\ 1/5 \\ 1 \\ 0 \end{pmatrix} + x_4 \begin{pmatrix} -7/5 \\ -1/5 \\ 0 \\ 1 \end{pmatrix}$$

- $\text{nullity}(B) = \dim(\text{Null}(B))$

# How to solve $BX = 0$ when $B$ is row-reduced echelon?

- Consider  $B = \begin{pmatrix} 1 & 0 & \frac{3}{5} & \frac{7}{5} \\ 0 & 1 & -\frac{1}{5} & \frac{1}{5} \\ 0 & 0 & 0 & 0 \end{pmatrix}$ , a row-reduced echelon matrix.
- The corresponding homogeneous system can be written as

$$x_1 + \frac{3}{5}x_3 + \frac{7}{5}x_4 = 0$$

$$x_2 - \frac{1}{5}x_3 + \frac{1}{5}x_4 = 0$$

- The solutions of the system are given by

$$\begin{pmatrix} x_1 \\ x_2 \\ x_3 \\ x_4 \end{pmatrix} = \begin{pmatrix} -\frac{3}{5}x_3 - \frac{7}{5}x_4 \\ \frac{1}{5}x_3 - \frac{1}{5}x_4 \\ x_3 \\ x_4 \end{pmatrix} = x_3 \begin{pmatrix} -3/5 \\ 1/5 \\ 1 \\ 0 \end{pmatrix} + x_4 \begin{pmatrix} -7/5 \\ -1/5 \\ 0 \\ 1 \end{pmatrix}$$

- $\text{nullity}(B) = \dim(\text{Null}(B)) = \text{the number of free variables.}$

# Isomorphism of vector spaces

## Definition

A linear map  $T : V \rightarrow W$  is said to be an **isomorphism**

# Isomorphism of vector spaces

## Definition

A linear map  $T : V \rightarrow W$  is said to be an **isomorphism** if there is a linear map  $S : W \rightarrow V$  such that

# Isomorphism of vector spaces

## Definition

A linear map  $T : V \rightarrow W$  is said to be an **isomorphism** if there is a linear map  $S : W \rightarrow V$  such that  $S \circ T = 1_V : V \rightarrow V$  (identity map)



# Isomorphism of vector spaces

## Definition

A linear map  $T : V \rightarrow W$  is said to be an **isomorphism** if there is a linear map  $S : W \rightarrow V$  such that  $S \circ T = 1_V : V \rightarrow V$  (identity map) and  $T \circ S = 1_W : W \rightarrow W$ .

# Isomorphism of vector spaces

## Definition

A linear map  $T : V \rightarrow W$  is said to be an **isomorphism** if there is a linear map  $S : W \rightarrow V$  such that

$S \circ T = 1_V : V \rightarrow V$  (identity map) and  $T \circ S = 1_W : W \rightarrow W$ .

If  $T : V \rightarrow W$  is an isomorphism,

# Isomorphism of vector spaces

## Definition

A linear map  $T : V \rightarrow W$  is said to be an **isomorphism** if there is a linear map  $S : W \rightarrow V$  such that

$S \circ T = 1_V : V \rightarrow V$  (identity map) and  $T \circ S = 1_W : W \rightarrow W$ .

If  $T : V \rightarrow W$  is an isomorphism, we say that  $V$  and  $W$  are isomorphic, and we write  $V \cong W$ .

# Isomorphism of vector spaces

## Definition

A linear map  $T : V \rightarrow W$  is said to be an **isomorphism** if there is a linear map  $S : W \rightarrow V$  such that

$S \circ T = 1_V : V \rightarrow V$  (identity map) and  $T \circ S = 1_W : W \rightarrow W$ .

If  $T : V \rightarrow W$  is an isomorphism, we say that  $V$  and  $W$  are isomorphic, and we write  $V \cong W$ .

## Example

Let  $A$  be an  $n \times n$  matrix over  $\mathbb{R}$ .

# Isomorphism of vector spaces

## Definition

A linear map  $T : V \rightarrow W$  is said to be an **isomorphism** if there is a linear map  $S : W \rightarrow V$  such that

$S \circ T = 1_V : V \rightarrow V$  (identity map) and  $T \circ S = 1_W : W \rightarrow W$ .

If  $T : V \rightarrow W$  is an isomorphism, we say that  $V$  and  $W$  are isomorphic, and we write  $V \cong W$ .

## Example

Let  $A$  be an  $n \times n$  matrix over  $\mathbb{R}$ . Consider  $A : \mathbb{R}^n \rightarrow \mathbb{R}^n$  as a linear map.

# Isomorphism of vector spaces

## Definition

A linear map  $T : V \rightarrow W$  is said to be an **isomorphism** if there is a linear map  $S : W \rightarrow V$  such that

$S \circ T = 1_V : V \rightarrow V$  (identity map) and  $T \circ S = 1_W : W \rightarrow W$ .

If  $T : V \rightarrow W$  is an isomorphism, we say that  $V$  and  $W$  are isomorphic, and we write  $V \cong W$ .

## Example

Let  $A$  be an  $n \times n$  matrix over  $\mathbb{R}$ . Consider  $A : \mathbb{R}^n \rightarrow \mathbb{R}^n$  as a linear map. When is it an isomorphism?

# Isomorphism of vector spaces

## Definition

A linear map  $T : V \rightarrow W$  is said to be an **isomorphism** if there is a linear map  $S : W \rightarrow V$  such that

$S \circ T = 1_V : V \rightarrow V$  (identity map) and  $T \circ S = 1_W : W \rightarrow W$ .

If  $T : V \rightarrow W$  is an isomorphism, we say that  $V$  and  $W$  are isomorphic, and we write  $V \cong W$ .

## Example

Let  $A$  be an  $n \times n$  matrix over  $\mathbb{R}$ . Consider  $A : \mathbb{R}^n \rightarrow \mathbb{R}^n$  as a linear map. When is it an isomorphism?

**Answer:** When there is an inverse linear map  $B : \mathbb{R}^n \rightarrow \mathbb{R}^n$  such that  $A \circ B = 1_{\mathbb{R}^n}$  and  $B \circ A = 1_{\mathbb{R}^n}$ ,

# Isomorphism of vector spaces

## Definition

A linear map  $T : V \rightarrow W$  is said to be an **isomorphism** if there is a linear map  $S : W \rightarrow V$  such that

$S \circ T = 1_V : V \rightarrow V$  (identity map) and  $T \circ S = 1_W : W \rightarrow W$ .

If  $T : V \rightarrow W$  is an isomorphism, we say that  $V$  and  $W$  are isomorphic, and we write  $V \cong W$ .

## Example

Let  $A$  be an  $n \times n$  matrix over  $\mathbb{R}$ . Consider  $A : \mathbb{R}^n \rightarrow \mathbb{R}^n$  as a linear map. When is it an isomorphism?

**Answer:** When there is an inverse linear map  $B : \mathbb{R}^n \rightarrow \mathbb{R}^n$  such that  $A \circ B = 1_{\mathbb{R}^n}$  and  $B \circ A = 1_{\mathbb{R}^n}$ , i.e., when there is an  $n \times n$  matrix  $B$  over  $\mathbb{R}$



# Isomorphism of vector spaces

## Definition

A linear map  $T : V \rightarrow W$  is said to be an **isomorphism** if there is a linear map  $S : W \rightarrow V$  such that

$S \circ T = 1_V : V \rightarrow V$  (identity map) and  $T \circ S = 1_W : W \rightarrow W$ .

If  $T : V \rightarrow W$  is an isomorphism, we say that  $V$  and  $W$  are isomorphic, and we write  $V \cong W$ .

## Example

Let  $A$  be an  $n \times n$  matrix over  $\mathbb{R}$ . Consider  $A : \mathbb{R}^n \rightarrow \mathbb{R}^n$  as a linear map. When is it an isomorphism?

**Answer:** When there is an inverse linear map  $B : \mathbb{R}^n \rightarrow \mathbb{R}^n$  such that  $A \circ B = 1_{\mathbb{R}^n}$  and  $B \circ A = 1_{\mathbb{R}^n}$ , i.e., when there is an  $n \times n$  matrix  $B$  over  $\mathbb{R}$  such that  $AB = I_n$

# Isomorphism of vector spaces

## Definition

A linear map  $T : V \rightarrow W$  is said to be an **isomorphism** if there is a linear map  $S : W \rightarrow V$  such that

$S \circ T = 1_V : V \rightarrow V$  (identity map) and  $T \circ S = 1_W : W \rightarrow W$ .

If  $T : V \rightarrow W$  is an isomorphism, we say that  $V$  and  $W$  are isomorphic, and we write  $V \cong W$ .

## Example

Let  $A$  be an  $n \times n$  matrix over  $\mathbb{R}$ . Consider  $A : \mathbb{R}^n \rightarrow \mathbb{R}^n$  as a linear map. When is it an isomorphism?

**Answer:** When there is an inverse linear map  $B : \mathbb{R}^n \rightarrow \mathbb{R}^n$  such that  $A \circ B = 1_{\mathbb{R}^n}$  and  $B \circ A = 1_{\mathbb{R}^n}$ , i.e., when there is an  $n \times n$  matrix  $B$  over  $\mathbb{R}$  such that  $AB = I_n$  and  $BA = I_n$ ,

# Isomorphism of vector spaces

## Definition

A linear map  $T : V \rightarrow W$  is said to be an **isomorphism** if there is a linear map  $S : W \rightarrow V$  such that

$S \circ T = 1_V : V \rightarrow V$  (identity map) and  $T \circ S = 1_W : W \rightarrow W$ .

If  $T : V \rightarrow W$  is an isomorphism, we say that  $V$  and  $W$  are isomorphic, and we write  $V \cong W$ .

## Example

Let  $A$  be an  $n \times n$  matrix over  $\mathbb{R}$ . Consider  $A : \mathbb{R}^n \rightarrow \mathbb{R}^n$  as a linear map. When is it an isomorphism?

**Answer:** When there is an inverse linear map  $B : \mathbb{R}^n \rightarrow \mathbb{R}^n$  such that  $A \circ B = 1_{\mathbb{R}^n}$  and  $B \circ A = 1_{\mathbb{R}^n}$ , i.e., when there is an  $n \times n$  matrix  $B$  over  $\mathbb{R}$  such that  $AB = I_n$  and  $BA = I_n$ , i.e., when  $A$  is an invertible matrix.

# Isomorphism of vector spaces

## Theorem

*Let  $T : V \rightarrow W$  be a linear map.*

# Isomorphism of vector spaces

## Theorem

*Let  $T : V \rightarrow W$  be a linear map. The following are equivalent:*

# Isomorphism of vector spaces

## Theorem

*Let  $T : V \rightarrow W$  be a linear map. The following are equivalent:*

- 1  *$T$  is an isomorphism.*

# Isomorphism of vector spaces

## Theorem

*Let  $T : V \rightarrow W$  be a linear map. The following are equivalent:*

- 1  *$T$  is an isomorphism.*
- 2  *$T$  is bijective (i.e., as a set map, it is injective and surjective).*

# Isomorphism of vector spaces

## Theorem

*Let  $T : V \rightarrow W$  be a linear map. The following are equivalent:*

- ❶  *$T$  is an isomorphism.*
- ❷  *$T$  is bijective (i.e., as a set map, it is injective and surjective).*

## Proof.

(1)  $\Rightarrow$  (2):



# Isomorphism of vector spaces

## Theorem

*Let  $T : V \rightarrow W$  be a linear map. The following are equivalent:*

- ①  *$T$  is an isomorphism.*
- ②  *$T$  is bijective (i.e., as a set map, it is injective and surjective).*

## Proof.

(1)  $\Rightarrow$  (2): Since  $T$  is an isomorphism, there is a linear map  $S : W \rightarrow V$

# Isomorphism of vector spaces

## Theorem

*Let  $T : V \rightarrow W$  be a linear map. The following are equivalent:*

- ①  *$T$  is an isomorphism.*
- ②  *$T$  is bijective (i.e., as a set map, it is injective and surjective).*

## Proof.

(1)  $\Rightarrow$  (2): Since  $T$  is an isomorphism, there is a linear map  $S : W \rightarrow V$  such that  $S \circ T = 1_V$

# Isomorphism of vector spaces

## Theorem

*Let  $T : V \rightarrow W$  be a linear map. The following are equivalent:*

- ①  *$T$  is an isomorphism.*
- ②  *$T$  is bijective (i.e., as a set map, it is injective and surjective).*

## Proof.

(1)  $\Rightarrow$  (2): Since  $T$  is an isomorphism, there is a linear map  $S : W \rightarrow V$  such that  $S \circ T = 1_V$  and  $T \circ S = 1_W$ .

# Isomorphism of vector spaces

## Theorem

*Let  $T : V \rightarrow W$  be a linear map. The following are equivalent:*

- 1  *$T$  is an isomorphism.*
- 2  *$T$  is bijective (i.e., as a set map, it is injective and surjective).*

## Proof.

(1)  $\Rightarrow$  (2): Since  $T$  is an isomorphism, there is a linear map  $S : W \rightarrow V$  such that  $S \circ T = 1_V$  and  $T \circ S = 1_W$ .

Let  $T(u) = T(v)$  for some  $u, v \in V$ .

# Isomorphism of vector spaces

## Theorem

*Let  $T : V \rightarrow W$  be a linear map. The following are equivalent:*

- ❶  *$T$  is an isomorphism.*
- ❷  *$T$  is bijective (i.e., as a set map, it is injective and surjective).*

## Proof.

(1)  $\Rightarrow$  (2): Since  $T$  is an isomorphism, there is a linear map  $S : W \rightarrow V$  such that  $S \circ T = 1_V$  and  $T \circ S = 1_W$ .

Let  $T(u) = T(v)$  for some  $u, v \in V$ .

Apply  $S$  on this equality,

# Isomorphism of vector spaces

## Theorem

*Let  $T : V \rightarrow W$  be a linear map. The following are equivalent:*

- ❶  *$T$  is an isomorphism.*
- ❷  *$T$  is bijective (i.e., as a set map, it is injective and surjective).*

## Proof.

(1)  $\Rightarrow$  (2): Since  $T$  is an isomorphism, there is a linear map  $S : W \rightarrow V$  such that  $S \circ T = 1_V$  and  $T \circ S = 1_W$ .

Let  $T(u) = T(v)$  for some  $u, v \in V$ .

Apply  $S$  on this equality, to get  $u = v$ .

# Isomorphism of vector spaces

## Theorem

*Let  $T : V \rightarrow W$  be a linear map. The following are equivalent:*

- ❶  *$T$  is an isomorphism.*
- ❷  *$T$  is bijective (i.e., as a set map, it is injective and surjective).*

## Proof.

(1)  $\Rightarrow$  (2): Since  $T$  is an isomorphism, there is a linear map  $S : W \rightarrow V$  such that  $S \circ T = 1_V$  and  $T \circ S = 1_W$ .

Let  $T(u) = T(v)$  for some  $u, v \in V$ .

Apply  $S$  on this equality, to get  $u = v$ . So  $T$  is injective.

# Isomorphism of vector spaces

## Theorem

*Let  $T : V \rightarrow W$  be a linear map. The following are equivalent:*

- 1  *$T$  is an isomorphism.*
- 2  *$T$  is bijective (i.e., as a set map, it is injective and surjective).*

## Proof.

(1)  $\Rightarrow$  (2): Since  $T$  is an isomorphism, there is a linear map  $S : W \rightarrow V$  such that  $S \circ T = 1_V$  and  $T \circ S = 1_W$ .

Let  $T(u) = T(v)$  for some  $u, v \in V$ .

Apply  $S$  on this equality, to get  $u = v$ . So  $T$  is injective.

For surjectivity, note that  $T(S(w)) = w$  for every  $w \in W$ .



# Isomorphism of vector spaces

## Theorem

*Let  $T : V \rightarrow W$  be a linear map. The following are equivalent:*

- 1  *$T$  is an isomorphism.*
- 2  *$T$  is bijective (i.e., as a set map, it is injective and surjective).*

## Proof.

(1)  $\Rightarrow$  (2): Since  $T$  is an isomorphism, there is a linear map  $S : W \rightarrow V$  such that  $S \circ T = 1_V$  and  $T \circ S = 1_W$ .

Let  $T(u) = T(v)$  for some  $u, v \in V$ .

Apply  $S$  on this equality, to get  $u = v$ . So  $T$  is injective.

For surjectivity, note that  $T(S(w)) = w$  for every  $w \in W$ .

(2)  $\Rightarrow$  (1):

# Isomorphism of vector spaces

## Theorem

*Let  $T : V \rightarrow W$  be a linear map. The following are equivalent:*

- 1  *$T$  is an isomorphism.*
- 2  *$T$  is bijective (i.e., as a set map, it is injective and surjective).*

## Proof.

(1)  $\Rightarrow$  (2): Since  $T$  is an isomorphism, there is a linear map  $S : W \rightarrow V$  such that  $S \circ T = 1_V$  and  $T \circ S = 1_W$ .

Let  $T(u) = T(v)$  for some  $u, v \in V$ .

Apply  $S$  on this equality, to get  $u = v$ . So  $T$  is injective.

For surjectivity, note that  $T(S(w)) = w$  for every  $w \in W$ .

(2)  $\Rightarrow$  (1): Since  $T$  is bijective,

# Isomorphism of vector spaces

## Theorem

*Let  $T : V \rightarrow W$  be a linear map. The following are equivalent:*

- ①  *$T$  is an isomorphism.*
- ②  *$T$  is bijective (i.e., as a set map, it is injective and surjective).*

## Proof.

(1)  $\Rightarrow$  (2): Since  $T$  is an isomorphism, there is a linear map  $S : W \rightarrow V$  such that  $S \circ T = 1_V$  and  $T \circ S = 1_W$ .

Let  $T(u) = T(v)$  for some  $u, v \in V$ .

Apply  $S$  on this equality, to get  $u = v$ . So  $T$  is injective.

For surjectivity, note that  $T(S(w)) = w$  for every  $w \in W$ .

(2)  $\Rightarrow$  (1): Since  $T$  is bijective, there is an inverse SET map  $S : W \rightarrow V$  such that  $S \circ T = 1_V$

# Isomorphism of vector spaces

## Theorem

*Let  $T : V \rightarrow W$  be a linear map. The following are equivalent:*

- 1  $T$  is an isomorphism.
- 2  $T$  is bijective (i.e., as a set map, it is injective and surjective).

## Proof.

(1)  $\Rightarrow$  (2): Since  $T$  is an isomorphism, there is a linear map  $S : W \rightarrow V$  such that  $S \circ T = 1_V$  and  $T \circ S = 1_W$ .

Let  $T(u) = T(v)$  for some  $u, v \in V$ .

Apply  $S$  on this equality, to get  $u = v$ . So  $T$  is injective.

For surjectivity, note that  $T(S(w)) = w$  for every  $w \in W$ .

(2)  $\Rightarrow$  (1): Since  $T$  is bijective, there is an inverse SET map  $S : W \rightarrow V$  such that  $S \circ T = 1_V$  and  $T \circ S = 1_W$ .

# Isomorphism of vector spaces

## Theorem

Let  $T : V \rightarrow W$  be a linear map. The following are equivalent:

- 1  $T$  is an isomorphism.
- 2  $T$  is bijective (i.e., as a set map, it is injective and surjective).

## Proof.

(1)  $\Rightarrow$  (2): Since  $T$  is an isomorphism, there is a linear map  $S : W \rightarrow V$  such that  $S \circ T = 1_V$  and  $T \circ S = 1_W$ .

Let  $T(u) = T(v)$  for some  $u, v \in V$ .

Apply  $S$  on this equality, to get  $u = v$ . So  $T$  is injective.

For surjectivity, note that  $T(S(w)) = w$  for every  $w \in W$ .

(2)  $\Rightarrow$  (1): Since  $T$  is bijective, there is an inverse SET map  $S : W \rightarrow V$  such that  $S \circ T = 1_V$  and  $T \circ S = 1_W$ .

All we need to show that  $S : W \rightarrow V$  is a linear map.

# Isomorphism of vector spaces

## Theorem

Let  $T : V \rightarrow W$  be a linear map. The following are equivalent:

- 1  $T$  is an isomorphism.
- 2  $T$  is bijective (i.e., as a set map, it is injective and surjective).

## Proof.

(1)  $\Rightarrow$  (2): Since  $T$  is an isomorphism, there is a linear map  $S : W \rightarrow V$  such that  $S \circ T = 1_V$  and  $T \circ S = 1_W$ .

Let  $T(u) = T(v)$  for some  $u, v \in V$ .

Apply  $S$  on this equality, to get  $u = v$ . So  $T$  is injective.

For surjectivity, note that  $T(S(w)) = w$  for every  $w \in W$ .

(2)  $\Rightarrow$  (1): Since  $T$  is bijective, there is an inverse SET map  $S : W \rightarrow V$  such that  $S \circ T = 1_V$  and  $T \circ S = 1_W$ .

All we need to show that  $S : W \rightarrow V$  is a linear map.

You may try on your own.

# Isomorphism of vector spaces

## Theorem

Let  $T : V \rightarrow W$  be a linear map. The following are equivalent:

- 1  $T$  is an isomorphism.
- 2  $T$  is bijective (i.e., as a set map, it is injective and surjective).

## Proof.

(1)  $\Rightarrow$  (2): Since  $T$  is an isomorphism, there is a linear map  $S : W \rightarrow V$  such that  $S \circ T = 1_V$  and  $T \circ S = 1_W$ .

Let  $T(u) = T(v)$  for some  $u, v \in V$ .

Apply  $S$  on this equality, to get  $u = v$ . So  $T$  is injective.

For surjectivity, note that  $T(S(w)) = w$  for every  $w \in W$ .

(2)  $\Rightarrow$  (1): Since  $T$  is bijective, there is an inverse SET map  $S : W \rightarrow V$  such that  $S \circ T = 1_V$  and  $T \circ S = 1_W$ .

All we need to show that  $S : W \rightarrow V$  is a linear map.

You may try on your own. Otherwise, see the next slide.

# Proof of the theorem contd...

(2)  $\Rightarrow$  (1):



# Proof of the theorem contd...

(2)  $\Rightarrow$  (1):

...

Let  $w_1, w_2 \in W$ .

# Proof of the theorem contd...

(2)  $\Rightarrow$  (1):

...

Let  $w_1, w_2 \in W$ . Want to show  $S(w_1 + w_2) = S(w_1) + S(w_2)$ .

# Proof of the theorem contd...

(2)  $\Rightarrow$  (1):

...

Let  $w_1, w_2 \in W$ . Want to show  $S(w_1 + w_2) = S(w_1) + S(w_2)$ .

Set  $v_1 := S(w_1)$  and  $v_2 := S(w_2)$ .

# Proof of the theorem contd...

(2)  $\Rightarrow$  (1):

...

Let  $w_1, w_2 \in W$ . Want to show  $S(w_1 + w_2) = S(w_1) + S(w_2)$ .

Set  $v_1 := S(w_1)$  and  $v_2 := S(w_2)$ .

Hence, since  $T$  is inverse of  $S$  (as a set map), it follows that

$$T(v_1) = w_1$$

# Proof of the theorem contd...

(2)  $\Rightarrow$  (1):

...

Let  $w_1, w_2 \in W$ . Want to show  $S(w_1 + w_2) = S(w_1) + S(w_2)$ .

Set  $v_1 := S(w_1)$  and  $v_2 := S(w_2)$ .

Hence, since  $T$  is inverse of  $S$  (as a set map), it follows that

$T(v_1) = w_1$  and  $T(v_2) = w_2$ .

# Proof of the theorem contd...

(2)  $\Rightarrow$  (1):

...

Let  $w_1, w_2 \in W$ . Want to show  $S(w_1 + w_2) = S(w_1) + S(w_2)$ .

Set  $v_1 := S(w_1)$  and  $v_2 := S(w_2)$ .

Hence, since  $T$  is inverse of  $S$  (as a set map), it follows that

$T(v_1) = w_1$  and  $T(v_2) = w_2$ . So  $T(v_1 + v_2) = w_1 + w_2$  because  $T$  is linear.

# Proof of the theorem contd...

(2)  $\Rightarrow$  (1):

...

Let  $w_1, w_2 \in W$ . Want to show  $S(w_1 + w_2) = S(w_1) + S(w_2)$ .

Set  $v_1 := S(w_1)$  and  $v_2 := S(w_2)$ .

Hence, since  $T$  is inverse of  $S$  (as a set map), it follows that

$T(v_1) = w_1$  and  $T(v_2) = w_2$ . So  $T(v_1 + v_2) = w_1 + w_2$  because  $T$  is linear. Therefore  $S(w_1 + w_2) = v_1 + v_2 = S(w_1) + S(w_2)$ .

# Proof of the theorem contd...

(2)  $\Rightarrow$  (1):

...

Let  $w_1, w_2 \in W$ . Want to show  $S(w_1 + w_2) = S(w_1) + S(w_2)$ .

Set  $v_1 := S(w_1)$  and  $v_2 := S(w_2)$ .

Hence, since  $T$  is inverse of  $S$  (as a set map), it follows that

$T(v_1) = w_1$  and  $T(v_2) = w_2$ . So  $T(v_1 + v_2) = w_1 + w_2$  because  $T$  is linear. Therefore  $S(w_1 + w_2) = v_1 + v_2 = S(w_1) + S(w_2)$ .

Similarly, one can prove that  $S(cw) = cS(w)$  for every scalar  $c \in \mathbb{R}$  and every vector  $w \in W$ .



# Conditions for a linear transformation to be isomorphism

## Theorem

*Let  $T : V \rightarrow V$  be a linear map*

# Conditions for a linear transformation to be isomorphism

## Theorem

*Let  $T : V \rightarrow V$  be a linear map (or **linear operator**),*

# Conditions for a linear transformation to be isomorphism

## Theorem

*Let  $T : V \rightarrow V$  be a linear map (or **linear operator**), where  $\dim(V) = n < \infty$ .*

# Conditions for a linear transformation to be isomorphism

## Theorem

*Let  $T : V \rightarrow V$  be a linear map (or **linear operator**), where  $\dim(V) = n < \infty$ . Then the following statements are equivalent:*

# Conditions for a linear transformation to be isomorphism

## Theorem

*Let  $T : V \rightarrow V$  be a linear map (or **linear operator**), where  $\dim(V) = n < \infty$ . Then the following statements are equivalent:*

- 1  *$T$  is an isomorphism (see the definition in the 1st slide).*

# Conditions for a linear transformation to be isomorphism

## Theorem

*Let  $T : V \rightarrow V$  be a linear map (or **linear operator**), where  $\dim(V) = n < \infty$ . Then the following statements are equivalent:*

- ①  *$T$  is an isomorphism (see the definition in the 1st slide).*
- ②  *$T$  is bijective (as a set map).*

# Conditions for a linear transformation to be isomorphism

## Theorem

*Let  $T : V \rightarrow V$  be a linear map (or **linear operator**), where  $\dim(V) = n < \infty$ . Then the following statements are equivalent:*

- 1  *$T$  is an isomorphism (see the definition in the 1st slide).*
- 2  *$T$  is bijective (as a set map).*
- 3  *$T$  is injective.*

# Conditions for a linear transformation to be isomorphism

## Theorem

*Let  $T : V \rightarrow V$  be a linear map (or **linear operator**), where  $\dim(V) = n < \infty$ . Then the following statements are equivalent:*

- 1  *$T$  is an isomorphism (see the definition in the 1st slide).*
- 2  *$T$  is bijective (as a set map).*
- 3  *$T$  is injective.*
- 4  *$\text{Ker}(T) = 0$ ,*



# Conditions for a linear transformation to be isomorphism

## Theorem

*Let  $T : V \rightarrow V$  be a linear map (or **linear operator**), where  $\dim(V) = n < \infty$ . Then the following statements are equivalent:*

- 1  *$T$  is an isomorphism (see the definition in the 1st slide).*
- 2  *$T$  is bijective (as a set map).*
- 3  *$T$  is injective.*
- 4  *$\text{Ker}(T) = 0$ , i.e.,  $\{T(v) = 0 \Rightarrow v = 0\}$ ,*

# Conditions for a linear transformation to be isomorphism

## Theorem

*Let  $T : V \rightarrow V$  be a linear map (or **linear operator**), where  $\dim(V) = n < \infty$ . Then the following statements are equivalent:*

- 1  *$T$  is an isomorphism (see the definition in the 1st slide).*
- 2  *$T$  is bijective (as a set map).*
- 3  *$T$  is injective.*
- 4  *$\text{Ker}(T) = 0$ , i.e.,  $\{T(v) = 0 \Rightarrow v = 0\}$ , i.e.,  $\text{Null}(T) = 0$ .*

# Conditions for a linear transformation to be isomorphism

## Theorem

Let  $T : V \rightarrow V$  be a linear map (or **linear operator**), where  $\dim(V) = n < \infty$ . Then the following statements are equivalent:

- 1  $T$  is an isomorphism (see the definition in the 1st slide).
- 2  $T$  is bijective (as a set map).
- 3  $T$  is injective.
- 4  $\text{Ker}(T) = 0$ , i.e.,  $\{T(v) = 0 \Rightarrow v = 0\}$ , i.e.,  $\text{Null}(T) = 0$ .
- 5  $T$  is surjective.

# Conditions for a linear transformation to be isomorphism

## Theorem

Let  $T : V \rightarrow V$  be a linear map (or **linear operator**), where  $\dim(V) = n < \infty$ . Then the following statements are equivalent:

- 1  $T$  is an isomorphism (see the definition in the 1st slide).
- 2  $T$  is bijective (as a set map).
- 3  $T$  is injective.
- 4  $\text{Ker}(T) = 0$ , i.e.,  $\{T(v) = 0 \Rightarrow v = 0\}$ , i.e.,  $\text{Null}(T) = 0$ .
- 5  $T$  is surjective.

**Proof.**

# Conditions for a linear transformation to be isomorphism

## Theorem

Let  $T : V \rightarrow V$  be a linear map (or **linear operator**), where  $\dim(V) = n < \infty$ . Then the following statements are equivalent:

- 1  $T$  is an isomorphism (see the definition in the 1st slide).
- 2  $T$  is bijective (as a set map).
- 3  $T$  is injective.
- 4  $\text{Ker}(T) = 0$ , i.e.,  $\{T(v) = 0 \Rightarrow v = 0\}$ , i.e.,  $\text{Null}(T) = 0$ .
- 5  $T$  is surjective.

**Proof.** We already proved  $(1) \Leftrightarrow (2)$ .

# Conditions for a linear transformation to be isomorphism

## Theorem

Let  $T : V \rightarrow V$  be a linear map (or **linear operator**), where  $\dim(V) = n < \infty$ . Then the following statements are equivalent:

- 1  $T$  is an isomorphism (see the definition in the 1st slide).
- 2  $T$  is bijective (as a set map).
- 3  $T$  is injective.
- 4  $\text{Ker}(T) = 0$ , i.e.,  $\{T(v) = 0 \Rightarrow v = 0\}$ , i.e.,  $\text{Null}(T) = 0$ .
- 5  $T$  is surjective.

**Proof.** We already proved  $(1) \Leftrightarrow (2)$ . The following implications are trivial:

# Conditions for a linear transformation to be isomorphism

## Theorem

Let  $T : V \rightarrow V$  be a linear map (or **linear operator**), where  $\dim(V) = n < \infty$ . Then the following statements are equivalent:

- 1  $T$  is an isomorphism (see the definition in the 1st slide).
- 2  $T$  is bijective (as a set map).
- 3  $T$  is injective.
- 4  $\text{Ker}(T) = 0$ , i.e.,  $\{T(v) = 0 \Rightarrow v = 0\}$ , i.e.,  $\text{Null}(T) = 0$ .
- 5  $T$  is surjective.

**Proof.** We already proved  $(1) \Leftrightarrow (2)$ . The following implications are trivial:  $(2) \Rightarrow (3)$

# Conditions for a linear transformation to be isomorphism

## Theorem

Let  $T : V \rightarrow V$  be a linear map (or **linear operator**), where  $\dim(V) = n < \infty$ . Then the following statements are equivalent:

- ①  $T$  is an isomorphism (see the definition in the 1st slide).
- ②  $T$  is bijective (as a set map).
- ③  $T$  is injective.
- ④  $\text{Ker}(T) = 0$ , i.e.,  $\{T(v) = 0 \Rightarrow v = 0\}$ , i.e.,  $\text{Null}(T) = 0$ .
- ⑤  $T$  is surjective.

**Proof.** We already proved  $(1) \Leftrightarrow (2)$ . The following implications are trivial:  $(2) \Rightarrow (3) \Rightarrow (4)$ .



# Conditions for a linear transformation to be isomorphism

## Theorem

Let  $T : V \rightarrow V$  be a linear map (or **linear operator**), where  $\dim(V) = n < \infty$ . Then the following statements are equivalent:

- 1  $T$  is an isomorphism (see the definition in the 1st slide).
- 2  $T$  is bijective (as a set map).
- 3  $T$  is injective.
- 4  $\text{Ker}(T) = 0$ , i.e.,  $\{T(v) = 0 \Rightarrow v = 0\}$ , i.e.,  $\text{Null}(T) = 0$ .
- 5  $T$  is surjective.

**Proof.** We already proved  $(1) \Leftrightarrow (2)$ . The following implications are trivial:  $(2) \Rightarrow (3) \Rightarrow (4)$ .  
 $(4) \Rightarrow (5)$ :

# Conditions for a linear transformation to be isomorphism

## Theorem

Let  $T : V \rightarrow V$  be a linear map (or **linear operator**), where  $\dim(V) = n < \infty$ . Then the following statements are equivalent:

- 1  $T$  is an isomorphism (see the definition in the 1st slide).
- 2  $T$  is bijective (as a set map).
- 3  $T$  is injective.
- 4  $\text{Ker}(T) = 0$ , i.e.,  $\{T(v) = 0 \Rightarrow v = 0\}$ , i.e.,  $\text{Null}(T) = 0$ .
- 5  $T$  is surjective.

**Proof.** We already proved  $(1) \Leftrightarrow (2)$ . The following implications are trivial:  $(2) \Rightarrow (3) \Rightarrow (4)$ .

$(4) \Rightarrow (5)$ : Since  $\text{Null}(T) = 0$ ,

# Conditions for a linear transformation to be isomorphism

## Theorem

Let  $T : V \rightarrow V$  be a linear map (or **linear operator**), where  $\dim(V) = n < \infty$ . Then the following statements are equivalent:

- 1  $T$  is an isomorphism (see the definition in the 1st slide).
- 2  $T$  is bijective (as a set map).
- 3  $T$  is injective.
- 4  $\text{Ker}(T) = 0$ , i.e.,  $\{T(v) = 0 \Rightarrow v = 0\}$ , i.e.,  $\text{Null}(T) = 0$ .
- 5  $T$  is surjective.

**Proof.** We already proved  $(1) \Leftrightarrow (2)$ . The following implications are trivial:  $(2) \Rightarrow (3) \Rightarrow (4)$ .

$(4) \Rightarrow (5)$ : Since  $\text{Null}(T) = 0$ ,  $\text{nullity}(T) = \dim(\text{Null}(T)) = 0$ .

# Conditions for a linear transformation to be isomorphism

## Theorem

Let  $T : V \rightarrow V$  be a linear map (or **linear operator**), where  $\dim(V) = n < \infty$ . Then the following statements are equivalent:

- 1  $T$  is an isomorphism (see the definition in the 1st slide).
- 2  $T$  is bijective (as a set map).
- 3  $T$  is injective.
- 4  $\text{Ker}(T) = 0$ , i.e.,  $\{T(v) = 0 \Rightarrow v = 0\}$ , i.e.,  $\text{Null}(T) = 0$ .
- 5  $T$  is surjective.

**Proof.** We already proved  $(1) \Leftrightarrow (2)$ . The following implications are trivial:  $(2) \Rightarrow (3) \Rightarrow (4)$ .

$(4) \Rightarrow (5)$ : Since  $\text{Null}(T) = 0$ ,  $\text{nullity}(T) = \dim(\text{Null}(T)) = 0$ . Hence, by Rank-Nullity Theorem,  $\text{rank}(T) = \dim(V)$ .

# Conditions for a linear transformation to be isomorphism

## Theorem

Let  $T : V \rightarrow V$  be a linear map (or **linear operator**), where  $\dim(V) = n < \infty$ . Then the following statements are equivalent:

- 1  $T$  is an isomorphism (see the definition in the 1st slide).
- 2  $T$  is bijective (as a set map).
- 3  $T$  is injective.
- 4  $\text{Ker}(T) = 0$ , i.e.,  $\{T(v) = 0 \Rightarrow v = 0\}$ , i.e.,  $\text{Null}(T) = 0$ .
- 5  $T$  is surjective.

**Proof.** We already proved  $(1) \Leftrightarrow (2)$ . The following implications are trivial:  $(2) \Rightarrow (3) \Rightarrow (4)$ .

$(4) \Rightarrow (5)$ : Since  $\text{Null}(T) = 0$ ,  $\text{nullity}(T) = \dim(\text{Null}(T)) = 0$ .

Hence, by Rank-Nullity Theorem,  $\text{rank}(T) = \dim(V)$ .

So  $\text{Image}(T) = V$ ,

# Conditions for a linear transformation to be isomorphism

## Theorem

Let  $T : V \rightarrow V$  be a linear map (or **linear operator**), where  $\dim(V) = n < \infty$ . Then the following statements are equivalent:

- 1  $T$  is an isomorphism (see the definition in the 1st slide).
- 2  $T$  is bijective (as a set map).
- 3  $T$  is injective.
- 4  $\text{Ker}(T) = 0$ , i.e.,  $\{T(v) = 0 \Rightarrow v = 0\}$ , i.e.,  $\text{Null}(T) = 0$ .
- 5  $T$  is surjective.

**Proof.** We already proved  $(1) \Leftrightarrow (2)$ . The following implications are trivial:  $(2) \Rightarrow (3) \Rightarrow (4)$ .

$(4) \Rightarrow (5)$ : Since  $\text{Null}(T) = 0$ ,  $\text{nullity}(T) = \dim(\text{Null}(T)) = 0$ .

Hence, by Rank-Nullity Theorem,  $\text{rank}(T) = \dim(V)$ .

So  $\text{Image}(T) = V$ , i.e.,  $T$  is surjective.

# Conditions for a linear transformation to be isomorphism

## Theorem

Let  $T : V \rightarrow V$  be a linear map (or **linear operator**), where  $\dim(V) = n < \infty$ . Then the following statements are equivalent:

- 1  $T$  is an isomorphism (see the definition in the 1st slide).
- 2  $T$  is bijective (as a set map).
- 3  $T$  is injective.
- 4  $\text{Ker}(T) = 0$ , i.e.,  $\{T(v) = 0 \Rightarrow v = 0\}$ , i.e.,  $\text{Null}(T) = 0$ .
- 5  $T$  is surjective.

**Proof.** We already proved  $(1) \Leftrightarrow (2)$ . The following implications are trivial:  $(2) \Rightarrow (3) \Rightarrow (4)$ .

$(4) \Rightarrow (5)$ : Since  $\text{Null}(T) = 0$ ,  $\text{nullity}(T) = \dim(\text{Null}(T)) = 0$ .

Hence, by Rank-Nullity Theorem,  $\text{rank}(T) = \dim(V)$ .

So  $\text{Image}(T) = V$ , i.e.,  $T$  is surjective.

$(5) \Rightarrow (2)$ :

# Conditions for a linear transformation to be isomorphism

## Theorem

Let  $T : V \rightarrow V$  be a linear map (or **linear operator**), where  $\dim(V) = n < \infty$ . Then the following statements are equivalent:

- 1  $T$  is an isomorphism (see the definition in the 1st slide).
- 2  $T$  is bijective (as a set map).
- 3  $T$  is injective.
- 4  $\text{Ker}(T) = 0$ , i.e.,  $\{T(v) = 0 \Rightarrow v = 0\}$ , i.e.,  $\text{Null}(T) = 0$ .
- 5  $T$  is surjective.

**Proof.** We already proved  $(1) \Leftrightarrow (2)$ . The following implications are trivial:  $(2) \Rightarrow (3) \Rightarrow (4)$ .

$(4) \Rightarrow (5)$ : Since  $\text{Null}(T) = 0$ ,  $\text{nullity}(T) = \dim(\text{Null}(T)) = 0$ .

Hence, by Rank-Nullity Theorem,  $\text{rank}(T) = \dim(V)$ .

So  $\text{Image}(T) = V$ , i.e.,  $T$  is surjective.

$(5) \Rightarrow (2)$ : Since  $T$  is surjective,



# Conditions for a linear transformation to be isomorphism

## Theorem

Let  $T : V \rightarrow V$  be a linear map (or **linear operator**), where  $\dim(V) = n < \infty$ . Then the following statements are equivalent:

- 1  $T$  is an isomorphism (see the definition in the 1st slide).
- 2  $T$  is bijective (as a set map).
- 3  $T$  is injective.
- 4  $\text{Ker}(T) = 0$ , i.e.,  $\{T(v) = 0 \Rightarrow v = 0\}$ , i.e.,  $\text{Null}(T) = 0$ .
- 5  $T$  is surjective.

**Proof.** We already proved  $(1) \Leftrightarrow (2)$ . The following implications are trivial:  $(2) \Rightarrow (3) \Rightarrow (4)$ .

$(4) \Rightarrow (5)$ : Since  $\text{Null}(T) = 0$ ,  $\text{nullity}(T) = \dim(\text{Null}(T)) = 0$ .

Hence, by Rank-Nullity Theorem,  $\text{rank}(T) = \dim(V)$ .

So  $\text{Image}(T) = V$ , i.e.,  $T$  is surjective.

$(5) \Rightarrow (2)$ : Since  $T$  is surjective,  $\text{rank}(T) = \dim(V)$ ,

# Conditions for a linear transformation to be isomorphism

## Theorem

Let  $T : V \rightarrow V$  be a linear map (or **linear operator**), where  $\dim(V) = n < \infty$ . Then the following statements are equivalent:

- 1  $T$  is an isomorphism (see the definition in the 1st slide).
- 2  $T$  is bijective (as a set map).
- 3  $T$  is injective.
- 4  $\text{Ker}(T) = 0$ , i.e.,  $\{T(v) = 0 \Rightarrow v = 0\}$ , i.e.,  $\text{Null}(T) = 0$ .
- 5  $T$  is surjective.

**Proof.** We already proved  $(1) \Leftrightarrow (2)$ . The following implications are trivial:  $(2) \Rightarrow (3) \Rightarrow (4)$ .

$(4) \Rightarrow (5)$ : Since  $\text{Null}(T) = 0$ ,  $\text{nullity}(T) = \dim(\text{Null}(T)) = 0$ . Hence, by Rank-Nullity Theorem,  $\text{rank}(T) = \dim(V)$ .

So  $\text{Image}(T) = V$ , i.e.,  $T$  is surjective.

$(5) \Rightarrow (2)$ : Since  $T$  is surjective,  $\text{rank}(T) = \dim(V)$ , hence  $\text{nullity}(T) = 0$ ,

# Conditions for a linear transformation to be isomorphism

## Theorem

Let  $T : V \rightarrow V$  be a linear map (or **linear operator**), where  $\dim(V) = n < \infty$ . Then the following statements are equivalent:

- 1  $T$  is an isomorphism (see the definition in the 1st slide).
- 2  $T$  is bijective (as a set map).
- 3  $T$  is injective.
- 4  $\text{Ker}(T) = 0$ , i.e.,  $\{T(v) = 0 \Rightarrow v = 0\}$ , i.e.,  $\text{Null}(T) = 0$ .
- 5  $T$  is surjective.

**Proof.** We already proved  $(1) \Leftrightarrow (2)$ . The following implications are trivial:  $(2) \Rightarrow (3) \Rightarrow (4)$ .

$(4) \Rightarrow (5)$ : Since  $\text{Null}(T) = 0$ ,  $\text{nullity}(T) = \dim(\text{Null}(T)) = 0$ . Hence, by Rank-Nullity Theorem,  $\text{rank}(T) = \dim(V)$ .

So  $\text{Image}(T) = V$ , i.e.,  $T$  is surjective.

$(5) \Rightarrow (2)$ : Since  $T$  is surjective,  $\text{rank}(T) = \dim(V)$ , hence  $\text{nullity}(T) = 0$ , i.e.,  $\text{Ker}(T) = 0$ .

# Conditions for a linear transformation to be isomorphism

## Theorem

Let  $T : V \rightarrow V$  be a linear map (or **linear operator**), where  $\dim(V) = n < \infty$ . Then the following statements are equivalent:

- 1  $T$  is an isomorphism (see the definition in the 1st slide).
- 2  $T$  is bijective (as a set map).
- 3  $T$  is injective.
- 4  $\text{Ker}(T) = 0$ , i.e.,  $\{T(v) = 0 \Rightarrow v = 0\}$ , i.e.,  $\text{Null}(T) = 0$ .
- 5  $T$  is surjective.

**Proof.** We already proved  $(1) \Leftrightarrow (2)$ . The following implications are trivial:  $(2) \Rightarrow (3) \Rightarrow (4)$ .

$(4) \Rightarrow (5)$ : Since  $\text{Null}(T) = 0$ ,  $\text{nullity}(T) = \dim(\text{Null}(T)) = 0$ .

Hence, by Rank-Nullity Theorem,  $\text{rank}(T) = \dim(V)$ .

So  $\text{Image}(T) = V$ , i.e.,  $T$  is surjective.

$(5) \Rightarrow (2)$ : Since  $T$  is surjective,  $\text{rank}(T) = \dim(V)$ , hence  $\text{nullity}(T) = 0$ , i.e.,  $\text{Ker}(T) = 0$ . Then, by linearity,  $T$  is injective.

# Conditions for a square matrix to be invertible

## Theorem

*Let  $A$  be an  $n \times n$  matrix over  $\mathbb{R}$ .*

# Conditions for a square matrix to be invertible

## Theorem

*Let  $A$  be an  $n \times n$  matrix over  $\mathbb{R}$ . The following are equivalent:*

# Conditions for a square matrix to be invertible

## Theorem

*Let  $A$  be an  $n \times n$  matrix over  $\mathbb{R}$ . The following are equivalent:*

- 1  *$A$  is invertible.*

# Conditions for a square matrix to be invertible

## Theorem

*Let  $A$  be an  $n \times n$  matrix over  $\mathbb{R}$ . The following are equivalent:*

- 1  *$A$  is invertible.*
- 2 *The homogeneous system  $AX = 0$  has only the trivial solution.*



# Conditions for a square matrix to be invertible

## Theorem

*Let  $A$  be an  $n \times n$  matrix over  $\mathbb{R}$ . The following are equivalent:*

- 1  *$A$  is invertible.*
- 2 *The homogeneous system  $AX = 0$  has only the trivial solution.*
- 3 *For every  $b \in \mathbb{R}^n$ , the system  $AX = b$  has a solution.*

# Conditions for a square matrix to be invertible

## Theorem

*Let  $A$  be an  $n \times n$  matrix over  $\mathbb{R}$ . The following are equivalent:*

- 1  *$A$  is invertible.*
- 2 *The homogeneous system  $AX = 0$  has only the trivial solution.*
- 3 *For every  $b \in \mathbb{R}^n$ , the system  $AX = b$  has a solution.*
- 4  *$\mathbb{R}^n$  is spanned by the column vectors of  $A$ .*

# Conditions for a square matrix to be invertible

## Theorem

*Let  $A$  be an  $n \times n$  matrix over  $\mathbb{R}$ . The following are equivalent:*

- 1  *$A$  is invertible.*
- 2 *The homogeneous system  $AX = 0$  has only the trivial solution.*
- 3 *For every  $b \in \mathbb{R}^n$ , the system  $AX = b$  has a solution.*
- 4  *$\mathbb{R}^n$  is spanned by the column vectors of  $A$ .*
- 5 *The column vectors of  $A$  are linearly independent.*

# Conditions for a square matrix to be invertible

## Theorem

*Let  $A$  be an  $n \times n$  matrix over  $\mathbb{R}$ . The following are equivalent:*

- 1  *$A$  is invertible.*
- 2 *The homogeneous system  $AX = 0$  has only the trivial solution.*
- 3 *For every  $b \in \mathbb{R}^n$ , the system  $AX = b$  has a solution.*
- 4  *$\mathbb{R}^n$  is spanned by the column vectors of  $A$ .*
- 5 *The column vectors of  $A$  are linearly independent.*
- 6  *$\mathbb{R}^n$  is spanned by the row vectors of  $A$ .*

# Conditions for a square matrix to be invertible

## Theorem

*Let  $A$  be an  $n \times n$  matrix over  $\mathbb{R}$ . The following are equivalent:*

- 1  *$A$  is invertible.*
- 2 *The homogeneous system  $AX = 0$  has only the trivial solution.*
- 3 *For every  $b \in \mathbb{R}^n$ , the system  $AX = b$  has a solution.*
- 4  *$\mathbb{R}^n$  is spanned by the column vectors of  $A$ .*
- 5 *The column vectors of  $A$  are linearly independent.*
- 6  *$\mathbb{R}^n$  is spanned by the row vectors of  $A$ .*
- 7 *The row vectors of  $A$  are linearly independent.*

# Conditions for a square matrix to be invertible

## Theorem

*Let  $A$  be an  $n \times n$  matrix over  $\mathbb{R}$ . The following are equivalent:*

- ①  *$A$  is invertible.*
- ② *The homogeneous system  $AX = 0$  has only the trivial solution.*
- ③ *For every  $b \in \mathbb{R}^n$ , the system  $AX = b$  has a solution.*
- ④  *$\mathbb{R}^n$  is spanned by the column vectors of  $A$ .*
- ⑤ *The column vectors of  $A$  are linearly independent.*
- ⑥  *$\mathbb{R}^n$  is spanned by the row vectors of  $A$ .*
- ⑦ *The row vectors of  $A$  are linearly independent.*

**Proof.**

# Conditions for a square matrix to be invertible

## Theorem

*Let  $A$  be an  $n \times n$  matrix over  $\mathbb{R}$ . The following are equivalent:*

- ①  *$A$  is invertible.*
- ② *The homogeneous system  $AX = 0$  has only the trivial solution.*
- ③ *For every  $b \in \mathbb{R}^n$ , the system  $AX = b$  has a solution.*
- ④  *$\mathbb{R}^n$  is spanned by the column vectors of  $A$ .*
- ⑤ *The column vectors of  $A$  are linearly independent.*
- ⑥  *$\mathbb{R}^n$  is spanned by the row vectors of  $A$ .*
- ⑦ *The row vectors of  $A$  are linearly independent.*

**Proof.** Consider  $A$  as a linear map  $A : \mathbb{R}^n \rightarrow \mathbb{R}^n$ .

# Conditions for a square matrix to be invertible

## Theorem

*Let  $A$  be an  $n \times n$  matrix over  $\mathbb{R}$ . The following are equivalent:*

- ①  *$A$  is invertible.*
- ② *The homogeneous system  $AX = 0$  has only the trivial solution.*
- ③ *For every  $b \in \mathbb{R}^n$ , the system  $AX = b$  has a solution.*
- ④  *$\mathbb{R}^n$  is spanned by the column vectors of  $A$ .*
- ⑤ *The column vectors of  $A$  are linearly independent.*
- ⑥  *$\mathbb{R}^n$  is spanned by the row vectors of  $A$ .*
- ⑦ *The row vectors of  $A$  are linearly independent.*

**Proof.** Consider  $A$  as a linear map  $A : \mathbb{R}^n \rightarrow \mathbb{R}^n$ . Then (2) is same as  $\text{Null}(A) = 0$ .



# Conditions for a square matrix to be invertible

## Theorem

*Let  $A$  be an  $n \times n$  matrix over  $\mathbb{R}$ . The following are equivalent:*

- ①  *$A$  is invertible.*
- ② *The homogeneous system  $AX = 0$  has only the trivial solution.*
- ③ *For every  $b \in \mathbb{R}^n$ , the system  $AX = b$  has a solution.*
- ④  *$\mathbb{R}^n$  is spanned by the column vectors of  $A$ .*
- ⑤ *The column vectors of  $A$  are linearly independent.*
- ⑥  *$\mathbb{R}^n$  is spanned by the row vectors of  $A$ .*
- ⑦ *The row vectors of  $A$  are linearly independent.*

**Proof.** Consider  $A$  as a linear map  $A : \mathbb{R}^n \rightarrow \mathbb{R}^n$ . Then (2) is same as  $\text{Null}(A) = 0$ . Moreover (3) is same as  $A$  is surjective.

# Conditions for a square matrix to be invertible

## Theorem

*Let  $A$  be an  $n \times n$  matrix over  $\mathbb{R}$ . The following are equivalent:*

- 1  *$A$  is invertible.*
- 2 *The homogeneous system  $AX = 0$  has only the trivial solution.*
- 3 *For every  $b \in \mathbb{R}^n$ , the system  $AX = b$  has a solution.*
- 4  *$\mathbb{R}^n$  is spanned by the column vectors of  $A$ .*
- 5 *The column vectors of  $A$  are linearly independent.*
- 6  *$\mathbb{R}^n$  is spanned by the row vectors of  $A$ .*
- 7 *The row vectors of  $A$  are linearly independent.*

**Proof.** Consider  $A$  as a linear map  $A : \mathbb{R}^n \rightarrow \mathbb{R}^n$ . Then (2) is same as  $\text{Null}(A) = 0$ . Moreover (3) is same as  $A$  is surjective. Thus, by the previous theorem, we have (1), (2) and (3) are equivalent.

# Conditions for a square matrix to be invertible

## Theorem

*Let  $A$  be an  $n \times n$  matrix over  $\mathbb{R}$ . The following are equivalent:*

- ①  *$A$  is invertible.*
- ② *The homogeneous system  $AX = 0$  has only the trivial solution.*
- ③ *For every  $b \in \mathbb{R}^n$ , the system  $AX = b$  has a solution.*
- ④  *$\mathbb{R}^n$  is spanned by the column vectors of  $A$ .*
- ⑤ *The column vectors of  $A$  are linearly independent.*
- ⑥  *$\mathbb{R}^n$  is spanned by the row vectors of  $A$ .*
- ⑦ *The row vectors of  $A$  are linearly independent.*

**Proof.** Consider  $A$  as a linear map  $A : \mathbb{R}^n \rightarrow \mathbb{R}^n$ . Then (2) is same as  $\text{Null}(A) = 0$ . Moreover (3) is same as  $A$  is surjective. Thus, by the previous theorem, we have (1), (2) and (3) are equivalent.

Since  $AX$  is nothing but a linear combination of column vectors of  $A$ ,

# Conditions for a square matrix to be invertible

## Theorem

*Let  $A$  be an  $n \times n$  matrix over  $\mathbb{R}$ . The following are equivalent:*

- ①  *$A$  is invertible.*
- ② *The homogeneous system  $AX = 0$  has only the trivial solution.*
- ③ *For every  $b \in \mathbb{R}^n$ , the system  $AX = b$  has a solution.*
- ④  *$\mathbb{R}^n$  is spanned by the column vectors of  $A$ .*
- ⑤ *The column vectors of  $A$  are linearly independent.*
- ⑥  *$\mathbb{R}^n$  is spanned by the row vectors of  $A$ .*
- ⑦ *The row vectors of  $A$  are linearly independent.*

**Proof.** Consider  $A$  as a linear map  $A : \mathbb{R}^n \rightarrow \mathbb{R}^n$ . Then (2) is same as  $\text{Null}(A) = 0$ . Moreover (3) is same as  $A$  is surjective. Thus, by the previous theorem, we have (1), (2) and (3) are equivalent. Since  $AX$  is nothing but a linear combination of column vectors of  $A$ , it follows that (3) and (4) are equivalent.

# Conditions for a square matrix to be invertible contd...

## Theorem

*Let  $A$  be an  $n \times n$  matrix over  $\mathbb{R}$ . The following are equivalent:*

- ①  *$A$  is invertible.*
- ② *The homogeneous system  $AX = 0$  has only the trivial solution.*
- ③ *For every  $b \in \mathbb{R}^n$ , the system  $AX = b$  has a solution.*
- ④  *$\mathbb{R}^n$  is spanned by the column vectors of  $A$ .*
- ⑤ *The column vectors of  $A$  are linearly independent.*
- ⑥  *$\mathbb{R}^n$  is spanned by the row vectors of  $A$ .*
- ⑦ *The row vectors of  $A$  are linearly independent.*

**Proof.** We already proved  $(1) \Leftrightarrow (2) \Leftrightarrow (3) \Leftrightarrow (4)$ .

# Conditions for a square matrix to be invertible contd...

## Theorem

*Let  $A$  be an  $n \times n$  matrix over  $\mathbb{R}$ . The following are equivalent:*

- ①  *$A$  is invertible.*
- ② *The homogeneous system  $AX = 0$  has only the trivial solution.*
- ③ *For every  $b \in \mathbb{R}^n$ , the system  $AX = b$  has a solution.*
- ④  *$\mathbb{R}^n$  is spanned by the column vectors of  $A$ .*
- ⑤ *The column vectors of  $A$  are linearly independent.*
- ⑥  *$\mathbb{R}^n$  is spanned by the row vectors of  $A$ .*
- ⑦ *The row vectors of  $A$  are linearly independent.*

**Proof.** We already proved  $(1) \Leftrightarrow (2) \Leftrightarrow (3) \Leftrightarrow (4)$ .

$(4) \Leftrightarrow (5)$  and  $(6) \Leftrightarrow (7)$ :

# Conditions for a square matrix to be invertible contd...

## Theorem

*Let  $A$  be an  $n \times n$  matrix over  $\mathbb{R}$ . The following are equivalent:*

- ①  *$A$  is invertible.*
- ② *The homogeneous system  $AX = 0$  has only the trivial solution.*
- ③ *For every  $b \in \mathbb{R}^n$ , the system  $AX = b$  has a solution.*
- ④  *$\mathbb{R}^n$  is spanned by the column vectors of  $A$ .*
- ⑤ *The column vectors of  $A$  are linearly independent.*
- ⑥  *$\mathbb{R}^n$  is spanned by the row vectors of  $A$ .*
- ⑦ *The row vectors of  $A$  are linearly independent.*

**Proof.** We already proved  $(1) \Leftrightarrow (2) \Leftrightarrow (3) \Leftrightarrow (4)$ .

$(4) \Leftrightarrow (5)$  and  $(6) \Leftrightarrow (7)$ : Since  $\dim(\mathbb{R}^n) = n$ ,

# Conditions for a square matrix to be invertible contd...

## Theorem

*Let  $A$  be an  $n \times n$  matrix over  $\mathbb{R}$ . The following are equivalent:*

- ①  *$A$  is invertible.*
- ② *The homogeneous system  $AX = 0$  has only the trivial solution.*
- ③ *For every  $b \in \mathbb{R}^n$ , the system  $AX = b$  has a solution.*
- ④  *$\mathbb{R}^n$  is spanned by the column vectors of  $A$ .*
- ⑤ *The column vectors of  $A$  are linearly independent.*
- ⑥  *$\mathbb{R}^n$  is spanned by the row vectors of  $A$ .*
- ⑦ *The row vectors of  $A$  are linearly independent.*

**Proof.** We already proved  $(1) \Leftrightarrow (2) \Leftrightarrow (3) \Leftrightarrow (4)$ .

$(4) \Leftrightarrow (5)$  and  $(6) \Leftrightarrow (7)$ : Since  $\dim(\mathbb{R}^n) = n$ , any collection of  $n$  vectors in  $\mathbb{R}^n$  is linearly independent



# Conditions for a square matrix to be invertible contd...

## Theorem

*Let  $A$  be an  $n \times n$  matrix over  $\mathbb{R}$ . The following are equivalent:*

- ①  *$A$  is invertible.*
- ② *The homogeneous system  $AX = 0$  has only the trivial solution.*
- ③ *For every  $b \in \mathbb{R}^n$ , the system  $AX = b$  has a solution.*
- ④  *$\mathbb{R}^n$  is spanned by the column vectors of  $A$ .*
- ⑤ *The column vectors of  $A$  are linearly independent.*
- ⑥  *$\mathbb{R}^n$  is spanned by the row vectors of  $A$ .*
- ⑦ *The row vectors of  $A$  are linearly independent.*

**Proof.** We already proved  $(1) \Leftrightarrow (2) \Leftrightarrow (3) \Leftrightarrow (4)$ .

$(4) \Leftrightarrow (5)$  and  $(6) \Leftrightarrow (7)$ : Since  $\dim(\mathbb{R}^n) = n$ , any collection of  $n$  vectors in  $\mathbb{R}^n$  is linearly independent if and only if it spans  $\mathbb{R}^n$ .

# Conditions for a square matrix to be invertible contd...

## Theorem

*Let  $A$  be an  $n \times n$  matrix over  $\mathbb{R}$ . The following are equivalent:*

- ①  *$A$  is invertible.*
- ② *The homogeneous system  $AX = 0$  has only the trivial solution.*
- ③ *For every  $b \in \mathbb{R}^n$ , the system  $AX = b$  has a solution.*
- ④  *$\mathbb{R}^n$  is spanned by the column vectors of  $A$ .*
- ⑤ *The column vectors of  $A$  are linearly independent.*
- ⑥  *$\mathbb{R}^n$  is spanned by the row vectors of  $A$ .*
- ⑦ *The row vectors of  $A$  are linearly independent.*

**Proof.** We already proved  $(1) \Leftrightarrow (2) \Leftrightarrow (3) \Leftrightarrow (4)$ .

$(4) \Leftrightarrow (5)$  and  $(6) \Leftrightarrow (7)$ : Since  $\dim(\mathbb{R}^n) = n$ , any collection of  $n$  vectors in  $\mathbb{R}^n$  is linearly independent if and only if it spans  $\mathbb{R}^n$ .

$(4) \Leftrightarrow (6)$ :

# Conditions for a square matrix to be invertible contd...

## Theorem

*Let  $A$  be an  $n \times n$  matrix over  $\mathbb{R}$ . The following are equivalent:*

- ①  *$A$  is invertible.*
- ② *The homogeneous system  $AX = 0$  has only the trivial solution.*
- ③ *For every  $b \in \mathbb{R}^n$ , the system  $AX = b$  has a solution.*
- ④  *$\mathbb{R}^n$  is spanned by the column vectors of  $A$ .*
- ⑤ *The column vectors of  $A$  are linearly independent.*
- ⑥  *$\mathbb{R}^n$  is spanned by the row vectors of  $A$ .*
- ⑦ *The row vectors of  $A$  are linearly independent.*

**Proof.** We already proved  $(1) \Leftrightarrow (2) \Leftrightarrow (3) \Leftrightarrow (4)$ .

$(4) \Leftrightarrow (5)$  and  $(6) \Leftrightarrow (7)$ : Since  $\dim(\mathbb{R}^n) = n$ , any collection of  $n$  vectors in  $\mathbb{R}^n$  is linearly independent if and only if it spans  $\mathbb{R}^n$ .

$(4) \Leftrightarrow (6)$ : It follows from the above equivalences “ $(4) \Leftrightarrow (5)$  and  $(6) \Leftrightarrow (7)$ ”

# Conditions for a square matrix to be invertible contd...

## Theorem

*Let  $A$  be an  $n \times n$  matrix over  $\mathbb{R}$ . The following are equivalent:*

- ①  *$A$  is invertible.*
- ② *The homogeneous system  $AX = 0$  has only the trivial solution.*
- ③ *For every  $b \in \mathbb{R}^n$ , the system  $AX = b$  has a solution.*
- ④  *$\mathbb{R}^n$  is spanned by the column vectors of  $A$ .*
- ⑤ *The column vectors of  $A$  are linearly independent.*
- ⑥  *$\mathbb{R}^n$  is spanned by the row vectors of  $A$ .*
- ⑦ *The row vectors of  $A$  are linearly independent.*

**Proof.** We already proved  $(1) \Leftrightarrow (2) \Leftrightarrow (3) \Leftrightarrow (4)$ .

$(4) \Leftrightarrow (5)$  and  $(6) \Leftrightarrow (7)$ : Since  $\dim(\mathbb{R}^n) = n$ , any collection of  $n$  vectors in  $\mathbb{R}^n$  is linearly independent if and only if it spans  $\mathbb{R}^n$ .

$(4) \Leftrightarrow (6)$ : It follows from the above equivalences “ $(4) \Leftrightarrow (5)$  and  $(6) \Leftrightarrow (7)$ ” and the fact that  $\text{column rank}(A) = \text{row rank}(A)$ .

# Thank You!