

Differential Equations (MA 1150)

Sukumar

Lecture 3

April 17, 2020

Definition An ODE of order n is called **linear** if it can be written as

$$a_0(x)y^{(n)} + a_1(x)y^{(n-1)} + \cdots + a_n(x)y = b(x) \text{ where } a_0(x) \neq 0.$$

Definition An ODE of order n is called **linear** if it can be written as

$$a_0(x)y^{(n)} + a_1(x)y^{(n-1)} + \cdots + a_n(x)y = b(x) \text{ where } a_0(x) \neq 0.$$

- ▶ Linear $Ay = b$
- ▶ Homogeneous $Ay = 0$ – Find all solutions
- ▶ Non-homogenous $Ay = b$ – Find one solution

Recall: Linear first order non-homogeneous differential equations

Consider the most **general form** of linear first order differential equation. By definition, this looks like

$$p(x)y' + q(x)y = h(x), \text{ where } p(x) \neq 0.$$

Recall: Linear first order non-homogeneous differential equations

Consider the most **general form** of linear first order differential equation. By definition, this looks like

$$p(x)y' + q(x)y = h(x), \text{ where } p(x) \neq 0.$$

Dividing this by $p(x)$, let us rewrite this in **the standard form**

$$y' + a(x)y = f(x).$$

Assume that $a(x)$ and $f(x)$ are defined on the interval $I = (x_0 - \epsilon, x_0 + \epsilon)$.

Recall: Linear first order non-homogeneous differential equations

Consider the most **general form** of linear first order differential equation. By definition, this looks like

$$p(x)y' + q(x)y = h(x), \text{ where } p(x) \neq 0.$$

Dividing this by $p(x)$, let us rewrite this in **the standard form**

$$y' + a(x)y = f(x).$$

Assume that $a(x)$ and $f(x)$ are defined on the interval $I = (x_0 - \epsilon, x_0 + \epsilon)$. The homogeneous equation / **complementary equation** of it is

$$y' + a(x)y = 0.$$

Solving first order linear ODE

To find all solutions of the given

$$y' + a(x)y = f(x).$$

Solving first order linear ODE

To find all solutions of the given

$$y' + a(x)y = f(x).$$

1. Complementary function (CF) - All solutions of homogeneous.
Find solution of complementary equation $y' + a(x)y = 0$.

Solving first order linear ODE

To find all solutions of the given

$$y' + a(x)y = f(x).$$

1. Complementary function (CF) - All solutions of homogeneous.
Find solution of complementary equation $y' + a(x)y = 0$.
2. Particular integral (PI) - One solution of non-homogeneous.
Find one solution of $y' + a(x)y = f(x)$.

Then the general solution of $y' + a(x)y = f(x)$ is

$$y = CF + PI$$

Solving first order linear ODE

To find all solutions of the given

$$y' + a(x)y = f(x).$$

1. Complementary function (CF) - All solutions of homogeneous.
Find solution of complementary equation $y' + a(x)y = 0$.
2. Particular integral (PI) - One solution of non-homogeneous.
Find one solution of $y' + a(x)y = f(x)$.

Then the general solution of $y' + a(x)y = f(x)$ is

$$y = CF + PI$$

Recall you have done the same in linear algebra in solving $Ay = b$. For example solving $u + v = 3$ for (u, v)

Complementary function

Solving homogeneous equation

Note $y(x) = 0$ is always a solution, called **trivial solution**.

Complementary function

Solving homogeneous equation

Note $y(x) = 0$ is always a solution, called **trivial solution**.

Finding non-trivial solution is easy as we can separate the variables.

Complementary function

Solving homogeneous equation

Note $y(x) = 0$ is always a solution, called **trivial solution**.

Finding non-trivial solution is easy as we can separate the variables.

$$y' + a(x)y = 0$$

$$\frac{dy}{dx} = -a(x)y$$

$$\frac{dy}{y} = -a(x)dx, \text{ by separating variables}$$

$$\ln y = - \int a(x)dx + c, \text{ on integration}$$

$$y = Ce^{-\int a(x)dx}, \text{ on integration and renaming } e^c \text{ as } C$$

Hence the CF is $y_h(x) = Ce^{-\int a(x)dx}$

Complementary function

Solving homogeneous equation

Note $y(x) = 0$ is always a solution, called **trivial solution**.

Finding non-trivial solution is easy as we can separate the variables.

$$y' + a(x)y = 0$$

$$\frac{dy}{dx} = -a(x)y$$

$$\frac{dy}{y} = -a(x)dx, \text{ by separating variables}$$

$$\ln y = - \int a(x)dx + c, \text{ on integration}$$

$$y = Ce^{-\int a(x)dx}, \text{ on integration and renaming } e^c \text{ as } C$$

Hence the CF is $y_h(x) = Ce^{-\int a(x)dx}$

Complementary function

Solving homogeneous equation

Note $y(x) = 0$ is always a solution, called **trivial solution**.

Finding non-trivial solution is easy as we can separate the variables.

$$y' + a(x)y = 0$$

$$\frac{dy}{dx} = -a(x)y$$

$$\frac{dy}{y} = -a(x)dx, \text{ by separating variables}$$

$$\ln y = - \int a(x)dx + c, \text{ on integration}$$

$$y = Ce^{-\int a(x)dx}, \text{ on integration and renaming } e^c \text{ as } C$$

Hence the CF is $y_h(x) = Ce^{-\int a(x)dx}$

If y_h is a solution then cy_h a constant multiple as well

Complementary function

Solving homogeneous equation

Note $y(x) = 0$ is always a solution, called **trivial solution**.

Finding non-trivial solution is easy as we can separate the variables.

$$y' + a(x)y = 0$$

$$\frac{dy}{dx} = -a(x)y$$

$$\frac{dy}{y} = -a(x)dx, \text{ by separating variables}$$

$$\ln y = - \int a(x)dx + c, \text{ on integration}$$

$$y = Ce^{-\int a(x)dx}, \text{ on integration and renaming } e^c \text{ as } C$$

Hence the CF is $y_h(x) = Ce^{-\int a(x)dx}$

If y_h is a solution then cy_h a constant multiple as well

If the co-efficient function is a constant (say α) then $y_h(x) = Ce^{-\alpha x}$.

Particular Integral

Solving non-homogeneous equation

Let $y_h(x)$ be a solution to the homogeneous equation.

Particular Integral

Solving non-homogeneous equation

Let $y_h(x)$ be a solution to the homogeneous equation. We look for a solution of the type $y_p(x) = u(x)y_h(x)$. Substituting this into the non-homogeneous differential equation we get

$$u'y_h + uy'_h + auy_h = f(x).$$

Since $y'_h + a(x)y_h = 0$, we get

$$u'y_h = f(x).$$

$$u(x) = \int_{x_0}^x \frac{f(s)}{y_h(s)} ds + c$$

Hence a particular integral is

$$y_p(x) = y_h(x) \left(\int_{x_0}^x \frac{f(s)}{y_h(s)} ds + c \right).$$

The general solution of non-homogeneous equation is

$$\begin{aligned}y(x) &= CF + PI \\&= y_h(x) + y_p(x) \\y(x) &= \alpha y_h(x) + y_h(s) \left(\int_{x_0}^x \frac{f(s)}{y_h(s)} ds \right)\end{aligned}$$

Example

(1) Solve the ODE $y' - 2xy = 1$.

Note that $y' - 2xy = 0$ has a solution $y_1(x) = e^{x^2}$.

The solution of ODE is $y = uy_1$, where

$$u'y_1 = 1 \implies u(x) = \int_0^x e^{s^2} ds + C$$

and this implies that

$$y(x) = e^{x^2} \left(\int_0^x e^{s^2} ds + C \right).$$

(2) Solve the ODE $y' - 2xy = 1$, where $y(0) = y_0$. Write the solution of ODE as

$$y(x) = e^{x^2} \left(\int_0^x e^{s^2} ds + C \right).$$

Then $y(0) = y_0$ gives $C = y_0$.

Definition. An ordinary differential equation of order n is an equation

$$F(x, y, y', y^{(2)}, \dots, y^{(n)}) = 0,$$

where F is a function of $(n + 2)$ -variables.

Solution of an ODE

Definition. Suppose we are given an ODE of order n

Let $x_0 \in \mathbb{R}$ and suppose there is a function $y(x)$ which is defined in a small neighborhood $(x_0 - \epsilon, x_0 + \epsilon)$, and which is n times differentiable in this interval.

If $F(x, y, y', y^{(2)}, \dots, y^{(n)}) = 0$, then we say that the function $y(x)$ is a **solution to the ODE** in the interval $(x_0 - \epsilon, x_0 + \epsilon)$.

(a) In view of the above definition, we can ask **what is the largest open interval around x_0** on which the given **ODE has a solution**

(b) If $y(x)$ is a solution to the ODE around x_0 , then the largest interval around x_0 in which $y(x)$ is defined is called the **interval of validity** for the solution y .

Example

(1) The function

$$y = \frac{x^2}{3} + \frac{1}{x}$$

satisfies

$$xy' + y = x^2$$

on $(-\infty, 0) \cup (0, \infty)$.

(2) For IVP

$$xy' + y = x^2, \text{ where } y(1) = \frac{4}{3}$$

the interval of validity of $y(x)$ is $(0, \infty)$.

(3) For IVP

$$xy' + y = x^2, \text{ where } y(-1) = -\frac{4}{3}$$

the interval of validity of $y(x)$ is $(-\infty, 0)$.

Definition of continuous function!

Definition of continuous function!

Definition and geometrical meaning of partial derivatives!

Theorem. If $G = G(x, y)$ has continuous partial derivatives G_x and G_y , then

$$G(x, y) = c \quad (c=\text{constant}), \quad (1)$$

is an implicit solution of the differential equation

$$G_x(x, y) dx + G_y(x, y) dy = 0. \quad (2)$$

Theorem. If $G = G(x, y)$ has continuous partial derivatives G_x and G_y , then

$$G(x, y) = c \quad (c=\text{constant}), \quad (1)$$

is an implicit solution of the differential equation

$$G_x(x, y) dx + G_y(x, y) dy = 0. \quad (2)$$

Proof. Assuming y as a function of x and differentiating (1) w.r.t x yields

$$G_x(x, y) + G_y(x, y) \frac{dy}{dx} = 0.$$

Theorem. If $G = G(x, y)$ has continuous partial derivatives G_x and G_y , then

$$G(x, y) = c \quad (c=\text{constant}), \quad (1)$$

is an implicit solution of the differential equation

$$G_x(x, y) dx + G_y(x, y) dy = 0. \quad (2)$$

Proof. Assuming y as a function of x and differentiating (1) w.r.t x yields

$$G_x(x, y) + G_y(x, y) \frac{dy}{dx} = 0.$$

Similarly, regarding x as a function of y and differentiating (1) w.r.t y yields

$$G_x(x, y) \frac{dx}{dy} + G_y(x, y) = 0.$$

Theorem. If $G = G(x, y)$ has continuous partial derivatives G_x and G_y , then

$$G(x, y) = c \quad (c=\text{constant}), \quad (1)$$

is an implicit solution of the differential equation

$$G_x(x, y) dx + G_y(x, y) dy = 0. \quad (2)$$

Proof. Assuming y as a function of x and differentiating (1) w.r.t x yields

$$G_x(x, y) + G_y(x, y) \frac{dy}{dx} = 0.$$

Similarly, regarding x as a function of y and differentiating (1) w.r.t y yields

$$G_x(x, y) \frac{dx}{dy} + G_y(x, y) = 0.$$

Thus (1) is an implicit solution of (2).

Exact ODE

Definition A first order ODE written in the form

$$M(x, y) + N(x, y) \frac{dy}{dx} = 0 \quad (3)$$

is said to be **exact** on open rectangle \mathcal{R} if

Exact ODE

Definition A first order ODE written in the form

$$M(x, y) + N(x, y) \frac{dy}{dx} = 0 \quad (3)$$

is said to be **exact** on open rectangle \mathcal{R} if there exists a function $G = G(x, y)$ such that $\frac{\partial G}{\partial x}$ and $\frac{\partial G}{\partial y}$ are continuous and

$$\frac{\partial G}{\partial x} = M(x, y) \quad \text{and} \quad \frac{\partial G}{\partial y} = N(x, y) \quad (4)$$

for all (x, y) in \mathcal{R} .

Exact ODE

Definition A first order ODE written in the form

$$M(x, y) + N(x, y) \frac{dy}{dx} = 0 \quad (3)$$

is said to be **exact** on open rectangle \mathcal{R} if there exists a function $G = G(x, y)$ such that $\frac{\partial G}{\partial x}$ and $\frac{\partial G}{\partial y}$ are continuous and

$$\frac{\partial G}{\partial x} = M(x, y) \quad \text{and} \quad \frac{\partial G}{\partial y} = N(x, y) \quad (4)$$

for all (x, y) in \mathcal{R} .

Question 1. Given an equation (3), how can we determine whether it's exact?

Exact ODE

Definition A first order ODE written in the form

$$M(x, y) + N(x, y) \frac{dy}{dx} = 0 \quad (3)$$

is said to be **exact** on open rectangle \mathcal{R} if there exists a function $G = G(x, y)$ such that $\frac{\partial G}{\partial x}$ and $\frac{\partial G}{\partial y}$ are continuous and

$$\frac{\partial G}{\partial x} = M(x, y) \quad \text{and} \quad \frac{\partial G}{\partial y} = N(x, y) \quad (4)$$

for all (x, y) in \mathcal{R} .

Question 1. Given an equation (3), how can we determine whether it's exact?

Question 2. If (3) is exact, how do we find a function G satisfying (4)?

When is an ODE exact ?

Theorem.(The Exactness Condition) Consider the ODE

$$M(x, y) + N(x, y) \frac{dy}{dx} = 0. \quad (5)$$

When is an ODE exact ?

Theorem.(The Exactness Condition) Consider the ODE

$$M(x, y) + N(x, y) \frac{dy}{dx} = 0. \quad (5)$$

Assume that functions M , N , $\frac{\partial M}{\partial y}$, $\frac{\partial N}{\partial x}$ are continuous on an open rectangle $\mathcal{R} := \{(x, y) \in \mathbb{R}^2 : (x, y) \in (a, b) \times (c, d)\}$.

When is an ODE exact ?

Theorem.(The Exactness Condition) Consider the ODE

$$M(x, y) + N(x, y) \frac{dy}{dx} = 0. \quad (5)$$

Assume that functions M , N , $\frac{\partial M}{\partial y}$, $\frac{\partial N}{\partial x}$ are continuous on an open rectangle $\mathcal{R} := \{(x, y) \in \mathbb{R}^2 : (x, y) \in (a, b) \times (c, d)\}$.

Then (5) is an exact ODE on open rectangle \mathcal{R} if and only if M and N satisfies the condition

$$\frac{\partial M}{\partial y} = \frac{\partial N}{\partial x}$$

for all $(x, y) \in \mathcal{R}$.

When is an ODE exact ?

Theorem.(The Exactness Condition) Consider the ODE

$$M(x, y) + N(x, y) \frac{dy}{dx} = 0. \quad (5)$$

Assume that functions M , N , $\frac{\partial M}{\partial y}$, $\frac{\partial N}{\partial x}$ are continuous on an open rectangle $\mathcal{R} := \{(x, y) \in \mathbb{R}^2 : (x, y) \in (a, b) \times (c, d)\}$.

Then (5) is an exact ODE on open rectangle \mathcal{R} if and only if M and N satisfies the condition

$$\frac{\partial M}{\partial y} = \frac{\partial N}{\partial x}$$

for all $(x, y) \in \mathcal{R}$.

In other words, there exists a function $G = G(x, y)$ such that

When is an ODE exact ?

Theorem.(The Exactness Condition) Consider the ODE

$$M(x, y) + N(x, y) \frac{dy}{dx} = 0. \quad (5)$$

Assume that functions M , N , $\frac{\partial M}{\partial y}$, $\frac{\partial N}{\partial x}$ are continuous on an open rectangle $\mathcal{R} := \{(x, y) \in \mathbb{R}^2 : (x, y) \in (a, b) \times (c, d)\}$.

Then (5) is an exact ODE on open rectangle \mathcal{R} if and only if M and N satisfies the condition

$$\frac{\partial M}{\partial y} = \frac{\partial N}{\partial x}$$

for all $(x, y) \in \mathcal{R}$.

In other words, there exists a function $G = G(x, y)$ such that $\frac{\partial G}{\partial x}$ and $\frac{\partial G}{\partial y}$ are continuous and

$$\frac{\partial G}{\partial x} = M(x, y) \quad \text{and} \quad \frac{\partial G}{\partial y} = N(x, y).$$

Exact Equations

Which of the following ODE's are exact?

Exact Equations

Which of the following ODE's are exact?

▶ $(2x + 3) + (2y - 2) \frac{dy}{dx} = 0.$

▶ $\frac{dy}{dx} = \frac{ax + by}{bx + cy}.$

▶ $\left(\frac{y}{x} + 6x\right) + (\ln x - 2) \frac{dy}{dx} = 0, \text{ where } x, y > 0.$

▶ $(3x^2y + 2xy + y^3) + (x^2 + y^2) \frac{dy}{dx} = 0.$

Exact Equations

Solve $(2x + 3) + (2y - 2)\frac{dy}{dx} = 0$.

Exact Equations

Solve $(2x + 3) + (2y - 2)\frac{dy}{dx} = 0$.

The ODE is exact, so we need to find $G(x, y)$ such that

$$\frac{\partial G}{\partial x} = 2x + 3 \text{ and } \frac{\partial G}{\partial y} = 2y - 2.$$

Exact Equations

Solve $(2x + 3) + (2y - 2)\frac{dy}{dx} = 0$.

The ODE is exact, so we need to find $G(x, y)$ such that

$$\frac{\partial G}{\partial x} = 2x + 3 \text{ and } \frac{\partial G}{\partial y} = 2y - 2.$$

Integrating first equation w.r.t. x gives

$$G(x, y) = x^2 + 3x + h(y).$$

Exact Equations

Solve $(2x + 3) + (2y - 2)\frac{dy}{dx} = 0$.

The ODE is exact, so we need to find $G(x, y)$ such that

$$\frac{\partial G}{\partial x} = 2x + 3 \text{ and } \frac{\partial G}{\partial y} = 2y - 2.$$

Integrating first equation w.r.t. x gives

$$G(x, y) = x^2 + 3x + h(y).$$

This gives

$$\frac{\partial G}{\partial y} = \frac{dh}{dy} = 2y - 2 \implies h(y) = y^2 - 2y + c_1.$$

Exact Equations

Solve $(2x + 3) + (2y - 2)\frac{dy}{dx} = 0$.

The ODE is exact, so we need to find $G(x, y)$ such that

$$\frac{\partial G}{\partial x} = 2x + 3 \text{ and } \frac{\partial G}{\partial y} = 2y - 2.$$

Integrating first equation w.r.t. x gives

$$G(x, y) = x^2 + 3x + h(y).$$

This gives

$$\frac{\partial G}{\partial y} = \frac{dh}{dy} = 2y - 2 \implies h(y) = y^2 - 2y + c_1.$$

Therefore, an implicit solution to ODE is

$$G(x, y) = x^2 + 3x + y^2 - 2y + c.$$

Exact Equations

Solve

$$(3x^2 + 6xy^2) + (6x^2y + 4y^3)\frac{dy}{dx} = 0.$$

Exact Equations

Solve

$$(3x^2 + 6xy^2) + (6x^2y + 4y^3)\frac{dy}{dx} = 0.$$

Here $M(x, y) = 3x^2 + 6xy^2$ and $N(x, y) = 6x^2y + 4y^3$.

Exact Equations

Solve

$$(3x^2 + 6xy^2) + (6x^2y + 4y^3)\frac{dy}{dx} = 0.$$

Here $M(x, y) = 3x^2 + 6xy^2$ and $N(x, y) = 6x^2y + 4y^3$.

Notice that $\frac{\partial M}{\partial y} = 12xy \equiv 12xy = \frac{\partial N}{\partial x}$.

Exact Equations

Solve

$$(3x^2 + 6xy^2) + (6x^2y + 4y^3)\frac{dy}{dx} = 0.$$

Here $M(x, y) = 3x^2 + 6xy^2$ and $N(x, y) = 6x^2y + 4y^3$.

Notice that $\frac{\partial M}{\partial y} = 12xy \equiv 12xy = \frac{\partial N}{\partial x}$.

The ODE is exact, so we need to find $G(x, y)$ such that

$$\frac{\partial G}{\partial x} = 3x^2 + 6xy^2 \text{ and } \frac{\partial G}{\partial y} = 6x^2y + 4y^3.$$

Exact Equations

Solve

$$(3x^2 + 6xy^2) + (6x^2y + 4y^3)\frac{dy}{dx} = 0.$$

Here $M(x, y) = 3x^2 + 6xy^2$ and $N(x, y) = 6x^2y + 4y^3$.

Notice that $\frac{\partial M}{\partial y} = 12xy \equiv 12xy = \frac{\partial N}{\partial x}$.

The ODE is exact, so we need to find $G(x, y)$ such that

$$\frac{\partial G}{\partial x} = 3x^2 + 6xy^2 \text{ and } \frac{\partial G}{\partial y} = 6x^2y + 4y^3.$$

The solution to ODE is

$$G(x, y) = x^3 + 3x^2y^2 + y^4 + c.$$

Exact Equations

Solve $\left(\frac{y}{x} + 6x\right) + (\ln x - 2) \frac{dy}{dx} = 0$, where $x, y > 0$.

Exact Equations

Solve $\left(\frac{y}{x} + 6x\right) + (\ln x - 2) \frac{dy}{dx} = 0$, where $x, y > 0$.

Check that given ODE is exact. We need to find $G(x, y)$ such that

$$\frac{\partial G}{\partial x} = \frac{y}{x} + 6x \text{ and } \frac{\partial G}{\partial y} = \ln x - 2.$$

Exact Equations

Solve $\left(\frac{y}{x} + 6x\right) + (\ln x - 2) \frac{dy}{dx} = 0$, where $x, y > 0$.

Check that given ODE is exact. We need to find $G(x, y)$ such that

$$\frac{\partial G}{\partial x} = \frac{y}{x} + 6x \text{ and } \frac{\partial G}{\partial y} = \ln x - 2.$$

Integrating first equation w.r.t. x gives

$$G(x, y) = y \ln |x| + 3x^2 + h(y).$$

Exact Equations

Solve $\left(\frac{y}{x} + 6x\right) + (\ln x - 2) \frac{dy}{dx} = 0$, where $x, y > 0$.

Check that given ODE is exact. We need to find $G(x, y)$ such that

$$\frac{\partial G}{\partial x} = \frac{y}{x} + 6x \text{ and } \frac{\partial G}{\partial y} = \ln x - 2.$$

Integrating first equation w.r.t. x gives

$$G(x, y) = y \ln |x| + 3x^2 + h(y).$$

Differentiate w.r.t. y to obtain

$$\frac{\partial G}{\partial y} = \ln |x| + \frac{dh}{dy} = \ln x - 2 \implies h(y) = -2y + c.$$

Exact Equations

Solve $\left(\frac{y}{x} + 6x\right) + (\ln x - 2) \frac{dy}{dx} = 0$, where $x, y > 0$.

Check that given ODE is exact. We need to find $G(x, y)$ such that

$$\frac{\partial G}{\partial x} = \frac{y}{x} + 6x \text{ and } \frac{\partial G}{\partial y} = \ln x - 2.$$

Integrating first equation w.r.t. x gives

$$G(x, y) = y \ln |x| + 3x^2 + h(y).$$

Differentiate w.r.t. y to obtain

$$\frac{\partial G}{\partial y} = \ln |x| + \frac{dh}{dy} = \ln x - 2 \implies h(y) = -2y + c.$$

Therefore, an implicit solution to ODE is

$$G(x, y) = y \ln |x| + 3x^2 - 2y + c.$$

Method of integrating factor

Example. Solve

$$(3x^2y + 2xy + y^3) + (x^2 + y^2)\frac{dy}{dx} = 0.$$

Method of integrating factor

Example. Solve

$$(3x^2y + 2xy + y^3) + (x^2 + y^2)\frac{dy}{dx} = 0.$$

Here $M = 3x^2y + 2xy + y^3$, and $N = x^2 + y^2$.

Method of integrating factor

Example. Solve

$$(3x^2y + 2xy + y^3) + (x^2 + y^2)\frac{dy}{dx} = 0.$$

Here $M = 3x^2y + 2xy + y^3$, and $N = x^2 + y^2$. We have

$$\frac{\partial M}{\partial y} = 3x^2 + 2x + 3y^2, \text{ and } \frac{\partial N}{\partial x} = 2x.$$

Method of integrating factor

Example. Solve

$$(3x^2y + 2xy + y^3) + (x^2 + y^2)\frac{dy}{dx} = 0.$$

Here $M = 3x^2y + 2xy + y^3$, and $N = x^2 + y^2$. We have

$$\frac{\partial M}{\partial y} = 3x^2 + 2x + 3y^2, \text{ and } \frac{\partial N}{\partial x} = 2x.$$

Therefore note that **ODE is not exact**.

Method of integrating factor

Example. Solve

$$(3x^2y + 2xy + y^3) + (x^2 + y^2)\frac{dy}{dx} = 0.$$

Here $M = 3x^2y + 2xy + y^3$, and $N = x^2 + y^2$. We have

$$\frac{\partial M}{\partial y} = 3x^2 + 2x + 3y^2, \text{ and } \frac{\partial N}{\partial x} = 2x.$$

Therefore note that **ODE is not exact**.

Question. Can we multiply the ODE by a function $\mu(x, y)$ so that it becomes exact.

Integrating factors

Definition. We say that a function $\mu(x, y)$ is an integrating factor of ODE

$$M(x, y) + N(x, y) \frac{dy}{dx} = 0 \quad (6)$$

Integrating factors

Definition. We say that a function $\mu(x, y)$ is an integrating factor of ODE

$$M(x, y) + N(x, y) \frac{dy}{dx} = 0 \quad (6)$$

if

$$\mu(x, y)M(x, y) + \mu(x, y)N(x, y) \frac{dy}{dx} = 0. \quad (7)$$

is exact.

Integrating factors

Definition. We say that a function $\mu(x, y)$ is an integrating factor of ODE

$$M(x, y) + N(x, y) \frac{dy}{dx} = 0 \quad (6)$$

if

$$\mu(x, y)M(x, y) + \mu(x, y)N(x, y) \frac{dy}{dx} = 0. \quad (7)$$

is exact.

Question. What if the ODE was already exact?

Integrating factors

Definition. We say that a function $\mu(x, y)$ is an integrating factor of ODE

$$M(x, y) + N(x, y) \frac{dy}{dx} = 0 \quad (6)$$

if

$$\mu(x, y)M(x, y) + \mu(x, y)N(x, y) \frac{dy}{dx} = 0. \quad (7)$$

is exact.

Question. What if the ODE was already exact?

The Exactness Theorem says that (7) is exact on an open rectangle \mathcal{R} if μM , μN , $\frac{\partial}{\partial y}(\mu M)$, and $\frac{\partial}{\partial x}(\mu N)$ are continuous

Integrating factors

Definition. We say that a function $\mu(x, y)$ is an integrating factor of ODE

$$M(x, y) + N(x, y) \frac{dy}{dx} = 0 \quad (6)$$

if

$$\mu(x, y)M(x, y) + \mu(x, y)N(x, y) \frac{dy}{dx} = 0. \quad (7)$$

is exact.

Question. What if the ODE was already exact?

The Exactness Theorem says that (7) is exact on an open rectangle \mathcal{R} if μM , μN , $\frac{\partial}{\partial y}(\mu M)$, and $\frac{\partial}{\partial x}(\mu N)$ are continuous and

$$\frac{\partial}{\partial y}(\mu M) = \frac{\partial}{\partial x}(\mu N)$$

on \mathcal{R} .

Integrating factors

Thus

$$\frac{\partial}{\partial y}(\mu M) = \frac{\partial}{\partial x}(\mu N)$$

gives

Integrating factors

Thus

$$\frac{\partial}{\partial y}(\mu M) = \frac{\partial}{\partial x}(\mu N)$$

gives

$$\mu_y M + \mu M_y = \mu_x N + \mu N_x.$$

Integrating factors

Thus

$$\frac{\partial}{\partial y}(\mu M) = \frac{\partial}{\partial x}(\mu N)$$

gives

$$\mu_y M + \mu M_y = \mu_x N + \mu N_x.$$

It's useful to rewrite the last equation as

$$\mu(M_y - N_x) = \mu_x N - \mu_y M, \tag{8}$$

Integrating factors

Thus

$$\frac{\partial}{\partial y}(\mu M) = \frac{\partial}{\partial x}(\mu N)$$

gives

$$\mu_y M + \mu M_y = \mu_x N + \mu N_x.$$

It's useful to rewrite the last equation as

$$\mu(M_y - N_x) = \mu_x N - \mu_y M, \quad (8)$$

Case study We will divide the discussion in three cases.

Finding the integrating factors

Case 1 If we assume that $\mu = \mu(x)$ is independent of y , then

Finding the integrating factors

Case 1 If we assume that $\mu = \mu(x)$ is independent of y , then

$$\begin{aligned}\mu(M_y - N_x) &= \frac{\partial \mu}{\partial x} N - \frac{\partial \mu}{\partial y} M \\ \implies \mu(M_y - N_x) &= \frac{\partial \mu}{\partial x} N \\ \implies \frac{1}{\mu} \frac{\partial \mu}{\partial x} &= \frac{M_y - N_x}{N} := p(x)\end{aligned}$$

If $\frac{M_y - N_x}{N}$ is a function of x only, say $p(x)$, then

$$\mu(x) = e^{\int p(x) dx}$$

is an integrating factor of ODE $M(x, y) + N(x, y) \frac{dy}{dx} = 0$ on \mathcal{R} .

Finding the integrating factors

Case 1 If we assume that $\mu = \mu(x)$ is independent of y , then

$$\mu(M_y - N_x) = \frac{\partial \mu}{\partial x} N - \frac{\partial \mu}{\partial y} M$$

$$\implies \frac{1}{\mu} \frac{\partial \mu}{\partial x} = \frac{M_y - N_x}{N} := p(x)$$

If $\frac{M_y - N_x}{N}$ is a function of x only, say $p(x)$, then

$$\mu(x) = e^{\int p(x) dx}$$

is an integrating factor of ODE $M(x, y) + N(x, y) \frac{dy}{dx} = 0$ on \mathcal{R} .

Finding the integrating factors

Case 1 If we assume that $\mu = \mu(x)$ is independent of y , then

$$\begin{aligned}\mu(M_y - N_x) &= \frac{\partial \mu}{\partial x} N - \frac{\partial \mu}{\partial y} M \\ \implies \mu(M_y - N_x) &= \frac{\partial \mu}{\partial x} N \\ \implies \frac{1}{\mu} \frac{\partial \mu}{\partial x} &= \frac{M_y - N_x}{N} := p(x)\end{aligned}$$

Finding the integrating factors

Case 1 If we assume that $\mu = \mu(x)$ is independent of y , then

$$\begin{aligned}\mu(M_y - N_x) &= \frac{\partial \mu}{\partial x} N - \frac{\partial \mu}{\partial y} M \\ \implies \mu(M_y - N_x) &= \frac{\partial \mu}{\partial x} N \\ \implies \frac{1}{\mu} \frac{\partial \mu}{\partial x} &= \frac{M_y - N_x}{N} := p(x)\end{aligned}$$

If $\frac{M_y - N_x}{N}$ is a function of x only, say $p(x)$, then

$$\mu(x) = e^{\int p(x) dx}$$

is an integrating factor of ODE $M(x, y) + N(x, y) \frac{dy}{dx} = 0$ on \mathcal{R} .

Finding the integrating factors

Case 2 If we assume that $\mu = \mu(y)$ is independent of x ,

Finding the integrating factors

Case 2 If we assume that $\mu = \mu(y)$ is independent of x , then

$$\begin{aligned}\mu(M_y - N_x) &= \frac{\partial \mu}{\partial x} N - \frac{\partial \mu}{\partial y} M \\ \implies \mu(M_y - N_x) &= -\frac{\partial \mu}{\partial y} M \\ \implies \frac{1}{\mu} \frac{\partial \mu}{\partial y} &= \frac{N_x - M_y}{M} := q(y)\end{aligned}$$

If $\frac{N_x - M_y}{M}$ is a function of y only, say $q(y)$, then

$$\mu = e^{\int q(y) dy}$$

is an integrating factor ODE $M(x, y) + N(x, y) \frac{dy}{dx} = 0$ on \mathcal{R} .

Finding the integrating factors

Case 2 If we assume that $\mu = \mu(y)$ is independent of x , then

$$\mu(M_y - N_x) = \frac{\partial \mu}{\partial x} N - \frac{\partial \mu}{\partial y} M$$

$$\implies \frac{1}{\mu} \frac{\partial \mu}{\partial y} = \frac{N_x - M_y}{M} := q(y)$$

If $\frac{N_x - M_y}{M}$ is a function of y only, say $q(y)$, then

$$\mu = e^{\int q(y) dy}$$

is an integrating factor ODE $M(x, y) + N(x, y) \frac{dy}{dx} = 0$ on \mathcal{R} .

Finding the integrating factors

Case 2 If we assume that $\mu = \mu(y)$ is independent of x , then

$$\begin{aligned}\mu(M_y - N_x) &= \frac{\partial \mu}{\partial x} N - \frac{\partial \mu}{\partial y} M \\ \implies \mu(M_y - N_x) &= -\frac{\partial \mu}{\partial y} M \\ \implies \frac{1}{\mu} \frac{\partial \mu}{\partial y} &= \frac{N_x - M_y}{M} := q(y)\end{aligned}$$

Finding the integrating factors

Case 2 If we assume that $\mu = \mu(y)$ is independent of x , then

$$\begin{aligned}\mu(M_y - N_x) &= \frac{\partial \mu}{\partial x} N - \frac{\partial \mu}{\partial y} M \\ \implies \mu(M_y - N_x) &= -\frac{\partial \mu}{\partial y} M \\ \implies \frac{1}{\mu} \frac{\partial \mu}{\partial y} &= \frac{N_x - M_y}{M} := q(y)\end{aligned}$$

If $\frac{N_x - M_y}{M}$ is a function of y only, say $q(y)$,

Finding the integrating factors

Case 2 If we assume that $\mu = \mu(y)$ is independent of x , then

$$\begin{aligned}\mu(M_y - N_x) &= \frac{\partial \mu}{\partial x} N - \frac{\partial \mu}{\partial y} M \\ \implies \mu(M_y - N_x) &= -\frac{\partial \mu}{\partial y} M \\ \implies \frac{1}{\mu} \frac{\partial \mu}{\partial y} &= \frac{N_x - M_y}{M} := q(y)\end{aligned}$$

If $\frac{N_x - M_y}{M}$ is a function of y only, say $q(y)$, then

$$\mu = e^{\int q(y) dy}$$

is an integrating factor ODE $M(x, y) + N(x, y) \frac{dy}{dx} = 0$ on \mathcal{R} .

Finding the integrating factors

Case 3 If we assume that $\mu(x, y) = P(x)Q(y)$, then

Finding the integrating factors

Case 3 If we assume that $\mu(x, y) = P(x)Q(y)$, then

$$\mu(M_y - N_x) = \frac{\partial \mu}{\partial x} N - \frac{\partial \mu}{\partial y} M$$

Finding the integrating factors

Case 3 If we assume that $\mu(x, y) = P(x)Q(y)$, then

$$\begin{aligned}\mu(M_y - N_x) &= \frac{\partial \mu}{\partial x} N - \frac{\partial \mu}{\partial y} M \\ \implies P(x)Q(y)(M_y - N_x) &= P'(x)Q(y)N - P(x)Q'(y)M\end{aligned}$$

Finding the integrating factors

Case 3 If we assume that $\mu(x, y) = P(x)Q(y)$, then

$$\begin{aligned}\mu(M_y - N_x) &= \frac{\partial \mu}{\partial x} N - \frac{\partial \mu}{\partial y} M \\ \implies P(x)Q(y)(M_y - N_x) &= P'(x)Q(y)N - P(x)Q'(y)M\end{aligned}$$

This implies that

$$M_y - N_x = \frac{P'}{P}N - \frac{Q'}{Q}M.$$

Finding the integrating factors

Case 3 If we assume that $\mu(x, y) = P(x)Q(y)$, then

$$\begin{aligned}\mu(M_y - N_x) &= \frac{\partial \mu}{\partial x} N - \frac{\partial \mu}{\partial y} M \\ \implies P(x)Q(y)(M_y - N_x) &= P'(x)Q(y)N - P(x)Q'(y)M\end{aligned}$$

This implies that

$$M_y - N_x = \frac{P'}{P} N - \frac{Q'}{Q} M.$$

If

$$M_y - N_x = p(x)N - q(y)M, \text{ where } \frac{P'}{P} = p(x), \frac{Q'}{Q} = q(y),$$

Finding the integrating factors

Case 3 If we assume that $\mu(x, y) = P(x)Q(y)$, then

$$\begin{aligned}\mu(M_y - N_x) &= \frac{\partial \mu}{\partial x} N - \frac{\partial \mu}{\partial y} M \\ \implies P(x)Q(y)(M_y - N_x) &= P'(x)Q(y)N - P(x)Q'(y)M\end{aligned}$$

This implies that

$$M_y - N_x = \frac{P'}{P}N - \frac{Q'}{Q}M.$$

If

$$M_y - N_x = p(x)N - q(y)M, \text{ where } \frac{P'}{P} = p(x), \frac{Q'}{Q} = q(y),$$

then

$$P(x) = e^{\int p(x)dx}, \text{ and } Q(y) = e^{\int q(y)dy}.$$

Finding the integrating factors

Case 3 If we assume that $\mu(x, y) = P(x)Q(y)$, then

$$\begin{aligned}\mu(M_y - N_x) &= \frac{\partial \mu}{\partial x} N - \frac{\partial \mu}{\partial y} M \\ \implies P(x)Q(y)(M_y - N_x) &= P'(x)Q(y)N - P(x)Q'(y)M\end{aligned}$$

This implies that

$$M_y - N_x = \frac{P'}{P}N - \frac{Q'}{Q}M.$$

If

$$M_y - N_x = p(x)N - q(y)M, \text{ where } \frac{P'}{P} = p(x), \frac{Q'}{Q} = q(y),$$

then

$$P(x) = e^{\int p(x)dx}, \text{ and } Q(y) = e^{\int q(y)dy}.$$

Thus

$$\mu(x, y) = P(x)Q(y) = e^{\int p(x)dx} e^{\int q(y)dy}$$

Integrating factors

Theorem Let M , N , M_y , and N_x be continuous on an open rectangle \mathcal{R} . Consider the ODE

$$M(x, y) + N(x, y) \frac{dy}{dx} = 0. \quad (9)$$

Integrating factors

Theorem Let M , N , M_y , and N_x be continuous on an open rectangle \mathcal{R} . Consider the ODE

$$M(x, y) + N(x, y) \frac{dy}{dx} = 0. \quad (9)$$

(a) If $\frac{M_y - N_x}{N}$ is a function of x only, say $p(x)$, then

$$\mu(x) = e^{\int p(x) dx}$$

is an integrating factor of ODE (9).

Integrating factors

Theorem Let M , N , M_y , and N_x be continuous on an open rectangle \mathcal{R} . Consider the ODE

$$M(x, y) + N(x, y) \frac{dy}{dx} = 0. \quad (9)$$

(a) If $\frac{M_y - N_x}{N}$ is a function of x only, say $p(x)$, then

$$\mu(x) = e^{\int p(x) dx}$$

is an integrating factor of ODE (9).

(b) If $\frac{N_x - M_y}{M}$ is a function of y only, say $q(y)$, then

$$\mu(y) = e^{\int q(y) dy}$$

is an integrating factor of ODE (9).

Integrating factors

Theorem Let M , N , M_y , and N_x be continuous on an open rectangle \mathcal{R} . Consider the ODE

$$M(x, y) + N(x, y) \frac{dy}{dx} = 0. \quad (9)$$

(a) If $\frac{M_y - N_x}{N}$ is a function of x only, say $p(x)$, then

$$\mu(x) = e^{\int p(x) dx}$$

is an integrating factor of ODE (9).

(b) If $\frac{N_x - M_y}{M}$ is a function of y only, say $q(y)$, then

$$\mu(y) = e^{\int q(y) dy}$$

is an integrating factor of ODE (9).

(c) If $M_y - N_x = p(x)N - q(y)M$, where $\frac{P'}{P} = p(x)$, $\frac{Q'}{Q} = q(y)$, then

$$\mu(x, y) = P(x)Q(y) = e^{\int p(x) dx} e^{\int q(y) dy}$$

is an integrating factor of ODE (9).

Integrating factors

Question Is it true that Integrating factor is exactly one of the following?

Integrating factors

Question Is it true that Integrating factor is exactly one of the following?

- ▶ functions of x only, that is, $\mu(x)$;

Integrating factors

Question Is it true that Integrating factor is exactly one of the following?

- ▶ functions of x only, that is, $\mu(x)$;
- ▶ functions of y only, that is, $\mu(y)$,

Integrating factors

Question Is it true that Integrating factor is exactly one of the following?

- ▶ functions of x only, that is, $\mu(x)$;
- ▶ functions of y only, that is, $\mu(y)$,
- ▶ functions of x and y , but $\mu(x, y) = P(x)Q(y)$, that is a pure product of functions of x and functions of y .

Integrating factors

Question Is it true that Integrating factor is exactly one of the following?

- ▶ functions of x only, that is, $\mu(x)$;
- ▶ functions of y only, that is, $\mu(y)$,
- ▶ functions of x and y , but $\mu(x, y) = P(x)Q(y)$, that is a pure product of functions of x and functions of y .

Question Is an Integrating factor unique (up to a constant) for a given ODE?

Example

Example Solve

$$\cos x \cos y \, dx + (\sin x \cos y - \sin x \sin y + y) \, dy = 0.$$

Example

Example Solve

$$\cos x \cos y \, dx + (\sin x \cos y - \sin x \sin y + y) \, dy = 0.$$

Here $M = \cos x \cos y$ and $N = \sin x \cos y - \sin x \sin y + y$

Example

Example Solve

$$\cos x \cos y \, dx + (\sin x \cos y - \sin x \sin y + y) \, dy = 0.$$

Here $M = \cos x \cos y$ and $N = \sin x \cos y - \sin x \sin y + y$

$$M_y - N_x = -\cos x \sin y - \cos x \cos y + \cos x \sin y$$

Example

Example Solve

$$\cos x \cos y \, dx + (\sin x \cos y - \sin x \sin y + y) \, dy = 0.$$

Here $M = \cos x \cos y$ and $N = \sin x \cos y - \sin x \sin y + y$

$$\begin{aligned} M_y - N_x &= -\cos x \sin y - \cos x \cos y + \cos x \sin y \\ \Rightarrow \frac{N_x - M_y}{M} &= 1 \end{aligned}$$

This implies that the integrating factor is $\mu = e^y$.

Example

Example Solve

$$\cos x \cos y \, dx + (\sin x \cos y - \sin x \sin y + y) \, dy = 0.$$

Here $M = \cos x \cos y$ and $N = \sin x \cos y - \sin x \sin y + y$

$$\begin{aligned} M_y - N_x &= -\cos x \sin y - \cos x \cos y + \cos x \sin y \\ \Rightarrow \frac{N_x - M_y}{M} &= 1 \end{aligned}$$

This implies that the integrating factor is $\mu = e^y$.

Thus

$$e^y \cos x \cos y \, dx + e^y (\sin x \cos y - \sin x \sin y + y) \, dy = 0$$

is exact.

Example

Example Solve

$$\cos x \cos y \, dx + (\sin x \cos y - \sin x \sin y + y) \, dy = 0.$$

Here $M = \cos x \cos y$ and $N = \sin x \cos y - \sin x \sin y + y$

$$\begin{aligned} M_y - N_x &= -\cos x \sin y - \cos x \cos y + \cos x \sin y \\ \Rightarrow \frac{N_x - M_y}{M} &= 1 \end{aligned}$$

This implies that the integrating factor is $\mu = e^y$.

Thus

$$e^y \cos x \cos y \, dx + e^y (\sin x \cos y - \sin x \sin y + y) \, dy = 0$$

is exact. So there exists $G(x, y)$ such that

$$\frac{\partial G}{\partial x} = e^y \cos x \cos y, \text{ and } \frac{\partial G}{\partial y} = e^y (\sin x \cos y - \sin x \sin y + y)$$

Example (continued ...)

Since $\frac{\partial G}{\partial x} = e^y \cos x \cos y$, integrating w.r.t. x , we get

$$G(x, y) = e^y \sin x \cos y + h(y).$$

Example (continued ...)

Since $\frac{\partial G}{\partial x} = e^y \cos x \cos y$, integrating w.r.t. x , we get

$$\begin{aligned} G(x, y) &= e^y \sin x \cos y + h(y). \\ \implies \frac{\partial G}{\partial y} &= e^y \sin x \cos y - e^y \sin x \sin y + \frac{dh}{dy} \end{aligned}$$

Example (continued ...)

Since $\frac{\partial G}{\partial x} = e^y \cos x \cos y$, integrating w.r.t. x , we get

$$\begin{aligned} G(x, y) &= e^y \sin x \cos y + h(y). \\ \implies \frac{\partial G}{\partial y} &= e^y \sin x \cos y - e^y \sin x \sin y + \frac{dh}{dy} \end{aligned}$$

Example (continued ...)

Since $\frac{\partial G}{\partial x} = e^y \cos x \cos y$, integrating w.r.t. x , we get

$$\begin{aligned} G(x, y) &= e^y \sin x \cos y + h(y). \\ \implies \frac{\partial G}{\partial y} &= e^y \sin x \cos y - e^y \sin x \sin y + \frac{dh}{dy} \end{aligned}$$

But

$$\frac{\partial G}{\partial y} = e^y (\sin x \cos y - \sin x \sin y + y)$$

Example (continued ...)

Since $\frac{\partial G}{\partial x} = e^y \cos x \cos y$, integrating w.r.t. x , we get

$$\begin{aligned} G(x, y) &= e^y \sin x \cos y + h(y). \\ \implies \frac{\partial G}{\partial y} &= e^y \sin x \cos y - e^y \sin x \sin y + \frac{dh}{dy} \end{aligned}$$

But

$$\frac{\partial G}{\partial y} = e^y (\sin x \cos y - \sin x \sin y + y)$$

Therefore $\frac{dh}{dy} = ye^y$,

Example (continued ...)

Since $\frac{\partial G}{\partial x} = e^y \cos x \cos y$, integrating w.r.t. x , we get

$$\begin{aligned} G(x, y) &= e^y \sin x \cos y + h(y). \\ \implies \frac{\partial G}{\partial y} &= e^y \sin x \cos y - e^y \sin x \sin y + \frac{dh}{dy} \end{aligned}$$

But

$$\frac{\partial G}{\partial y} = e^y (\sin x \cos y - \sin x \sin y + y)$$

Therefore $\frac{dh}{dy} = ye^y$, and hence $h(y) = ye^y - e^y + c$.

Example (continued ...)

Since $\frac{\partial G}{\partial x} = e^y \cos x \cos y$, integrating w.r.t. x , we get

$$\begin{aligned} G(x, y) &= e^y \sin x \cos y + h(y). \\ \implies \frac{\partial G}{\partial y} &= e^y \sin x \cos y - e^y \sin x \sin y + \frac{dh}{dy} \end{aligned}$$

But

$$\frac{\partial G}{\partial y} = e^y (\sin x \cos y - \sin x \sin y + y)$$

Therefore $\frac{dh}{dy} = ye^y$, and hence $h(y) = ye^y - e^y + c$.

Thus

$$G(x, y) = e^y (\sin x \cos y + y - 1) + c$$

is an implicit solution of ODE.

Example

Example Solve

$$(3x^2y^3 - y^2 + y)dx + (-xy + 2x)dy = 0$$

Example

Example Solve

$$(3x^2y^3 - y^2 + y)dx + (-xy + 2x)dy = 0$$

Here $M(x, y) = 3x^2y^3 - y^2 + y$ and $N(x, y) = -xy + 2x$. Note that

$$M_y - N_x = 9x^2y^2 - 2y + 1 - 2 = 9x^2y^2 - y - 1.$$

Example

Example Solve

$$(3x^2y^3 - y^2 + y)dx + (-xy + 2x)dy = 0$$

Here $M(x, y) = 3x^2y^3 - y^2 + y$ and $N(x, y) = -xy + 2x$. Note that

$$M_y - N_x = 9x^2y^2 - 2y + 1 - 2 = 9x^2y^2 - y - 1.$$

Observe that

$$\frac{-M_y + N_x}{M} \neq q(y), \text{ and } \frac{M_y - N_x}{N} \neq p(x).$$

Example

Example Solve

$$(3x^2y^3 - y^2 + y)dx + (-xy + 2x)dy = 0$$

Here $M(x, y) = 3x^2y^3 - y^2 + y$ and $N(x, y) = -xy + 2x$. Note that

$$M_y - N_x = 9x^2y^2 - 2y + 1 - 2 = 9x^2y^2 - y - 1.$$

Observe that

$$\frac{-M_y + N_x}{M} \neq q(y), \text{ and } \frac{M_y - N_x}{N} \neq p(x).$$

Can we write

$$M_y - N_x = p(x)N - q(y)M$$

for some $p(x)$ and $q(y)$?

Example (continued ...)

Choose $p(x) = \frac{-2}{x}$ and $q(y) = \frac{-3}{y}$. Then

Example (continued ...)

Choose $p(x) = \frac{-2}{x}$ and $q(y) = \frac{-3}{y}$. Then

$$M_y - N_x = p(x)N - q(y).$$

The integrating factor is then given by

$$\mu(x, y) = e^{\int \frac{-2}{x} dx} e^{\int \frac{-3}{y} dy} = \frac{1}{x^2 y^3}.$$

Example (continued ...)

Choose $p(x) = \frac{-2}{x}$ and $q(y) = \frac{-3}{y}$. Then

$$M_y - N_x = p(x)N - q(y).$$

The integrating factor is then given by

$$\mu(x, y) = e^{\int \frac{-2}{x} dx} e^{\int \frac{-3}{y} dy} = \frac{1}{x^2 y^3}.$$

Example (continued ...)

Choose $p(x) = \frac{-2}{x}$ and $q(y) = \frac{-3}{y}$. Then

$$M_y - N_x = p(x)N - q(y).$$

The integrating factor is then given by

$$\mu(x, y) = e^{\int \frac{-2}{x} dx} e^{\int \frac{-3}{y} dy} = \frac{1}{x^2 y^3}.$$

We get an exact ODE

$$\frac{1}{x^2 y^3} [(3x^2 y^3 - y^2 + y)dx + (-xy + 2x)dy] = 0.$$