

# Lectures 3 - 4 - 5

## Vector space, basis and dimension

Dipankar Ghosh  
(IIT Hyderabad)

January, 2020

- Linear Algebra is the study of 'vector spaces' and the 'maps' between them.

- Linear Algebra is the study of 'vector spaces' and the 'maps' between them.
- For now, we keep  $\mathbb{R}^n$  as an example of a vector space.


- Linear Algebra is the study of 'vector spaces' and the 'maps' between them.
- For now, we keep  $\mathbb{R}^n$  as an example of a vector space.
- Essentially, a vector space means a collection of objects,

- Linear Algebra is the study of 'vector spaces' and the 'maps' between them.
- For now, we keep  $\mathbb{R}^n$  as an example of a vector space.
- Essentially, a vector space means a collection of objects, we call them vectors,

- Linear Algebra is the study of 'vector spaces' and the 'maps' between them.
- For now, we keep  $\mathbb{R}^n$  as an example of a vector space.
- Essentially, a vector space means a collection of objects, we call them vectors, where we can add two vectors, and what we get is a vector;


- Linear Algebra is the study of ‘vector spaces’ and the ‘maps’ between them.
- For now, we keep  $\mathbb{R}^n$  as an example of a vector space.
- Essentially, a vector space means a collection of objects, we call them vectors, where we can add two vectors, and what we get is a vector; we can multiply a vector by a scalar, and what we get is a vector.

# Vector addition in $\mathbb{R}^2$







# Vector addition in $\mathbb{R}^2$




# Vector addition in $\mathbb{R}^2$




# Vector addition in $\mathbb{R}^2$




# Vector addition in $\mathbb{R}^2$




# Vector addition in $\mathbb{R}^2$




# Vector addition in $\mathbb{R}^2$



# Vector addition in $\mathbb{R}^2$




# Scalar multiplication in $\mathbb{R}^2$







# Scalar multiplication in $\mathbb{R}^2$




# Scalar multiplication in $\mathbb{R}^2$



# Scalar multiplication in $\mathbb{R}^2$



# Scalar multiplication in $\mathbb{R}^2$



# Definition of Vector Space

A set  $V$  of objects (called vectors)

# Definition of Vector Space

A set  $V$  of objects (called vectors) along with vector addition '+' and scalar multiplication '·'

# Definition of Vector Space

A set  $V$  of objects (called vectors) along with vector addition '+' and scalar multiplication ' $\cdot$ ' is said to be a vector space over a field  $\mathbb{F}$  (say,  $\mathbb{F} = \mathbb{R}$ , the set of real numbers) if the following hold:

# Definition of Vector Space

A set  $V$  of objects (called vectors) along with vector addition '+' and scalar multiplication ' $\cdot$ ' is said to be a vector space over a field  $\mathbb{F}$  (say,  $\mathbb{F} = \mathbb{R}$ , the set of real numbers) if the following hold:

- 1  $V$  is closed under '+', i.e.  $x + y \in V$  for all  $x, y \in V$ .



# Definition of Vector Space

A set  $V$  of objects (called vectors) along with vector addition '+' and scalar multiplication ' $\cdot$ ' is said to be a vector space over a field  $\mathbb{F}$  (say,  $\mathbb{F} = \mathbb{R}$ , the set of real numbers) if the following hold:

- 1  $V$  is closed under '+', i.e.  $x + y \in V$  for all  $x, y \in V$ .
- 2 Addition is commutative, i.e.  $x + y = y + x$  for all  $x, y \in V$ .

# Definition of Vector Space

A set  $V$  of objects (called vectors) along with vector addition '+' and scalar multiplication ' $\cdot$ ' is said to be a vector space over a field  $\mathbb{F}$  (say,  $\mathbb{F} = \mathbb{R}$ , the set of real numbers) if the following hold:

- 1  $V$  is closed under '+', i.e.  $x + y \in V$  for all  $x, y \in V$ .
- 2 Addition is commutative, i.e.  $x + y = y + x$  for all  $x, y \in V$ .
- 3 Addition is associative, i.e.  $(x + y) + z = x + (y + z)$  for all  $x, y, z \in V$ .

# Definition of Vector Space

A set  $V$  of objects (called vectors) along with vector addition '+' and scalar multiplication ' $\cdot$ ' is said to be a vector space over a field  $\mathbb{F}$  (say,  $\mathbb{F} = \mathbb{R}$ , the set of real numbers) if the following hold:

- 1  $V$  is closed under '+', i.e.  $x + y \in V$  for all  $x, y \in V$ .
- 2 Addition is commutative, i.e.  $x + y = y + x$  for all  $x, y \in V$ .
- 3 Addition is associative, i.e.  $(x + y) + z = x + (y + z)$  for all  $x, y, z \in V$ .
- 4 Additive identity, i.e. there is  $0 \in V$  such that  $x + 0 = x$  for all  $x \in V$ .

# Definition of Vector Space

A set  $V$  of objects (called vectors) along with vector addition '+' and scalar multiplication ' $\cdot$ ' is said to be a vector space over a field  $\mathbb{F}$  (say,  $\mathbb{F} = \mathbb{R}$ , the set of real numbers) if the following hold:

- 1  $V$  is closed under '+', i.e.  $x + y \in V$  for all  $x, y \in V$ .
- 2 Addition is commutative, i.e.  $x + y = y + x$  for all  $x, y \in V$ .
- 3 Addition is associative, i.e.  $(x + y) + z = x + (y + z)$  for all  $x, y, z \in V$ .
- 4 Additive identity, i.e. there is  $0 \in V$  such that  $x + 0 = x$  for all  $x \in V$ .
- 5 Additive inverse, i.e. for every  $x \in V$ , there is  $-x \in V$  such that  $x + (-x) = 0$ .

# Definition of Vector Space

A set  $V$  of objects (called vectors) along with vector addition '+' and scalar multiplication ' $\cdot$ ' is said to be a vector space over a field  $\mathbb{F}$  (say,  $\mathbb{F} = \mathbb{R}$ , the set of real numbers) if the following hold:

- 1  $V$  is closed under '+', i.e.  $x + y \in V$  for all  $x, y \in V$ .
- 2 Addition is commutative, i.e.  $x + y = y + x$  for all  $x, y \in V$ .
- 3 Addition is associative, i.e.  $(x + y) + z = x + (y + z)$  for all  $x, y, z \in V$ .
- 4 Additive identity, i.e. there is  $0 \in V$  such that  $x + 0 = x$  for all  $x \in V$ .
- 5 Additive inverse, i.e. for every  $x \in V$ , there is  $-x \in V$  such that  $x + (-x) = 0$ .
- 6  $V$  is closed under ' $\cdot$ ', i.e.  $c \cdot x \in V$  for all  $c \in \mathbb{F}$  and  $x \in V$ .

# Definition of Vector Space

A set  $V$  of objects (called vectors) along with vector addition '+' and scalar multiplication ' $\cdot$ ' is said to be a vector space over a field  $\mathbb{F}$  (say,  $\mathbb{F} = \mathbb{R}$ , the set of real numbers) if the following hold:

- 1  $V$  is closed under '+', i.e.  $x + y \in V$  for all  $x, y \in V$ .
- 2 Addition is commutative, i.e.  $x + y = y + x$  for all  $x, y \in V$ .
- 3 Addition is associative, i.e.  $(x + y) + z = x + (y + z)$  for all  $x, y, z \in V$ .
- 4 Additive identity, i.e. there is  $0 \in V$  such that  $x + 0 = x$  for all  $x \in V$ .
- 5 Additive inverse, i.e. for every  $x \in V$ , there is  $-x \in V$  such that  $x + (-x) = 0$ .
- 6  $V$  is closed under ' $\cdot$ ', i.e.  $c \cdot x \in V$  for all  $c \in \mathbb{F}$  and  $x \in V$ .
- 7  $1 \cdot x = x$  for all  $x \in V$ .

# Definition of Vector Space

A set  $V$  of objects (called vectors) along with vector addition '+' and scalar multiplication ' $\cdot$ ' is said to be a vector space over a field  $\mathbb{F}$  (say,  $\mathbb{F} = \mathbb{R}$ , the set of real numbers) if the following hold:

- 1  $V$  is closed under '+', i.e.  $x + y \in V$  for all  $x, y \in V$ .
- 2 Addition is commutative, i.e.  $x + y = y + x$  for all  $x, y \in V$ .
- 3 Addition is associative, i.e.  $(x + y) + z = x + (y + z)$  for all  $x, y, z \in V$ .
- 4 Additive identity, i.e. there is  $0 \in V$  such that  $x + 0 = x$  for all  $x \in V$ .
- 5 Additive inverse, i.e. for every  $x \in V$ , there is  $-x \in V$  such that  $x + (-x) = 0$ .
- 6  $V$  is closed under ' $\cdot$ ', i.e.  $c \cdot x \in V$  for all  $c \in \mathbb{F}$  and  $x \in V$ .
- 7  $1 \cdot x = x$  for all  $x \in V$ .
- 8  $a \cdot (b \cdot x) = (ab) \cdot x$  for all  $a, b \in \mathbb{R}$  and  $x \in V$ .

# Definition of Vector Space

A set  $V$  of objects (called vectors) along with vector addition '+' and scalar multiplication ' $\cdot$ ' is said to be a vector space over a field  $\mathbb{F}$  (say,  $\mathbb{F} = \mathbb{R}$ , the set of real numbers) if the following hold:

- 1  $V$  is closed under '+', i.e.  $x + y \in V$  for all  $x, y \in V$ .
- 2 Addition is commutative, i.e.  $x + y = y + x$  for all  $x, y \in V$ .
- 3 Addition is associative, i.e.  $(x + y) + z = x + (y + z)$  for all  $x, y, z \in V$ .
- 4 Additive identity, i.e. there is  $0 \in V$  such that  $x + 0 = x$  for all  $x \in V$ .
- 5 Additive inverse, i.e. for every  $x \in V$ , there is  $-x \in V$  such that  $x + (-x) = 0$ .
- 6  $V$  is closed under ' $\cdot$ ', i.e.  $c \cdot x \in V$  for all  $c \in \mathbb{F}$  and  $x \in V$ .
- 7  $1 \cdot x = x$  for all  $x \in V$ .
- 8  $a \cdot (b \cdot x) = (ab) \cdot x$  for all  $a, b \in \mathbb{R}$  and  $x \in V$ .
- 9  $a \cdot (x + y) = a \cdot x + a \cdot y$  for all  $a \in \mathbb{R}$  and  $x, y \in V$ .



# Definition of Vector Space

A set  $V$  of objects (called vectors) along with vector addition '+' and scalar multiplication ' $\cdot$ ' is said to be a vector space over a field  $\mathbb{F}$  (say,  $\mathbb{F} = \mathbb{R}$ , the set of real numbers) if the following hold:

- 1  $V$  is closed under '+', i.e.  $x + y \in V$  for all  $x, y \in V$ .
- 2 Addition is commutative, i.e.  $x + y = y + x$  for all  $x, y \in V$ .
- 3 Addition is associative, i.e.  $(x + y) + z = x + (y + z)$  for all  $x, y, z \in V$ .
- 4 Additive identity, i.e. there is  $0 \in V$  such that  $x + 0 = x$  for all  $x \in V$ .
- 5 Additive inverse, i.e. for every  $x \in V$ , there is  $-x \in V$  such that  $x + (-x) = 0$ .
- 6  $V$  is closed under ' $\cdot$ ', i.e.  $c \cdot x \in V$  for all  $c \in \mathbb{F}$  and  $x \in V$ .
- 7  $1 \cdot x = x$  for all  $x \in V$ .
- 8  $a \cdot (b \cdot x) = (ab) \cdot x$  for all  $a, b \in \mathbb{R}$  and  $x \in V$ .
- 9  $a \cdot (x + y) = a \cdot x + a \cdot y$  for all  $a \in \mathbb{R}$  and  $x, y \in V$ .
- 10  $(a + b) \cdot x = a \cdot x + b \cdot x$  for all  $a, b \in \mathbb{R}$  and  $x \in V$ .

- 1 The elements of  $\mathbb{F}$  are called **scalars**,

- 1 The elements of  $\mathbb{F}$  are called **scalars**, and the elements of  $V$  are called **vectors**.

# Remarks on vector spaces

- 1 The elements of  $\mathbb{F}$  are called **scalars**, and the elements of  $V$  are called **vectors**.
- 2 The first five properties are nothing but the properties of abelian group,

# Remarks on vector spaces

- 1 The elements of  $\mathbb{F}$  are called **scalars**, and the elements of  $V$  are called **vectors**.
- 2 The first five properties are nothing but the properties of abelian group, i.e.  $(V, +)$  is an abelian group.

- 1 The elements of  $\mathbb{F}$  are called **scalars**, and the elements of  $V$  are called **vectors**.
- 2 The first five properties are nothing but the properties of abelian group, i.e.  $(V, +)$  is an abelian group.
- 3 We simply write  $cx$  instead of  $c \cdot x$  for  $c \in \mathbb{F}$  and  $x \in V$  when there is no confusion.

- 1 The elements of  $\mathbb{F}$  are called **scalars**, and the elements of  $V$  are called **vectors**.
- 2 The first five properties are nothing but the properties of abelian group, i.e.  $(V, +)$  is an abelian group.
- 3 We simply write  $cx$  instead of  $c \cdot x$  for  $c \in \mathbb{F}$  and  $x \in V$  when there is no confusion.
- 4 From now, we work over the field  $\mathbb{R}$ .

# Examples of vector spaces

(1) The  $n$ -tuple space,  $V = \mathbb{R}^n$ .



# Examples of vector spaces

(1) The  $n$ -tuple space,  $V = \mathbb{R}^n$ .

$$\begin{pmatrix} x_1 \\ x_2 \\ \vdots \\ x_n \end{pmatrix} + \begin{pmatrix} y_1 \\ y_2 \\ \vdots \\ y_n \end{pmatrix} = \begin{pmatrix} x_1 + y_1 \\ x_2 + y_2 \\ \vdots \\ x_n + y_n \end{pmatrix} \quad \text{and} \quad c \cdot \begin{pmatrix} x_1 \\ x_2 \\ \vdots \\ x_n \end{pmatrix} = \begin{pmatrix} cx_1 \\ cx_2 \\ \vdots \\ cx_n \end{pmatrix}$$

# Examples of vector spaces

(1) The  $n$ -tuple space,  $V = \mathbb{R}^n$ .

$$\begin{pmatrix} x_1 \\ x_2 \\ \vdots \\ x_n \end{pmatrix} + \begin{pmatrix} y_1 \\ y_2 \\ \vdots \\ y_n \end{pmatrix} = \begin{pmatrix} x_1 + y_1 \\ x_2 + y_2 \\ \vdots \\ x_n + y_n \end{pmatrix} \quad \text{and} \quad c \cdot \begin{pmatrix} x_1 \\ x_2 \\ \vdots \\ x_n \end{pmatrix} = \begin{pmatrix} cx_1 \\ cx_2 \\ \vdots \\ cx_n \end{pmatrix}$$

(2) The space  $\mathbb{R}^{m \times n}$  of all  $m \times n$  matrices

$$\begin{pmatrix} x_{11} & x_{12} & \cdots & x_{1n} \\ x_{21} & x_{22} & \cdots & x_{2n} \\ \vdots & \vdots & \ddots & \vdots \\ x_{m1} & x_{m2} & \cdots & x_{mn} \end{pmatrix} \quad \text{where } x_{ij} \in \mathbb{R}.$$

# Examples of vector spaces

(1) The  $n$ -tuple space,  $V = \mathbb{R}^n$ .

$$\begin{pmatrix} x_1 \\ x_2 \\ \vdots \\ x_n \end{pmatrix} + \begin{pmatrix} y_1 \\ y_2 \\ \vdots \\ y_n \end{pmatrix} = \begin{pmatrix} x_1 + y_1 \\ x_2 + y_2 \\ \vdots \\ x_n + y_n \end{pmatrix} \quad \text{and} \quad c \cdot \begin{pmatrix} x_1 \\ x_2 \\ \vdots \\ x_n \end{pmatrix} = \begin{pmatrix} cx_1 \\ cx_2 \\ \vdots \\ cx_n \end{pmatrix}$$

(2) The space  $\mathbb{R}^{m \times n}$  of all  $m \times n$  matrices

$$\begin{pmatrix} x_{11} & x_{12} & \cdots & x_{1n} \\ x_{21} & x_{22} & \cdots & x_{2n} \\ \vdots & \vdots & \ddots & \vdots \\ x_{m1} & x_{m2} & \cdots & x_{mn} \end{pmatrix} \quad \text{where } x_{ij} \in \mathbb{R}.$$

Component wise addition and component wise scalar multiplication.

# Examples

(3) Let  $S$  be any non-empty set.

# Examples

- (3) Let  $S$  be any non-empty set. Let  $V$  be the set of all functions from  $S$  into  $\mathbb{R}$ .

- (3) Let  $S$  be any non-empty set. Let  $V$  be the set of all functions from  $S$  into  $\mathbb{R}$ . The sum  $f + g$  of two vectors  $f$  and  $g$  in  $V$  is defined to be

$$(f + g)(s) := f(s) + g(s) \text{ for all } s \in S.$$

- (3) Let  $S$  be any non-empty set. Let  $V$  be the set of all functions from  $S$  into  $\mathbb{R}$ . The sum  $f + g$  of two vectors  $f$  and  $g$  in  $V$  is defined to be

$$(f + g)(s) := f(s) + g(s) \text{ for all } s \in S.$$

The scalar multiplication  $c \cdot f$  (for  $c \in \mathbb{R}$ ) is defined by

$$(c \cdot f)(s) := c f(s) \text{ for all } s \in S.$$

- (3) Let  $S$  be any non-empty set. Let  $V$  be the set of all functions from  $S$  into  $\mathbb{R}$ . The sum  $f + g$  of two vectors  $f$  and  $g$  in  $V$  is defined to be

$$(f + g)(s) := f(s) + g(s) \text{ for all } s \in S.$$

The scalar multiplication  $c \cdot f$  (for  $c \in \mathbb{R}$ ) is defined by

$$(c \cdot f)(s) := c f(s) \text{ for all } s \in S.$$

Is  $V$  a vector space?



- (3) Let  $S$  be any non-empty set. Let  $V$  be the set of all functions from  $S$  into  $\mathbb{R}$ . The sum  $f + g$  of two vectors  $f$  and  $g$  in  $V$  is defined to be

$$(f + g)(s) := f(s) + g(s) \text{ for all } s \in S.$$

The scalar multiplication  $c \cdot f$  (for  $c \in \mathbb{R}$ ) is defined by

$$(c \cdot f)(s) := c f(s) \text{ for all } s \in S.$$

Is  $V$  a vector space? Answer: Yes.

# Examples

- (3) Let  $S$  be any non-empty set. Let  $V$  be the set of all functions from  $S$  into  $\mathbb{R}$ . The sum  $f + g$  of two vectors  $f$  and  $g$  in  $V$  is defined to be

$$(f + g)(s) := f(s) + g(s) \text{ for all } s \in S.$$

The scalar multiplication  $c \cdot f$  (for  $c \in \mathbb{R}$ ) is defined by

$$(c \cdot f)(s) := c f(s) \text{ for all } s \in S.$$

Is  $V$  a vector space? Answer: Yes.

- (4) The set  $\mathbb{R}[x]$  of all polynomials

# Examples

- (3) Let  $S$  be any non-empty set. Let  $V$  be the set of all functions from  $S$  into  $\mathbb{R}$ . The sum  $f + g$  of two vectors  $f$  and  $g$  in  $V$  is defined to be

$$(f + g)(s) := f(s) + g(s) \text{ for all } s \in S.$$

The scalar multiplication  $c \cdot f$  (for  $c \in \mathbb{R}$ ) is defined by

$$(c \cdot f)(s) := c f(s) \text{ for all } s \in S.$$

Is  $V$  a vector space? Answer: Yes.

- (4) The set  $\mathbb{R}[x]$  of all polynomials  $a_0 + a_1x + \cdots + a_mx^m$ ,

# Examples

- (3) Let  $S$  be any non-empty set. Let  $V$  be the set of all functions from  $S$  into  $\mathbb{R}$ . The sum  $f + g$  of two vectors  $f$  and  $g$  in  $V$  is defined to be

$$(f + g)(s) := f(s) + g(s) \text{ for all } s \in S.$$

The scalar multiplication  $c \cdot f$  (for  $c \in \mathbb{R}$ ) is defined by

$$(c \cdot f)(s) := c f(s) \text{ for all } s \in S.$$

Is  $V$  a vector space? Answer: Yes.

- (4) The set  $\mathbb{R}[x]$  of all polynomials  $a_0 + a_1x + \cdots + a_mx^m$ , where  $a_i \in \mathbb{R}$ ,

# Examples

- (3) Let  $S$  be any non-empty set. Let  $V$  be the set of all functions from  $S$  into  $\mathbb{R}$ . The sum  $f + g$  of two vectors  $f$  and  $g$  in  $V$  is defined to be

$$(f + g)(s) := f(s) + g(s) \text{ for all } s \in S.$$

The scalar multiplication  $c \cdot f$  (for  $c \in \mathbb{R}$ ) is defined by

$$(c \cdot f)(s) := c f(s) \text{ for all } s \in S.$$

Is  $V$  a vector space? Answer: Yes.

- (4) The set  $\mathbb{R}[x]$  of all polynomials  $a_0 + a_1x + \cdots + a_mx^m$ , where  $a_i \in \mathbb{R}$ ,  $x$  is an indeterminate

# Examples

- (3) Let  $S$  be any non-empty set. Let  $V$  be the set of all functions from  $S$  into  $\mathbb{R}$ . The sum  $f + g$  of two vectors  $f$  and  $g$  in  $V$  is defined to be

$$(f + g)(s) := f(s) + g(s) \text{ for all } s \in S.$$

The scalar multiplication  $c \cdot f$  (for  $c \in \mathbb{R}$ ) is defined by

$$(c \cdot f)(s) := c f(s) \text{ for all } s \in S.$$

Is  $V$  a vector space? Answer: Yes.

- (4) The set  $\mathbb{R}[x]$  of all polynomials  $a_0 + a_1x + \cdots + a_mx^m$ , where  $a_i \in \mathbb{R}$ ,  $x$  is an indeterminate and  $m$  varies over non-negative integers.

- (3) Let  $S$  be any non-empty set. Let  $V$  be the set of all functions from  $S$  into  $\mathbb{R}$ . The sum  $f + g$  of two vectors  $f$  and  $g$  in  $V$  is defined to be

$$(f + g)(s) := f(s) + g(s) \text{ for all } s \in S.$$

The scalar multiplication  $c \cdot f$  (for  $c \in \mathbb{R}$ ) is defined by

$$(c \cdot f)(s) := c f(s) \text{ for all } s \in S.$$

Is  $V$  a vector space? Answer: Yes.

- (4) The set  $\mathbb{R}[x]$  of all polynomials  $a_0 + a_1x + \cdots + a_mx^m$ , where  $a_i \in \mathbb{R}$ ,  $x$  is an indeterminate and  $m$  varies over non-negative integers. The vector addition and scalar multiplication are defined in obvious way.

# Examples

- (3) Let  $S$  be any non-empty set. Let  $V$  be the set of all functions from  $S$  into  $\mathbb{R}$ . The sum  $f + g$  of two vectors  $f$  and  $g$  in  $V$  is defined to be

$$(f + g)(s) := f(s) + g(s) \text{ for all } s \in S.$$

The scalar multiplication  $c \cdot f$  (for  $c \in \mathbb{R}$ ) is defined by

$$(c \cdot f)(s) := c f(s) \text{ for all } s \in S.$$

Is  $V$  a vector space? Answer: Yes.

- (4) The set  $\mathbb{R}[x]$  of all polynomials  $a_0 + a_1x + \cdots + a_mx^m$ , where  $a_i \in \mathbb{R}$ ,  $x$  is an indeterminate and  $m$  varies over non-negative integers. The vector addition and scalar multiplication are defined in obvious way. Then  $\mathbb{R}[x]$  is a vector space over  $\mathbb{R}$ .



(5) The set  $\mathbb{R}^{n \times n}$  of all  $n \times n$  matrices

(5) The set  $\mathbb{R}^{n \times n}$  of all  $n \times n$  matrices with **vector addition**:

$$\begin{pmatrix} x_{11} & x_{12} & \cdots & x_{1n} \\ x_{21} & x_{22} & \cdots & x_{2n} \\ \vdots & \vdots & \ddots & \vdots \\ x_{n1} & x_{n2} & \cdots & x_{nn} \end{pmatrix} \times \begin{pmatrix} y_{11} & y_{12} & \cdots & y_{1n} \\ y_{21} & y_{22} & \cdots & y_{2n} \\ \vdots & \vdots & \ddots & \vdots \\ y_{n1} & y_{n2} & \cdots & y_{nn} \end{pmatrix} \\ = \begin{pmatrix} \text{---} & \text{---} & \text{---} \\ \text{---} & (\sum_{k=1}^n x_{ik} y_{kj}) & \text{---} \\ \text{---} & \text{---} & \text{---} \end{pmatrix} \quad (\text{matrix multiplication})$$

(5) The set  $\mathbb{R}^{n \times n}$  of all  $n \times n$  matrices with **vector addition**:

$$\begin{pmatrix} x_{11} & x_{12} & \cdots & x_{1n} \\ x_{21} & x_{22} & \cdots & x_{2n} \\ \vdots & \vdots & \ddots & \vdots \\ x_{n1} & x_{n2} & \cdots & x_{nn} \end{pmatrix} \times \begin{pmatrix} y_{11} & y_{12} & \cdots & y_{1n} \\ y_{21} & y_{22} & \cdots & y_{2n} \\ \vdots & \vdots & \ddots & \vdots \\ y_{n1} & y_{n2} & \cdots & y_{nn} \end{pmatrix} \\ = \begin{pmatrix} \text{---} & \text{---} & \text{---} \\ \text{---} & (\sum_{k=1}^n x_{ik} y_{kj}) & \text{---} \\ \text{---} & \text{---} & \text{---} \end{pmatrix} \quad (\text{matrix multiplication})$$

and scalar multiplication as before.

(5) The set  $\mathbb{R}^{n \times n}$  of all  $n \times n$  matrices with **vector addition**:

$$\begin{pmatrix} x_{11} & x_{12} & \cdots & x_{1n} \\ x_{21} & x_{22} & \cdots & x_{2n} \\ \vdots & \vdots & \ddots & \vdots \\ x_{n1} & x_{n2} & \cdots & x_{nn} \end{pmatrix} \times \begin{pmatrix} y_{11} & y_{12} & \cdots & y_{1n} \\ y_{21} & y_{22} & \cdots & y_{2n} \\ \vdots & \vdots & \ddots & \vdots \\ y_{n1} & y_{n2} & \cdots & y_{nn} \end{pmatrix} \\ = \begin{pmatrix} \text{---} & \text{---} & \text{---} \\ \text{---} & (\sum_{k=1}^n x_{ik} y_{kj}) & \text{---} \\ \text{---} & \text{---} & \text{---} \end{pmatrix} \quad (\text{matrix multiplication})$$

and scalar multiplication as before.

Is  $V$  a vector space?

(5) The set  $\mathbb{R}^{n \times n}$  of all  $n \times n$  matrices with **vector addition**:

$$\begin{pmatrix} x_{11} & x_{12} & \cdots & x_{1n} \\ x_{21} & x_{22} & \cdots & x_{2n} \\ \vdots & \vdots & \ddots & \vdots \\ x_{n1} & x_{n2} & \cdots & x_{nn} \end{pmatrix} \times \begin{pmatrix} y_{11} & y_{12} & \cdots & y_{1n} \\ y_{21} & y_{22} & \cdots & y_{2n} \\ \vdots & \vdots & \ddots & \vdots \\ y_{n1} & y_{n2} & \cdots & y_{nn} \end{pmatrix} \\ = \begin{pmatrix} \text{---} & \text{---} & \text{---} \\ \text{---} & (\sum_{k=1}^n x_{ik} y_{kj}) & \text{---} \\ \text{---} & \text{---} & \text{---} \end{pmatrix} \quad (\text{matrix multiplication})$$

and scalar multiplication as before.

Is  $V$  a vector space? **Answer: No.**

# $\mathbb{R}^{n \times n}$ with matrix multiplication ' $\times$ ' is not a vector space

## Reasons:

- The operation ' $\times$ ' is not **commutative**,

## Reasons:

- The operation ' $\times$ ' is not **commutative**, because

$$\begin{pmatrix} 1 & 0 \\ 0 & 0 \end{pmatrix} \times \begin{pmatrix} 0 & 1 \\ 0 & 0 \end{pmatrix} = \begin{pmatrix} 0 & 1 \\ 0 & 0 \end{pmatrix} \neq \begin{pmatrix} 0 & 0 \\ 0 & 0 \end{pmatrix} = \begin{pmatrix} 0 & 0 \\ 0 & 0 \end{pmatrix} \times \begin{pmatrix} 0 & 1 \\ 0 & 0 \end{pmatrix}$$

## Reasons:

- The operation ' $\times$ ' is not **commutative**, because

$$\begin{pmatrix} 1 & 0 \\ 0 & 0 \end{pmatrix} \times \begin{pmatrix} 0 & 1 \\ 0 & 0 \end{pmatrix} = \begin{pmatrix} 0 & 1 \\ 0 & 0 \end{pmatrix} \neq \begin{pmatrix} 0 & 0 \\ 0 & 0 \end{pmatrix} = \begin{pmatrix} 0 & 0 \\ 0 & 0 \end{pmatrix} \times \begin{pmatrix} 0 & 1 \\ 0 & 0 \end{pmatrix}$$

- Another reason is that every matrix does not necessarily have multiplicative inverse.



## Reasons:

- The operation ' $\times$ ' is not **commutative**, because

$$\begin{pmatrix} 1 & 0 \\ 0 & 0 \end{pmatrix} \times \begin{pmatrix} 0 & 1 \\ 0 & 0 \end{pmatrix} = \begin{pmatrix} 0 & 1 \\ 0 & 0 \end{pmatrix} \neq \begin{pmatrix} 0 & 0 \\ 0 & 0 \end{pmatrix} = \begin{pmatrix} 0 & 1 \\ 0 & 0 \end{pmatrix} \times \begin{pmatrix} 1 & 0 \\ 0 & 0 \end{pmatrix}$$

- Another reason is that every matrix does not necessarily have multiplicative inverse. Note that  $\begin{pmatrix} 1 & 0 \\ 0 & 1 \end{pmatrix}$  is the identity element under the operation  $\times$ .

## Reasons:

- The operation ' $\times$ ' is not **commutative**, because

$$\begin{pmatrix} 1 & 0 \\ 0 & 0 \end{pmatrix} \times \begin{pmatrix} 0 & 1 \\ 0 & 0 \end{pmatrix} = \begin{pmatrix} 0 & 1 \\ 0 & 0 \end{pmatrix} \neq \begin{pmatrix} 0 & 0 \\ 0 & 0 \end{pmatrix} = \begin{pmatrix} 0 & 0 \\ 0 & 0 \end{pmatrix} \times \begin{pmatrix} 1 & 0 \\ 0 & 0 \end{pmatrix}$$

- Another reason is that every matrix does not necessarily have multiplicative inverse. Note that  $\begin{pmatrix} 1 & 0 \\ 0 & 1 \end{pmatrix}$  is the identity element under the operation  $\times$ . But  $\begin{pmatrix} 1 & 0 \\ 0 & 0 \end{pmatrix}$  does not have inverse under the operation  $\times$

## Reasons:

- The operation ' $\times$ ' is not **commutative**, because

$$\begin{pmatrix} 1 & 0 \\ 0 & 0 \end{pmatrix} \times \begin{pmatrix} 0 & 1 \\ 0 & 0 \end{pmatrix} = \begin{pmatrix} 0 & 1 \\ 0 & 0 \end{pmatrix} \neq \begin{pmatrix} 0 & 0 \\ 0 & 0 \end{pmatrix} = \begin{pmatrix} 0 & 1 \\ 0 & 0 \end{pmatrix} \times \begin{pmatrix} 1 & 0 \\ 0 & 0 \end{pmatrix}$$

- Another reason is that every matrix does not necessarily have multiplicative inverse. Note that  $\begin{pmatrix} 1 & 0 \\ 0 & 1 \end{pmatrix}$  is the identity element under the operation  $\times$ . But  $\begin{pmatrix} 1 & 0 \\ 0 & 0 \end{pmatrix}$  does not have inverse under the operation  $\times$  as there does not exist a matrix  $A$  such that

$$\begin{pmatrix} 1 & 0 \\ 0 & 0 \end{pmatrix} \times A = \begin{pmatrix} 1 & 0 \\ 0 & 1 \end{pmatrix}.$$

# Linear combination: Definition and examples

## Definition

A vector  $\beta$  in  $V$  is said to be a **linear combination** of vectors  $\alpha_1, \alpha_2$  and  $\alpha_r$  in  $V$

# Linear combination: Definition and examples

## Definition

A vector  $\beta$  in  $V$  is said to be a **linear combination** of vectors  $\alpha_1, \alpha_2$  and  $\alpha_r$  in  $V$  if


$$\beta = c_1\alpha_1 + c_2\alpha_2 + \cdots + c_r\alpha_r \text{ for some } c_1, c_2, \dots, c_r \in \mathbb{R}.$$

# Linear combination: Definition and examples

## Definition

A vector  $\beta$  in  $V$  is said to be a **linear combination** of vectors  $\alpha_1, \alpha_2$  and  $\alpha_r$  in  $V$  if

$$\beta = c_1\alpha_1 + c_2\alpha_2 + \cdots + c_r\alpha_r \text{ for some } c_1, c_2, \dots, c_r \in \mathbb{R}.$$




# Linear combination: Definition and examples

## Definition

A vector  $\beta$  in  $V$  is said to be a **linear combination** of vectors  $\alpha_1, \alpha_2$  and  $\alpha_r$  in  $V$  if

$$\beta = c_1\alpha_1 + c_2\alpha_2 + \cdots + c_r\alpha_r \text{ for some } c_1, c_2, \dots, c_r \in \mathbb{R}.$$




In this figure, the **linear combinations of  $\{u\}$**  are the vectors on the **dotted line**;

# Linear combination: Definition and examples

## Definition

A vector  $\beta$  in  $V$  is said to be a **linear combination** of vectors  $\alpha_1, \alpha_2$  and  $\alpha_r$  in  $V$  if

$$\beta = c_1\alpha_1 + c_2\alpha_2 + \cdots + c_r\alpha_r \text{ for some } c_1, c_2, \dots, c_r \in \mathbb{R}.$$



In this figure, the **linear combinations of  $\{u\}$**  are the vectors on the **dotted line**; while the **linear combinations of  $\{u, v\}$**  yield  $\mathbb{R}^2$ .



# More on linear combinations

- 1 Suppose  $\alpha_1, \alpha_2, \dots, \alpha_r$  and  $\beta$  are given vectors in  $V$ .

# More on linear combinations

- 1 Suppose  $\alpha_1, \alpha_2, \dots, \alpha_r$  and  $\beta$  are given vectors in  $V$ .
- 2 Is  $\beta$  a linear combination of  $\alpha_1, \alpha_2, \dots, \alpha_r$ ?

# More on linear combinations

- 1 Suppose  $\alpha_1, \alpha_2, \dots, \alpha_r$  and  $\beta$  are given vectors in  $V$ .
- 2 Is  $\beta$  a linear combination of  $\alpha_1, \alpha_2, \dots, \alpha_r$ ?
- 3 If yes, then how many ways are there to write that lin. comb.?

# More on linear combinations

- 1 Suppose  $\alpha_1, \alpha_2, \dots, \alpha_r$  and  $\beta$  are given vectors in  $V$ .
- 2 Is  $\beta$  a linear combination of  $\alpha_1, \alpha_2, \dots, \alpha_r$ ?
- 3 If yes, then how many ways are there to write that lin. comb.?
- 4 We find the answers to these questions as we proceed further.

# More on linear combinations

- 1 Suppose  $\alpha_1, \alpha_2, \dots, \alpha_r$  and  $\beta$  are given vectors in  $V$ .
- 2 Is  $\beta$  a linear combination of  $\alpha_1, \alpha_2, \dots, \alpha_r$ ?
- 3 If yes, then how many ways are there to write that lin. comb.?
- 4 We find the answers to these questions as we proceed further.
- 5 Let us see some examples here:

# More on linear combinations

- 1 Suppose  $\alpha_1, \alpha_2, \dots, \alpha_r$  and  $\beta$  are given vectors in  $V$ .
- 2 Is  $\beta$  a linear combination of  $\alpha_1, \alpha_2, \dots, \alpha_r$ ?
- 3 If yes, then how many ways are there to write that lin. comb.?
- 4 We find the answers to these questions as we proceed further.
- 5 Let us see some examples here:
- 6 In  $\mathbb{R}^2$ , the vector  $\begin{pmatrix} 1 \\ 2 \end{pmatrix}$

# More on linear combinations

- 1 Suppose  $\alpha_1, \alpha_2, \dots, \alpha_r$  and  $\beta$  are given vectors in  $V$ .
- 2 Is  $\beta$  a linear combination of  $\alpha_1, \alpha_2, \dots, \alpha_r$ ?
- 3 If yes, then how many ways are there to write that lin. comb.?
- 4 We find the answers to these questions as we proceed further.
- 5 Let us see some examples here:
- 6 In  $\mathbb{R}^2$ , the vector  $\begin{pmatrix} 1 \\ 2 \end{pmatrix} = 1 \cdot \begin{pmatrix} 1 \\ 0 \end{pmatrix} + 2 \cdot \begin{pmatrix} 0 \\ 1 \end{pmatrix}$

# More on linear combinations

- 1 Suppose  $\alpha_1, \alpha_2, \dots, \alpha_r$  and  $\beta$  are given vectors in  $V$ .
- 2 Is  $\beta$  a linear combination of  $\alpha_1, \alpha_2, \dots, \alpha_r$ ?
- 3 If yes, then how many ways are there to write that lin. comb.?
- 4 We find the answers to these questions as we proceed further.
- 5 Let us see some examples here:
- 6 In  $\mathbb{R}^2$ , the vector  $\begin{pmatrix} 1 \\ 2 \end{pmatrix} = 1 \cdot \begin{pmatrix} 1 \\ 0 \end{pmatrix} + 2 \cdot \begin{pmatrix} 0 \\ 1 \end{pmatrix}$  (Unique way).



# More on linear combinations

- 1 Suppose  $\alpha_1, \alpha_2, \dots, \alpha_r$  and  $\beta$  are given vectors in  $V$ .
- 2 Is  $\beta$  a linear combination of  $\alpha_1, \alpha_2, \dots, \alpha_r$ ?
- 3 If yes, then how many ways are there to write that lin. comb.?
- 4 We find the answers to these questions as we proceed further.
- 5 Let us see some examples here:
- 6 In  $\mathbb{R}^2$ , the vector  $\begin{pmatrix} 1 \\ 2 \end{pmatrix} = 1 \cdot \begin{pmatrix} 1 \\ 0 \end{pmatrix} + 2 \cdot \begin{pmatrix} 0 \\ 1 \end{pmatrix}$  (Unique way).
- 7  $\begin{pmatrix} 1 \\ 2 \end{pmatrix}$

# More on linear combinations

- 1 Suppose  $\alpha_1, \alpha_2, \dots, \alpha_r$  and  $\beta$  are given vectors in  $V$ .
- 2 Is  $\beta$  a linear combination of  $\alpha_1, \alpha_2, \dots, \alpha_r$ ?
- 3 If yes, then how many ways are there to write that lin. comb.?
- 4 We find the answers to these questions as we proceed further.
- 5 Let us see some examples here:
- 6 In  $\mathbb{R}^2$ , the vector  $\begin{pmatrix} 1 \\ 2 \end{pmatrix} = 1 \cdot \begin{pmatrix} 1 \\ 0 \end{pmatrix} + 2 \cdot \begin{pmatrix} 0 \\ 1 \end{pmatrix}$  (Unique way).
- 7  $\begin{pmatrix} 1 \\ 2 \end{pmatrix} = 0 \cdot \begin{pmatrix} 1 \\ 0 \end{pmatrix} + 1 \cdot \begin{pmatrix} 0 \\ 1 \end{pmatrix} + 1 \cdot \begin{pmatrix} 1 \\ 1 \end{pmatrix}$

# More on linear combinations

- 1 Suppose  $\alpha_1, \alpha_2, \dots, \alpha_r$  and  $\beta$  are given vectors in  $V$ .
- 2 Is  $\beta$  a linear combination of  $\alpha_1, \alpha_2, \dots, \alpha_r$ ?
- 3 If yes, then how many ways are there to write that lin. comb.?
- 4 We find the answers to these questions as we proceed further.
- 5 Let us see some examples here:
- 6 In  $\mathbb{R}^2$ , the vector  $\begin{pmatrix} 1 \\ 2 \end{pmatrix} = 1 \cdot \begin{pmatrix} 1 \\ 0 \end{pmatrix} + 2 \cdot \begin{pmatrix} 0 \\ 1 \end{pmatrix}$  (Unique way).
- 7  $\begin{pmatrix} 1 \\ 2 \end{pmatrix} = 0 \cdot \begin{pmatrix} 1 \\ 0 \end{pmatrix} + 1 \cdot \begin{pmatrix} 0 \\ 1 \end{pmatrix} + 1 \cdot \begin{pmatrix} 1 \\ 1 \end{pmatrix} = 2 \cdot \begin{pmatrix} 1 \\ 0 \end{pmatrix} + 3 \cdot \begin{pmatrix} 0 \\ 1 \end{pmatrix} - 1 \cdot \begin{pmatrix} 1 \\ 1 \end{pmatrix}.$

# More on linear combinations

- 1 Suppose  $\alpha_1, \alpha_2, \dots, \alpha_r$  and  $\beta$  are given vectors in  $V$ .
- 2 Is  $\beta$  a linear combination of  $\alpha_1, \alpha_2, \dots, \alpha_r$ ?
- 3 If yes, then how many ways are there to write that lin. comb.?
- 4 We find the answers to these questions as we proceed further.
- 5 Let us see some examples here:
- 6 In  $\mathbb{R}^2$ , the vector  $\begin{pmatrix} 1 \\ 2 \end{pmatrix} = 1 \cdot \begin{pmatrix} 1 \\ 0 \end{pmatrix} + 2 \cdot \begin{pmatrix} 0 \\ 1 \end{pmatrix}$  (Unique way).
- 7  $\begin{pmatrix} 1 \\ 2 \end{pmatrix} = 0 \cdot \begin{pmatrix} 1 \\ 0 \end{pmatrix} + 1 \cdot \begin{pmatrix} 0 \\ 1 \end{pmatrix} + 1 \cdot \begin{pmatrix} 1 \\ 1 \end{pmatrix} = 2 \cdot \begin{pmatrix} 1 \\ 0 \end{pmatrix} + 3 \cdot \begin{pmatrix} 0 \\ 1 \end{pmatrix} - 1 \cdot \begin{pmatrix} 1 \\ 1 \end{pmatrix}$ .
- 8 So the vector  $\begin{pmatrix} 1 \\ 2 \end{pmatrix}$  can be written as linear combinations of  $\begin{pmatrix} 1 \\ 0 \end{pmatrix}$ ,  $\begin{pmatrix} 0 \\ 1 \end{pmatrix}$  and  $\begin{pmatrix} 1 \\ 1 \end{pmatrix}$  in infinitely many ways.

# Subspaces: Definition and examples

## Definition

Let  $V$  be a vector space over  $\mathbb{R}$ .

# Subspaces: Definition and examples

## Definition

Let  $V$  be a vector space over  $\mathbb{R}$ . A subspace of  $V$  is a subset  $W$  of  $V$

# Subspaces: Definition and examples

## Definition

Let  $V$  be a vector space over  $\mathbb{R}$ . A subspace of  $V$  is a subset  $W$  of  $V$  which is itself a vector space over  $\mathbb{R}$

# Subspaces: Definition and examples

## Definition


Let  $V$  be a vector space over  $\mathbb{R}$ . A subspace of  $V$  is a subset  $W$  of  $V$  which is itself a vector space over  $\mathbb{R}$  with the same operations of vector addition and scalar multiplication on  $V$ .



# Subspaces: Definition and examples

## Definition


Let  $V$  be a vector space over  $\mathbb{R}$ . A subspace of  $V$  is a subset  $W$  of  $V$  which is itself a vector space over  $\mathbb{R}$  with the same operations of vector addition and scalar multiplication on  $V$ .



# Subspaces: Definition and examples

## Definition

Let  $V$  be a vector space over  $\mathbb{R}$ . A subspace of  $V$  is a subset  $W$  of  $V$  which is itself a vector space over  $\mathbb{R}$  with the same operations of vector addition and scalar multiplication on  $V$ .




Both red and blue lines are subspaces of  $\mathbb{R}^2$ .

# Subspaces: Definition and examples

## Definition

Let  $V$  be a vector space over  $\mathbb{R}$ . A subspace of  $V$  is a subset  $W$  of  $V$  which is itself a vector space over  $\mathbb{R}$  with the same operations of vector addition and scalar multiplication on  $V$ .




Both red and blue lines are subspaces of  $\mathbb{R}^2$ . In fact, any line passing through the origin is a subspace of  $\mathbb{R}^2$ .

# Subspaces: Definition and examples

## Definition

Let  $V$  be a vector space over  $\mathbb{R}$ . A subspace of  $V$  is a subset  $W$  of  $V$  which is itself a vector space over  $\mathbb{R}$  with the same operations of vector addition and scalar multiplication on  $V$ .




Both red and blue lines are subspaces of  $\mathbb{R}^2$ . In fact, any line passing through the origin is a subspace of  $\mathbb{R}^2$ .

For a subspace  $W$  of  $V$ , is it true that  $0 \in W$ ?

# Subspaces: Definition and examples

## Definition

Let  $V$  be a vector space over  $\mathbb{R}$ . A subspace of  $V$  is a subset  $W$  of  $V$  which is itself a vector space over  $\mathbb{R}$  with the same operations of vector addition and scalar multiplication on  $V$ .



Both red and blue lines are subspaces of  $\mathbb{R}^2$ . In fact, any line passing through the origin is a subspace of  $\mathbb{R}^2$ .

For a subspace  $W$  of  $V$ , is it true that  $0 \in W$ ? Answer: Yes.

# Criterion for a subset of a vector space to be a subspace

## Theorem

*Let  $W$  be a non-empty subset of a vector space  $V$  over  $\mathbb{R}$ .*

# Criterion for a subset of a vector space to be a subspace

## Theorem

*Let  $W$  be a non-empty subset of a vector space  $V$  over  $\mathbb{R}$ .  
Then  $W$  is a subspace of  $V$*

# Criterion for a subset of a vector space to be a subspace

## Theorem

*Let  $W$  be a non-empty subset of a vector space  $V$  over  $\mathbb{R}$ .*

*Then  $W$  is a subspace of  $V$*

*if and only if*

*for each pair of vectors  $\alpha, \beta \in W$  and each scalar  $c \in \mathbb{R}$ ,*



# Criterion for a subset of a vector space to be a subspace

## Theorem

*Let  $W$  be a non-empty subset of a vector space  $V$  over  $\mathbb{R}$ .*

*Then  $W$  is a subspace of  $V$*

*if and only if*

*for each pair of vectors  $\alpha, \beta \in W$  and each scalar  $c \in \mathbb{R}$ , the vector  $c\alpha + \beta$  belongs to  $W$ .*

# Criterion for a subset of a vector space to be a subspace

## Theorem

*Let  $W$  be a non-empty subset of a vector space  $V$  over  $\mathbb{R}$ .  
Then  $W$  is a subspace of  $V$*

*if and only if*

*for each pair of vectors  $\alpha, \beta \in W$  and each scalar  $c \in \mathbb{R}$ , the vector  $c\alpha + \beta$  belongs to  $W$ .*

**Proof.** Exercise!

# Criterion for a subset of a vector space to be a subspace

## Theorem

*Let  $W$  be a non-empty subset of a vector space  $V$  over  $\mathbb{R}$ .  
Then  $W$  is a subspace of  $V$*

*if and only if*

*for each pair of vectors  $\alpha, \beta \in W$  and each scalar  $c \in \mathbb{R}$ , the vector  $c\alpha + \beta$  belongs to  $W$ .*

**Proof.** Exercise!

Note that many properties of  $W$  will be inherited from  $V$ .

# Examples of subspaces

- The subset  $\{0\}$  consisting of the zero vector of  $V$

# Examples of subspaces

- The subset  $\{0\}$  consisting of the zero vector of  $V$  is a subspace of  $V$ .

# Examples of subspaces

- The subset  $\{0\}$  consisting of the zero vector of  $V$  is a subspace of  $V$ . It is also denoted by  $0$ .

# Examples of subspaces

- The subset  $\{0\}$  consisting of the zero vector of  $V$  is a subspace of  $V$ . It is also denoted by  $0$ .
- In  $\mathbb{R}^n$ , the set

$$\{(x_1, \dots, x_n) \in \mathbb{R}^n : x_1 = 0\}$$

# Examples of subspaces

- The subset  $\{0\}$  consisting of the zero vector of  $V$  is a subspace of  $V$ . It is also denoted by  $0$ .
- In  $\mathbb{R}^n$ , the set

$$\{(x_1, \dots, x_n) \in \mathbb{R}^n : x_1 = 0\}$$

is a subspace;



# Examples of subspaces

- The subset  $\{0\}$  consisting of the zero vector of  $V$  is a subspace of  $V$ . It is also denoted by  $0$ .
- In  $\mathbb{R}^n$ , the set

$$\{(x_1, \dots, x_n) \in \mathbb{R}^n : x_1 = 0\}$$

is a subspace; while the set

$$\{(x_1, \dots, x_n) \in \mathbb{R}^n : x_1 = 1\}$$

# Examples of subspaces

- The subset  $\{0\}$  consisting of the zero vector of  $V$  is a subspace of  $V$ . It is also denoted by  $0$ .
- In  $\mathbb{R}^n$ , the set

$$\{(x_1, \dots, x_n) \in \mathbb{R}^n : x_1 = 0\}$$

is a subspace; while the set

$$\{(x_1, \dots, x_n) \in \mathbb{R}^n : x_1 = 1\}$$

is NOT a subspace.

# Examples of subspaces

- The subset  $\{0\}$  consisting of the zero vector of  $V$  is a subspace of  $V$ . It is also denoted by  $0$ .
- In  $\mathbb{R}^n$ , the set

$$\{(x_1, \dots, x_n) \in \mathbb{R}^n : x_1 = 0\}$$

is a subspace; while the set

$$\{(x_1, \dots, x_n) \in \mathbb{R}^n : x_1 = 1\}$$

is NOT a subspace.

- The set of all 'symmetric matrices'

# Examples of subspaces

- The subset  $\{0\}$  consisting of the zero vector of  $V$  is a subspace of  $V$ . It is also denoted by  $0$ .
- In  $\mathbb{R}^n$ , the set

$$\{(x_1, \dots, x_n) \in \mathbb{R}^n : x_1 = 0\}$$

is a subspace; while the set

$$\{(x_1, \dots, x_n) \in \mathbb{R}^n : x_1 = 1\}$$

is NOT a subspace.

- The set of all 'symmetric matrices' forms a subspace of the space of all  $n \times n$  matrices.

# Examples of subspaces

- The subset  $\{0\}$  consisting of the zero vector of  $V$  is a subspace of  $V$ . It is also denoted by  $0$ .
- In  $\mathbb{R}^n$ , the set

$$\{(x_1, \dots, x_n) \in \mathbb{R}^n : x_1 = 0\}$$

is a subspace; while the set

$$\{(x_1, \dots, x_n) \in \mathbb{R}^n : x_1 = 1\}$$

is NOT a subspace.

- The set of all 'symmetric matrices' forms a subspace of the space of all  $n \times n$  matrices. Recall that an  $n \times n$  square matrix  $A$  is said to be symmetric if  $A_{ij} = A_{ji}$  for each  $i$  and  $j$ .

# Subspace spanned by a set

## Definition

Let  $S$  be a set of vectors in a vector space  $V$ .

# Subspace spanned by a set

## Definition

Let  $S$  be a set of vectors in a vector space  $V$ . The **subspace spanned** by (or generated by)  $S$  is defined to be the smallest subspace of  $V$  containing  $S$ .

# Subspace spanned by a set

## Definition

Let  $S$  be a set of vectors in a vector space  $V$ . The **subspace spanned** by (or generated by)  $S$  is defined to be the smallest subspace of  $V$  containing  $S$ . We denote this subspace by  $\text{Span}(S)$ .



# Subspace spanned by a set

## Definition

Let  $S$  be a set of vectors in a vector space  $V$ . The **subspace spanned** by (or generated by)  $S$  is defined to be the smallest subspace of  $V$  containing  $S$ . We denote this subspace by  $\text{Span}(S)$ . We write that  $S$  spans the subspace  $\text{Span}(S)$ .

# Subspace spanned by a set

## Definition

Let  $S$  be a set of vectors in a vector space  $V$ . The **subspace spanned** by (or generated by)  $S$  is defined to be the smallest subspace of  $V$  containing  $S$ . We denote this subspace by  $\text{Span}(S)$ . We write that  $S$  spans the subspace  $\text{Span}(S)$ .

## Theorem

*Let  $S$  be a set of vectors in a vector space  $V$ .*

# Subspace spanned by a set

## Definition

Let  $S$  be a set of vectors in a vector space  $V$ . The **subspace spanned** by (or generated by)  $S$  is defined to be the smallest subspace of  $V$  containing  $S$ . We denote this subspace by  $\text{Span}(S)$ . We write that  $S$  spans the subspace  $\text{Span}(S)$ .

## Theorem

*Let  $S$  be a set of vectors in a vector space  $V$ . The following subspaces are equal.*

# Subspace spanned by a set

## Definition

Let  $S$  be a set of vectors in a vector space  $V$ . The **subspace spanned** by (or generated by)  $S$  is defined to be the smallest subspace of  $V$  containing  $S$ . We denote this subspace by  $\text{Span}(S)$ . We write that  $S$  spans the subspace  $\text{Span}(S)$ .

## Theorem

*Let  $S$  be a set of vectors in a vector space  $V$ . The following subspaces are equal.*

- 1 *The intersection of all subspaces of  $V$  containing  $S$ .*

# Subspace spanned by a set

## Definition

Let  $S$  be a set of vectors in a vector space  $V$ . The **subspace spanned** by (or generated by)  $S$  is defined to be the smallest subspace of  $V$  containing  $S$ . We denote this subspace by  $\text{Span}(S)$ . We write that  $S$  spans the subspace  $\text{Span}(S)$ .

## Theorem

*Let  $S$  be a set of vectors in a vector space  $V$ . The following subspaces are equal.*

- ❶ *The intersection of all subspaces of  $V$  containing  $S$ .*
- ❷ *The set of all linear combinations of vectors in  $S$ , i.e.*

$$\{c_1 v_1 + \cdots + c_r v_r : c_i \in \mathbb{R}, v_i \in S\}.$$

# Subspace spanned by a set

## Definition

Let  $S$  be a set of vectors in a vector space  $V$ . The **subspace spanned** by (or generated by)  $S$  is defined to be the smallest subspace of  $V$  containing  $S$ . We denote this subspace by  $\text{Span}(S)$ . We write that  $S$  spans the subspace  $\text{Span}(S)$ .

## Theorem

*Let  $S$  be a set of vectors in a vector space  $V$ . The following subspaces are equal.*

- 1 *The intersection of all subspaces of  $V$  containing  $S$ .*
- 2 *The set of all linear combinations of vectors in  $S$ , i.e.*

$$\{c_1 v_1 + \cdots + c_r v_r : c_i \in \mathbb{R}, v_i \in S\}.$$

- 3 *The subspace spanned by  $S$  (i.e. the smallest subspace of  $V$  containing  $S$ ).*

# Subspace spanned by a set

## Theorem

*Let  $S$  be a subset of  $V$ . The following subspaces are equal.*

- ❶ *The intersection of all subspaces of  $V$  containing  $S$ .*
- ❷ *The set of all linear combinations of vectors in  $S$ , i.e.  $\{c_1 v_1 + \cdots + c_r v_r : c_i \in \mathbb{R}, v_i \in S\}$ .*
- ❸ *The subspace spanned by  $S$ , i.e. the smallest subspace of  $V$  containing  $S$ .*

# Subspace spanned by a set

## Theorem

*Let  $S$  be a subset of  $V$ . The following subspaces are equal.*

- ❶ *The intersection of all subspaces of  $V$  containing  $S$ .*
- ❷ *The set of all linear combinations of vectors in  $S$ , i.e.  $\{c_1 v_1 + \cdots + c_r v_r : c_i \in \mathbb{R}, v_i \in S\}$ .*
- ❸ *The subspace spanned by  $S$ , i.e. the smallest subspace of  $V$  containing  $S$ .*

*Proof.* Let  $W_1, W_2$  and  $W_3$  be the subspaces described as in (1), (2) and (3) respectively.



# Subspace spanned by a set

## Theorem

*Let  $S$  be a subset of  $V$ . The following subspaces are equal.*

- ❶ *The intersection of all subspaces of  $V$  containing  $S$ .*
- ❷ *The set of all linear combinations of vectors in  $S$ , i.e.  $\{c_1 v_1 + \cdots + c_r v_r : c_i \in \mathbb{R}, v_i \in S\}$ .*
- ❸ *The subspace spanned by  $S$ , i.e. the smallest subspace of  $V$  containing  $S$ .*

*Proof.* Let  $W_1, W_2$  and  $W_3$  be the subspaces described as in (1), (2) and (3) respectively. Clearly,  $W_1$  is contained in any subspace of  $V$  containing  $S$ .

# Subspace spanned by a set

## Theorem

*Let  $S$  be a subset of  $V$ . The following subspaces are equal.*

- ❶ *The intersection of all subspaces of  $V$  containing  $S$ .*
- ❷ *The set of all linear combinations of vectors in  $S$ , i.e.  $\{c_1 v_1 + \cdots + c_r v_r : c_i \in \mathbb{R}, v_i \in S\}$ .*
- ❸ *The subspace spanned by  $S$ , i.e. the smallest subspace of  $V$  containing  $S$ .*

*Proof.* Let  $W_1, W_2$  and  $W_3$  be the subspaces described as in (1), (2) and (3) respectively. Clearly,  $W_1$  is contained in any subspace of  $V$  containing  $S$ . Since  $W_1$  is a subspace,  $W_1$  is the smallest subspace of  $V$  containing  $S$ ,

# Subspace spanned by a set

## Theorem

*Let  $S$  be a subset of  $V$ . The following subspaces are equal.*

- ❶ *The intersection of all subspaces of  $V$  containing  $S$ .*
- ❷ *The set of all linear combinations of vectors in  $S$ , i.e.  $\{c_1 v_1 + \cdots + c_r v_r : c_i \in \mathbb{R}, v_i \in S\}$ .*
- ❸ *The subspace spanned by  $S$ , i.e. the smallest subspace of  $V$  containing  $S$ .*

*Proof.* Let  $W_1, W_2$  and  $W_3$  be the subspaces described as in (1), (2) and (3) respectively. Clearly,  $W_1$  is contained in any subspace of  $V$  containing  $S$ . Since  $W_1$  is a subspace,  $W_1$  is the smallest subspace of  $V$  containing  $S$ , i.e.  $W_1 = W_3$ .

# Subspace spanned by a set

## Theorem

*Let  $S$  be a subset of  $V$ . The following subspaces are equal.*

- ❶ *The intersection of all subspaces of  $V$  containing  $S$ .*
- ❷ *The set of all linear combinations of vectors in  $S$ , i.e.  $\{c_1 v_1 + \cdots + c_r v_r : c_i \in \mathbb{R}, v_i \in S\}$ .*
- ❸ *The subspace spanned by  $S$ , i.e. the smallest subspace of  $V$  containing  $S$ .*

*Proof.* Let  $W_1, W_2$  and  $W_3$  be the subspaces described as in (1), (2) and (3) respectively. Clearly,  $W_1$  is contained in any subspace of  $V$  containing  $S$ . Since  $W_1$  is a subspace,  $W_1$  is the smallest subspace of  $V$  containing  $S$ , i.e.  $W_1 = W_3$ .

Note that  $W_2$  is a subspace containing  $S$ .

# Subspace spanned by a set

## Theorem

*Let  $S$  be a subset of  $V$ . The following subspaces are equal.*

- ❶ *The intersection of all subspaces of  $V$  containing  $S$ .*
- ❷ *The set of all linear combinations of vectors in  $S$ , i.e.  $\{c_1 v_1 + \cdots + c_r v_r : c_i \in \mathbb{R}, v_i \in S\}$ .*
- ❸ *The subspace spanned by  $S$ , i.e. the smallest subspace of  $V$  containing  $S$ .*

*Proof.* Let  $W_1, W_2$  and  $W_3$  be the subspaces described as in (1), (2) and (3) respectively. Clearly,  $W_1$  is contained in any subspace of  $V$  containing  $S$ . Since  $W_1$  is a subspace,  $W_1$  is the smallest subspace of  $V$  containing  $S$ , i.e.  $W_1 = W_3$ .

Note that  $W_2$  is a subspace containing  $S$ . So  $W_1 \subseteq W_2$ .

# Subspace spanned by a set

## Theorem

*Let  $S$  be a subset of  $V$ . The following subspaces are equal.*

- ① The intersection of all subspaces of  $V$  containing  $S$ .*
- ② The set of all linear combinations of vectors in  $S$ , i.e.  $\{c_1 v_1 + \cdots + c_r v_r : c_i \in \mathbb{R}, v_i \in S\}$ .*
- ③ The subspace spanned by  $S$ , i.e. the smallest subspace of  $V$  containing  $S$ .*

*Proof.* Let  $W_1, W_2$  and  $W_3$  be the subspaces described as in (1), (2) and (3) respectively. Clearly,  $W_1$  is contained in any subspace of  $V$  containing  $S$ . Since  $W_1$  is a subspace,  $W_1$  is the smallest subspace of  $V$  containing  $S$ , i.e.  $W_1 = W_3$ .

Note that  $W_2$  is a subspace containing  $S$ . So  $W_1 \subseteq W_2$ . Notice that any subspace of  $V$  containing  $S$  also contains all linear combinations of vectors in  $S$ .

# Subspace spanned by a set

## Theorem

*Let  $S$  be a subset of  $V$ . The following subspaces are equal.*

- ❶ *The intersection of all subspaces of  $V$  containing  $S$ .*
- ❷ *The set of all linear combinations of vectors in  $S$ , i.e.  $\{c_1 v_1 + \cdots + c_r v_r : c_i \in \mathbb{R}, v_i \in S\}$ .*
- ❸ *The subspace spanned by  $S$ , i.e. the smallest subspace of  $V$  containing  $S$ .*

*Proof.* Let  $W_1, W_2$  and  $W_3$  be the subspaces described as in (1), (2) and (3) respectively. Clearly,  $W_1$  is contained in any subspace of  $V$  containing  $S$ . Since  $W_1$  is a subspace,  $W_1$  is the smallest subspace of  $V$  containing  $S$ , i.e.  $W_1 = W_3$ .

Note that  $W_2$  is a subspace containing  $S$ . So  $W_1 \subseteq W_2$ . Notice that any subspace of  $V$  containing  $S$  also contains all linear combinations of vectors in  $S$ . Hence it follows that  $W_2 \subseteq W_1$ .

# Subspace spanned by a set

## Theorem

*Let  $S$  be a subset of  $V$ . The following subspaces are equal.*

- ❶ *The intersection of all subspaces of  $V$  containing  $S$ .*
- ❷ *The set of all linear combinations of vectors in  $S$ , i.e.  $\{c_1 v_1 + \cdots + c_r v_r : c_i \in \mathbb{R}, v_i \in S\}$ .*
- ❸ *The subspace spanned by  $S$ , i.e. the smallest subspace of  $V$  containing  $S$ .*

*Proof.* Let  $W_1, W_2$  and  $W_3$  be the subspaces described as in (1), (2) and (3) respectively. Clearly,  $W_1$  is contained in any subspace of  $V$  containing  $S$ . Since  $W_1$  is a subspace,  $W_1$  is the smallest subspace of  $V$  containing  $S$ , i.e.  $W_1 = W_3$ .

Note that  $W_2$  is a subspace containing  $S$ . So  $W_1 \subseteq W_2$ . Notice that any subspace of  $V$  containing  $S$  also contains all linear combinations of vectors in  $S$ . Hence it follows that  $W_2 \subseteq W_1$ . Therefore  $W_1 = W_2$ .



# Subspace spanned by a set

## Theorem

*Let  $S$  be a subset of  $V$ . The following subspaces are equal.*

- ❶ *The intersection of all subspaces of  $V$  containing  $S$ .*
- ❷ *The set of all linear combinations of vectors in  $S$ , i.e.  $\{c_1 v_1 + \cdots + c_r v_r : c_i \in \mathbb{R}, v_i \in S\}$ .*
- ❸ *The subspace spanned by  $S$ , i.e. the smallest subspace of  $V$  containing  $S$ .*

*Proof.* Let  $W_1, W_2$  and  $W_3$  be the subspaces described as in (1), (2) and (3) respectively. Clearly,  $W_1$  is contained in any subspace of  $V$  containing  $S$ . Since  $W_1$  is a subspace,  $W_1$  is the smallest subspace of  $V$  containing  $S$ , i.e.  $W_1 = W_3$ .

Note that  $W_2$  is a subspace containing  $S$ . So  $W_1 \subseteq W_2$ . Notice that any subspace of  $V$  containing  $S$  also contains all linear combinations of vectors in  $S$ . Hence it follows that  $W_2 \subseteq W_1$ . Therefore  $W_1 = W_2$ . Thus  $W_1 = W_2 = W_3$ .

# Examples of subspaces spanned by subsets

- 1 Let  $V = \mathbb{R}^{2 \times 2}$  be the space of all  $2 \times 2$  matrices over  $\mathbb{R}$ .

# Examples of subspaces spanned by subsets

- ① Let  $V = \mathbb{R}^{2 \times 2}$  be the space of all  $2 \times 2$  matrices over  $\mathbb{R}$ . Set

$$S := \left\{ \begin{pmatrix} 1 & 0 \\ 0 & 0 \end{pmatrix}, \begin{pmatrix} 0 & 0 \\ 0 & 1 \end{pmatrix}, \begin{pmatrix} 0 & 1 \\ 1 & 0 \end{pmatrix} \right\}.$$

# Examples of subspaces spanned by subsets

- ① Let  $V = \mathbb{R}^{2 \times 2}$  be the space of all  $2 \times 2$  matrices over  $\mathbb{R}$ . Set

$$S := \left\{ \begin{pmatrix} 1 & 0 \\ 0 & 0 \end{pmatrix}, \begin{pmatrix} 0 & 0 \\ 0 & 1 \end{pmatrix}, \begin{pmatrix} 0 & 1 \\ 1 & 0 \end{pmatrix} \right\}.$$

The subspace spanned by  $S$  is the subspace of all  $2 \times 2$  symmetric matrices over  $\mathbb{R}$ .

# Examples of subspaces spanned by subsets

- ❶ Let  $V = \mathbb{R}^{2 \times 2}$  be the space of all  $2 \times 2$  matrices over  $\mathbb{R}$ . Set

$$S := \left\{ \begin{pmatrix} 1 & 0 \\ 0 & 0 \end{pmatrix}, \begin{pmatrix} 0 & 0 \\ 0 & 1 \end{pmatrix}, \begin{pmatrix} 0 & 1 \\ 1 & 0 \end{pmatrix} \right\}.$$

The subspace spanned by  $S$  is the subspace of all  $2 \times 2$  symmetric matrices over  $\mathbb{R}$ .

- ❷ Let  $V = \mathbb{R}[x]$  (set of all polynomials).

# Examples of subspaces spanned by subsets

- ① Let  $V = \mathbb{R}^{2 \times 2}$  be the space of all  $2 \times 2$  matrices over  $\mathbb{R}$ . Set

$$S := \left\{ \begin{pmatrix} 1 & 0 \\ 0 & 0 \end{pmatrix}, \begin{pmatrix} 0 & 0 \\ 0 & 1 \end{pmatrix}, \begin{pmatrix} 0 & 1 \\ 1 & 0 \end{pmatrix} \right\}.$$

The subspace spanned by  $S$  is the subspace of all  $2 \times 2$  symmetric matrices over  $\mathbb{R}$ .

- ② Let  $V = \mathbb{R}[x]$  (set of all polynomials). Set

$$S := \{f_n(x) = x^n : n = 0, 1, 2, \dots\}.$$

# Examples of subspaces spanned by subsets

- ❶ Let  $V = \mathbb{R}^{2 \times 2}$  be the space of all  $2 \times 2$  matrices over  $\mathbb{R}$ . Set

$$S := \left\{ \begin{pmatrix} 1 & 0 \\ 0 & 0 \end{pmatrix}, \begin{pmatrix} 0 & 0 \\ 0 & 1 \end{pmatrix}, \begin{pmatrix} 0 & 1 \\ 1 & 0 \end{pmatrix} \right\}.$$


The subspace spanned by  $S$  is the subspace of all  $2 \times 2$  symmetric matrices over  $\mathbb{R}$ .

- ❷ Let  $V = \mathbb{R}[x]$  (set of all polynomials). Set

$$S := \{f_n(x) = x^n : n = 0, 1, 2, \dots\}.$$

Then the subspace spanned by  $S$  is  $V$  itself.


# Examples of subspaces spanned by subsets



In this figure,




# Examples of subspaces spanned by subsets



In this figure,  
the subspace spanned by  $\{u\}$  can be described by the dotted line;

# Examples of subspaces spanned by subsets



In this figure,  
the subspace spanned by  $\{u\}$  can be described by the dotted line;  
while the subspace spanned by  $\{u, v\}$  is  $\mathbb{R}^2$ .

# Linearly dependent or independent

Throughout, let  $V$  be a vector space over  $\mathbb{R}$ .

# Linearly dependent or independent

Throughout, let  $V$  be a vector space over  $\mathbb{R}$ .

## Definition

A subset  $S$  of vectors in  $V$  is said to be **linearly dependent**

# Linearly dependent or independent

Throughout, let  $V$  be a vector space over  $\mathbb{R}$ .

## Definition

A subset  $S$  of vectors in  $V$  is said to be **linearly dependent** if there exists vectors  $v_1, v_2, \dots, v_r$  in  $S$  and

# Linearly dependent or independent

Throughout, let  $V$  be a vector space over  $\mathbb{R}$ .

## Definition

A subset  $S$  of vectors in  $V$  is said to be **linearly dependent** if there exists vectors  $v_1, v_2, \dots, v_r$  in  $S$  and scalars  $c_1, c_2, \dots, c_r$  in  $\mathbb{R}$ ,

# Linearly dependent or independent

Throughout, let  $V$  be a vector space over  $\mathbb{R}$ .

## Definition

A subset  $S$  of vectors in  $V$  is said to be **linearly dependent** if there exists vectors  $v_1, v_2, \dots, v_r$  in  $S$  and scalars  $c_1, c_2, \dots, c_r$  in  $\mathbb{R}$ , NOT ALL of which are 0,

# Linearly dependent or independent

Throughout, let  $V$  be a vector space over  $\mathbb{R}$ .

## Definition

A subset  $S$  of vectors in  $V$  is said to be **linearly dependent** if there exists vectors  $v_1, v_2, \dots, v_r$  in  $S$  and scalars  $c_1, c_2, \dots, c_r$  in  $\mathbb{R}$ , NOT ALL of which are 0, such that

$$c_1 v_1 + \dots + c_r v_r = 0.$$



# Linearly dependent or independent

Throughout, let  $V$  be a vector space over  $\mathbb{R}$ .

## Definition

A subset  $S$  of vectors in  $V$  is said to be **linearly dependent** if there exists vectors  $v_1, v_2, \dots, v_r$  in  $S$  and scalars  $c_1, c_2, \dots, c_r$  in  $\mathbb{R}$ , NOT ALL of which are 0, such that

$$c_1 v_1 + \dots + c_r v_r = 0.$$

## Definition

A set  $S$  which is not linearly dependent is called **linearly independent**.

# Linearly dependent or independent

Throughout, let  $V$  be a vector space over  $\mathbb{R}$ .

## Definition

A subset  $S$  of vectors in  $V$  is said to be **linearly dependent** if there exists vectors  $v_1, v_2, \dots, v_r$  in  $S$  and scalars  $c_1, c_2, \dots, c_r$  in  $\mathbb{R}$ , NOT ALL of which are 0, such that

$$c_1 v_1 + \dots + c_r v_r = 0.$$

## Definition

A set  $S$  which is not linearly dependent is called **linearly independent**.

If  $S = \{v_1, \dots, v_n\}$ , we say that  $v_1, \dots, v_n$  are linearly dependent (or independent) instead of saying that  $S$  is so.

# Remarks on linearly dependent vectors

- 1 Any set containing the  $0$  vector

# Remarks on linearly dependent vectors

- 1 Any set containing the  $0$  vector is linearly dependent.

# Remarks on linearly dependent vectors

- ① Any set containing the 0 vector is linearly dependent.
- ② A set  $S$  of vectors is linearly dependent if and only if there exists a non-trivial relation of some vectors of  $S$ :

$$c_1 v_1 + \cdots + c_r v_r = 0, \text{ where at least one } c_i \neq 0.$$

# Remarks on linearly dependent vectors

- ① Any set containing the 0 vector is linearly dependent.
- ② A set  $S$  of vectors is linearly dependent if and only if there exists a non-trivial relation of some vectors of  $S$ :

$$c_1 v_1 + \cdots + c_r v_r = 0, \text{ where at least one } c_i \neq 0.$$

This is equivalent to say that there exists at least one vector  $v \in S$

# Remarks on linearly dependent vectors

- ① Any set containing the 0 vector is linearly dependent.
- ② A set  $S$  of vectors is linearly dependent if and only if there exists a non-trivial relation of some vectors of  $S$ :

$$c_1 v_1 + \cdots + c_r v_r = 0, \text{ where at least one } c_i \neq 0.$$

This is equivalent to say that there exists at least one vector  $v \in S$  which belongs to the subspace spanned by  $S \setminus \{v\}$ .

# Remarks on linearly dependent vectors

- ① Any set containing the 0 vector is linearly dependent.
- ② A set  $S$  of vectors is linearly dependent if and only if there exists a non-trivial relation of some vectors of  $S$ :

$$c_1 v_1 + \cdots + c_r v_r = 0, \text{ where at least one } c_i \neq 0.$$

This is equivalent to say that there exists at least one vector  $v \in S$  which belongs to the subspace spanned by  $S \setminus \{v\}$ .

- ③ Any set containing a linearly dependent subset



# Remarks on linearly dependent vectors

- ① Any set containing the 0 vector is linearly dependent.
- ② A set  $S$  of vectors is linearly dependent if and only if there exists a non-trivial relation of some vectors of  $S$ :

$$c_1 v_1 + \cdots + c_r v_r = 0, \text{ where at least one } c_i \neq 0.$$

This is equivalent to say that there exists at least one vector  $v \in S$  which belongs to the subspace spanned by  $S \setminus \{v\}$ .

- ③ Any set containing a linearly dependent subset is again linearly dependent.

# Remarks on linearly independent vectors

- 1 Every non-zero vector  $v$  in  $V$

# Remarks on linearly independent vectors

- 1 Every non-zero vector  $v$  in  $V$  is linearly independent.

# Remarks on linearly independent vectors

- ① Every non-zero vector  $v$  in  $V$  is linearly independent.
- ② A finite set  $\{v_1, \dots, v_r\}$  is linearly independent

# Remarks on linearly independent vectors

- ① Every non-zero vector  $v$  in  $V$  is linearly independent.
- ② A finite set  $\{v_1, \dots, v_r\}$  is linearly independent if and only if
$$c_1 v_1 + \dots + c_r v_r = 0 \implies c_i = 0 \text{ for all } 1 \leq i \leq r.$$

# Remarks on linearly independent vectors

- ① Every non-zero vector  $v$  in  $V$  is linearly independent.
- ② A finite set  $\{v_1, \dots, v_r\}$  is linearly independent if and only if
$$c_1 v_1 + \dots + c_r v_r = 0 \implies c_i = 0 \text{ for all } 1 \leq i \leq r.$$
- ③ A set  $S$  of vectors is linearly independent if and only if every finite subset of  $S$  is linearly independent,

# Remarks on linearly independent vectors

- ① Every non-zero vector  $v$  in  $V$  is linearly independent.
- ② A finite set  $\{v_1, \dots, v_r\}$  is linearly independent if and only if
$$c_1 v_1 + \dots + c_r v_r = 0 \implies c_i = 0 \text{ for all } 1 \leq i \leq r.$$
- ③ A set  $S$  of vectors is linearly independent if and only if every finite subset of  $S$  is linearly independent, i.e., if and only if for every subset  $\{v_1, \dots, v_r\} \subseteq S$ ,

$$c_1 v_1 + \dots + c_r v_r = 0$$

# Remarks on linearly independent vectors

- ① Every non-zero vector  $v$  in  $V$  is linearly independent.
- ② A finite set  $\{v_1, \dots, v_r\}$  is linearly independent if and only if
$$c_1 v_1 + \dots + c_r v_r = 0 \implies c_i = 0 \text{ for all } 1 \leq i \leq r.$$
- ③ A set  $S$  of vectors is linearly independent if and only if every finite subset of  $S$  is linearly independent, i.e., if and only if for every subset  $\{v_1, \dots, v_r\} \subseteq S$ ,
$$c_1 v_1 + \dots + c_r v_r = 0 \implies c_i = 0 \text{ for all } 1 \leq i \leq r.$$



# Remarks on linearly independent vectors

- ① Every non-zero vector  $v$  in  $V$  is linearly independent.
- ② A finite set  $\{v_1, \dots, v_r\}$  is linearly independent if and only if
$$c_1 v_1 + \dots + c_r v_r = 0 \implies c_i = 0 \text{ for all } 1 \leq i \leq r.$$
- ③ A set  $S$  of vectors is linearly independent if and only if every finite subset of  $S$  is linearly independent, i.e., if and only if for every subset  $\{v_1, \dots, v_r\} \subseteq S$ ,
$$c_1 v_1 + \dots + c_r v_r = 0 \implies c_i = 0 \text{ for all } 1 \leq i \leq r.$$
- ④ Any subset of a linearly independent set

# Remarks on linearly independent vectors

- ① Every non-zero vector  $v$  in  $V$  is linearly independent.
- ② A finite set  $\{v_1, \dots, v_r\}$  is linearly independent if and only if
$$c_1 v_1 + \dots + c_r v_r = 0 \implies c_i = 0 \text{ for all } 1 \leq i \leq r.$$
- ③ A set  $S$  of vectors is linearly independent if and only if every finite subset of  $S$  is linearly independent, i.e., if and only if for every subset  $\{v_1, \dots, v_r\} \subseteq S$ ,
$$c_1 v_1 + \dots + c_r v_r = 0 \implies c_i = 0 \text{ for all } 1 \leq i \leq r.$$
- ④ Any subset of a linearly independent set is linearly independent.

# Adjoining a vector to a linearly independent set

## Lemma

*Let  $S$  be a linearly independent subset of a vector space  $V$ .*

# Adjoining a vector to a linearly independent set

## Lemma

*Let  $S$  be a linearly independent subset of a vector space  $V$ .  
Suppose  $v \notin \text{Span}(S) := \{c_1 v_1 + \cdots + c_r v_r : v_i \in S, c_i \in \mathbb{R}\}$ .*

# Adjoining a vector to a linearly independent set

## Lemma

*Let  $S$  be a linearly independent subset of a vector space  $V$ .  
Suppose  $v \notin \text{Span}(S) := \{c_1 v_1 + \cdots + c_r v_r : v_i \in S, c_i \in \mathbb{R}\}$ .  
Then  $S \cup \{v\}$  is linearly independent.*

# Adjoining a vector to a linearly independent set

## Lemma

*Let  $S$  be a linearly independent subset of a vector space  $V$ .  
Suppose  $v \notin \text{Span}(S) := \{c_1 v_1 + \cdots + c_r v_r : v_i \in S, c_i \in \mathbb{R}\}$ .  
Then  $S \cup \{v\}$  is linearly independent.*

## Proof.

Let  $c_1 v_1 + \cdots + c_r v_r + cv = 0$  for some vectors  $v_1, \dots, v_r \in S$  and scalars  $c_1, \dots, c_r, c$ .

# Adjoining a vector to a linearly independent set

## Lemma

*Let  $S$  be a linearly independent subset of a vector space  $V$ .  
Suppose  $v \notin \text{Span}(S) := \{c_1 v_1 + \cdots + c_r v_r : v_i \in S, c_i \in \mathbb{R}\}$ .  
Then  $S \cup \{v\}$  is linearly independent.*

## Proof.

Let  $c_1 v_1 + \cdots + c_r v_r + cv = 0$  for some vectors  $v_1, \dots, v_r \in S$  and scalars  $c_1, \dots, c_r, c$ .

If  $c \neq 0$ ,

# Adjoining a vector to a linearly independent set

## Lemma

*Let  $S$  be a linearly independent subset of a vector space  $V$ .  
Suppose  $v \notin \text{Span}(S) := \{c_1 v_1 + \cdots + c_r v_r : v_i \in S, c_i \in \mathbb{R}\}$ .  
Then  $S \cup \{v\}$  is linearly independent.*

## Proof.

Let  $c_1 v_1 + \cdots + c_r v_r + cv = 0$  for some vectors  $v_1, \dots, v_r \in S$  and scalars  $c_1, \dots, c_r, c$ .

If  $c \neq 0$ , then it follows that  $v \in \text{Span}(S)$ ,



# Adjoining a vector to a linearly independent set

## Lemma

*Let  $S$  be a linearly independent subset of a vector space  $V$ .  
Suppose  $v \notin \text{Span}(S) := \{c_1 v_1 + \cdots + c_r v_r : v_i \in S, c_i \in \mathbb{R}\}$ .  
Then  $S \cup \{v\}$  is linearly independent.*

## Proof.

Let  $c_1 v_1 + \cdots + c_r v_r + cv = 0$  for some vectors  $v_1, \dots, v_r \in S$  and scalars  $c_1, \dots, c_r, c$ .

If  $c \neq 0$ , then it follows that  $v \in \text{Span}(S)$ , which is a contradiction.

# Adjoining a vector to a linearly independent set

## Lemma

*Let  $S$  be a linearly independent subset of a vector space  $V$ .  
Suppose  $v \notin \text{Span}(S) := \{c_1 v_1 + \cdots + c_r v_r : v_i \in S, c_i \in \mathbb{R}\}$ .  
Then  $S \cup \{v\}$  is linearly independent.*

## Proof.

Let  $c_1 v_1 + \cdots + c_r v_r + cv = 0$  for some vectors  $v_1, \dots, v_r \in S$  and scalars  $c_1, \dots, c_r, c$ .

If  $c \neq 0$ , then it follows that  $v \in \text{Span}(S)$ , which is a contradiction.  
Therefore  $c = 0$ ,

# Adjoining a vector to a linearly independent set

## Lemma

*Let  $S$  be a linearly independent subset of a vector space  $V$ .  
Suppose  $v \notin \text{Span}(S) := \{c_1 v_1 + \cdots + c_r v_r : v_i \in S, c_i \in \mathbb{R}\}$ .  
Then  $S \cup \{v\}$  is linearly independent.*

## Proof.

Let  $c_1 v_1 + \cdots + c_r v_r + cv = 0$  for some vectors  $v_1, \dots, v_r \in S$  and scalars  $c_1, \dots, c_r, c$ .

If  $c \neq 0$ , then it follows that  $v \in \text{Span}(S)$ , which is a contradiction.  
Therefore  $c = 0$ , and hence  $c_1 v_1 + \cdots + c_r v_r = 0$ .

# Adjoining a vector to a linearly independent set

## Lemma


*Let  $S$  be a linearly independent subset of a vector space  $V$ .  
Suppose  $v \notin \text{Span}(S) := \{c_1 v_1 + \cdots + c_r v_r : v_i \in S, c_i \in \mathbb{R}\}$ .  
Then  $S \cup \{v\}$  is linearly independent.*

## Proof.


Let  $c_1 v_1 + \cdots + c_r v_r + cv = 0$  for some vectors  $v_1, \dots, v_r \in S$  and scalars  $c_1, \dots, c_r, c$ .

If  $c \neq 0$ , then it follows that  $v \in \text{Span}(S)$ , which is a contradiction.  
Therefore  $c = 0$ , and hence  $c_1 v_1 + \cdots + c_r v_r = 0$ . Since  $S$  is linearly independent, it follows that  $c_i = 0$  for all  $1 \leq i \leq r$ .  $\square$


# Vectors in $\mathbb{R}^2$ plane




# Vectors in $\mathbb{R}^2$ plane



# Vectors in $\mathbb{R}^2$ plane




# Vectors in $\mathbb{R}^2$ plane







# Vectors in $\mathbb{R}^2$ plane




# Vectors in $\mathbb{R}^2$ plane




# Vectors in $\mathbb{R}^2$ plane




# Vectors in $\mathbb{R}^2$ plane




# Vectors in $\mathbb{R}^2$ plane




# Vectors in $\mathbb{R}^2$ plane



# Vectors in $\mathbb{R}^2$ plane




# Vectors in $\mathbb{R}^2$ plane





# Vectors in $\mathbb{R}^2$ plane



# Basis of a vector space

## Definition

A set  $S$  of vectors in  $V$  is called a **basis** of  $V$  if

# Basis of a vector space

## Definition

A set  $S$  of vectors in  $V$  is called a **basis** of  $V$  if  
(i)  $S$  is linearly independent, and

# Basis of a vector space

## Definition

A set  $S$  of vectors in  $V$  is called a **basis** of  $V$  if

- (i)  $S$  is linearly independent, and
- (ii) it spans the space  $V$  (i.e., the subspace spanned by  $S$  is  $V$ ).

# Basis of a vector space

## Definition

A set  $S$  of vectors in  $V$  is called a **basis** of  $V$  if

- (i)  $S$  is linearly independent, and
- (ii) it spans the space  $V$  (i.e., the subspace spanned by  $S$  is  $V$ ).

The space  $V$  is said to be **finite dimensional** if it has a finite basis.

## Definition

A set  $S$  of vectors in  $V$  is called a **basis** of  $V$  if

- (i)  $S$  is linearly independent, and
- (ii) it spans the space  $V$  (i.e., the subspace spanned by  $S$  is  $V$ ).

The space  $V$  is said to be **finite dimensional** if it has a finite basis. If  $V$  does not have a finite basis, then  $V$  is said to be **infinite dimensional**.

# Every vector space has a basis

What is the guarantee that a basis exists?

# Every vector space has a basis

What is the guarantee that a basis exists?

We can prove the existence at least when  $V$  is generated (or spanned) by finitely many vectors.



# Every vector space has a basis

What is the guarantee that a basis exists?

We can prove the existence at least when  $V$  is generated (or spanned) by finitely many vectors. How?

# Every vector space has a basis

What is the guarantee that a basis exists?

We can prove the existence at least when  $V$  is generated (or spanned) by finitely many vectors. How?

Start with a finite spanning set  $S$ .

# Every vector space has a basis

What is the guarantee that a basis exists?

We can prove the existence at least when  $V$  is generated (or spanned) by finitely many vectors. How?

Start with a finite spanning set  $S$ . Then check whether it is linearly independent.

# Every vector space has a basis

What is the guarantee that a basis exists?

We can prove the existence at least when  $V$  is generated (or spanned) by finitely many vectors. How?

Start with a finite spanning set  $S$ . Then check whether it is linearly independent. If yes, then we are done.

# Every vector space has a basis

What is the guarantee that a basis exists?

We can prove the existence at least when  $V$  is generated (or spanned) by finitely many vectors. How?

Start with a finite spanning set  $S$ . Then check whether it is linearly independent. If yes, then we are done. Otherwise  $S$  should be linearly dependent, and

# Every vector space has a basis

What is the guarantee that a basis exists?

We can prove the existence at least when  $V$  is generated (or spanned) by finitely many vectors. How?

Start with a finite spanning set  $S$ . Then check whether it is linearly independent. If yes, then we are done. Otherwise  $S$  should be linearly dependent, and there is  $v \in S$  such that  $v$  belongs to the subspace spanned by  $S \setminus \{v\}$ .

# Every vector space has a basis

What is the guarantee that a basis exists?

We can prove the existence at least when  $V$  is generated (or spanned) by finitely many vectors. How?

Start with a finite spanning set  $S$ . Then check whether it is linearly independent. If yes, then we are done. Otherwise  $S$  should be linearly dependent, and there is  $v \in S$  such that  $v$  belongs to the subspace spanned by  $S \setminus \{v\}$ . One can prove that  $S \setminus \{v\}$  spans  $V$ .

# Every vector space has a basis

What is the guarantee that a basis exists?

We can prove the existence at least when  $V$  is generated (or spanned) by finitely many vectors. How?

Start with a finite spanning set  $S$ . Then check whether it is linearly independent. If yes, then we are done. Otherwise  $S$  should be linearly dependent, and there is  $v \in S$  such that  $v$  belongs to the subspace spanned by  $S \setminus \{v\}$ . One can prove that  $S \setminus \{v\}$  spans  $V$ . Repeat the process till we get a linearly independent subset of  $S$  which spans  $V$ .



# Every vector space has a basis

What is the guarantee that a basis exists?

We can prove the existence at least when  $V$  is generated (or spanned) by finitely many vectors. How?

Start with a finite spanning set  $S$ . Then check whether it is linearly independent. If yes, then we are done. Otherwise  $S$  should be linearly dependent, and there is  $v \in S$  such that  $v$  belongs to the subspace spanned by  $S \setminus \{v\}$ . One can prove that  $S \setminus \{v\}$  spans  $V$ . Repeat the process till we get a linearly independent subset of  $S$  which spans  $V$ .

For vector space which is not finitely generated,

# Every vector space has a basis

What is the guarantee that a basis exists?

We can prove the existence at least when  $V$  is generated (or spanned) by finitely many vectors. How?

Start with a finite spanning set  $S$ . Then check whether it is linearly independent. If yes, then we are done. Otherwise  $S$  should be linearly dependent, and there is  $v \in S$  such that  $v$  belongs to the subspace spanned by  $S \setminus \{v\}$ . One can prove that  $S \setminus \{v\}$  spans  $V$ . Repeat the process till we get a linearly independent subset of  $S$  which spans  $V$ .

For vector space which is not finitely generated, we need the axiom of choice.

# Every vector space has a basis

What is the guarantee that a basis exists?

We can prove the existence at least when  $V$  is generated (or spanned) by finitely many vectors. How?

Start with a finite spanning set  $S$ . Then check whether it is linearly independent. If yes, then we are done. Otherwise  $S$  should be linearly dependent, and there is  $v \in S$  such that  $v$  belongs to the subspace spanned by  $S \setminus \{v\}$ . One can prove that  $S \setminus \{v\}$  spans  $V$ . Repeat the process till we get a linearly independent subset of  $S$  which spans  $V$ .

For vector space which is not finitely generated, we need the axiom of choice. We will not do that in this course.

# An example of a basis of $\mathbb{R}^2$

The set  $\left\{ u = \begin{pmatrix} 1 \\ 2 \end{pmatrix}, v = \begin{pmatrix} 2 \\ 1 \end{pmatrix} \right\}$  forms a basis of  $\mathbb{R}^2$ .

# An example of a basis of $\mathbb{R}^2$

The set  $\left\{ u = \begin{pmatrix} 1 \\ 2 \end{pmatrix}, v = \begin{pmatrix} 2 \\ 1 \end{pmatrix} \right\}$  forms a basis of  $\mathbb{R}^2$ .

Indeed, geometrically, it can be observed that  $u, v$  are linearly independent,

# An example of a basis of $\mathbb{R}^2$

The set  $\left\{ u = \begin{pmatrix} 1 \\ 2 \end{pmatrix}, v = \begin{pmatrix} 2 \\ 1 \end{pmatrix} \right\}$  forms a basis of  $\mathbb{R}^2$ .

Indeed, geometrically, it can be observed that  $u, v$  are linearly independent, and  $\{u, v\}$  spans  $\mathbb{R}^2$ .

# An example of a basis of $\mathbb{R}^2$

The set  $\left\{ u = \begin{pmatrix} 1 \\ 2 \end{pmatrix}, v = \begin{pmatrix} 2 \\ 1 \end{pmatrix} \right\}$  forms a basis of  $\mathbb{R}^2$ .

Indeed, geometrically, it can be observed that  $u, v$  are linearly independent, and  $\{u, v\}$  spans  $\mathbb{R}^2$ .

Or directly, we see that for EVERY vector  $\begin{pmatrix} a \\ b \end{pmatrix} \in \mathbb{R}^2$ ,

# An example of a basis of $\mathbb{R}^2$

The set  $\left\{ u = \begin{pmatrix} 1 \\ 2 \end{pmatrix}, v = \begin{pmatrix} 2 \\ 1 \end{pmatrix} \right\}$  forms a basis of  $\mathbb{R}^2$ .

Indeed, geometrically, it can be observed that  $u, v$  are linearly independent, and  $\{u, v\}$  spans  $\mathbb{R}^2$ .

Or directly, we see that for EVERY vector  $\begin{pmatrix} a \\ b \end{pmatrix} \in \mathbb{R}^2$ , the system

$$x \begin{pmatrix} 1 \\ 2 \end{pmatrix} + y \begin{pmatrix} 2 \\ 1 \end{pmatrix} = \begin{pmatrix} a \\ b \end{pmatrix} \quad \text{i.e.,} \quad \begin{cases} x + 2y = a \\ 2x + y = b \end{cases} \quad \text{or} \quad \begin{pmatrix} 1 & 2 \\ 2 & 1 \end{pmatrix} \begin{pmatrix} x \\ y \end{pmatrix} = \begin{pmatrix} a \\ b \end{pmatrix}$$

has a UNIQUE solution in  $x, y$



# An example of a basis of $\mathbb{R}^2$

The set  $\left\{ u = \begin{pmatrix} 1 \\ 2 \end{pmatrix}, v = \begin{pmatrix} 2 \\ 1 \end{pmatrix} \right\}$  forms a basis of  $\mathbb{R}^2$ .

Indeed, geometrically, it can be observed that  $u, v$  are linearly independent, and  $\{u, v\}$  spans  $\mathbb{R}^2$ .

Or directly, we see that for EVERY vector  $\begin{pmatrix} a \\ b \end{pmatrix} \in \mathbb{R}^2$ , the system

$$x \begin{pmatrix} 1 \\ 2 \end{pmatrix} + y \begin{pmatrix} 2 \\ 1 \end{pmatrix} = \begin{pmatrix} a \\ b \end{pmatrix} \quad \text{i.e.,} \quad \begin{cases} x + 2y = a \\ 2x + y = b \end{cases} \quad \text{or} \quad \begin{pmatrix} 1 & 2 \\ 2 & 1 \end{pmatrix} \begin{pmatrix} x \\ y \end{pmatrix} = \begin{pmatrix} a \\ b \end{pmatrix}$$

has a UNIQUE solution in  $x, y$  because the coefficient matrix  $\begin{pmatrix} 1 & 2 \\ 2 & 1 \end{pmatrix}$  is invertible.

# An example of a basis of $\mathbb{R}^2$

The set  $\left\{ u = \begin{pmatrix} 1 \\ 2 \end{pmatrix}, v = \begin{pmatrix} 2 \\ 1 \end{pmatrix} \right\}$  forms a basis of  $\mathbb{R}^2$ .

Indeed, geometrically, it can be observed that  $u, v$  are linearly independent, and  $\{u, v\}$  spans  $\mathbb{R}^2$ .

Or directly, we see that for EVERY vector  $\begin{pmatrix} a \\ b \end{pmatrix} \in \mathbb{R}^2$ , the system

$$x \begin{pmatrix} 1 \\ 2 \end{pmatrix} + y \begin{pmatrix} 2 \\ 1 \end{pmatrix} = \begin{pmatrix} a \\ b \end{pmatrix} \quad \text{i.e.,} \quad \begin{cases} x + 2y = a \\ 2x + y = b \end{cases} \quad \text{or} \quad \begin{pmatrix} 1 & 2 \\ 2 & 1 \end{pmatrix} \begin{pmatrix} x \\ y \end{pmatrix} = \begin{pmatrix} a \\ b \end{pmatrix}$$

has a UNIQUE solution in  $x, y$  because the coefficient matrix

$\begin{pmatrix} 1 & 2 \\ 2 & 1 \end{pmatrix}$  is invertible.

So every vector in  $\mathbb{R}^2$  can be written as a linear combination of  $\{u, v\}$ ,

# An example of a basis of $\mathbb{R}^2$

The set  $\left\{ u = \begin{pmatrix} 1 \\ 2 \end{pmatrix}, v = \begin{pmatrix} 2 \\ 1 \end{pmatrix} \right\}$  forms a basis of  $\mathbb{R}^2$ .

Indeed, geometrically, it can be observed that  $u, v$  are linearly independent, and  $\{u, v\}$  spans  $\mathbb{R}^2$ .

Or directly, we see that for EVERY vector  $\begin{pmatrix} a \\ b \end{pmatrix} \in \mathbb{R}^2$ , the system

$$x \begin{pmatrix} 1 \\ 2 \end{pmatrix} + y \begin{pmatrix} 2 \\ 1 \end{pmatrix} = \begin{pmatrix} a \\ b \end{pmatrix} \quad \text{i.e.,} \quad \begin{cases} x + 2y = a \\ 2x + y = b \end{cases} \quad \text{or} \quad \begin{pmatrix} 1 & 2 \\ 2 & 1 \end{pmatrix} \begin{pmatrix} x \\ y \end{pmatrix} = \begin{pmatrix} a \\ b \end{pmatrix}$$

has a UNIQUE solution in  $x, y$  because the coefficient matrix  $\begin{pmatrix} 1 & 2 \\ 2 & 1 \end{pmatrix}$  is invertible.

So every vector in  $\mathbb{R}^2$  can be written as a linear combination of  $\{u, v\}$ , hence it spans the space  $\mathbb{R}^2$ .

# An example of a basis of $\mathbb{R}^2$

The set  $\left\{ u = \begin{pmatrix} 1 \\ 2 \end{pmatrix}, v = \begin{pmatrix} 2 \\ 1 \end{pmatrix} \right\}$  forms a basis of  $\mathbb{R}^2$ .

Indeed, geometrically, it can be observed that  $u, v$  are linearly independent, and  $\{u, v\}$  spans  $\mathbb{R}^2$ .

Or directly, we see that for EVERY vector  $\begin{pmatrix} a \\ b \end{pmatrix} \in \mathbb{R}^2$ , the system

$$x \begin{pmatrix} 1 \\ 2 \end{pmatrix} + y \begin{pmatrix} 2 \\ 1 \end{pmatrix} = \begin{pmatrix} a \\ b \end{pmatrix} \quad \text{i.e.,} \quad \begin{cases} x + 2y = a \\ 2x + y = b \end{cases} \quad \text{or} \quad \begin{pmatrix} 1 & 2 \\ 2 & 1 \end{pmatrix} \begin{pmatrix} x \\ y \end{pmatrix} = \begin{pmatrix} a \\ b \end{pmatrix}$$

has a UNIQUE solution in  $x, y$  because the coefficient matrix  $\begin{pmatrix} 1 & 2 \\ 2 & 1 \end{pmatrix}$  is invertible.

So every vector in  $\mathbb{R}^2$  can be written as a linear combination of  $\{u, v\}$ , hence it spans the space  $\mathbb{R}^2$ .

Moreover, when  $a = b = 0$ , then the system has THE trivial solution  $x = y = 0$ .

# An example of a basis of $\mathbb{R}^2$

The set  $\left\{ u = \begin{pmatrix} 1 \\ 2 \end{pmatrix}, v = \begin{pmatrix} 2 \\ 1 \end{pmatrix} \right\}$  forms a basis of  $\mathbb{R}^2$ .

Indeed, geometrically, it can be observed that  $u, v$  are linearly independent, and  $\{u, v\}$  spans  $\mathbb{R}^2$ .

Or directly, we see that for EVERY vector  $\begin{pmatrix} a \\ b \end{pmatrix} \in \mathbb{R}^2$ , the system

$$x \begin{pmatrix} 1 \\ 2 \end{pmatrix} + y \begin{pmatrix} 2 \\ 1 \end{pmatrix} = \begin{pmatrix} a \\ b \end{pmatrix} \quad \text{i.e.,} \quad \begin{cases} x + 2y = a \\ 2x + y = b \end{cases} \quad \text{or} \quad \begin{pmatrix} 1 & 2 \\ 2 & 1 \end{pmatrix} \begin{pmatrix} x \\ y \end{pmatrix} = \begin{pmatrix} a \\ b \end{pmatrix}$$

has a UNIQUE solution in  $x, y$  because the coefficient matrix  $\begin{pmatrix} 1 & 2 \\ 2 & 1 \end{pmatrix}$  is invertible.

So every vector in  $\mathbb{R}^2$  can be written as a linear combination of  $\{u, v\}$ , hence it spans the space  $\mathbb{R}^2$ .

Moreover, when  $a = b = 0$ , then the system has THE trivial solution  $x = y = 0$ . Thus  $\{u, v\}$  is linearly independent as well.

# Dimension of a vector space

## Theorem

*Let  $V$  be a vector space (over  $\mathbb{R}$ ). Then*

# Dimension of a vector space

## Theorem

*Let  $V$  be a vector space (over  $\mathbb{R}$ ). Then*

- 1  *$V$  has a basis, and*

# Dimension of a vector space

## Theorem

*Let  $V$  be a vector space (over  $\mathbb{R}$ ). Then*

- ①  $V$  has a basis, and*
- ② all the bases of  $V$  have the same cardinality,*



# Dimension of a vector space

## Theorem

*Let  $V$  be a vector space (over  $\mathbb{R}$ ). Then*

- ①  *$V$  has a basis, and*
- ② *all the bases of  $V$  have the same cardinality, i.e., if  $S_1$  and  $S_2$  are any two bases of  $V$ ,*

# Dimension of a vector space

## Theorem

*Let  $V$  be a vector space (over  $\mathbb{R}$ ). Then*

- ①  *$V$  has a basis, and*
- ② *all the bases of  $V$  have the same cardinality, i.e., if  $S_1$  and  $S_2$  are any two bases of  $V$ , then there exists a bijection from  $S_1$  to  $S_2$ .*

# Dimension of a vector space

## Theorem

*Let  $V$  be a vector space (over  $\mathbb{R}$ ). Then*

- ①  *$V$  has a basis, and*
- ② *all the bases of  $V$  have the same cardinality, i.e., if  $S_1$  and  $S_2$  are any two bases of  $V$ , then there exists a bijection from  $S_1$  to  $S_2$ . Particularly, if  $S_1$  and  $S_2$  are finite, then they have the same number of vectors.*

# Dimension of a vector space

## Theorem

*Let  $V$  be a vector space (over  $\mathbb{R}$ ). Then*

- ①  *$V$  has a basis, and*
- ② *all the bases of  $V$  have the same cardinality, i.e., if  $S_1$  and  $S_2$  are any two bases of  $V$ , then there exists a bijection from  $S_1$  to  $S_2$ . Particularly, if  $S_1$  and  $S_2$  are finite, then they have the same number of vectors.*

## Definition

- ① The cardinality of a basis of the vector space  $V$  is called the **dimension** of  $V$ .

# Dimension of a vector space

## Theorem

*Let  $V$  be a vector space (over  $\mathbb{R}$ ). Then*

- 1  *$V$  has a basis, and*
- 2 *all the bases of  $V$  have the same cardinality, i.e., if  $S_1$  and  $S_2$  are any two bases of  $V$ , then there exists a bijection from  $S_1$  to  $S_2$ . Particularly, if  $S_1$  and  $S_2$  are finite, then they have the same number of vectors.*

## Definition

- 1 The cardinality of a basis of the vector space  $V$  is called the **dimension** of  $V$ .
- 2 The dimension of  $V$  is denoted by  $\dim(V)$ .

# Standard basis of $\mathbb{R}^n$

① In  $\mathbb{R}^n$ ,

# Standard basis of $\mathbb{R}^n$

① In  $\mathbb{R}^n$ , let  $S$  be the subset consisting of the vectors:

$$e_1 = (1, 0, 0, \dots, 0)$$

$$e_2 = (0, 1, 0, \dots, 0)$$

$$\vdots$$

$$e_n = (0, 0, 0, \dots, 1).$$

# Standard basis of $\mathbb{R}^n$

- ① In  $\mathbb{R}^n$ , let  $S$  be the subset consisting of the vectors:

$$e_1 = (1, 0, 0, \dots, 0)$$

$$e_2 = (0, 1, 0, \dots, 0)$$

$$\vdots$$

$$e_n = (0, 0, 0, \dots, 1).$$

- ② Note that any vector  $v = (x_1, \dots, x_n) \in \mathbb{R}^n$  can be written as a linear combination  $x_1 e_1 + \dots + x_n e_n$ .



# Standard basis of $\mathbb{R}^n$

- ① In  $\mathbb{R}^n$ , let  $S$  be the subset consisting of the vectors:

$$e_1 = (1, 0, 0, \dots, 0)$$

$$e_2 = (0, 1, 0, \dots, 0)$$

$$\vdots$$

$$e_n = (0, 0, 0, \dots, 1).$$

- ② Note that any vector  $v = (x_1, \dots, x_n) \in \mathbb{R}^n$  can be written as a linear combination  $x_1 e_1 + \dots + x_n e_n$ . So  $S$  spans  $\mathbb{R}^n$ .

# Standard basis of $\mathbb{R}^n$

- ① In  $\mathbb{R}^n$ , let  $S$  be the subset consisting of the vectors:

$$e_1 = (1, 0, 0, \dots, 0)$$

$$e_2 = (0, 1, 0, \dots, 0)$$

$$\vdots$$

$$e_n = (0, 0, 0, \dots, 1).$$

- ② Note that any vector  $v = (x_1, \dots, x_n) \in \mathbb{R}^n$  can be written as a linear combination  $x_1 e_1 + \dots + x_n e_n$ . So  $S$  spans  $\mathbb{R}^n$ .
- ③ Is  $S$  linearly independent?

# Standard basis of $\mathbb{R}^n$

- ① In  $\mathbb{R}^n$ , let  $S$  be the subset consisting of the vectors:

$$e_1 = (1, 0, 0, \dots, 0)$$

$$e_2 = (0, 1, 0, \dots, 0)$$

$$\vdots$$

$$e_n = (0, 0, 0, \dots, 1).$$

- ② Note that any vector  $v = (x_1, \dots, x_n) \in \mathbb{R}^n$  can be written as a linear combination  $x_1 e_1 + \dots + x_n e_n$ . So  $S$  spans  $\mathbb{R}^n$ .
- ③ Is  $S$  linearly independent? Answer: Yes.

# Standard basis of $\mathbb{R}^n$

- ① In  $\mathbb{R}^n$ , let  $S$  be the subset consisting of the vectors:

$$e_1 = (1, 0, 0, \dots, 0)$$

$$e_2 = (0, 1, 0, \dots, 0)$$

$$\vdots$$

$$e_n = (0, 0, 0, \dots, 1).$$

- ② Note that any vector  $v = (x_1, \dots, x_n) \in \mathbb{R}^n$  can be written as a linear combination  $x_1 e_1 + \dots + x_n e_n$ . So  $S$  spans  $\mathbb{R}^n$ .
- ③ Is  $S$  linearly independent? Answer: Yes. Why?

# Standard basis of $\mathbb{R}^n$

- ① In  $\mathbb{R}^n$ , let  $S$  be the subset consisting of the vectors:

$$e_1 = (1, 0, 0, \dots, 0)$$

$$e_2 = (0, 1, 0, \dots, 0)$$

$$\vdots$$

$$e_n = (0, 0, 0, \dots, 1).$$

- ② Note that any vector  $v = (x_1, \dots, x_n) \in \mathbb{R}^n$  can be written as a linear combination  $x_1 e_1 + \dots + x_n e_n$ . So  $S$  spans  $\mathbb{R}^n$ .
- ③ Is  $S$  linearly independent? Answer: Yes. Why?  
(Because  $x_1 e_1 + \dots + x_n e_n = 0$

# Standard basis of $\mathbb{R}^n$

- ① In  $\mathbb{R}^n$ , let  $S$  be the subset consisting of the vectors:

$$e_1 = (1, 0, 0, \dots, 0)$$

$$e_2 = (0, 1, 0, \dots, 0)$$

$$\vdots$$

$$e_n = (0, 0, 0, \dots, 1).$$

- ② Note that any vector  $v = (x_1, \dots, x_n) \in \mathbb{R}^n$  can be written as a linear combination  $x_1 e_1 + \dots + x_n e_n$ . So  $S$  spans  $\mathbb{R}^n$ .
- ③ Is  $S$  linearly independent? Answer: Yes. Why?  
(Because  $x_1 e_1 + \dots + x_n e_n = 0 \implies$  every  $x_i = 0$ .)

# Standard basis of $\mathbb{R}^n$

- ① In  $\mathbb{R}^n$ , let  $S$  be the subset consisting of the vectors:

$$e_1 = (1, 0, 0, \dots, 0)$$

$$e_2 = (0, 1, 0, \dots, 0)$$

$$\vdots$$

$$e_n = (0, 0, 0, \dots, 1).$$

- ② Note that any vector  $v = (x_1, \dots, x_n) \in \mathbb{R}^n$  can be written as a linear combination  $x_1 e_1 + \dots + x_n e_n$ . So  $S$  spans  $\mathbb{R}^n$ .
- ③ Is  $S$  linearly independent? Answer: Yes. Why?  
(Because  $x_1 e_1 + \dots + x_n e_n = 0 \implies$  every  $x_i = 0$ .)
- ④ Therefore  $S$  is a basis of  $\mathbb{R}^n$ .

# Standard basis of $\mathbb{R}^n$

- ① In  $\mathbb{R}^n$ , let  $S$  be the subset consisting of the vectors:

$$e_1 = (1, 0, 0, \dots, 0)$$

$$e_2 = (0, 1, 0, \dots, 0)$$

$$\vdots$$

$$e_n = (0, 0, 0, \dots, 1).$$

- ② Note that any vector  $v = (x_1, \dots, x_n) \in \mathbb{R}^n$  can be written as a linear combination  $x_1 e_1 + \dots + x_n e_n$ . So  $S$  spans  $\mathbb{R}^n$ .
- ③ Is  $S$  linearly independent? Answer: Yes. Why?  
(Because  $x_1 e_1 + \dots + x_n e_n = 0 \implies$  every  $x_i = 0$ .)
- ④ Therefore  $S$  is a basis of  $\mathbb{R}^n$ .
- ⑤ This particular basis is called the **standard basis** of  $\mathbb{R}^n$ .



# Standard basis of $\mathbb{R}^n$

- ① In  $\mathbb{R}^n$ , let  $S$  be the subset consisting of the vectors:

$$e_1 = (1, 0, 0, \dots, 0)$$

$$e_2 = (0, 1, 0, \dots, 0)$$

$$\vdots$$

$$e_n = (0, 0, 0, \dots, 1).$$

- ② Note that any vector  $v = (x_1, \dots, x_n) \in \mathbb{R}^n$  can be written as a linear combination  $x_1 e_1 + \dots + x_n e_n$ . So  $S$  spans  $\mathbb{R}^n$ .
- ③ Is  $S$  linearly independent? Answer: Yes. Why?  
(Because  $x_1 e_1 + \dots + x_n e_n = 0 \implies$  every  $x_i = 0$ .)
- ④ Therefore  $S$  is a basis of  $\mathbb{R}^n$ .
- ⑤ This particular basis is called the **standard basis** of  $\mathbb{R}^n$ .
- ⑥ Thus  $\dim(\mathbb{R}^n)$

# Standard basis of $\mathbb{R}^n$

- ① In  $\mathbb{R}^n$ , let  $S$  be the subset consisting of the vectors:

$$e_1 = (1, 0, 0, \dots, 0)$$

$$e_2 = (0, 1, 0, \dots, 0)$$

$$\vdots$$

$$e_n = (0, 0, 0, \dots, 1).$$

- ② Note that any vector  $v = (x_1, \dots, x_n) \in \mathbb{R}^n$  can be written as a linear combination  $x_1 e_1 + \dots + x_n e_n$ . So  $S$  spans  $\mathbb{R}^n$ .
- ③ Is  $S$  linearly independent? Answer: Yes. Why?  
(Because  $x_1 e_1 + \dots + x_n e_n = 0 \implies$  every  $x_i = 0$ .)
- ④ Therefore  $S$  is a basis of  $\mathbb{R}^n$ .
- ⑤ This particular basis is called the **standard basis** of  $\mathbb{R}^n$ .
- ⑥ Thus  $\dim(\mathbb{R}^n) = n$ .

# Basis of $\mathbb{R}[x]$

For  $\mathbb{R}[x]$ , the set of all polynomials over  $\mathbb{R}$ ,

For  $\mathbb{R}[x]$ , the set of all polynomials over  $\mathbb{R}$ , the subset

$$S = \{x^n : n = 0, 1, 2, \dots\}$$

For  $\mathbb{R}[x]$ , the set of all polynomials over  $\mathbb{R}$ , the subset

$$S = \{x^n : n = 0, 1, 2, \dots\}$$

forms a basis.

For  $\mathbb{R}[x]$ , the set of all polynomials over  $\mathbb{R}$ , the subset

$$S = \{x^n : n = 0, 1, 2, \dots\}$$

forms a basis. Thus  $\dim(\mathbb{R}[x])$

For  $\mathbb{R}[x]$ , the set of all polynomials over  $\mathbb{R}$ , the subset

$$S = \{x^n : n = 0, 1, 2, \dots\}$$

forms a basis. Thus  $\dim(\mathbb{R}[x]) = \infty$ .

# Theorems related to the dimension of a vector space

- In the rest of the slides, it is proved that if  $V$  is a finite dimensional vector space, then any two bases of  $V$  have the same number of elements.



# Theorems related to the dimension of a vector space

- In the rest of the slides, it is proved that if  $V$  is a finite dimensional vector space, then any two bases of  $V$  have the same number of elements.
- Interested students can read it on their own

# On spanning set of $V$

## Lemma

*Suppose  $\{v_1, v_2, \dots, v_n\}$  spans  $V$ .*

# On spanning set of $V$

## Lemma

*Suppose  $\{v_1, v_2, \dots, v_n\}$  spans  $V$ . Let  $u \neq 0$  is a vector in  $V$ .*

# On spanning set of $V$

## Lemma

*Suppose  $\{v_1, v_2, \dots, v_n\}$  spans  $V$ . Let  $u \neq 0$  is a vector in  $V$ . Then some  $v_i$  can be replaced by  $u$  to get another spanning set of  $V$ ,*

# On spanning set of $V$

## Lemma

*Suppose  $\{v_1, v_2, \dots, v_n\}$  spans  $V$ . Let  $u \neq 0$  is a vector in  $V$ . Then some  $v_i$  can be replaced by  $u$  to get another spanning set of  $V$ , i.e., if necessary, then after renaming the vectors  $\{v_1, v_2, \dots, v_n\}$ ,*

# On spanning set of $V$

## Lemma

*Suppose  $\{v_1, v_2, \dots, v_n\}$  spans  $V$ . Let  $u \neq 0$  is a vector in  $V$ . Then some  $v_i$  can be replaced by  $u$  to get another spanning set of  $V$ , i.e., if necessary, then after renaming the vectors  $\{v_1, v_2, \dots, v_n\}$ , we obtain that  $\{u, v_2, \dots, v_n\}$  spans  $V$ .*

# On spanning set of $V$

## Lemma

*Suppose  $\{v_1, v_2, \dots, v_n\}$  spans  $V$ . Let  $u \neq 0$  is a vector in  $V$ . Then some  $v_i$  can be replaced by  $u$  to get another spanning set of  $V$ , i.e., if necessary, then after renaming the vectors  $\{v_1, v_2, \dots, v_n\}$ , we obtain that  $\{u, v_2, \dots, v_n\}$  spans  $V$ .*

*Proof.* Since  $\{v_1, v_2, \dots, v_n\}$  spans  $V$ ,

# On spanning set of $V$

## Lemma

*Suppose  $\{v_1, v_2, \dots, v_n\}$  spans  $V$ . Let  $u \neq 0$  is a vector in  $V$ . Then some  $v_i$  can be replaced by  $u$  to get another spanning set of  $V$ , i.e., if necessary, then after renaming the vectors  $\{v_1, v_2, \dots, v_n\}$ , we obtain that  $\{u, v_2, \dots, v_n\}$  spans  $V$ .*

*Proof.* Since  $\{v_1, v_2, \dots, v_n\}$  spans  $V$ ,  $u$  can be written as

$$u = c_1 v_1 + c_2 v_2 + \dots + c_n v_n \text{ for some } c_1, \dots, c_n \in \mathbb{R}. \quad (1)$$



# On spanning set of $V$

## Lemma

*Suppose  $\{v_1, v_2, \dots, v_n\}$  spans  $V$ . Let  $u \neq 0$  is a vector in  $V$ . Then some  $v_i$  can be replaced by  $u$  to get another spanning set of  $V$ , i.e., if necessary, then after renaming the vectors  $\{v_1, v_2, \dots, v_n\}$ , we obtain that  $\{u, v_2, \dots, v_n\}$  spans  $V$ .*

*Proof.* Since  $\{v_1, v_2, \dots, v_n\}$  spans  $V$ ,  $u$  can be written as

$$u = c_1 v_1 + c_2 v_2 + \dots + c_n v_n \text{ for some } c_1, \dots, c_n \in \mathbb{R}. \quad (1)$$

Since  $u \neq 0$ , at least one  $c_i \neq 0$ .

# On spanning set of $V$

## Lemma

*Suppose  $\{v_1, v_2, \dots, v_n\}$  spans  $V$ . Let  $u \neq 0$  is a vector in  $V$ . Then some  $v_i$  can be replaced by  $u$  to get another spanning set of  $V$ , i.e., if necessary, then after renaming the vectors  $\{v_1, v_2, \dots, v_n\}$ , we obtain that  $\{u, v_2, \dots, v_n\}$  spans  $V$ .*

*Proof.* Since  $\{v_1, v_2, \dots, v_n\}$  spans  $V$ ,  $u$  can be written as

$$u = c_1 v_1 + c_2 v_2 + \dots + c_n v_n \text{ for some } c_1, \dots, c_n \in \mathbb{R}. \quad (1)$$

Since  $u \neq 0$ , at least one  $c_i \neq 0$ . So (1) yields that  $v_i =$

$$(1/c_i)u + (c_1/c_i)v_1 + \dots + (c_{i-1}/c_i)v_{i-1} + (c_{i+1}/c_i)v_{i+1} + \dots + (c_n/c_i)v_n.$$

# On spanning set of $V$

## Lemma

*Suppose  $\{v_1, v_2, \dots, v_n\}$  spans  $V$ . Let  $u \neq 0$  is a vector in  $V$ . Then some  $v_i$  can be replaced by  $u$  to get another spanning set of  $V$ , i.e., if necessary, then after renaming the vectors  $\{v_1, v_2, \dots, v_n\}$ , we obtain that  $\{u, v_2, \dots, v_n\}$  spans  $V$ .*

*Proof.* Since  $\{v_1, v_2, \dots, v_n\}$  spans  $V$ ,  $u$  can be written as

$$u = c_1 v_1 + c_2 v_2 + \dots + c_n v_n \text{ for some } c_1, \dots, c_n \in \mathbb{R}. \quad (1)$$

Since  $u \neq 0$ , at least one  $c_i \neq 0$ . So (1) yields that  $v_i =$

$$(1/c_i)u + (c_1/c_i)v_1 + \dots + (c_{i-1}/c_i)v_{i-1} + (c_{i+1}/c_i)v_{i+1} + \dots + (c_n/c_i)v_n.$$

Since every  $v \in V$  is a linear combination of  $v_1, v_2, \dots, v_n$ , using the expression of  $v_i$  in that linear combination,

# On spanning set of $V$

## Lemma

*Suppose  $\{v_1, v_2, \dots, v_n\}$  spans  $V$ . Let  $u \neq 0$  is a vector in  $V$ . Then some  $v_i$  can be replaced by  $u$  to get another spanning set of  $V$ , i.e., if necessary, then after renaming the vectors  $\{v_1, v_2, \dots, v_n\}$ , we obtain that  $\{u, v_2, \dots, v_n\}$  spans  $V$ .*

*Proof.* Since  $\{v_1, v_2, \dots, v_n\}$  spans  $V$ ,  $u$  can be written as

$$u = c_1 v_1 + c_2 v_2 + \dots + c_n v_n \text{ for some } c_1, \dots, c_n \in \mathbb{R}. \quad (1)$$

Since  $u \neq 0$ , at least one  $c_i \neq 0$ . So (1) yields that  $v_i =$

$$(1/c_i)u + (c_1/c_i)v_1 + \dots + (c_{i-1}/c_i)v_{i-1} + (c_{i+1}/c_i)v_{i+1} + \dots + (c_n/c_i)v_n.$$

Since every  $v \in V$  is a linear combination of  $v_1, v_2, \dots, v_n$ , using the expression of  $v_i$  in that linear combination, it follows that  $v$  can be written as a linear combination of  $v_1, \dots, v_{i-1}, u, v_{i+1}, \dots, v_n$ .

# On spanning set of $V$

## Lemma

*Suppose  $\{v_1, v_2, \dots, v_n\}$  spans  $V$ . Let  $u \neq 0$  is a vector in  $V$ . Then some  $v_i$  can be replaced by  $u$  to get another spanning set of  $V$ , i.e., if necessary, then after renaming the vectors  $\{v_1, v_2, \dots, v_n\}$ , we obtain that  $\{u, v_2, \dots, v_n\}$  spans  $V$ .*

*Proof.* Since  $\{v_1, v_2, \dots, v_n\}$  spans  $V$ ,  $u$  can be written as

$$u = c_1 v_1 + c_2 v_2 + \dots + c_n v_n \text{ for some } c_1, \dots, c_n \in \mathbb{R}. \quad (1)$$

Since  $u \neq 0$ , at least one  $c_i \neq 0$ . So (1) yields that  $v_i =$

$$(1/c_i)u + (c_1/c_i)v_1 + \dots + (c_{i-1}/c_i)v_{i-1} + (c_{i+1}/c_i)v_{i+1} + \dots + (c_n/c_i)v_n.$$

Since every  $v \in V$  is a linear combination of  $v_1, v_2, \dots, v_n$ , using the expression of  $v_i$  in that linear combination, it follows that  $v$  can be written as a linear combination of  $v_1, \dots, v_{i-1}, u, v_{i+1}, \dots, v_n$ .

Thus  $\{v_1, \dots, v_{i-1}, u, v_{i+1}, \dots, v_n\}$  spans  $V$ .



# Linearly independent set has less vectors than spanning set

## Theorem

*Suppose  $V = \text{Span}\{v_1, v_2, \dots, v_n\}$ ,*

# Linearly independent set has less vectors than spanning set

## Theorem

*Suppose  $V = \text{Span}\{v_1, v_2, \dots, v_n\}$ , and  $\{u_1, u_2, \dots, u_m\}$  is a linearly independent subset of  $V$ .*

# Linearly independent set has less vectors than spanning set

## Theorem

*Suppose  $V = \text{Span}\{v_1, v_2, \dots, v_n\}$ , and  $\{u_1, u_2, \dots, u_m\}$  is a linearly independent subset of  $V$ . Then  $m \leq n$ .*



# Linearly independent set has less vectors than spanning set

## Theorem

*Suppose  $V = \text{Span}\{v_1, v_2, \dots, v_n\}$ , and  $\{u_1, u_2, \dots, u_m\}$  is a linearly independent subset of  $V$ . Then  $m \leq n$ .*

*Proof.* If possible, let  $n < m$ .

# Linearly independent set has less vectors than spanning set

## Theorem

*Suppose  $V = \text{Span}\{v_1, v_2, \dots, v_n\}$ , and  $\{u_1, u_2, \dots, u_m\}$  is a linearly independent subset of  $V$ . Then  $m \leq n$ .*

*Proof.* If possible, let  $n < m$ . Note that  $u_i \neq 0$ .

# Linearly independent set has less vectors than spanning set

## Theorem

*Suppose  $V = \text{Span}\{v_1, v_2, \dots, v_n\}$ , and  $\{u_1, u_2, \dots, u_m\}$  is a linearly independent subset of  $V$ . Then  $m \leq n$ .*

*Proof.* If possible, let  $n < m$ . Note that  $u_i \neq 0$ . So, by renaming the vectors  $v_1, \dots, v_n$ , we have  $\{u_1, v_2, v_3, \dots, v_n\}$  spans  $V$ .

# Linearly independent set has less vectors than spanning set

## Theorem

*Suppose  $V = \text{Span}\{v_1, v_2, \dots, v_n\}$ , and  $\{u_1, u_2, \dots, u_m\}$  is a linearly independent subset of  $V$ . Then  $m \leq n$ .*

*Proof.* If possible, let  $n < m$ . Note that  $u_i \neq 0$ . So, by renaming the vectors  $v_1, \dots, v_n$ , we have  $\{u_1, v_2, v_3, \dots, v_n\}$  spans  $V$ . In the 2nd step, since  $u_2 \in V = \text{Span}\{u_1, v_2, v_3, \dots, v_n\}$ ,

# Linearly independent set has less vectors than spanning set

## Theorem

*Suppose  $V = \text{Span}\{v_1, v_2, \dots, v_n\}$ , and  $\{u_1, u_2, \dots, u_m\}$  is a linearly independent subset of  $V$ . Then  $m \leq n$ .*

*Proof.* If possible, let  $n < m$ . Note that  $u_i \neq 0$ . So, by renaming the vectors  $v_1, \dots, v_n$ , we have  $\{u_1, v_2, v_3, \dots, v_n\}$  spans  $V$ .

In the 2nd step, since  $u_2 \in V = \text{Span}\{u_1, v_2, v_3, \dots, v_n\}$ ,

$$u_2 = b_1 u_1 + b_2 v_2 + b_3 v_3 + \dots + b_n v_n \quad \text{for some } b_i \in \mathbb{R}.$$

# Linearly independent set has less vectors than spanning set

## Theorem

*Suppose  $V = \text{Span}\{v_1, v_2, \dots, v_n\}$ , and  $\{u_1, u_2, \dots, u_m\}$  is a linearly independent subset of  $V$ . Then  $m \leq n$ .*

*Proof.* If possible, let  $n < m$ . Note that  $u_i \neq 0$ . So, by renaming the vectors  $v_1, \dots, v_n$ , we have  $\{u_1, v_2, v_3, \dots, v_n\}$  spans  $V$ .

In the 2nd step, since  $u_2 \in V = \text{Span}\{u_1, v_2, v_3, \dots, v_n\}$ ,

$$u_2 = b_1 u_1 + b_2 v_2 + b_3 v_3 + \dots + b_n v_n \quad \text{for some } b_i \in \mathbb{R}.$$

Then at least one of  $\{b_2, \dots, b_n\}$  is non-zero.

# Linearly independent set has less vectors than spanning set

## Theorem

*Suppose  $V = \text{Span}\{v_1, v_2, \dots, v_n\}$ , and  $\{u_1, u_2, \dots, u_m\}$  is a linearly independent subset of  $V$ . Then  $m \leq n$ .*

*Proof.* If possible, let  $n < m$ . Note that  $u_i \neq 0$ . So, by renaming the vectors  $v_1, \dots, v_n$ , we have  $\{u_1, v_2, v_3, \dots, v_n\}$  spans  $V$ .

In the 2nd step, since  $u_2 \in V = \text{Span}\{u_1, v_2, v_3, \dots, v_n\}$ ,

$$u_2 = b_1 u_1 + b_2 v_2 + b_3 v_3 + \dots + b_n v_n \quad \text{for some } b_i \in \mathbb{R}.$$

Then at least one of  $\{b_2, \dots, b_n\}$  is non-zero.

Hence, if necessary, by renaming the vectors  $v_2, \dots, v_n$ , we have that  $\{u_1, u_2, v_3, \dots, v_n\}$  spans  $V$ .

# Linearly independent set has less vectors than spanning set

## Theorem

*Suppose  $V = \text{Span}\{v_1, v_2, \dots, v_n\}$ , and  $\{u_1, u_2, \dots, u_m\}$  is a linearly independent subset of  $V$ . Then  $m \leq n$ .*

*Proof.* If possible, let  $n < m$ . Note that  $u_i \neq 0$ . So, by renaming the vectors  $v_1, \dots, v_n$ , we have  $\{u_1, v_2, v_3, \dots, v_n\}$  spans  $V$ .

In the 2nd step, since  $u_2 \in V = \text{Span}\{u_1, v_2, v_3, \dots, v_n\}$ ,

$$u_2 = b_1 u_1 + b_2 v_2 + b_3 v_3 + \dots + b_n v_n \quad \text{for some } b_i \in \mathbb{R}.$$

Then at least one of  $\{b_2, \dots, b_n\}$  is non-zero.

Hence, if necessary, by renaming the vectors  $v_2, \dots, v_n$ , we have that  $\{u_1, u_2, v_3, \dots, v_n\}$  spans  $V$ .

Continuing in this way, after  $n$  steps, we obtain that  $\{u_1, u_2, \dots, u_n\}$  spans  $V$ .



# Linearly independent set has less vectors than spanning set

## Theorem

*Suppose  $V = \text{Span}\{v_1, v_2, \dots, v_n\}$ , and  $\{u_1, u_2, \dots, u_m\}$  is a linearly independent subset of  $V$ . Then  $m \leq n$ .*

*Proof.* If possible, let  $n < m$ . Note that  $u_i \neq 0$ . So, by renaming the vectors  $v_1, \dots, v_n$ , we have  $\{u_1, v_2, v_3, \dots, v_n\}$  spans  $V$ .

In the 2nd step, since  $u_2 \in V = \text{Span}\{u_1, v_2, v_3, \dots, v_n\}$ ,

$$u_2 = b_1 u_1 + b_2 v_2 + b_3 v_3 + \dots + b_n v_n \quad \text{for some } b_i \in \mathbb{R}.$$

Then at least one of  $\{b_2, \dots, b_n\}$  is non-zero.

Hence, if necessary, by renaming the vectors  $v_2, \dots, v_n$ , we have that  $\{u_1, u_2, v_3, \dots, v_n\}$  spans  $V$ .

Continuing in this way, after  $n$  steps, we obtain that  $\{u_1, u_2, \dots, u_n\}$  spans  $V$ . Hence

$$u_{n+1} \in V = \text{Span}\{u_1, u_2, \dots, u_n\}.$$

# Linearly independent set has less vectors than spanning set

## Theorem

*Suppose  $V = \text{Span}\{v_1, v_2, \dots, v_n\}$ , and  $\{u_1, u_2, \dots, u_m\}$  is a linearly independent subset of  $V$ . Then  $m \leq n$ .*

*Proof.* If possible, let  $n < m$ . Note that  $u_i \neq 0$ . So, by renaming the vectors  $v_1, \dots, v_n$ , we have  $\{u_1, v_2, v_3, \dots, v_n\}$  spans  $V$ .

In the 2nd step, since  $u_2 \in V = \text{Span}\{u_1, v_2, v_3, \dots, v_n\}$ ,

$$u_2 = b_1 u_1 + b_2 v_2 + b_3 v_3 + \dots + b_n v_n \quad \text{for some } b_i \in \mathbb{R}.$$

Then at least one of  $\{b_2, \dots, b_n\}$  is non-zero.

Hence, if necessary, by renaming the vectors  $v_2, \dots, v_n$ , we have that  $\{u_1, u_2, v_3, \dots, v_n\}$  spans  $V$ .

Continuing in this way, after  $n$  steps, we obtain that  $\{u_1, u_2, \dots, u_n\}$  spans  $V$ . Hence

$$u_{n+1} \in V = \text{Span}\{u_1, u_2, \dots, u_n\}.$$

Therefore  $\{u_1, u_2, \dots, u_{n+1}\}$  is linearly dependent,

# Linearly independent set has less vectors than spanning set

## Theorem

*Suppose  $V = \text{Span}\{v_1, v_2, \dots, v_n\}$ , and  $\{u_1, u_2, \dots, u_m\}$  is a linearly independent subset of  $V$ . Then  $m \leq n$ .*

*Proof.* If possible, let  $n < m$ . Note that  $u_i \neq 0$ . So, by renaming the vectors  $v_1, \dots, v_n$ , we have  $\{u_1, v_2, v_3, \dots, v_n\}$  spans  $V$ .

In the 2nd step, since  $u_2 \in V = \text{Span}\{u_1, v_2, v_3, \dots, v_n\}$ ,

$$u_2 = b_1 u_1 + b_2 v_2 + b_3 v_3 + \dots + b_n v_n \quad \text{for some } b_i \in \mathbb{R}.$$

Then at least one of  $\{b_2, \dots, b_n\}$  is non-zero.

Hence, if necessary, by renaming the vectors  $v_2, \dots, v_n$ , we have that  $\{u_1, u_2, v_3, \dots, v_n\}$  spans  $V$ .

Continuing in this way, after  $n$  steps, we obtain that  $\{u_1, u_2, \dots, u_n\}$  spans  $V$ . Hence

$$u_{n+1} \in V = \text{Span}\{u_1, u_2, \dots, u_n\}.$$

Therefore  $\{u_1, u_2, \dots, u_{n+1}\}$  is linearly dependent, which contradicts the assumption.

# Linearly independent set has less vectors than spanning set

## Theorem

*Suppose  $V = \text{Span}\{v_1, v_2, \dots, v_n\}$ , and  $\{u_1, u_2, \dots, u_m\}$  is a linearly independent subset of  $V$ . Then  $m \leq n$ .*

*Proof.* If possible, let  $n < m$ . Note that  $u_i \neq 0$ . So, by renaming the vectors  $v_1, \dots, v_n$ , we have  $\{u_1, v_2, v_3, \dots, v_n\}$  spans  $V$ .

In the 2nd step, since  $u_2 \in V = \text{Span}\{u_1, v_2, v_3, \dots, v_n\}$ ,

$$u_2 = b_1 u_1 + b_2 v_2 + b_3 v_3 + \dots + b_n v_n \quad \text{for some } b_i \in \mathbb{R}.$$

Then at least one of  $\{b_2, \dots, b_n\}$  is non-zero.

Hence, if necessary, by renaming the vectors  $v_2, \dots, v_n$ , we have that  $\{u_1, u_2, v_3, \dots, v_n\}$  spans  $V$ .

Continuing in this way, after  $n$  steps, we obtain that  $\{u_1, u_2, \dots, u_n\}$  spans  $V$ . Hence

$$u_{n+1} \in V = \text{Span}\{u_1, u_2, \dots, u_n\}.$$

Therefore  $\{u_1, u_2, \dots, u_{n+1}\}$  is linearly dependent, which contradicts the assumption. Hence  $m \leq n$ .

# Any two bases of $V$ have the same number of elements

## Corollary

*If  $V$  is a finite dimensional vector space,*

# Any two bases of $V$ have the same number of elements

## Corollary

*If  $V$  is a finite dimensional vector space, then any two bases of  $V$  have the same number of elements.*

# Any two bases of $V$ have the same number of elements

## Corollary

*If  $V$  is a finite dimensional vector space, then any two bases of  $V$  have the same number of elements.*

*Proof.* Since  $V$  is finite dimensional, it has a finite basis  $\{v_1, \dots, v_n\}$ .

# Any two bases of $V$ have the same number of elements

## Corollary

*If  $V$  is a finite dimensional vector space, then any two bases of  $V$  have the same number of elements.*

*Proof.* Since  $V$  is finite dimensional, it has a finite basis  $\{v_1, \dots, v_n\}$ .

If  $\{u_1, \dots, u_m\}$  is another basis of  $V$ ,



# Any two bases of $V$ have the same number of elements

## Corollary

*If  $V$  is a finite dimensional vector space, then any two bases of  $V$  have the same number of elements.*

*Proof.* Since  $V$  is finite dimensional, it has a finite basis  $\{v_1, \dots, v_n\}$ .

If  $\{u_1, \dots, u_m\}$  is another basis of  $V$ , then by the last theorem,  $m \leq n$ .

# Any two bases of $V$ have the same number of elements

## Corollary

*If  $V$  is a finite dimensional vector space, then any two bases of  $V$  have the same number of elements.*

*Proof.* Since  $V$  is finite dimensional, it has a finite basis  $\{v_1, \dots, v_n\}$ .

If  $\{u_1, \dots, u_m\}$  is another basis of  $V$ , then by the last theorem,  $m \leq n$ .

By the same argument,  $n \leq m$ .

# Any two bases of $V$ have the same number of elements

## Corollary

*If  $V$  is a finite dimensional vector space, then any two bases of  $V$  have the same number of elements.*

*Proof.* Since  $V$  is finite dimensional, it has a finite basis  $\{v_1, \dots, v_n\}$ .

If  $\{u_1, \dots, u_m\}$  is another basis of  $V$ , then by the last theorem,  $m \leq n$ .

By the same argument,  $n \leq m$ . Thus  $m = n$ . □

# Any two bases of $V$ have the same number of elements

## Corollary

*If  $V$  is a finite dimensional vector space, then any two bases of  $V$  have the same number of elements.*

*Proof.* Since  $V$  is finite dimensional, it has a finite basis  $\{v_1, \dots, v_n\}$ .

If  $\{u_1, \dots, u_m\}$  is another basis of  $V$ , then by the last theorem,  $m \leq n$ .

By the same argument,  $n \leq m$ . Thus  $m = n$ . □

# Consequences of the fact that a linearly independent set has less or equal number of vectors than a spanning set

## Corollary

*Let  $V$  be a finite dimensional vector space,*

# Consequences of the fact that a linearly independent set has less or equal number of vectors than a spanning set

## Corollary

*Let  $V$  be a finite dimensional vector space,  $d = \dim(V)$ .*

# Consequences of the fact that a linearly independent set has less or equal number of vectors than a spanning set

## Corollary

*Let  $V$  be a finite dimensional vector space,  $d = \dim(V)$ . Then*

- 1 any subset of  $V$  containing more than  $d$  vectors is linearly dependent.*

# Consequences of the fact that a linearly independent set has less or equal number of vectors than a spanning set

## Corollary

*Let  $V$  be a finite dimensional vector space,  $d = \dim(V)$ . Then*

- 1 any subset of  $V$  containing more than  $d$  vectors is linearly dependent. Thus a basis of  $V$  is a **maximal linearly independent subset** of  $V$ .*



# Consequences of the fact that a linearly independent set has less or equal number of vectors than a spanning set

## Corollary

*Let  $V$  be a finite dimensional vector space,  $d = \dim(V)$ . Then*

- ① any subset of  $V$  containing more than  $d$  vectors is linearly dependent. Thus a basis of  $V$  is a **maximal linearly independent subset** of  $V$ .*
- ② A subset of  $V$  containing fewer than  $d$  vectors cannot span  $V$ .*

# Consequences of the fact that a linearly independent set has less or equal number of vectors than a spanning set

## Corollary

*Let  $V$  be a finite dimensional vector space,  $d = \dim(V)$ . Then*

- ① any subset of  $V$  containing more than  $d$  vectors is linearly dependent. Thus a basis of  $V$  is a **maximal linearly independent subset** of  $V$ .*
- ② A subset of  $V$  containing fewer than  $d$  vectors cannot span  $V$ . Hence a basis of  $V$  can also be expressed as a **minimal spanning set** of  $V$ .*

# A linearly independent set can be extended to a basis

## Theorem

*Let  $S$  be a linearly independent subset of a finite dimensional vector space  $V$ .*

# A linearly independent set can be extended to a basis

## Theorem

*Let  $S$  be a linearly independent subset of a finite dimensional vector space  $V$ . Then  $S$  is finite,*

# A linearly independent set can be extended to a basis


## Theorem

*Let  $S$  be a linearly independent subset of a finite dimensional vector space  $V$ . Then  $S$  is finite, and it is part of a (finite) basis of  $V$ .*

# A linearly independent set can be extended to a basis

## Theorem


*Let  $S$  be a linearly independent subset of a finite dimensional vector space  $V$ . Then  $S$  is finite, and it is part of a (finite) basis of  $V$ .*



# A linearly independent set can be extended to a basis

## Theorem


*Let  $S$  be a linearly independent subset of a finite dimensional vector space  $V$ . Then  $S$  is finite, and it is part of a (finite) basis of  $V$ .*



# A linearly independent set can be extended to a basis

## Theorem

*Let  $S$  be a linearly independent subset of a finite dimensional vector space  $V$ . Then  $S$  is finite, and it is part of a (finite) basis of  $V$ .*






# A linearly independent set can be extended to a basis

## Theorem


*Let  $S$  be a linearly independent subset of a finite dimensional vector space  $V$ . Then  $S$  is finite, and it is part of a (finite) basis of  $V$ .*



# A linearly independent set can be extended to a basis

## Theorem


*Let  $S$  be a linearly independent subset of a finite dimensional vector space  $V$ . Then  $S$  is finite, and it is part of a (finite) basis of  $V$ .*



# A linearly independent set can be extended to a basis

## Theorem

*Let  $S$  be a linearly independent subset of a finite dimensional vector space  $V$ . Then  $S$  is finite, and it is part of a (finite) basis of  $V$ .*



# A linearly independent set can be extended to a basis

## Theorem

*Let  $S$  be a linearly independent subset of a finite dimensional vector space  $V$ . Then  $S$  is finite, and it is part of a (finite) basis of  $V$ .*

# A linearly independent set can be extended to a basis

## Theorem

*Let  $S$  be a linearly independent subset of a finite dimensional vector space  $V$ . Then  $S$  is finite, and it is part of a (finite) basis of  $V$ .*

*Proof.* Note that  $S$  contains at most  $\dim(V)$  elements.

# A linearly independent set can be extended to a basis

## Theorem

*Let  $S$  be a linearly independent subset of a finite dimensional vector space  $V$ . Then  $S$  is finite, and it is part of a (finite) basis of  $V$ .*

*Proof.* Note that  $S$  contains at most  $\dim(V)$  elements. So  $S$  is finite.

# A linearly independent set can be extended to a basis

## Theorem

*Let  $S$  be a linearly independent subset of a finite dimensional vector space  $V$ . Then  $S$  is finite, and it is part of a (finite) basis of  $V$ .*

*Proof.* Note that  $S$  contains at most  $\dim(V)$  elements. So  $S$  is finite.

If  $S$  spans  $V$ ,

# A linearly independent set can be extended to a basis

## Theorem

*Let  $S$  be a linearly independent subset of a finite dimensional vector space  $V$ . Then  $S$  is finite, and it is part of a (finite) basis of  $V$ .*

*Proof.* Note that  $S$  contains at most  $\dim(V)$  elements. So  $S$  is finite.

If  $S$  spans  $V$ , then  $S$  is a basis of  $V$ , and we are done.



# A linearly independent set can be extended to a basis

## Theorem

*Let  $S$  be a linearly independent subset of a finite dimensional vector space  $V$ . Then  $S$  is finite, and it is part of a (finite) basis of  $V$ .*

*Proof.* Note that  $S$  contains at most  $\dim(V)$  elements. So  $S$  is finite.

If  $S$  spans  $V$ , then  $S$  is a basis of  $V$ , and we are done. If  $\text{Span}(S) \neq V$ ,

# A linearly independent set can be extended to a basis

## Theorem

*Let  $S$  be a linearly independent subset of a finite dimensional vector space  $V$ . Then  $S$  is finite, and it is part of a (finite) basis of  $V$ .*

*Proof.* Note that  $S$  contains at most  $\dim(V)$  elements. So  $S$  is finite.

If  $S$  spans  $V$ , then  $S$  is a basis of  $V$ , and we are done. If  $\text{Span}(S) \neq V$ , then  $\exists v_1 \in V \setminus \text{Span}(S)$ .

# A linearly independent set can be extended to a basis

## Theorem

*Let  $S$  be a linearly independent subset of a finite dimensional vector space  $V$ . Then  $S$  is finite, and it is part of a (finite) basis of  $V$ .*

*Proof.* Note that  $S$  contains at most  $\dim(V)$  elements. So  $S$  is finite.

If  $S$  spans  $V$ , then  $S$  is a basis of  $V$ , and we are done. If  $\text{Span}(S) \neq V$ , then  $\exists v_1 \in V \setminus \text{Span}(S)$ . Hence,  $S_1 := S \cup \{v_1\}$  is linearly independent.

# A linearly independent set can be extended to a basis

## Theorem

*Let  $S$  be a linearly independent subset of a finite dimensional vector space  $V$ . Then  $S$  is finite, and it is part of a (finite) basis of  $V$ .*

*Proof.* Note that  $S$  contains at most  $\dim(V)$  elements. So  $S$  is finite.

If  $S$  spans  $V$ , then  $S$  is a basis of  $V$ , and we are done. If  $\text{Span}(S) \neq V$ , then  $\exists v_1 \in V \setminus \text{Span}(S)$ . Hence,  $S_1 := S \cup \{v_1\}$  is linearly independent. If  $\text{Span}(S_1) = V$ , then we are done.

# A linearly independent set can be extended to a basis

## Theorem

*Let  $S$  be a linearly independent subset of a finite dimensional vector space  $V$ . Then  $S$  is finite, and it is part of a (finite) basis of  $V$ .*

*Proof.* Note that  $S$  contains at most  $\dim(V)$  elements. So  $S$  is finite.

If  $S$  spans  $V$ , then  $S$  is a basis of  $V$ , and we are done. If  $\text{Span}(S) \neq V$ , then  $\exists v_1 \in V \setminus \text{Span}(S)$ . Hence,  $S_1 := S \cup \{v_1\}$  is linearly independent. If  $\text{Span}(S_1) = V$ , then we are done. Otherwise, if  $\text{Span}(S_1) \neq V$ ,

# A linearly independent set can be extended to a basis

## Theorem

*Let  $S$  be a linearly independent subset of a finite dimensional vector space  $V$ . Then  $S$  is finite, and it is part of a (finite) basis of  $V$ .*

*Proof.* Note that  $S$  contains at most  $\dim(V)$  elements. So  $S$  is finite.

If  $S$  spans  $V$ , then  $S$  is a basis of  $V$ , and we are done. If  $\text{Span}(S) \neq V$ , then  $\exists v_1 \in V \setminus \text{Span}(S)$ . Hence,  $S_1 := S \cup \{v_1\}$  is linearly independent. If  $\text{Span}(S_1) = V$ , then we are done. Otherwise, if  $\text{Span}(S_1) \neq V$ , there is  $v_2 \in V \setminus \text{Span}(S_1)$ ,

# A linearly independent set can be extended to a basis

## Theorem

*Let  $S$  be a linearly independent subset of a finite dimensional vector space  $V$ . Then  $S$  is finite, and it is part of a (finite) basis of  $V$ .*

*Proof.* Note that  $S$  contains at most  $\dim(V)$  elements. So  $S$  is finite.

If  $S$  spans  $V$ , then  $S$  is a basis of  $V$ , and we are done. If  $\text{Span}(S) \neq V$ , then  $\exists v_1 \in V \setminus \text{Span}(S)$ . Hence,  $S_1 := S \cup \{v_1\}$  is linearly independent. If  $\text{Span}(S_1) = V$ , then we are done. Otherwise, if  $\text{Span}(S_1) \neq V$ , there is  $v_2 \in V \setminus \text{Span}(S_1)$ , and hence  $S_2 := S \cup \{v_1, v_2\}$  is linearly independent.

# A linearly independent set can be extended to a basis

## Theorem

*Let  $S$  be a linearly independent subset of a finite dimensional vector space  $V$ . Then  $S$  is finite, and it is part of a (finite) basis of  $V$ .*

*Proof.* Note that  $S$  contains at most  $\dim(V)$  elements. So  $S$  is finite.

If  $S$  spans  $V$ , then  $S$  is a basis of  $V$ , and we are done. If  $\text{Span}(S) \neq V$ , then  $\exists v_1 \in V \setminus \text{Span}(S)$ . Hence,  $S_1 := S \cup \{v_1\}$  is linearly independent. If  $\text{Span}(S_1) = V$ , then we are done.

Otherwise, if  $\text{Span}(S_1) \neq V$ , there is  $v_2 \in V \setminus \text{Span}(S_1)$ , and hence  $S_2 := S \cup \{v_1, v_2\}$  is linearly independent.

This process stops after some finite steps because at most  $\dim(V)$  linearly independent vectors can be there in  $V$ .



# A linearly independent set can be extended to a basis

## Theorem

*Let  $S$  be a linearly independent subset of a finite dimensional vector space  $V$ . Then  $S$  is finite, and it is part of a (finite) basis of  $V$ .*

*Proof.* Note that  $S$  contains at most  $\dim(V)$  elements. So  $S$  is finite.

If  $S$  spans  $V$ , then  $S$  is a basis of  $V$ , and we are done. If  $\text{Span}(S) \neq V$ , then  $\exists v_1 \in V \setminus \text{Span}(S)$ . Hence,  $S_1 := S \cup \{v_1\}$  is linearly independent. If  $\text{Span}(S_1) = V$ , then we are done.

Otherwise, if  $\text{Span}(S_1) \neq V$ , there is  $v_2 \in V \setminus \text{Span}(S_1)$ , and hence  $S_2 := S \cup \{v_1, v_2\}$  is linearly independent.

This process stops after some finite steps because at most  $\dim(V)$  linearly independent vectors can be there in  $V$ .

So finally we obtain a set  $S \cup \{v_1, v_2, \dots, v_m\} \subset V$  which is linearly independent and spans  $V$ ,

# A linearly independent set can be extended to a basis

## Theorem

*Let  $S$  be a linearly independent subset of a finite dimensional vector space  $V$ . Then  $S$  is finite, and it is part of a (finite) basis of  $V$ .*

*Proof.* Note that  $S$  contains at most  $\dim(V)$  elements. So  $S$  is finite.

If  $S$  spans  $V$ , then  $S$  is a basis of  $V$ , and we are done. If  $\text{Span}(S) \neq V$ , then  $\exists v_1 \in V \setminus \text{Span}(S)$ . Hence,  $S_1 := S \cup \{v_1\}$  is linearly independent. If  $\text{Span}(S_1) = V$ , then we are done.

Otherwise, if  $\text{Span}(S_1) \neq V$ , there is  $v_2 \in V \setminus \text{Span}(S_1)$ , and hence  $S_2 := S \cup \{v_1, v_2\}$  is linearly independent.

This process stops after some finite steps because at most  $\dim(V)$  linearly independent vectors can be there in  $V$ .

So finally we obtain a set  $S \cup \{v_1, v_2, \dots, v_m\} \subset V$  which is linearly independent and spans  $V$ , i.e., it forms a basis of  $V$ .

# A proper subspace has less dimension

Let  $V$  be a finite dimensional vector space.

# A proper subspace has less dimension

Let  $V$  be a finite dimensional vector space.

## Theorem

*In  $V$ , every linearly independent set of vectors is part of a basis.*

# A proper subspace has less dimension

Let  $V$  be a finite dimensional vector space.

## Theorem

*In  $V$ , every linearly independent set of vectors is part of a basis.*

## Corollary

*A subspace  $W$  of  $V$  is PROPER*

# A proper subspace has less dimension

Let  $V$  be a finite dimensional vector space.

## Theorem

*In  $V$ , every linearly independent set of vectors is part of a basis.*

## Corollary

*A subspace  $W$  of  $V$  is PROPER if and only if  $\dim(W) < \dim(V)$ .*

# A proper subspace has less dimension

Let  $V$  be a finite dimensional vector space.

## Theorem

*In  $V$ , every linearly independent set of vectors is part of a basis.*

## Corollary

*A subspace  $W$  of  $V$  is PROPER if and only if  $\dim(W) < \dim(V)$ .*

The 'if' part is trivial,

# A proper subspace has less dimension

Let  $V$  be a finite dimensional vector space.

## Theorem

*In  $V$ , every linearly independent set of vectors is part of a basis.*

## Corollary

*A subspace  $W$  of  $V$  is PROPER if and only if  $\dim(W) < \dim(V)$ .*

The 'if' part is trivial, because if  $W = V$ , then  $\dim(W) = \dim(V)$ .



# A proper subspace has less dimension

Let  $V$  be a finite dimensional vector space.

## Theorem

*In  $V$ , every linearly independent set of vectors is part of a basis.*

## Corollary

*A subspace  $W$  of  $V$  is PROPER if and only if  $\dim(W) < \dim(V)$ .*

The 'if' part is trivial, because if  $W = V$ , then  $\dim(W) = \dim(V)$ .

*Proof of 'only if' part.*

# A proper subspace has less dimension

Let  $V$  be a finite dimensional vector space.

## Theorem

*In  $V$ , every linearly independent set of vectors is part of a basis.*

## Corollary

*A subspace  $W$  of  $V$  is PROPER if and only if  $\dim(W) < \dim(V)$ .*

The 'if' part is trivial, because if  $W = V$ , then  $\dim(W) = \dim(V)$ .

*Proof of 'only if' part.* Let  $W \subsetneq V$ .

# A proper subspace has less dimension

Let  $V$  be a finite dimensional vector space.

## Theorem

*In  $V$ , every linearly independent set of vectors is part of a basis.*

## Corollary

*A subspace  $W$  of  $V$  is PROPER if and only if  $\dim(W) < \dim(V)$ .*

The 'if' part is trivial, because if  $W = V$ , then  $\dim(W) = \dim(V)$ .

*Proof of 'only if' part.* Let  $W \subsetneq V$ . If  $W = 0$ , then we are done.

# A proper subspace has less dimension

Let  $V$  be a finite dimensional vector space.

## Theorem

*In  $V$ , every linearly independent set of vectors is part of a basis.*

## Corollary

*A subspace  $W$  of  $V$  is PROPER if and only if  $\dim(W) < \dim(V)$ .*

The 'if' part is trivial, because if  $W = V$ , then  $\dim(W) = \dim(V)$ .

*Proof of 'only if' part.* Let  $W \subsetneq V$ . If  $W = 0$ , then we are done.

Thus we may assume that  $\exists u \neq 0$  in  $W$ .

# A proper subspace has less dimension

Let  $V$  be a finite dimensional vector space.

## Theorem

*In  $V$ , every linearly independent set of vectors is part of a basis.*

## Corollary

*A subspace  $W$  of  $V$  is PROPER if and only if  $\dim(W) < \dim(V)$ .*

The 'if' part is trivial, because if  $W = V$ , then  $\dim(W) = \dim(V)$ .

*Proof of 'only if' part.* Let  $W \subsetneq V$ . If  $W = 0$ , then we are done.

Thus we may assume that  $\exists u \neq 0$  in  $W$ .

Then  $\{u\}$  can be extended to a finite basis (say  $S$ ) of  $W$ .

# A proper subspace has less dimension

Let  $V$  be a finite dimensional vector space.

## Theorem

*In  $V$ , every linearly independent set of vectors is part of a basis.*

## Corollary

*A subspace  $W$  of  $V$  is PROPER if and only if  $\dim(W) < \dim(V)$ .*

The 'if' part is trivial, because if  $W = V$ , then  $\dim(W) = \dim(V)$ .

*Proof of 'only if' part.* Let  $W \subsetneq V$ . If  $W = 0$ , then we are done.

Thus we may assume that  $\exists u \neq 0$  in  $W$ .

Then  $\{u\}$  can be extended to a finite basis (say  $S$ ) of  $W$ . So, in particular,  $W$  is finite dimensional.

# A proper subspace has less dimension

Let  $V$  be a finite dimensional vector space.

## Theorem

*In  $V$ , every linearly independent set of vectors is part of a basis.*

## Corollary

*A subspace  $W$  of  $V$  is PROPER if and only if  $\dim(W) < \dim(V)$ .*

The 'if' part is trivial, because if  $W = V$ , then  $\dim(W) = \dim(V)$ .

*Proof of 'only if' part.* Let  $W \subsetneq V$ . If  $W = 0$ , then we are done.

Thus we may assume that  $\exists u \neq 0$  in  $W$ .

Then  $\{u\}$  can be extended to a finite basis (say  $S$ ) of  $W$ . So, in particular,  $W$  is finite dimensional.

Since  $\text{Span}(S) = W \subsetneq V$ ,

# A proper subspace has less dimension

Let  $V$  be a finite dimensional vector space.

## Theorem

*In  $V$ , every linearly independent set of vectors is part of a basis.*

## Corollary

*A subspace  $W$  of  $V$  is PROPER if and only if  $\dim(W) < \dim(V)$ .*

The 'if' part is trivial, because if  $W = V$ , then  $\dim(W) = \dim(V)$ .

*Proof of 'only if' part.* Let  $W \subsetneq V$ . If  $W = 0$ , then we are done.

Thus we may assume that  $\exists u \neq 0$  in  $W$ .

Then  $\{u\}$  can be extended to a finite basis (say  $S$ ) of  $W$ . So, in particular,  $W$  is finite dimensional.

Since  $\text{Span}(S) = W \subsetneq V$ , there is a vector  $v \in V \setminus \text{Span}(S)$ .



# A proper subspace has less dimension

Let  $V$  be a finite dimensional vector space.

## Theorem

*In  $V$ , every linearly independent set of vectors is part of a basis.*

## Corollary

*A subspace  $W$  of  $V$  is PROPER if and only if  $\dim(W) < \dim(V)$ .*

The 'if' part is trivial, because if  $W = V$ , then  $\dim(W) = \dim(V)$ .

*Proof of 'only if' part.* Let  $W \subsetneq V$ . If  $W = 0$ , then we are done.

Thus we may assume that  $\exists u \neq 0$  in  $W$ .

Then  $\{u\}$  can be extended to a finite basis (say  $S$ ) of  $W$ . So, in particular,  $W$  is finite dimensional.

Since  $\text{Span}(S) = W \subsetneq V$ , there is a vector  $v \in V \setminus \text{Span}(S)$ .

Then  $S \cup \{v\}$  is a linearly independent subset of  $V$ .

# A proper subspace has less dimension

Let  $V$  be a finite dimensional vector space.

## Theorem

*In  $V$ , every linearly independent set of vectors is part of a basis.*

## Corollary

*A subspace  $W$  of  $V$  is PROPER if and only if  $\dim(W) < \dim(V)$ .*

The 'if' part is trivial, because if  $W = V$ , then  $\dim(W) = \dim(V)$ .

*Proof of 'only if' part.* Let  $W \subsetneq V$ . If  $W = 0$ , then we are done.

Thus we may assume that  $\exists u \neq 0$  in  $W$ .

Then  $\{u\}$  can be extended to a finite basis (say  $S$ ) of  $W$ . So, in particular,  $W$  is finite dimensional.

Since  $\text{Span}(S) = W \subsetneq V$ , there is a vector  $v \in V \setminus \text{Span}(S)$ .

Then  $S \cup \{v\}$  is a linearly independent subset of  $V$ .

Hence  $S \cup \{v\}$  can be extended to a basis of  $V$ .

# A proper subspace has less dimension

Let  $V$  be a finite dimensional vector space.

## Theorem

*In  $V$ , every linearly independent set of vectors is part of a basis.*

## Corollary

*A subspace  $W$  of  $V$  is PROPER if and only if  $\dim(W) < \dim(V)$ .*

The 'if' part is trivial, because if  $W = V$ , then  $\dim(W) = \dim(V)$ .

*Proof of 'only if' part.* Let  $W \subsetneq V$ . If  $W = 0$ , then we are done.

Thus we may assume that  $\exists u \neq 0$  in  $W$ .

Then  $\{u\}$  can be extended to a finite basis (say  $S$ ) of  $W$ . So, in particular,  $W$  is finite dimensional.


Since  $\text{Span}(S) = W \subsetneq V$ , there is a vector  $v \in V \setminus \text{Span}(S)$ .

Then  $S \cup \{v\}$  is a linearly independent subset of  $V$ .


Hence  $S \cup \{v\}$  can be extended to a basis of  $V$ .

Therefore  $\dim(W) < \dim(V)$ .


# Example: Proper subspaces of $\mathbb{R}^2$




## Example: Proper subspaces of $\mathbb{R}^2$




## Example: Proper subspaces of $\mathbb{R}^2$



## Example: Proper subspaces of $\mathbb{R}^2$



## Example: Proper subspaces of $\mathbb{R}^2$





# Sum of two subspaces

## Definition

Let  $W_1$  and  $W_2$  be two subspaces of  $V$ .

# Sum of two subspaces


## Definition

Let  $W_1$  and  $W_2$  be two subspaces of  $V$ . Then  
 $W_1 + W_2 := \{w_1 + w_2 : w_i \in W_i\}.$

# Sum of two subspaces

## Definition


Let  $W_1$  and  $W_2$  be two subspaces of  $V$ . Then  
 $W_1 + W_2 := \{w_1 + w_2 : w_i \in W_i\}.$



# Sum of two subspaces

## Definition


Let  $W_1$  and  $W_2$  be two subspaces of  $V$ . Then  
 $W_1 + W_2 := \{w_1 + w_2 : w_i \in W_i\}.$



# Sum of two subspaces

## Definition


Let  $W_1$  and  $W_2$  be two subspaces of  $V$ . Then  
 $W_1 + W_2 := \{w_1 + w_2 : w_i \in W_i\}.$



# Sum of two subspaces

## Definition


Let  $W_1$  and  $W_2$  be two subspaces of  $V$ . Then  
 $W_1 + W_2 := \{w_1 + w_2 : w_i \in W_i\}.$



# Sum of two subspaces

## Definition


Let  $W_1$  and  $W_2$  be two subspaces of  $V$ . Then  
 $W_1 + W_2 := \{w_1 + w_2 : w_i \in W_i\}.$



# Sum of two subspaces

## Definition

Let  $W_1$  and  $W_2$  be two subspaces of  $V$ . Then  
 $W_1 + W_2 := \{w_1 + w_2 : w_i \in W_i\}$ .





# Sum of two subspaces, and its dimension

## Theorem

*Let  $W_1$  and  $W_2$  be finite dimensional subspaces of  $V$ .*

# Sum of two subspaces, and its dimension

## Theorem

*Let  $W_1$  and  $W_2$  be finite dimensional subspaces of  $V$ . Then*

$$W_1 + W_2 := \{w_1 + w_2 : w_i \in W_i\}$$

*is a finite dimensional subspace of  $V$ ,*

# Sum of two subspaces, and its dimension

## Theorem

*Let  $W_1$  and  $W_2$  be finite dimensional subspaces of  $V$ . Then*

$$W_1 + W_2 := \{w_1 + w_2 : w_i \in W_i\}$$

*is a finite dimensional subspace of  $V$ , and*

$$\dim(W_1) + \dim(W_2) = \dim(W_1 \cap W_2) + \dim(W_1 + W_2).$$

# Sum of two subspaces, and its dimension

## Theorem

*Let  $W_1$  and  $W_2$  be finite dimensional subspaces of  $V$ . Then*

$$W_1 + W_2 := \{w_1 + w_2 : w_i \in W_i\}$$

*is a finite dimensional subspace of  $V$ , and*

$$\dim(W_1) + \dim(W_2) = \dim(W_1 \cap W_2) + \dim(W_1 + W_2).$$

*Proof.* Since  $W_1 \cap W_2 \subseteq W_1$ ,

# Sum of two subspaces, and its dimension

## Theorem

Let  $W_1$  and  $W_2$  be finite dimensional subspaces of  $V$ . Then

$$W_1 + W_2 := \{w_1 + w_2 : w_i \in W_i\}$$

is a finite dimensional subspace of  $V$ , and

$$\dim(W_1) + \dim(W_2) = \dim(W_1 \cap W_2) + \dim(W_1 + W_2).$$

*Proof.* Since  $W_1 \cap W_2 \subseteq W_1$ , it follows that  $W_1 \cap W_2$  has a finite basis  $\{u_1, \dots, u_r\}$ ,

# Sum of two subspaces, and its dimension

## Theorem

Let  $W_1$  and  $W_2$  be finite dimensional subspaces of  $V$ . Then

$$W_1 + W_2 := \{w_1 + w_2 : w_i \in W_i\}$$

is a finite dimensional subspace of  $V$ , and

$$\dim(W_1) + \dim(W_2) = \dim(W_1 \cap W_2) + \dim(W_1 + W_2).$$

*Proof.* Since  $W_1 \cap W_2 \subseteq W_1$ , it follows that  $W_1 \cap W_2$  has a finite basis  $\{u_1, \dots, u_r\}$ , which can be extended to a basis

$$\{u_1, \dots, u_r, v_1, \dots, v_m\} \text{ of } W_1$$

# Sum of two subspaces, and its dimension

## Theorem

Let  $W_1$  and  $W_2$  be finite dimensional subspaces of  $V$ . Then

$$W_1 + W_2 := \{w_1 + w_2 : w_i \in W_i\}$$

is a finite dimensional subspace of  $V$ , and

$$\dim(W_1) + \dim(W_2) = \dim(W_1 \cap W_2) + \dim(W_1 + W_2).$$

*Proof.* Since  $W_1 \cap W_2 \subseteq W_1$ , it follows that  $W_1 \cap W_2$  has a finite basis  $\{u_1, \dots, u_r\}$ , which can be extended to a basis

$$\{u_1, \dots, u_r, v_1, \dots, v_m\} \text{ of } W_1$$

and a basis

$$\{u_1, \dots, u_r, w_1, \dots, w_n\} \text{ of } W_2.$$

# Sum of two subspaces, and its dimension

## Theorem

Let  $W_1$  and  $W_2$  be finite dimensional subspaces of  $V$ . Then

$$W_1 + W_2 := \{w_1 + w_2 : w_i \in W_i\}$$

is a finite dimensional subspace of  $V$ , and

$$\dim(W_1) + \dim(W_2) = \dim(W_1 \cap W_2) + \dim(W_1 + W_2).$$

*Proof.* Since  $W_1 \cap W_2 \subseteq W_1$ , it follows that  $W_1 \cap W_2$  has a finite basis  $\{u_1, \dots, u_r\}$ , which can be extended to a basis

$$\{u_1, \dots, u_r, v_1, \dots, v_m\} \text{ of } W_1$$

and a basis

$$\{u_1, \dots, u_r, w_1, \dots, w_n\} \text{ of } W_2.$$

Show that  $\{u_1, \dots, u_r, v_1, \dots, v_m, w_1, \dots, w_n\}$  is a basis of  $W_1 + W_2$ .



# Ordered bases

- 1 Let  $V$  be a finite dimensional vector space over  $\mathbb{R}$ .

# Ordered bases

- 1 Let  $V$  be a finite dimensional vector space over  $\mathbb{R}$ .
- 2 An **ordered basis** of  $V$  is nothing but a basis  $\mathcal{B} = \{v_1, \dots, v_n\}$  with a specified ordering of its vectors.

# Ordered bases

- 1 Let  $V$  be a finite dimensional vector space over  $\mathbb{R}$ .
- 2 An **ordered basis** of  $V$  is nothing but a basis  $\mathcal{B} = \{v_1, \dots, v_n\}$  with a specified ordering of its vectors.
- 3 It essentially enables us to introduce coordinates in  $V$ .

# Ordered bases

- 1 Let  $V$  be a finite dimensional vector space over  $\mathbb{R}$ .
- 2 An **ordered basis** of  $V$  is nothing but a basis  $\mathcal{B} = \{v_1, \dots, v_n\}$  with a specified ordering of its vectors.
- 3 It essentially enables us to introduce coordinates in  $V$ .
- 4 Let  $\mathcal{B} = \{v_1, \dots, v_n\}$  be an ordered basis of  $V$ .

# Ordered bases

- 1 Let  $V$  be a finite dimensional vector space over  $\mathbb{R}$ .
- 2 An **ordered basis** of  $V$  is nothing but a basis  $\mathcal{B} = \{v_1, \dots, v_n\}$  with a specified ordering of its vectors.
- 3 It essentially enables us to introduce coordinates in  $V$ .
- 4 Let  $\mathcal{B} = \{v_1, \dots, v_n\}$  be an ordered basis of  $V$ .
- 5 Given  $v \in V$ ,

# Ordered bases

- 1 Let  $V$  be a finite dimensional vector space over  $\mathbb{R}$ .
- 2 An **ordered basis** of  $V$  is nothing but a basis  $\mathcal{B} = \{v_1, \dots, v_n\}$  with a specified ordering of its vectors.
- 3 It essentially enables us to introduce coordinates in  $V$ .
- 4 Let  $\mathcal{B} = \{v_1, \dots, v_n\}$  be an ordered basis of  $V$ .
- 5 Given  $v \in V$ , there is a UNIQUE  $n$ -tuple  $(x_1, \dots, x_n)$  of scalars such that  $v = x_1 v_1 + \dots + x_n v_n$ .

# Ordered bases

- 1 Let  $V$  be a finite dimensional vector space over  $\mathbb{R}$ .
- 2 An **ordered basis** of  $V$  is nothing but a basis  $\mathcal{B} = \{v_1, \dots, v_n\}$  with a specified ordering of its vectors.
- 3 It essentially enables us to introduce coordinates in  $V$ .
- 4 Let  $\mathcal{B} = \{v_1, \dots, v_n\}$  be an ordered basis of  $V$ .
- 5 Given  $v \in V$ , there is a UNIQUE  $n$ -tuple  $(x_1, \dots, x_n)$  of scalars such that  $v = x_1 v_1 + \dots + x_n v_n$ . Why?

# Ordered bases

- 1 Let  $V$  be a finite dimensional vector space over  $\mathbb{R}$ .
- 2 An **ordered basis** of  $V$  is nothing but a basis  $\mathcal{B} = \{v_1, \dots, v_n\}$  with a specified ordering of its vectors.
- 3 It essentially enables us to introduce coordinates in  $V$ .
- 4 Let  $\mathcal{B} = \{v_1, \dots, v_n\}$  be an ordered basis of  $V$ .
- 5 Given  $v \in V$ , there is a UNIQUE  $n$ -tuple  $(x_1, \dots, x_n)$  of scalars such that  $v = x_1 v_1 + \dots + x_n v_n$ . Why?
- 6 We call such  $x_i$  the  $i$ th **coordinate of  $v$  relative to  $\mathcal{B}$** .



# Ordered bases

- 1 Let  $V$  be a finite dimensional vector space over  $\mathbb{R}$ .
- 2 An **ordered basis** of  $V$  is nothing but a basis  $\mathcal{B} = \{v_1, \dots, v_n\}$  with a specified ordering of its vectors.
- 3 It essentially enables us to introduce coordinates in  $V$ .
- 4 Let  $\mathcal{B} = \{v_1, \dots, v_n\}$  be an ordered basis of  $V$ .
- 5 Given  $v \in V$ , there is a UNIQUE  $n$ -tuple  $(x_1, \dots, x_n)$  of scalars such that  $v = x_1 v_1 + \dots + x_n v_n$ . Why?
- 6 We call such  $x_i$  the  $i$ th **coordinate of  $v$  relative to  $\mathcal{B}$** .
- 7 We denote the **coordinate vector of  $v$  relative to  $\mathcal{B}$**  by

$$[v]_{\mathcal{B}} := \begin{bmatrix} x_1 \\ \vdots \\ x_n \end{bmatrix}.$$

# Ordered bases

- 1 Let  $V$  be a finite dimensional vector space over  $\mathbb{R}$ .
- 2 An **ordered basis** of  $V$  is nothing but a basis  $\mathcal{B} = \{v_1, \dots, v_n\}$  with a specified ordering of its vectors.
- 3 It essentially enables us to introduce coordinates in  $V$ .
- 4 Let  $\mathcal{B} = \{v_1, \dots, v_n\}$  be an ordered basis of  $V$ .
- 5 Given  $v \in V$ , there is a **UNIQUE**  $n$ -tuple  $(x_1, \dots, x_n)$  of scalars such that  $v = x_1 v_1 + \dots + x_n v_n$ . Why?
- 6 We call such  $x_i$  the  $i$ th **coordinate of  $v$  relative to  $\mathcal{B}$** .
- 7 We denote the **coordinate vector of  $v$  relative to  $\mathcal{B}$**  by

$$[v]_{\mathcal{B}} := \begin{bmatrix} x_1 \\ \vdots \\ x_n \end{bmatrix}.$$

- 8 On the other hand, every  $n$ -tuple  $(x_1, \dots, x_n) \in \mathbb{R}^n$  is the coordinate vector of some vector in  $V$  relative to  $\mathcal{B}$ ,

# Ordered bases

- 1 Let  $V$  be a finite dimensional vector space over  $\mathbb{R}$ .
- 2 An **ordered basis** of  $V$  is nothing but a basis  $\mathcal{B} = \{v_1, \dots, v_n\}$  with a specified ordering of its vectors.
- 3 It essentially enables us to introduce coordinates in  $V$ .
- 4 Let  $\mathcal{B} = \{v_1, \dots, v_n\}$  be an ordered basis of  $V$ .
- 5 Given  $v \in V$ , there is a UNIQUE  $n$ -tuple  $(x_1, \dots, x_n)$  of scalars such that  $v = x_1 v_1 + \dots + x_n v_n$ . Why?
- 6 We call such  $x_i$  the  $i$ th **coordinate of  $v$  relative to  $\mathcal{B}$** .
- 7 We denote the **coordinate vector of  $v$  relative to  $\mathcal{B}$**  by

$$[v]_{\mathcal{B}} := \begin{bmatrix} x_1 \\ \vdots \\ x_n \end{bmatrix}.$$

- 8 On the other hand, every  $n$ -tuple  $(x_1, \dots, x_n) \in \mathbb{R}^n$  is the coordinate vector of some vector in  $V$  relative to  $\mathcal{B}$ , namely the vector  $x_1 v_1 + \dots + x_n v_n$ .

# Correspondence between vectors in $V$ and $n$ -tuples in $\mathbb{R}^n$

- 1 Let  $u, v \in V$  and  $c \in \mathbb{R}$ .

# Correspondence between vectors in $V$ and $n$ -tuples in $\mathbb{R}^n$

- ① Let  $u, v \in V$  and  $c \in \mathbb{R}$ . It can be easily verified that

$$[u + v]_{\mathcal{B}} = [u]_{\mathcal{B}} + [v]_{\mathcal{B}} \quad \text{and} \quad [cu]_{\mathcal{B}} = c[u]_{\mathcal{B}}.$$

# Correspondence between vectors in $V$ and $n$ -tuples in $\mathbb{R}^n$

- ① Let  $u, v \in V$  and  $c \in \mathbb{R}$ . It can be easily verified that

$$[u + v]_{\mathcal{B}} = [u]_{\mathcal{B}} + [v]_{\mathcal{B}} \quad \text{and} \quad [cu]_{\mathcal{B}} = c[u]_{\mathcal{B}}.$$

- ② Thus an ordered basis  $\mathcal{B}$  of  $V$  determines a one-to-one correspondence

$$v \longleftrightarrow [v]_{\mathcal{B}}$$

# Correspondence between vectors in $V$ and $n$ -tuples in $\mathbb{R}^n$

- ① Let  $u, v \in V$  and  $c \in \mathbb{R}$ . It can be easily verified that

$$[u + v]_{\mathcal{B}} = [u]_{\mathcal{B}} + [v]_{\mathcal{B}} \quad \text{and} \quad [cu]_{\mathcal{B}} = c[u]_{\mathcal{B}}.$$

- ② Thus an ordered basis  $\mathcal{B}$  of  $V$  determines a one-to-one correspondence

$$v \longleftrightarrow [v]_{\mathcal{B}}$$

between the set of all vectors in  $V$  and the set of all  $n$ -tuples in  $\mathbb{R}^n$ ,

# Correspondence between vectors in $V$ and $n$ -tuples in $\mathbb{R}^n$

- ① Let  $u, v \in V$  and  $c \in \mathbb{R}$ . It can be easily verified that

$$[u + v]_{\mathcal{B}} = [u]_{\mathcal{B}} + [v]_{\mathcal{B}} \quad \text{and} \quad [cu]_{\mathcal{B}} = c[u]_{\mathcal{B}}.$$

- ② Thus an ordered basis  $\mathcal{B}$  of  $V$  determines a one-to-one correspondence

$$v \longleftrightarrow [v]_{\mathcal{B}}$$

between the set of all vectors in  $V$  and the set of all  $n$ -tuples in  $\mathbb{R}^n$ , and this correspondence respects both the vector additions and scalar multiplications.



# Correspondence between vectors in $V$ and $n$ -tuples in $\mathbb{R}^n$

- ① Let  $u, v \in V$  and  $c \in \mathbb{R}$ . It can be easily verified that

$$[u + v]_{\mathcal{B}} = [u]_{\mathcal{B}} + [v]_{\mathcal{B}} \quad \text{and} \quad [cu]_{\mathcal{B}} = c[u]_{\mathcal{B}}.$$

- ② Thus an ordered basis  $\mathcal{B}$  of  $V$  determines a one-to-one correspondence

$$v \longleftrightarrow [v]_{\mathcal{B}}$$

between the set of all vectors in  $V$  and the set of all  $n$ -tuples in  $\mathbb{R}^n$ , and this correspondence respects both the vector additions and scalar multiplications.

- ③ This also shows that any vector space  $V$  of dimension  $n$  over  $\mathbb{R}$  is isomorphic to  $\mathbb{R}^n$ .

# Change of coordinates matrix

- ① Let  $\mathcal{B} = \{v_1, \dots, v_n\}$ ,  $\mathcal{B}' = \{v'_1, \dots, v'_n\}$  be ordered bases of  $V$ .

# Change of coordinates matrix

- 1 Let  $\mathcal{B} = \{v_1, \dots, v_n\}$ ,  $\mathcal{B}' = \{v'_1, \dots, v'_n\}$  be ordered bases of  $V$ .
- 2 There exists an  $n \times n$  matrix  $P$  such that  $P[v]_{\mathcal{B}} = [v]_{\mathcal{B}'} \forall v \in V$ .

# Change of coordinates matrix

- 1 Let  $\mathcal{B} = \{v_1, \dots, v_n\}$ ,  $\mathcal{B}' = \{v'_1, \dots, v'_n\}$  be ordered bases of  $V$ .
- 2 There exists an  $n \times n$  matrix  $P$  such that  $P[v]_{\mathcal{B}} = [v]_{\mathcal{B}'} \forall v \in V$ .  
This  $P$  is called the **change of coordinates matrix** from  $\mathcal{B}$  to  $\mathcal{B}'$ .

# Change of coordinates matrix

- 1 Let  $\mathcal{B} = \{v_1, \dots, v_n\}$ ,  $\mathcal{B}' = \{v'_1, \dots, v'_n\}$  be ordered bases of  $V$ .
- 2 There exists an  $n \times n$  matrix  $P$  such that  $P[v]_{\mathcal{B}} = [v]_{\mathcal{B}'} \quad \forall v \in V$ .  
This  $P$  is called the **change of coordinates matrix** from  $\mathcal{B}$  to  $\mathcal{B}'$ .
- 3 Since  $\mathcal{B}'$  is a basis,  $\exists$  UNIQUE scalars  $P_{ij}$  ( $1 \leq i, j \leq n$ ) such that

$$v_j = \sum_{i=1}^n P_{ij} v'_i \quad \text{for every } 1 \leq j \leq n. \quad (2)$$

# Change of coordinates matrix

- 1 Let  $\mathcal{B} = \{v_1, \dots, v_n\}$ ,  $\mathcal{B}' = \{v'_1, \dots, v'_n\}$  be ordered bases of  $V$ .
- 2 There exists an  $n \times n$  matrix  $P$  such that  $P[v]_{\mathcal{B}} = [v]_{\mathcal{B}'} \quad \forall v \in V$ .  
This  $P$  is called the **change of coordinates matrix** from  $\mathcal{B}$  to  $\mathcal{B}'$ .
- 3 Since  $\mathcal{B}'$  is a basis,  $\exists$  UNIQUE scalars  $P_{ij}$  ( $1 \leq i, j \leq n$ ) such that

$$v_j = \sum_{i=1}^n P_{ij} v'_i \quad \text{for every } 1 \leq j \leq n. \quad (2)$$

- 4 Fix  $v \in V$ .

# Change of coordinates matrix

- 1 Let  $\mathcal{B} = \{v_1, \dots, v_n\}$ ,  $\mathcal{B}' = \{v'_1, \dots, v'_n\}$  be ordered bases of  $V$ .
- 2 There exists an  $n \times n$  matrix  $P$  such that  $P[v]_{\mathcal{B}} = [v]_{\mathcal{B}'} \quad \forall v \in V$ .  
This  $P$  is called the **change of coordinates matrix** from  $\mathcal{B}$  to  $\mathcal{B}'$ .
- 3 Since  $\mathcal{B}'$  is a basis,  $\exists$  UNIQUE scalars  $P_{ij}$  ( $1 \leq i, j \leq n$ ) such that

$$v_j = \sum_{i=1}^n P_{ij} v'_i \quad \text{for every } 1 \leq j \leq n. \quad (2)$$

- 4 Fix  $v \in V$ . Let  $[v]_{\mathcal{B}} = \begin{bmatrix} x_1 \\ \vdots \\ x_n \end{bmatrix}$

# Change of coordinates matrix

- 1 Let  $\mathcal{B} = \{v_1, \dots, v_n\}$ ,  $\mathcal{B}' = \{v'_1, \dots, v'_n\}$  be ordered bases of  $V$ .
- 2 There exists an  $n \times n$  matrix  $P$  such that  $P[v]_{\mathcal{B}} = [v]_{\mathcal{B}'} \quad \forall v \in V$ .  
This  $P$  is called the **change of coordinates matrix** from  $\mathcal{B}$  to  $\mathcal{B}'$ .
- 3 Since  $\mathcal{B}'$  is a basis,  $\exists$  UNIQUE scalars  $P_{ij}$  ( $1 \leq i, j \leq n$ ) such that

$$v_j = \sum_{i=1}^n P_{ij} v'_i \quad \text{for every } 1 \leq j \leq n. \quad (2)$$

- 4 Fix  $v \in V$ . Let  $[v]_{\mathcal{B}} = \begin{bmatrix} x_1 \\ \vdots \\ x_n \end{bmatrix}$  and  $[v]_{\mathcal{B}'} = \begin{bmatrix} x'_1 \\ \vdots \\ x'_n \end{bmatrix}$ .



# Change of coordinates matrix

- 1 Let  $\mathcal{B} = \{v_1, \dots, v_n\}$ ,  $\mathcal{B}' = \{v'_1, \dots, v'_n\}$  be ordered bases of  $V$ .
- 2 There exists an  $n \times n$  matrix  $P$  such that  $P[v]_{\mathcal{B}} = [v]_{\mathcal{B}'} \quad \forall v \in V$ .  
This  $P$  is called the **change of coordinates matrix** from  $\mathcal{B}$  to  $\mathcal{B}'$ .
- 3 Since  $\mathcal{B}'$  is a basis,  $\exists$  UNIQUE scalars  $P_{ij}$  ( $1 \leq i, j \leq n$ ) such that

$$v_j = \sum_{i=1}^n P_{ij} v'_i \quad \text{for every } 1 \leq j \leq n. \quad (2)$$

- 4 Fix  $v \in V$ . Let  $[v]_{\mathcal{B}} = \begin{bmatrix} x_1 \\ \vdots \\ x_n \end{bmatrix}$  and  $[v]_{\mathcal{B}'} = \begin{bmatrix} x'_1 \\ \vdots \\ x'_n \end{bmatrix}$ . Hence

$$v = \sum_{j=1}^n x_j v_j = \sum_{j=1}^n x_j \left( \sum_{i=1}^n P_{ij} v'_i \right) = \sum_{i=1}^n \left( \sum_{j=1}^n P_{ij} x_j \right) v'_i. \quad (3)$$

# Change of coordinates matrix

- 1 Let  $\mathcal{B} = \{v_1, \dots, v_n\}$ ,  $\mathcal{B}' = \{v'_1, \dots, v'_n\}$  be ordered bases of  $V$ .
- 2 There exists an  $n \times n$  matrix  $P$  such that  $P[v]_{\mathcal{B}} = [v]_{\mathcal{B}'} \quad \forall v \in V$ .  
This  $P$  is called the **change of coordinates matrix** from  $\mathcal{B}$  to  $\mathcal{B}'$ .
- 3 Since  $\mathcal{B}'$  is a basis,  $\exists$  UNIQUE scalars  $P_{ij}$  ( $1 \leq i, j \leq n$ ) such that

$$v_j = \sum_{i=1}^n P_{ij} v'_i \quad \text{for every } 1 \leq j \leq n. \quad (2)$$

- 4 Fix  $v \in V$ . Let  $[v]_{\mathcal{B}} = \begin{bmatrix} x_1 \\ \vdots \\ x_n \end{bmatrix}$  and  $[v]_{\mathcal{B}'} = \begin{bmatrix} x'_1 \\ \vdots \\ x'_n \end{bmatrix}$ . Hence

$$v = \sum_{j=1}^n x_j v_j = \sum_{j=1}^n x_j \left( \sum_{i=1}^n P_{ij} v'_i \right) = \sum_{i=1}^n \left( \sum_{j=1}^n P_{ij} x_j \right) v'_i. \quad (3)$$

- 5 It follows from the above equalities that

$$[v]_{\mathcal{B}'} = P[v]_{\mathcal{B}} \quad \text{for every vector } v \in V.$$

# Thank You!