

Differential Equations (MA 1150)

Sukumar

Lecture 8

May 09, 2020

Higher Order Linear Differential Equations

- The Wronskian and Abel's Formula

- Particular solution

- Higher order constant coefficient equations

Section 1

Higher Order Linear Differential Equations

Definition An n th order differential equation is said to be linear if it can be written in the form

$$y^{(n)} + p_1(x)y^{(n-1)} + \cdots + p_n(x)y = f(x).$$

Higher Order Linear Differential Equations

More generally, n th order linear differential equations can be written as

$$P_0(x)y^{(n)} + P_1(x)y^{(n-1)} + \cdots + P_n(x)y = F(x), \quad (1)$$

Higher Order Linear Differential Equations

More generally, n th order linear differential equations can be written as

$$P_0(x)y^{(n)} + P_1(x)y^{(n-1)} + \cdots + P_n(x)y = F(x), \quad (1)$$

Suppose $P_0(x) \neq 0$ for all x in some open interval (a, b) , then we can divide by $P_0(x)$ in (1) on (a, b) , and obtain the equation

$$y^{(n)} + p_1(x)y^{(n-1)} + \cdots + p_n(x)y = f(x).$$

where $p_i(x) = \frac{P_i(x)}{P_0(x)}$ and $f(x) = \frac{F(x)}{P_0(x)}$.

Convention The left side of n th order linear differential equation

$$P_0(x)y^{(n)} + P_1(x)y^{(n-1)} + \cdots + P_n(x)y = F(x), \quad (2)$$

is denoted by Ly ; that is,

Higher Order Linear Differential Equations

Convention The left side of n th order linear differential equation

$$P_0(x)y^{(n)} + P_1(x)y^{(n-1)} + \cdots + P_n(x)y = F(x), \quad (2)$$

is denoted by Ly ; that is,

$$Ly = P_0y^{(n)} + P_1y^{(n-1)} + \cdots + P_ny.$$

Higher Order Linear Differential Equations

Convention The left side of n th order linear differential equation

$$P_0(x)y^{(n)} + P_1(x)y^{(n-1)} + \cdots + P_n(x)y = F(x), \quad (2)$$

is denoted by Ly ; that is,

$$Ly = P_0y^{(n)} + P_1y^{(n-1)} + \cdots + P_ny.$$

We say that the equation $Ly = F$ is *normal* on (a, b) if P_0, P_1, \dots, P_n and F are continuous on (a, b) and P_0 has no zeros on (a, b) .

Higher Order Linear Differential Equations

Convention The left side of n th order linear differential equation

$$P_0(x)y^{(n)} + P_1(x)y^{(n-1)} + \cdots + P_n(x)y = F(x), \quad (2)$$

is denoted by Ly ; that is,

$$Ly = P_0y^{(n)} + P_1y^{(n-1)} + \cdots + P_ny.$$

We say that the equation $Ly = F$ is *normal* on (a, b) if P_0, P_1, \dots, P_n and F are continuous on (a, b) and P_0 has no zeros on (a, b) .

Homogeneous If $F \equiv 0$,

Higher Order Linear Differential Equations

Convention The left side of n th order linear differential equation

$$P_0(x)y^{(n)} + P_1(x)y^{(n-1)} + \cdots + P_n(x)y = F(x), \quad (2)$$

is denoted by Ly ; that is,

$$Ly = P_0y^{(n)} + P_1y^{(n-1)} + \cdots + P_ny.$$

We say that the equation $Ly = F$ is *normal* on (a, b) if P_0, P_1, \dots, P_n and F are continuous on (a, b) and P_0 has no zeros on (a, b) .

Homogeneous If $F \equiv 0$, otherwise **Nonhomogeneous**.

Theorem Suppose $Ly = F$ is normal on (a, b) and let $x_0 \in (a, b)$. Then the initial value problem

$$Ly = F, \quad y(x_0) = k_0, \quad y'(x_0) = k_1, \quad \dots, \quad y^{(n-1)}(x_0) = k_{n-1}$$

has a unique solution on (a, b) .

Theorem Suppose $Ly = F$ is normal on (a, b) and let $x_0 \in (a, b)$. Then the initial value problem

$$Ly = F, \quad y(x_0) = k_0, \quad y'(x_0) = k_1, \quad \dots, \quad y^{(n-1)}(x_0) = k_{n-1}$$

has a unique solution on (a, b) .

You may compare this with the 1st order and 2nd order linear ODE.

Higher Order Linear Differential Equations

$$Ly = P_0 y^{(n)} + P_1 y^{(n-1)} + \cdots + P_n y.$$

Higher Order Linear Differential Equations

$$Ly = P_0 y^{(n)} + P_1 y^{(n-1)} + \cdots + P_n y.$$

Theorem if y_1, y_2, \dots, y_n are solutions of $Ly = 0$ on (a, b) , then so is any linear combination of $\{y_1, y_2, \dots, y_n\}$.

Higher Order Linear Differential Equations

$$Ly = P_0y^{(n)} + P_1y^{(n-1)} + \cdots + P_ny.$$

Theorem if y_1, y_2, \dots, y_n are solutions of $Ly = 0$ on (a, b) , then so is any linear combination of $\{y_1, y_2, \dots, y_n\}$.

We say that $\{y_1, y_2, \dots, y_n\}$ is a fundamental set of solutions of $Ly = 0$ on (a, b) if every solution of $Ly = 0$ on (a, b) can be written as a linear combination of $\{y_1, y_2, \dots, y_n\}$.

Higher Order Linear Differential Equations

$$Ly = P_0y^{(n)} + P_1y^{(n-1)} + \cdots + P_ny.$$

Theorem if y_1, y_2, \dots, y_n are solutions of $Ly = 0$ on (a, b) , then so is any linear combination of $\{y_1, y_2, \dots, y_n\}$.

We say that $\{y_1, y_2, \dots, y_n\}$ is a fundamental set of solutions of $Ly = 0$ on (a, b) if every solution of $Ly = 0$ on (a, b) can be written as a linear combination of $\{y_1, y_2, \dots, y_n\}$.

Theorem If $Ly = 0$ is normal on (a, b) , then a set $\{y_1, y_2, \dots, y_n\}$ of n solutions of $Ly = 0$ on (a, b) is a fundamental set if and only if it's linearly independent on (a, b) .

Subsection 1

The Wronskian and Abel's Formula

The Wronskian and Abel's Formula

Let y_1 and y_2 be solutions of

$$y'' + p(x)y' + q(x)y = 0 \tag{3}$$

on (a, b) .

The Wronskian and Abel's Formula

Let y_1 and y_2 be solutions of

$$y'' + p(x)y' + q(x)y = 0 \quad (3)$$

on (a, b) .

Definition: The Wronskian of y_1 and y_2 is

$$W = \begin{vmatrix} y_1 & y_2 \\ y_1' & y_2' \end{vmatrix} = y_1 y_2' - y_1' y_2.$$

The Wronskian and Abel's Formula

Let y_1 and y_2 be solutions of

$$y'' + p(x)y' + q(x)y = 0 \quad (3)$$

on (a, b) .

Definition: The Wronskian of y_1 and y_2 is

$$W = \begin{vmatrix} y_1 & y_2 \\ y_1' & y_2' \end{vmatrix} = y_1 y_2' - y_1' y_2.$$

Remark: y_1 and y_2 are linearly independent on (a, b) if and only if W has no zeros on (a, b) .

The Wronskian and Abel's Formula

Let y_1 and y_2 be solutions of

$$y'' + p(x)y' + q(x)y = 0 \quad (3)$$

on (a, b) .

Definition: The Wronskian of y_1 and y_2 is

$$W = \begin{vmatrix} y_1 & y_2 \\ y_1' & y_2' \end{vmatrix} = y_1 y_2' - y_1' y_2.$$

Remark: y_1 and y_2 are linearly independent on (a, b) if and only if W has no zeros on (a, b) .

Abel's formula

$$W(x) = W(x_0)e^{-\int_{x_0}^x p(t) dt}, \quad a < x < b.$$

Higher Order Linear Differential Equations

Consider the following equation:

$$Ly = P_0y^{(n)} + P_1y^{(n-1)} + \cdots + P_ny = 0$$

Higher Order Linear Differential Equations

Consider the following equation:

$$Ly = P_0 y^{(n)} + P_1 y^{(n-1)} + \cdots + P_n y = 0$$

Theorem if y_1, y_2, \dots, y_n are solutions of $Ly = 0$ on (a, b) , then so is any linear combination of $\{y_1, y_2, \dots, y_n\}$.

Higher Order Linear Differential Equations

Consider the following equation:

$$Ly = P_0 y^{(n)} + P_1 y^{(n-1)} + \cdots + P_n y = 0$$

Theorem if y_1, y_2, \dots, y_n are solutions of $Ly = 0$ on (a, b) , then so is any linear combination of $\{y_1, y_2, \dots, y_n\}$.

We say that $\{y_1, y_2, \dots, y_n\}$ is a fundamental set of solutions of $Ly = 0$ on (a, b) if every solution of $Ly = 0$ on (a, b) can be written as a linear combination of $\{y_1, y_2, \dots, y_n\}$.

Higher Order Linear Differential Equations

Consider the following equation:

$$Ly = P_0 y^{(n)} + P_1 y^{(n-1)} + \cdots + P_n y = 0$$

Theorem if y_1, y_2, \dots, y_n are solutions of $Ly = 0$ on (a, b) , then so is any linear combination of $\{y_1, y_2, \dots, y_n\}$.

We say that $\{y_1, y_2, \dots, y_n\}$ is a fundamental set of solutions of $Ly = 0$ on (a, b) if every solution of $Ly = 0$ on (a, b) can be written as a linear combination of $\{y_1, y_2, \dots, y_n\}$.

Theorem If $Ly = 0$ is normal on (a, b) , then a set $\{y_1, y_2, \dots, y_n\}$ of n solutions of $Ly = 0$ on (a, b) is a fundamental set if and only if it is linearly independent on (a, b) .

Higher Order Linear Differential Equations

Theorem If $Ly = 0$ is normal on (a, b) , then a set $\{y_1, y_2, \dots, y_n\}$ of n solutions of $Ly = 0$ on (a, b) is a fundamental set if and only if it's linearly independent on (a, b) .

Higher Order Linear Differential Equations

Theorem If $Ly = 0$ is normal on (a, b) , then a set $\{y_1, y_2, \dots, y_n\}$ of n solutions of $Ly = 0$ on (a, b) is a fundamental set if and only if it's linearly independent on (a, b) .

To check linear independence of $\{y_1, y_2, \dots, y_n\}$ on (a, b) , where y_1, y_2, \dots, y_n are solutions of n th order equation $Ly = 0$. (Assume that $Ly = 0$ is normal).

Higher Order Linear Differential Equations

Theorem If $Ly = 0$ is normal on (a, b) , then a set $\{y_1, y_2, \dots, y_n\}$ of n solutions of $Ly = 0$ on (a, b) is a fundamental set if and only if it's linearly independent on (a, b) .

To check linear independence of $\{y_1, y_2, \dots, y_n\}$ on (a, b) , where y_1, y_2, \dots, y_n are solutions of n th order equation $Ly = 0$. (Assume that $Ly = 0$ is normal).

Suppose that there are constants c_1, c_2, \dots, c_n such that

$$c_1 y_1 + c_2 y_2 + \dots + c_n y_n = 0, \quad a < x < b.$$

Higher Order Linear Differential Equations

Theorem If $Ly = 0$ is normal on (a, b) , then a set $\{y_1, y_2, \dots, y_n\}$ of n solutions of $Ly = 0$ on (a, b) is a fundamental set if and only if it's linearly independent on (a, b) .

To check linear independence of $\{y_1, y_2, \dots, y_n\}$ on (a, b) , where y_1, y_2, \dots, y_n are solutions of n th order equation $Ly = 0$. (Assume that $Ly = 0$ is normal).

Suppose that there are constants c_1, c_2, \dots, c_n such that

$$c_1 y_1 + c_2 y_2 + \dots + c_n y_n = 0, \quad a < x < b.$$

Differentiating $n - 1$ times leads to the $n \times n$ system of equations

$$\begin{aligned} c_1 y_1(x) + c_2 y_2(x) + \dots + c_n y_n(x) &= 0 \\ c_1 y_1'(x) + c_2 y_2'(x) + \dots + c_n y_n'(x) &= 0 \\ &\vdots \\ c_1 y_1^{(n-1)}(x) + c_2 y_2^{(n-1)}(x) + \dots + c_n y_n^{(n-1)}(x) &= 0 \end{aligned} \tag{4}$$

Wronskian of $\{y_1, y_2, \dots, y_n\}$.

For a fixed x , let

$$W(x) = \begin{vmatrix} y_1(x) & y_2(x) & \cdots & y_n(x) \\ y_1'(x) & y_2'(x) & \cdots & y_n'(x) \\ \vdots & \vdots & \ddots & \vdots \\ y_1^{(n-1)}(x) & y_2^{(n-1)}(x) & \cdots & y_n^{(n-1)}(x) \end{vmatrix}.$$

Wronskian of $\{y_1, y_2, \dots, y_n\}$.

For a fixed x , let

$$W(x) = \begin{vmatrix} y_1(x) & y_2(x) & \cdots & y_n(x) \\ y_1'(x) & y_2'(x) & \cdots & y_n'(x) \\ \vdots & \vdots & \ddots & \vdots \\ y_1^{(n-1)}(x) & y_2^{(n-1)}(x) & \cdots & y_n^{(n-1)}(x) \end{vmatrix}.$$

We call this determinant the Wronskian of $\{y_1, y_2, \dots, y_n\}$.

Wronskian of $\{y_1, y_2, \dots, y_n\}$.

For a fixed x , let

$$W(x) = \begin{vmatrix} y_1(x) & y_2(x) & \cdots & y_n(x) \\ y_1'(x) & y_2'(x) & \cdots & y_n'(x) \\ \vdots & \vdots & \ddots & \vdots \\ y_1^{(n-1)}(x) & y_2^{(n-1)}(x) & \cdots & y_n^{(n-1)}(x) \end{vmatrix}.$$

We call this determinant the Wronskian of $\{y_1, y_2, \dots, y_n\}$.

If $W(x) \neq 0$ for some x in (a, b) then the system (4) has only the trivial solution $c_1 = c_2 = \cdots = c_n = 0$.

Wronskian of $\{y_1, y_2, \dots, y_n\}$.

For a fixed x , let

$$W(x) = \begin{vmatrix} y_1(x) & y_2(x) & \cdots & y_n(x) \\ y_1'(x) & y_2'(x) & \cdots & y_n'(x) \\ \vdots & \vdots & \ddots & \vdots \\ y_1^{(n-1)}(x) & y_2^{(n-1)}(x) & \cdots & y_n^{(n-1)}(x) \end{vmatrix}.$$

We call this determinant the Wronskian of $\{y_1, y_2, \dots, y_n\}$.

If $W(x) \neq 0$ for some x in (a, b) then the system (4) has only the trivial solution $c_1 = c_2 = \cdots = c_n = 0$.

Thus using previous theorem we conclude that

$$y = c_1 y_1 + c_2 y_2 + \cdots + c_n y_n$$

is the general solution of $Ly = 0$ on (a, b) .

Wronskian of $\{y_1, y_2, \dots, y_n\}$.

The Wronskian of $\{y_1, y_2, \dots, y_n\}$ is

$$W(y_1, y_2, \dots, y_n; x) = \begin{vmatrix} y_1(x) & y_2(x) & \cdots & y_n(x) \\ y_1'(x) & y_2'(x) & \cdots & y_n'(x) \\ \vdots & \vdots & \ddots & \vdots \\ y_1^{(n-1)}(x) & y_2^{(n-1)}(x) & \cdots & y_n^{(n-1)}(x) \end{vmatrix}.$$

Wronskian of $\{y_1, y_2, \dots, y_n\}$.

The Wronskian of $\{y_1, y_2, \dots, y_n\}$ is

$$W(y_1, y_2, \dots, y_n; x) = \begin{vmatrix} y_1(x) & y_2(x) & \cdots & y_n(x) \\ y_1'(x) & y_2'(x) & \cdots & y_n'(x) \\ \vdots & \vdots & \ddots & \vdots \\ y_1^{(n-1)}(x) & y_2^{(n-1)}(x) & \cdots & y_n^{(n-1)}(x) \end{vmatrix}.$$

Equivalent formulation

$$W(y_1, y_2, \dots, y_n; x) = W(y_1, y_2, \dots, y_n; x_0) \cdot$$

Wronskian of $\{y_1, y_2, \dots, y_n\}$.

The Wronskian of $\{y_1, y_2, \dots, y_n\}$ is

$$W(y_1, y_2, \dots, y_n; x) = \begin{vmatrix} y_1(x) & y_2(x) & \cdots & y_n(x) \\ y_1'(x) & y_2'(x) & \cdots & y_n'(x) \\ \vdots & \vdots & \ddots & \vdots \\ y_1^{(n-1)}(x) & y_2^{(n-1)}(x) & \cdots & y_n^{(n-1)}(x) \end{vmatrix}.$$

Equivalent formulation

$$W(y_1, y_2, \dots, y_n; x) = W(y_1, y_2, \dots, y_n; x_0) \cdot \text{????}$$

Wronskian of $\{y_1, y_2, \dots, y_n\}$.

Theorem Suppose the homogeneous linear n th order equation

$$P_0(x)y^{(n)} + P_1(x)y^{(n-1)} + \dots + P_n(x)y = 0 \quad (5)$$

is normal on (a, b) , let y_1, y_2, \dots, y_n be solutions of (5) on (a, b) , and let x_0 be in (a, b) .

Wronskian of $\{y_1, y_2, \dots, y_n\}$.

Theorem Suppose the homogeneous linear n th order equation

$$P_0(x)y^{(n)} + P_1(x)y^{n-1} + \dots + P_n(x)y = 0 \quad (5)$$

is normal on (a, b) , let y_1, y_2, \dots, y_n be solutions of (5) on (a, b) , and let x_0 be in (a, b) .

Then the Wronskian of $\{y_1, y_2, \dots, y_n\}$ is given by

Wronskian of $\{y_1, y_2, \dots, y_n\}$.

Theorem Suppose the homogeneous linear n th order equation

$$P_0(x)y^{(n)} + P_1(x)y^{n-1} + \dots + P_n(x)y = 0 \quad (5)$$

is normal on (a, b) , let y_1, y_2, \dots, y_n be solutions of (5) on (a, b) , and let x_0 be in (a, b) .

Then the Wronskian of $\{y_1, y_2, \dots, y_n\}$ is given by

$$W(x) = W(x_0)$$

Wronskian of $\{y_1, y_2, \dots, y_n\}$.

Theorem Suppose the homogeneous linear n th order equation

$$P_0(x)y^{(n)} + P_1(x)y^{n-1} + \dots + P_n(x)y = 0 \quad (5)$$

is normal on (a, b) , let y_1, y_2, \dots, y_n be solutions of (5) on (a, b) , and let x_0 be in (a, b) .

Then the Wronskian of $\{y_1, y_2, \dots, y_n\}$ is given by

$$W(x) = W(x_0) \exp \left\{ - \int_{x_0}^x \frac{P_1(t)}{P_0(t)} dt \right\}, \quad a < x < b. \quad (6)$$

Wronskian of $\{y_1, y_2, \dots, y_n\}$.

Theorem Suppose the homogeneous linear n th order equation

$$P_0(x)y^{(n)} + P_1(x)y^{n-1} + \dots + P_n(x)y = 0 \quad (5)$$

is normal on (a, b) , let y_1, y_2, \dots, y_n be solutions of (5) on (a, b) , and let x_0 be in (a, b) .

Then the Wronskian of $\{y_1, y_2, \dots, y_n\}$ is given by

$$W(x) = W(x_0) \exp \left\{ - \int_{x_0}^x \frac{P_1(t)}{P_0(t)} dt \right\}, \quad a < x < b. \quad (6)$$

Therefore, either W has no zeros in (a, b) or $W \equiv 0$ on (a, b) .

Wronskian of $\{y_1, y_2, \dots, y_n\}$.

Theorem Suppose the homogeneous linear n th order equation

$$P_0(x)y^{(n)} + P_1(x)y^{n-1} + \dots + P_n(x)y = 0 \quad (5)$$

is normal on (a, b) , let y_1, y_2, \dots, y_n be solutions of (5) on (a, b) , and let x_0 be in (a, b) .

Then the Wronskian of $\{y_1, y_2, \dots, y_n\}$ is given by

$$W(x) = W(x_0) \exp \left\{ - \int_{x_0}^x \frac{P_1(t)}{P_0(t)} dt \right\}, \quad a < x < b. \quad (6)$$

Therefore, either W has no zeros in (a, b) or $W \equiv 0$ on (a, b) .

Proof Similar to homogeneous linear 2nd order ODE.

Wronskian of $\{y_1, y_2, \dots, y_n\}$.

Theorem Suppose the homogeneous linear n th order equation

$$P_0(x)y^{(n)} + P_1(x)y^{n-1} + \dots + P_n(x)y = 0 \quad (5)$$

is normal on (a, b) , let y_1, y_2, \dots, y_n be solutions of (5) on (a, b) , and let x_0 be in (a, b) .

Then the Wronskian of $\{y_1, y_2, \dots, y_n\}$ is given by

$$W(x) = W(x_0) \exp \left\{ - \int_{x_0}^x \frac{P_1(t)}{P_0(t)} dt \right\}, \quad a < x < b. \quad (6)$$

Therefore, either W has no zeros in (a, b) or $W \equiv 0$ on (a, b) .

Proof Similar to homogeneous linear 2nd order ODE.

Compare this with homogeneous linear 2nd order ODE.

Subsection 2

Particular solution

Particular solution.

Theorem Suppose $Ly = F$ is normal on (a, b) . Let y_p be a particular solution of $Ly = F$ on (a, b) , and let $\{y_1, y_2, \dots, y_n\}$ be a fundamental set of solutions of the complementary equation $Ly = 0$ on (a, b) .

Particular solution.

Theorem Suppose $Ly = F$ is normal on (a, b) . Let y_p be a particular solution of $Ly = F$ on (a, b) , and let $\{y_1, y_2, \dots, y_n\}$ be a fundamental set of solutions of the complementary equation $Ly = 0$ on (a, b) .

Then y is a solution of $Ly = F$ on (a, b) if and only if

Particular solution.

Theorem Suppose $Ly = F$ is normal on (a, b) . Let y_p be a particular solution of $Ly = F$ on (a, b) , and let $\{y_1, y_2, \dots, y_n\}$ be a fundamental set of solutions of the complementary equation $Ly = 0$ on (a, b) .

Then y is a solution of $Ly = F$ on (a, b) if and only if

$$y = y_p + c_1y_1 + c_2y_2 + \cdots + c_ny_n,$$

where c_1, c_2, \dots, c_n are constants.

Particular solution.

Theorem Suppose $Ly = F$ is normal on (a, b) . Let y_p be a particular solution of $Ly = F$ on (a, b) , and let $\{y_1, y_2, \dots, y_n\}$ be a fundamental set of solutions of the complementary equation $Ly = 0$ on (a, b) .

Then y is a solution of $Ly = F$ on (a, b) if and only if

$$y = y_p + c_1y_1 + c_2y_2 + \cdots + c_ny_n,$$

where c_1, c_2, \dots, c_n are constants.

Theorem[The Principle of Superposition] Suppose for each $i = 1, 2, \dots, r$, the function y_{p_i} is a particular solution of $Ly = F_i$ on (a, b) .

Particular solution.

Theorem Suppose $Ly = F$ is normal on (a, b) . Let y_p be a particular solution of $Ly = F$ on (a, b) , and let $\{y_1, y_2, \dots, y_n\}$ be a fundamental set of solutions of the complementary equation $Ly = 0$ on (a, b) .

Then y is a solution of $Ly = F$ on (a, b) if and only if

$$y = y_p + c_1y_1 + c_2y_2 + \cdots + c_ny_n,$$

where c_1, c_2, \dots, c_n are constants.

Theorem[The Principle of Superposition] Suppose for each $i = 1, 2, \dots, r$, the function y_{p_i} is a particular solution of $Ly = F_i$ on (a, b) .

Then

$$y_p = y_{p_1} + y_{p_2} + \cdots + y_{p_r}$$

is a particular solution on (a, b) of

$$Ly = F_1(x) + F_2(x) + \cdots + F_r(x).$$

Particular solution.

Theorem Suppose $Ly = F$ is normal on (a, b) . Let y_p be a particular solution of $Ly = F$ on (a, b) , and let $\{y_1, y_2, \dots, y_n\}$ be a fundamental set of solutions of the complementary equation $Ly = 0$ on (a, b) .

Then y is a solution of $Ly = F$ on (a, b) if and only if

$$y = y_p + c_1y_1 + c_2y_2 + \cdots + c_ny_n,$$

where c_1, c_2, \dots, c_n are constants.

Theorem[The Principle of Superposition] Suppose for each $i = 1, 2, \dots, r$, the function y_{p_i} is a particular solution of $Ly = F_i$ on (a, b) .

Then

$$y_p = y_{p_1} + y_{p_2} + \cdots + y_{p_r}$$

is a particular solution on (a, b) of

$$Ly = F_1(x) + F_2(x) + \cdots + F_r(x).$$

Compare these with homogeneous linear 2nd order ODE.

Subsection 3

Higher order constant coefficient equations

Higher order constant coefficient equations

If a_0, a_1, \dots, a_n are constants and $a_0 \neq 0$, then

$$a_0 y^{(n)} + a_1 y^{(n-1)} + \dots + a_n y = 0. \quad (7)$$

is said to be a *higher order constant coefficient equation*.

Higher order constant coefficient equations

If a_0, a_1, \dots, a_n are constants and $a_0 \neq 0$, then

$$a_0 y^{(n)} + a_1 y^{(n-1)} + \dots + a_n y = 0. \quad (7)$$

is said to be a *higher order constant coefficient equation*.

We may take open interval to be $(a, b) = (-\infty, \infty)$.

Higher order constant coefficient equations

If a_0, a_1, \dots, a_n are constants and $a_0 \neq 0$, then

$$a_0 y^{(n)} + a_1 y^{(n-1)} + \dots + a_n y = 0. \quad (7)$$

is said to be a *higher order constant coefficient equation*.

We may take open interval to be $(a, b) = (-\infty, \infty)$.

Similar to 2nd order ODE: we call

$$p(m) = a_0 m^n + a_1 m^{n-1} + \dots + a_n \quad (8)$$

Higher order constant coefficient equations

If a_0, a_1, \dots, a_n are constants and $a_0 \neq 0$, then

$$a_0 y^{(n)} + a_1 y^{(n-1)} + \dots + a_n y = 0. \quad (7)$$

is said to be a *higher order constant coefficient equation*.

We may take open interval to be $(a, b) = (-\infty, \infty)$.

Similar to 2nd order ODE: we call

$$p(m) = a_0 m^n + a_1 m^{n-1} + \dots + a_n \quad (8)$$

the *characteristic polynomial* of (7).

Higher order constant coefficient equations

If k is a positive integer, let D^k stand for the k -th derivative operator; that is

$$D^k y = y^{(k)}.$$

Higher order constant coefficient equations

If k is a positive integer, let D^k stand for the k -th derivative operator; that is

$$D^k y = y^{(k)}.$$

If

$$q(m) = b_0 m^n + b_1 m^{n-1} + \cdots + b_n$$

is an arbitrary polynomial,

Higher order constant coefficient equations

If k is a positive integer, let D^k stand for the k -th derivative operator; that is

$$D^k y = y^{(k)}.$$

If

$$q(m) = b_0 m^n + b_1 m^{n-1} + \cdots + b_n$$

is an arbitrary polynomial, define the operator

$$q(D) = b_0 D^n + b_1 D^{n-1} + \cdots + b_n$$

such that

Higher order constant coefficient equations

If k is a positive integer, let D^k stand for the k -th derivative operator; that is

$$D^k y = y^{(k)}.$$

If

$$q(m) = b_0 m^n + b_1 m^{n-1} + \cdots + b_n$$

is an arbitrary polynomial, define the operator

$$q(D) = b_0 D^n + b_1 D^{n-1} + \cdots + b_n$$

such that

$$q(D)y = (b_0 D^n + b_1 D^{n-1} + \cdots + b_n)y = b_0 y^{(n)} + b_1 y^{(n-1)} + \cdots + b_n y$$

Higher order constant coefficient equations

If k is a positive integer, let D^k stand for the k -th derivative operator; that is

$$D^k y = y^{(k)}.$$

If

$$q(m) = b_0 m^n + b_1 m^{n-1} + \cdots + b_n$$

is an arbitrary polynomial, define the operator

$$q(D) = b_0 D^n + b_1 D^{n-1} + \cdots + b_n$$

such that

$$q(D)y = (b_0 D^n + b_1 D^{n-1} + \cdots + b_n)y = b_0 y^{(n)} + b_1 y^{(n-1)} + \cdots + b_n y$$

whenever y is a function with n derivatives.

Higher order constant coefficient equations

If k is a positive integer, let D^k stand for the k -th derivative operator; that is

$$D^k y = y^{(k)}.$$

If

$$q(m) = b_0 m^n + b_1 m^{n-1} + \cdots + b_n$$

is an arbitrary polynomial, define the operator

$$q(D) = b_0 D^n + b_1 D^{n-1} + \cdots + b_n$$

such that

$$q(D)y = (b_0 D^n + b_1 D^{n-1} + \cdots + b_n)y = b_0 y^{(n)} + b_1 y^{(n-1)} + \cdots + b_n y$$

whenever y is a function with n derivatives.

We call $q(D)$ a *polynomial* operator.

Higher order constant coefficient equations

Recall $p(m) = a_0 m^n + a_1 m^{n-1} + \cdots + a_n$ is the *characteristic polynomial*, we have

$$p(D) = a_0 D^n + a_1 D^{n-1} + \cdots + a_n,$$

Higher order constant coefficient equations

Recall $p(m) = a_0 m^n + a_1 m^{n-1} + \cdots + a_n$ is the *characteristic polynomial*, we have

$$p(D) = a_0 D^n + a_1 D^{n-1} + \cdots + a_n,$$

Hence

$$a_0 y^{(n)} + a_1 y^{(n-1)} + \cdots + a_n y = 0. \tag{9}$$

can be written as $p(D)y = 0$.

Higher order constant coefficient equations

Recall $p(m) = a_0m^n + a_1m^{n-1} + \cdots + a_n$ is the *characteristic polynomial*, we have

$$p(D) = a_0D^n + a_1D^{n-1} + \cdots + a_n,$$

Hence

$$a_0y^{(n)} + a_1y^{(n-1)} + \cdots + a_ny = 0. \quad (9)$$

can be written as $p(D)y = 0$.

If m is a constant then

$$\begin{aligned} p(D)e^{mx} &= (a_0D^n e^{mx} + a_1D^{n-1} e^{mx} + \cdots + a_n e^{mx}) \\ &= (a_0m^n + a_1m^{n-1} + \cdots + a_n)e^{mx}; \end{aligned}$$

Higher order constant coefficient equations

Recall $p(m) = a_0 m^n + a_1 m^{n-1} + \cdots + a_n$ is the *characteristic polynomial*, we have

$$p(D) = a_0 D^n + a_1 D^{n-1} + \cdots + a_n,$$

Hence

$$a_0 y^{(n)} + a_1 y^{(n-1)} + \cdots + a_n y = 0. \quad (9)$$

can be written as $p(D)y = 0$.

If m is a constant then

$$\begin{aligned} p(D)e^{mx} &= (a_0 D^n e^{mx} + a_1 D^{n-1} e^{mx} + \cdots + a_n e^{mx}) \\ &= (a_0 m^n + a_1 m^{n-1} + \cdots + a_n) e^{mx}; \end{aligned}$$

that is

$$p(D)(e^{mx}) = p(m)e^{mx}.$$

Higher order constant coefficient equations

Recall $p(m) = a_0 m^n + a_1 m^{n-1} + \cdots + a_n$ is the *characteristic polynomial*, we have

$$p(D) = a_0 D^n + a_1 D^{n-1} + \cdots + a_n,$$

Hence

$$a_0 y^{(n)} + a_1 y^{(n-1)} + \cdots + a_n y = 0. \quad (9)$$

can be written as $p(D)y = 0$.

If m is a constant then

$$\begin{aligned} p(D)e^{mx} &= (a_0 D^n e^{mx} + a_1 D^{n-1} e^{mx} + \cdots + a_n e^{mx}) \\ &= (a_0 m^n + a_1 m^{n-1} + \cdots + a_n) e^{mx}; \end{aligned}$$

that is

$$p(D)(e^{mx}) = p(m)e^{mx}.$$

This shows that $y = e^{mx}$ is a solution of (11)

Higher order constant coefficient equations

Recall $p(m) = a_0 m^n + a_1 m^{n-1} + \cdots + a_n$ is the *characteristic polynomial*, we have

$$p(D) = a_0 D^n + a_1 D^{n-1} + \cdots + a_n,$$

Hence

$$a_0 y^{(n)} + a_1 y^{(n-1)} + \cdots + a_n y = 0. \quad (9)$$

can be written as $p(D)y = 0$.

If m is a constant then

$$\begin{aligned} p(D)e^{mx} &= (a_0 D^n e^{mx} + a_1 D^{n-1} e^{mx} + \cdots + a_n e^{mx}) \\ &= (a_0 m^n + a_1 m^{n-1} + \cdots + a_n) e^{mx}; \end{aligned}$$

that is

$$p(D)(e^{mx}) = p(m)e^{mx}.$$

This shows that $y = e^{mx}$ is a solution of (11) if $p(m) = 0$.

Higher order constant coefficient equations

Example Find the general solution of

$$y''' - 6y'' + 11y' - 6y = 0. \quad (10)$$

Higher order constant coefficient equations

Example Find the general solution of

$$y''' - 6y'' + 11y' - 6y = 0. \quad (10)$$

Solution

The characteristic polynomial is

$$p(m) = m^3 - 6m^2 + 11m - 6 = (m - 1)(m - 2)(m - 3).$$

Higher order constant coefficient equations

Example Find the general solution of

$$y''' - 6y'' + 11y' - 6y = 0. \quad (10)$$

Solution

The characteristic polynomial is

$$p(m) = m^3 - 6m^2 + 11m - 6 = (m - 1)(m - 2)(m - 3).$$

Therefore $\{e^x, e^{2x}, e^{3x}\}$ is a set of solutions.

Higher order constant coefficient equations

Example Find the general solution of

$$y''' - 6y'' + 11y' - 6y = 0. \quad (10)$$

Solution

The characteristic polynomial is

$$p(m) = m^3 - 6m^2 + 11m - 6 = (m - 1)(m - 2)(m - 3).$$

Therefore $\{e^x, e^{2x}, e^{3x}\}$ is a set of solutions.

$$W(x) = \begin{vmatrix} e^x & e^{2x} & e^{3x} \\ e^x & 2e^{2x} & 3e^{3x} \\ e^x & 4e^{2x} & 9e^{3x} \end{vmatrix} =$$

Higher order constant coefficient equations

Example Find the general solution of

$$y''' - 6y'' + 11y' - 6y = 0. \quad (10)$$

Solution

The characteristic polynomial is

$$p(m) = m^3 - 6m^2 + 11m - 6 = (m - 1)(m - 2)(m - 3).$$

Therefore $\{e^x, e^{2x}, e^{3x}\}$ is a set of solutions.

$$W(x) = \begin{vmatrix} e^x & e^{2x} & e^{3x} \\ e^x & 2e^{2x} & 3e^{3x} \\ e^x & 4e^{2x} & 9e^{3x} \end{vmatrix} = e^{6x} \begin{vmatrix} 1 & 1 & 1 \\ 1 & 2 & 3 \\ 1 & 4 & 9 \end{vmatrix} =$$

Higher order constant coefficient equations

Example Find the general solution of

$$y''' - 6y'' + 11y' - 6y = 0. \quad (10)$$

Solution

The characteristic polynomial is

$$p(m) = m^3 - 6m^2 + 11m - 6 = (m - 1)(m - 2)(m - 3).$$

Therefore $\{e^x, e^{2x}, e^{3x}\}$ is a set of solutions.

$$W(x) = \begin{vmatrix} e^x & e^{2x} & e^{3x} \\ e^x & 2e^{2x} & 3e^{3x} \\ e^x & 4e^{2x} & 9e^{3x} \end{vmatrix} = e^{6x} \begin{vmatrix} 1 & 1 & 1 \\ 1 & 2 & 3 \\ 1 & 4 & 9 \end{vmatrix} = 2e^{6x} \neq 0.$$

Higher order constant coefficient equations

Example Find the general solution of

$$y''' - 6y'' + 11y' - 6y = 0. \quad (10)$$

Solution

The characteristic polynomial is

$$p(m) = m^3 - 6m^2 + 11m - 6 = (m - 1)(m - 2)(m - 3).$$

Therefore $\{e^x, e^{2x}, e^{3x}\}$ is a set of solutions.

$$W(x) = \begin{vmatrix} e^x & e^{2x} & e^{3x} \\ e^x & 2e^{2x} & 3e^{3x} \\ e^x & 4e^{2x} & 9e^{3x} \end{vmatrix} = e^{6x} \begin{vmatrix} 1 & 1 & 1 \\ 1 & 2 & 3 \\ 1 & 4 & 9 \end{vmatrix} = 2e^{6x} \neq 0.$$

Therefore the general solution is

$$y = c_1 e^x + c_2 e^{2x} + c_3 e^{3x}.$$

Higher order constant coefficient equations

Example Solve the initial value problem

$$y''' - 6y'' + 11y' - 6y = 0, \quad y(0) = 4, \quad y'(0) = 5, \quad y''(0) = 9. \quad (11)$$

Higher order constant coefficient equations

Example Solve the initial value problem

$$y''' - 6y'' + 11y' - 6y = 0, \quad y(0) = 4, \quad y'(0) = 5, \quad y''(0) = 9. \quad (11)$$

Solution Differentiating $y = c_1 e^x + c_2 e^{2x} + c_3 e^{3x}$ two times gives

$$\begin{aligned} y' &= c_1 e^x + 2c_2 e^{2x} + 3c_3 e^{3x} \\ y'' &= c_1 e^x + 4c_2 e^{2x} + 9c_3 e^{3x}. \end{aligned} \quad (12)$$

Higher order constant coefficient equations

Example Solve the initial value problem

$$y''' - 6y'' + 11y' - 6y = 0, \quad y(0) = 4, \quad y'(0) = 5, \quad y''(0) = 9. \quad (11)$$

Solution Differentiating $y = c_1 e^x + c_2 e^{2x} + c_3 e^{3x}$ two times gives

$$\begin{aligned} y' &= c_1 e^x + 2c_2 e^{2x} + 3c_3 e^{3x} \\ y'' &= c_1 e^x + 4c_2 e^{2x} + 9c_3 e^{3x}. \end{aligned} \quad (12)$$

Set $x = 0$ and using the initial conditions,

Higher order constant coefficient equations

Example Solve the initial value problem

$$y''' - 6y'' + 11y' - 6y = 0, \quad y(0) = 4, \quad y'(0) = 5, \quad y''(0) = 9. \quad (11)$$

Solution Differentiating $y = c_1 e^x + c_2 e^{2x} + c_3 e^{3x}$ two times gives

$$\begin{aligned} y' &= c_1 e^x + 2c_2 e^{2x} + 3c_3 e^{3x} \\ y'' &= c_1 e^x + 4c_2 e^{2x} + 9c_3 e^{3x}. \end{aligned} \quad (12)$$

Set $x = 0$ and using the initial conditions, to get $c_1 = 4$, $c_2 = -1$, $c_3 = 1$.

Higher order constant coefficient equations

Example Solve the initial value problem

$$y''' - 6y'' + 11y' - 6y = 0, \quad y(0) = 4, \quad y'(0) = 5, \quad y''(0) = 9. \quad (11)$$

Solution Differentiating $y = c_1 e^x + c_2 e^{2x} + c_3 e^{3x}$ two times gives

$$\begin{aligned} y' &= c_1 e^x + 2c_2 e^{2x} + 3c_3 e^{3x} \\ y'' &= c_1 e^x + 4c_2 e^{2x} + 9c_3 e^{3x}. \end{aligned} \quad (12)$$

Set $x = 0$ and using the initial conditions, to get $c_1 = 4$, $c_2 = -1$, $c_3 = 1$.

Therefore solution to IVP is

$$y = 4e^x - e^{2x} + e^{3x}.$$

Higher order constant coefficient equations

Example find the general solution of

$$y''' - y'' + y' - y = 0. \quad (13)$$

Higher order constant coefficient equations

Example find the general solution of

$$y''' - y'' + y' - y = 0. \quad (13)$$

Solution

The characteristic polynomial is

$$p(m) = m^3 - m^2 + m - 1 = (m - 1)(m^2 + 1).$$

Higher order constant coefficient equations

Example find the general solution of

$$y''' - y'' + y' - y = 0. \quad (13)$$

Solution

The characteristic polynomial is

$$p(m) = m^3 - m^2 + m - 1 = (m - 1)(m^2 + 1).$$

Rewrite (13) as

$$(D - 1)(D^2 + 1)y = 0,$$

Higher order constant coefficient equations

Example find the general solution of

$$y''' - y'' + y' - y = 0. \quad (13)$$

Solution

The characteristic polynomial is

$$p(m) = m^3 - m^2 + m - 1 = (m - 1)(m^2 + 1).$$

Rewrite (13) as

$$(D - 1)(D^2 + 1)y = 0,$$

which implies that any solution of $(D^2 + 1)y = 0$ is a solution of (13).

Higher order constant coefficient equations

Example find the general solution of

$$y''' - y'' + y' - y = 0. \quad (13)$$

Solution

The characteristic polynomial is

$$p(m) = m^3 - m^2 + m - 1 = (m - 1)(m^2 + 1).$$

Rewrite (13) as

$$(D - 1)(D^2 + 1)y = 0,$$

which implies that any solution of $(D^2 + 1)y = 0$ is a solution of (13).

Therefore $y_1 = \cos x$ and $y_2 = \sin x$ are solutions of (13).

Higher order constant coefficient equations

Example find the general solution of

$$y''' - y'' + y' - y = 0. \quad (13)$$

Solution

The characteristic polynomial is

$$p(m) = m^3 - m^2 + m - 1 = (m - 1)(m^2 + 1).$$

Rewrite (13) as

$$(D - 1)(D^2 + 1)y = 0,$$

which implies that any solution of $(D^2 + 1)y = 0$ is a solution of (13).

Therefore $y_1 = \cos x$ and $y_2 = \sin x$ are solutions of (13).

Now rewrite (13) as

$$(D^2 + 1)(D - 1)y = 0,$$

which implies that any solution of $(D - 1)y = 0$ is a solution of (13).

Higher order constant coefficient equations

Example find the general solution of

$$y''' - y'' + y' - y = 0. \quad (13)$$

Solution

The characteristic polynomial is

$$p(m) = m^3 - m^2 + m - 1 = (m - 1)(m^2 + 1).$$

Rewrite (13) as

$$(D - 1)(D^2 + 1)y = 0,$$

which implies that any solution of $(D^2 + 1)y = 0$ is a solution of (13).

Therefore $y_1 = \cos x$ and $y_2 = \sin x$ are solutions of (13).

Now rewrite (13) as

$$(D^2 + 1)(D - 1)y = 0,$$

which implies that any solution of $(D - 1)y = 0$ is a solution of (13).

Therefore $y_3 = e^x$ is solution of (13).

Higher order constant coefficient equations

Example (continued...)

Example (continued...)

The Wronskian of $\{e^x, \cos x, \sin x\}$ is

$$W(x) = \begin{vmatrix} \cos x & \sin x & e^x \\ -\sin x & \cos x & e^x \\ -\cos x & -\sin x & e^x \end{vmatrix}.$$

Higher order constant coefficient equations

Example (continued...)

The Wronskian of $\{e^x, \cos x, \sin x\}$ is

$$W(x) = \begin{vmatrix} \cos x & \sin x & e^x \\ -\sin x & \cos x & e^x \\ -\cos x & -\sin x & e^x \end{vmatrix}.$$

Note that

$$W(0) = \begin{vmatrix} 1 & 0 & 1 \\ 0 & 1 & 1 \\ -1 & 0 & 1 \end{vmatrix} =$$

Example (continued...)

The Wronskian of $\{e^x, \cos x, \sin x\}$ is

$$W(x) = \begin{vmatrix} \cos x & \sin x & e^x \\ -\sin x & \cos x & e^x \\ -\cos x & -\sin x & e^x \end{vmatrix}.$$

Note that

$$W(0) = \begin{vmatrix} 1 & 0 & 1 \\ 0 & 1 & 1 \\ -1 & 0 & 1 \end{vmatrix} = 2,$$

Hence $\{\cos x, \sin x, e^x\}$ is linearly independent and

$$y = c_1 \cos x + c_2 \sin x + c_3 e^x$$

is the general solution of (13).

Higher order constant coefficient equations

Example find the general solution of

$$y^{(4)} - 16y = 0. \quad (14)$$

Higher order constant coefficient equations

Example find the general solution of

$$y^{(4)} - 16y = 0. \quad (14)$$

Solution

Higher order constant coefficient equations

Example find the general solution of

$$y^{(4)} - 16y = 0. \quad (14)$$

Solution

The characteristic polynomial is

$$p(m) = m^4 - 16 = (m^2 - 4)(m^2 + 4) = (m - 2)(m + 2)(m^2 + 4)$$

Higher order constant coefficient equations

Example find the general solution of

$$y^{(4)} - 16y = 0. \quad (14)$$

Solution

The characteristic polynomial is

$$p(m) = m^4 - 16 = (m^2 - 4)(m^2 + 4) = (m - 2)(m + 2)(m^2 + 4)$$

it can be written as

$$(D^2 + 4)(D + 2)(D - 2)y = 0$$

Higher order constant coefficient equations

Example find the general solution of

$$y^{(4)} - 16y = 0. \quad (14)$$

Solution

The characteristic polynomial is

$$p(m) = m^4 - 16 = (m^2 - 4)(m^2 + 4) = (m - 2)(m + 2)(m^2 + 4)$$

it can be written as

$$(D^2 + 4)(D + 2)(D - 2)y = 0$$

or

$$(D^2 + 4)(D - 2)(D + 2)y = 0$$

Higher order constant coefficient equations

Example find the general solution of

$$y^{(4)} - 16y = 0. \quad (14)$$

Solution

The characteristic polynomial is

$$p(m) = m^4 - 16 = (m^2 - 4)(m^2 + 4) = (m - 2)(m + 2)(m^2 + 4)$$

it can be written as

$$(D^2 + 4)(D + 2)(D - 2)y = 0$$

or

$$(D^2 + 4)(D - 2)(D + 2)y = 0$$

or

$$(D - 2)(D + 2)(D^2 + 4)y = 0.$$

Higher order constant coefficient equations

Example find the general solution of

$$y^{(4)} - 16y = 0. \quad (14)$$

Solution

The characteristic polynomial is

$$p(m) = m^4 - 16 = (m^2 - 4)(m^2 + 4) = (m - 2)(m + 2)(m^2 + 4)$$

it can be written as

$$(D^2 + 4)(D + 2)(D - 2)y = 0$$

or

$$(D^2 + 4)(D - 2)(D + 2)y = 0$$

or

$$(D - 2)(D + 2)(D^2 + 4)y = 0.$$

Therefore y is a solution if it is a solution of any of the three equations

$$(D - 2)y = 0, \quad (D + 2)y = 0, \quad (D^2 + 4)y = 0.$$

Higher order constant coefficient equations

Hence, $\{e^{2x}, e^{-2x}, \cos 2x, \sin 2x\}$ is a set of solutions.

Higher order constant coefficient equations

Hence, $\{e^{2x}, e^{-2x}, \cos 2x, \sin 2x\}$ is a set of solutions.

$$W(x) =$$

Higher order constant coefficient equations

Hence, $\{e^{2x}, e^{-2x}, \cos 2x, \sin 2x\}$ is a set of solutions.

$$W(x) = \begin{vmatrix} e^{2x} & e^{-2x} & \cos 2x & \sin 2x \end{vmatrix}$$

Higher order constant coefficient equations

Hence, $\{e^{2x}, e^{-2x}, \cos 2x, \sin 2x\}$ is a set of solutions.

$$W(x) = \begin{vmatrix} e^{2x} & e^{-2x} & \cos 2x & \sin 2x \\ 2e^{2x} & -2e^{-2x} & -2\sin 2x & 2\cos 2x \end{vmatrix}$$

Higher order constant coefficient equations

Hence, $\{e^{2x}, e^{-2x}, \cos 2x, \sin 2x\}$ is a set of solutions.

$$W(x) = \begin{vmatrix} e^{2x} & e^{-2x} & \cos 2x & \sin 2x \\ 2e^{2x} & -2e^{-2x} & -2\sin 2x & 2\cos 2x \\ 4e^{2x} & 4e^{-2x} & -4\cos 2x & -4\sin 2x \end{vmatrix}$$

Higher order constant coefficient equations

Hence, $\{e^{2x}, e^{-2x}, \cos 2x, \sin 2x\}$ is a set of solutions.

$$W(x) = \begin{vmatrix} e^{2x} & e^{-2x} & \cos 2x & \sin 2x \\ 2e^{2x} & -2e^{-2x} & -2\sin 2x & 2\cos 2x \\ 4e^{2x} & 4e^{-2x} & -4\cos 2x & -4\sin 2x \\ 8e^{2x} & -8e^{-2x} & 8\sin 2x & -8\cos 2x \end{vmatrix}.$$

Higher order constant coefficient equations

Hence, $\{e^{2x}, e^{-2x}, \cos 2x, \sin 2x\}$ is a set of solutions.

$$W(x) = \begin{vmatrix} e^{2x} & e^{-2x} & \cos 2x & \sin 2x \\ 2e^{2x} & -2e^{-2x} & -2\sin 2x & 2\cos 2x \\ 4e^{2x} & 4e^{-2x} & -4\cos 2x & -4\sin 2x \\ 8e^{2x} & -8e^{-2x} & 8\sin 2x & -8\cos 2x \end{vmatrix}.$$

Since

$$W(0) =$$

Higher order constant coefficient equations

Hence, $\{e^{2x}, e^{-2x}, \cos 2x, \sin 2x\}$ is a set of solutions.

$$W(x) = \begin{vmatrix} e^{2x} & e^{-2x} & \cos 2x & \sin 2x \\ 2e^{2x} & -2e^{-2x} & -2\sin 2x & 2\cos 2x \\ 4e^{2x} & 4e^{-2x} & -4\cos 2x & -4\sin 2x \\ 8e^{2x} & -8e^{-2x} & 8\sin 2x & -8\cos 2x \end{vmatrix}.$$

Since

$$W(0) = \begin{vmatrix} 1 & 1 & 1 & 0 \\ 2 & -2 & 0 & 2 \\ 4 & 4 & -4 & 0 \\ 8 & -8 & 0 & -8 \end{vmatrix} =$$

Higher order constant coefficient equations

Hence, $\{e^{2x}, e^{-2x}, \cos 2x, \sin 2x\}$ is a set of solutions.

$$W(x) = \begin{vmatrix} e^{2x} & e^{-2x} & \cos 2x & \sin 2x \\ 2e^{2x} & -2e^{-2x} & -2\sin 2x & 2\cos 2x \\ 4e^{2x} & 4e^{-2x} & -4\cos 2x & -4\sin 2x \\ 8e^{2x} & -8e^{-2x} & 8\sin 2x & -8\cos 2x \end{vmatrix}.$$

Since

$$W(0) = \begin{vmatrix} 1 & 1 & 1 & 0 \\ 2 & -2 & 0 & 2 \\ 4 & 4 & -4 & 0 \\ 8 & -8 & 0 & -8 \end{vmatrix} = -512 \neq 0,$$

Higher order constant coefficient equations

Hence, $\{e^{2x}, e^{-2x}, \cos 2x, \sin 2x\}$ is a set of solutions.

$$W(x) = \begin{vmatrix} e^{2x} & e^{-2x} & \cos 2x & \sin 2x \\ 2e^{2x} & -2e^{-2x} & -2\sin 2x & 2\cos 2x \\ 4e^{2x} & 4e^{-2x} & -4\cos 2x & -4\sin 2x \\ 8e^{2x} & -8e^{-2x} & 8\sin 2x & -8\cos 2x \end{vmatrix}.$$

Since

$$W(0) = \begin{vmatrix} 1 & 1 & 1 & 0 \\ 2 & -2 & 0 & 2 \\ 4 & 4 & -4 & 0 \\ 8 & -8 & 0 & -8 \end{vmatrix} = -512 \neq 0,$$

$\{e^{2x}, e^{-2x}, \cos 2x, \sin 2x\}$ is linearly independent, and

$$y_1 = c_1 e^{2x} + c_2 e^{-2x} + c_3 \cos 2x + c_4 \sin 2x$$

is the general solution of $y^{(4)} - 16y = 0$.

Example find the general solution of

$$y''' + 3y'' + 3y' + y = 0. \quad (15)$$

Example find the general solution of

$$y''' + 3y'' + 3y' + y = 0. \quad (15)$$

Solution

Higher order constant coefficient equations

Example find the general solution of

$$y''' + 3y'' + 3y' + y = 0. \quad (15)$$

Solution

The characteristic polynomial is

$$p(m) = m^3 + 3m^2 + 3m + 1 = (m + 1)^3.$$

Higher order constant coefficient equations

Example find the general solution of

$$y''' + 3y'' + 3y' + y = 0. \quad (15)$$

Solution

The characteristic polynomial is

$$p(m) = m^3 + 3m^2 + 3m + 1 = (m + 1)^3.$$

it can be written as

$$(D + 1)^3 y = 0.$$

Higher order constant coefficient equations

Example find the general solution of

$$y''' + 3y'' + 3y' + y = 0. \quad (15)$$

Solution

The characteristic polynomial is

$$p(m) = m^3 + 3m^2 + 3m + 1 = (m + 1)^3.$$

it can be written as

$$(D + 1)^3 y = 0.$$

Therefore the general solution of (15) is

$$y = e^{-x}(c_1 + c_2x + c_3x^2).$$

Higher order constant coefficient equations

Theorem: If $\omega \neq 0$ and m is a positive integer, then

$$\{e^{\lambda x} \cos \omega x, xe^{\lambda x} \cos \omega x, \dots, x^{m-1}e^{\lambda x} \cos \omega x, \\ e^{\lambda x} \sin \omega x, xe^{\lambda x} \sin \omega x, \dots, x^{m-1}e^{\lambda x} \sin \omega x\}$$

is a fundamental set of solutions of

$$[(D - \lambda)^2 + \omega^2]^m y = 0.$$

Higher order constant coefficient equations

Theorem: If $\omega \neq 0$ and m is a positive integer, then

$$\{e^{\lambda x} \cos \omega x, xe^{\lambda x} \cos \omega x, \dots, x^{m-1}e^{\lambda x} \cos \omega x, \\ e^{\lambda x} \sin \omega x, xe^{\lambda x} \sin \omega x, \dots, x^{m-1}e^{\lambda x} \sin \omega x\}$$

is a fundamental set of solutions of

$$[(D - \lambda)^2 + \omega^2]^m y = 0.$$

Example: Find the general solution of

$$(D^2 + 4D + 13)^3 y = 0. \tag{16}$$

Higher order constant coefficient equations

Theorem: If $\omega \neq 0$ and m is a positive integer, then

$$\left\{ e^{\lambda x} \cos \omega x, x e^{\lambda x} \cos \omega x, \dots, x^{m-1} e^{\lambda x} \cos \omega x, \right. \\ \left. e^{\lambda x} \sin \omega x, x e^{\lambda x} \sin \omega x, \dots, x^{m-1} e^{\lambda x} \sin \omega x \right\}$$

is a fundamental set of solutions of

$$[(D - \lambda)^2 + \omega^2]^m y = 0.$$

Example: Find the general solution of

$$(D^2 + 4D + 13)^3 y = 0. \tag{16}$$

We can re-write $D^2 + 4D + 13$ as

Higher order constant coefficient equations

Theorem: If $\omega \neq 0$ and m is a positive integer, then

$$\left\{ e^{\lambda x} \cos \omega x, x e^{\lambda x} \cos \omega x, \dots, x^{m-1} e^{\lambda x} \cos \omega x, \right. \\ \left. e^{\lambda x} \sin \omega x, x e^{\lambda x} \sin \omega x, \dots, x^{m-1} e^{\lambda x} \sin \omega x \right\}$$

is a fundamental set of solutions of

$$[(D - \lambda)^2 + \omega^2]^m y = 0.$$

Example: Find the general solution of

$$(D^2 + 4D + 13)^3 y = 0. \tag{16}$$

We can re-write $D^2 + 4D + 13$ as

$$[(D + 2)^2 + 9]^3 y = 0.$$

Higher order constant coefficient equations

Theorem: If $\omega \neq 0$ and m is a positive integer, then

$$\{e^{\lambda x} \cos \omega x, xe^{\lambda x} \cos \omega x, \dots, x^{m-1}e^{\lambda x} \cos \omega x, \\ e^{\lambda x} \sin \omega x, xe^{\lambda x} \sin \omega x, \dots, x^{m-1}e^{\lambda x} \sin \omega x\}$$

is a fundamental set of solutions of

$$[(D - \lambda)^2 + \omega^2]^m y = 0.$$

Example: Find the general solution of

$$(D^2 + 4D + 13)^3 y = 0. \tag{16}$$

We can re-write $D^2 + 4D + 13$ as

$$[(D + 2)^2 + 9]^3 y = 0.$$

Therefore the general solution would be

$$y = (c_1 + c_2x + c_3x^2)e^{-2x} \cos 3x + (c_4 + c_5x + c_6x^2)e^{-2x} \sin 3x.$$

Example: Find a particular solution of

$$y''' + 3y'' + 2y' - y = e^x(21 + 24x + 28x^2 + 5x^3). \quad (17)$$

Higher order constant coefficient equations

Example: Find a particular solution of

$$y''' + 3y'' + 2y' - y = e^x(21 + 24x + 28x^2 + 5x^3). \quad (17)$$

Solution.

Higher order constant coefficient equations

Example: Find a particular solution of

$$y''' + 3y'' + 2y' - y = e^x(21 + 24x + 28x^2 + 5x^3). \quad (17)$$

Solution. Let $y = u(x)e^x$ and

$$\begin{aligned} y &= ue^x, \\ y' &= e^x(u' + u), \end{aligned}$$

Higher order constant coefficient equations

Example: Find a particular solution of

$$y''' + 3y'' + 2y' - y = e^x(21 + 24x + 28x^2 + 5x^3). \quad (17)$$

Solution. Let $y = u(x)e^x$ and

$$\begin{aligned} y &= ue^x, \\ y' &= e^x(u' + u), \\ y'' &= e^x(u'' + 2u' + u), \end{aligned}$$

Higher order constant coefficient equations

Example: Find a particular solution of

$$y''' + 3y'' + 2y' - y = e^x(21 + 24x + 28x^2 + 5x^3). \quad (17)$$

Solution. Let $y = u(x)e^x$ and

$$\begin{aligned} y &= ue^x, \\ y' &= e^x(u' + u), \\ y'' &= e^x(u'' + 2u' + u), \\ y''' &= e^x(u''' + 3u'' + 3u' + u) \end{aligned}$$

Higher order constant coefficient equations

Example: Find a particular solution of

$$y''' + 3y'' + 2y' - y = e^x(21 + 24x + 28x^2 + 5x^3). \quad (17)$$

Solution. Let $y = u(x)e^x$ and

$$\begin{aligned} y &= ue^x, \\ y' &= e^x(u' + u), \\ y'' &= e^x(u'' + 2u' + u), \\ y''' &= e^x(u''' + 3u'' + 3u' + u) \end{aligned}$$

Higher order constant coefficient equations

Example: Find a particular solution of

$$y''' + 3y'' + 2y' - y = e^x(21 + 24x + 28x^2 + 5x^3). \quad (17)$$

Solution. Let $y = u(x)e^x$ and

$$\begin{aligned} y &= ue^x, \\ y' &= e^x(u' + u), \\ y'' &= e^x(u'' + 2u' + u), \\ y''' &= e^x(u''' + 3u'' + 3u' + u) \end{aligned}$$

Substitute into (17), and also canceling e^x gives

$$(u''' + 3u'' + 3u' + u) + 3(u'' + 2u' + u) + 2(u' + u) - u =$$

Higher order constant coefficient equations

Example: Find a particular solution of

$$y''' + 3y'' + 2y' - y = e^x(21 + 24x + 28x^2 + 5x^3). \quad (17)$$

Solution. Let $y = u(x)e^x$ and

$$\begin{aligned} y &= ue^x, \\ y' &= e^x(u' + u), \\ y'' &= e^x(u'' + 2u' + u), \\ y''' &= e^x(u''' + 3u'' + 3u' + u) \end{aligned}$$

Substitute into (17), and also canceling e^x gives

$$(u''' + 3u'' + 3u' + u) + 3(u'' + 2u' + u) + 2(u' + u) - u = 21 + 24x + 28x^2 + 5x^3,$$

Higher order constant coefficient equations

A particular solution is of the form

$$u_p = A + Bx + Cx^2 + Dx^3.$$

Higher order constant coefficient equations

A particular solution is of the form

$$u_p = A + Bx + Cx^2 + Dx^3.$$

Then

$$u'_p = B + 2Cx + 3Dx^2$$

Higher order constant coefficient equations

A particular solution is of the form

$$u_p = A + Bx + Cx^2 + Dx^3.$$

Then

$$u'_p = B + 2Cx + 3Dx^2$$

$$u''_p = 2C + 6Dx$$

Higher order constant coefficient equations

A particular solution is of the form

$$u_p = A + Bx + Cx^2 + Dx^3.$$

Then

$$\begin{aligned}u_p' &= B + 2Cx + 3Dx^2 \\u_p'' &= 2C + 6Dx \\u_p''' &= 6D.\end{aligned}$$

Higher order constant coefficient equations

A particular solution is of the form

$$u_p = A + Bx + Cx^2 + Dx^3.$$

Then

$$\begin{aligned}u_p' &= B + 2Cx + 3Dx^2 \\u_p'' &= 2C + 6Dx \\u_p''' &= 6D.\end{aligned}$$

Substituting this, we get

Higher order constant coefficient equations

A particular solution is of the form

$$u_p = A + Bx + Cx^2 + Dx^3.$$

Then

$$\begin{aligned}u_p' &= B + 2Cx + 3Dx^2 \\u_p'' &= 2C + 6Dx \\u_p''' &= 6D.\end{aligned}$$

Substituting this, we get

$$u_p''' + 6u_p'' + 11u_p' + 5u_p = 6D + 6(2C + 6Dx) + 11(B + 2Cx + 3Dx^2)$$

Higher order constant coefficient equations

A particular solution is of the form

$$u_p = A + Bx + Cx^2 + Dx^3.$$

Then

$$\begin{aligned}u_p' &= B + 2Cx + 3Dx^2 \\u_p'' &= 2C + 6Dx \\u_p''' &= 6D.\end{aligned}$$

Substituting this, we get

$$\begin{aligned}u_p''' + 6u_p'' + 11u_p' + 5u_p &= 6D + 6(2C + 6Dx) + 11(B + 2Cx + 3Dx^2) \\&\quad + 5(A + Bx + Cx^2 + Dx^3)\end{aligned}$$

Higher order constant coefficient equations

A particular solution is of the form

$$u_p = A + Bx + Cx^2 + Dx^3.$$

Then

$$\begin{aligned}u_p' &= B + 2Cx + 3Dx^2 \\u_p'' &= 2C + 6Dx \\u_p''' &= 6D.\end{aligned}$$

Substituting this, we get

$$\begin{aligned}u_p''' + 6u_p'' + 11u_p' + 5u_p &= 6D + 6(2C + 6Dx) + 11(B + 2Cx + 3Dx^2) \\&\quad + 5(A + Bx + Cx^2 + Dx^3) \\&= (5A + 11B + 12C + 6D) + (5B + 22C + 36D)x \\&\quad + (5C + 33D)x^2 + 5Dx^3.\end{aligned}$$

Higher order constant coefficient equations

A particular solution is of the form

$$u_p = A + Bx + Cx^2 + Dx^3.$$

Then

$$\begin{aligned}u_p' &= B + 2Cx + 3Dx^2 \\u_p'' &= 2C + 6Dx \\u_p''' &= 6D.\end{aligned}$$

Substituting this, we get

$$\begin{aligned}u_p''' + 6u_p'' + 11u_p' + 5u_p &= 6D + 6(2C + 6Dx) + 11(B + 2Cx + 3Dx^2) \\&\quad + 5(A + Bx + Cx^2 + Dx^3) \\&= (5A + 11B + 12C + 6D) + (5B + 22C + 36D)x \\&\quad + (5C + 33D)x^2 + 5Dx^3.\end{aligned}$$

Higher order constant coefficient equations

Comparing coefficients of like powers of x on both sides,

Higher order constant coefficient equations

Comparing coefficients of like powers of x on both sides,

$$5D = 5$$

Higher order constant coefficient equations

Comparing coefficients of like powers of x on both sides,

$$\begin{aligned}5D &= 5 \\ 5C + 33D &= 28\end{aligned}$$

Higher order constant coefficient equations

Comparing coefficients of like powers of x on both sides,

$$5D = 5$$

$$5C + 33D = 28$$

$$5B + 22C + 36D = 24$$

Higher order constant coefficient equations

Comparing coefficients of like powers of x on both sides,

$$5D = 5$$

$$5C + 33D = 28$$

$$5B + 22C + 36D = 24$$

$$5A + 11B + 12C + 6D = 21.$$

Higher order constant coefficient equations

Comparing coefficients of like powers of x on both sides,

$$5D = 5$$

$$5C + 33D = 28$$

$$5B + 22C + 36D = 24$$

$$5A + 11B + 12C + 6D = 21.$$

Higher order constant coefficient equations

Comparing coefficients of like powers of x on both sides,

$$5D = 5$$

$$5C + 33D = 28$$

$$5B + 22C + 36D = 24$$

$$5A + 11B + 12C + 6D = 21.$$

Solving these equations, we get

Higher order constant coefficient equations

Comparing coefficients of like powers of x on both sides,

$$5D = 5$$

$$5C + 33D = 28$$

$$5B + 22C + 36D = 24$$

$$5A + 11B + 12C + 6D = 21.$$

Solving these equations, we get

$$D = 1, C = -1, B = 2, A = 1$$

Higher order constant coefficient equations

Comparing coefficients of like powers of x on both sides,

$$\begin{aligned}5D &= 5 \\5C + 33D &= 28 \\5B + 22C + 36D &= 24 \\5A + 11B + 12C + 6D &= 21.\end{aligned}$$

Solving these equations, we get

$$D = 1, C = -1, B = 2, A = 1$$

Therefore

$$u_p = 1 + 2x - x^2 + x^3$$

Higher order constant coefficient equations

Comparing coefficients of like powers of x on both sides,

$$\begin{aligned}5D &= 5 \\5C + 33D &= 28 \\5B + 22C + 36D &= 24 \\5A + 11B + 12C + 6D &= 21.\end{aligned}$$

Solving these equations, we get

$$D = 1, C = -1, B = 2, A = 1$$

Therefore

$$u_p = 1 + 2x - x^2 + x^3$$

and hence

$$y_p = e^x u_p = e^x(1 + 2x - x^2 + x^3)$$

is a particular solution of (17)