

Differential Equations (MA 1150)

Sukumar

Lecture 6

May 01, 2020

Second order linear ODE

Fundamental solutions : Linearly independent solution

Second order ODE with Constant coefficients

Distinct real roots

Repeated real roots

Complex conjugate roots

Wronskian

Non-homogeneous second order linear ODE

Section 1

Second order linear ODE

Definition: A second order differential equation is said to be linear if it can be written as

$$y'' + p(x)y' + q(x)y = f(x). \quad (1)$$

Definition: A second order differential equation is said to be linear if it can be written as

$$y'' + p(x)y' + q(x)y = f(x). \quad (1)$$

Homogeneous: If $f(x) = 0$ in (1).

Definition: A second order differential equation is said to be linear if it can be written as

$$y'' + p(x)y' + q(x)y = f(x). \quad (1)$$

Homogeneous: If $f(x) = 0$ in (1).

Non-homogeneous: If $f(x) \neq 0$ in (1).

Linear Second Order ODE

Definition: A second order differential equation is said to be linear if it can be written as

$$y'' + p(x)y' + q(x)y = f(x). \quad (1)$$

Homogeneous: If $f(x) = 0$ in (1).

Non-homogeneous: If $f(x) \neq 0$ in (1).

Let us try to understand how to solve homogeneous linear second order ODE:

$$y'' + p(x)y' + q(x)y = 0.$$

Linear Second Order ODE

Theorem: Suppose $p(x)$ and $q(x)$ are continuous function on an open interval (a, b) , let x_0 be any point in (a, b) . Then the initial value problem

$$y'' + p(x)y' + q(x)y = 0, \quad y(x_0) = y_0, \quad y'(x_0) = y_1$$

has a unique solution on (a, b) .

Linear Second Order ODE

Theorem: Suppose $p(x)$ and $q(x)$ are continuous function on an open interval (a, b) , let x_0 be any point in (a, b) . Then the initial value problem

$$y'' + p(x)y' + q(x)y = 0, \quad y(x_0) = y_0, \quad y'(x_0) = y_1$$

has a unique solution on (a, b) .

Remark: The initial value problem

$$y'' + p(x)y' + q(x)y = 0, \quad y(x_0) = 0, \quad y'(x_0) = 0$$

has $y \equiv 0$ the only solution (Trivial solution).

Linear Second Order ODE

Theorem: Suppose $p(x)$ and $q(x)$ are continuous function on an open interval (a, b) , let x_0 be any point in (a, b) . Then the initial value problem

$$y'' + p(x)y' + q(x)y = 0, \quad y(x_0) = y_0, \quad y'(x_0) = y_1$$

has a **unique solution** on (a, b) .

Remark: The initial value problem

$$y'' + p(x)y' + q(x)y = 0, \quad y(x_0) = 0, \quad y'(x_0) = 0$$

has $y \equiv 0$ the only solution (Trivial solution).

Solve: The initial value problem

$$y'' - y = 0, \quad y(0) = 1, \quad y'(0) = 3.$$

Solve: The initial value problem

$$y'' - y = 0, \quad y(0) = 1, \quad y'(0) = 3. \quad (2)$$

Linear Second Order ODE

Solve: The initial value problem

$$y'' - y = 0, \quad y(0) = 1, \quad y'(0) = 3. \quad (2)$$

Step 1: Observe that $p(x) = 0$ and $q(x) = -1$ are continuous on $(-\infty, \infty)$.

Linear Second Order ODE

Solve: The initial value problem

$$y'' - y = 0, \quad y(0) = 1, \quad y'(0) = 3. \quad (2)$$

Step 1: Observe that $p(x) = 0$ and $q(x) = -1$ are continuous on $(-\infty, \infty)$.

Step 2: Verify $y_1 = e^x$ and $y_2 = e^{-x}$ are solutions of $y'' - y = 0$.

Linear Second Order ODE

Solve: The initial value problem

$$y'' - y = 0, \quad y(0) = 1, \quad y'(0) = 3. \quad (2)$$

Step 1: Observe that $p(x) = 0$ and $q(x) = -1$ are continuous on $(-\infty, \infty)$.

Step 2: Verify $y_1 = e^x$ and $y_2 = e^{-x}$ are solutions of $y'' - y = 0$.

Step 3: Verify that if c_1 and c_2 are arbitrary constants, $y = c_1 e^x + c_2 e^{-x}$ is a solution of (2) on $(-\infty, \infty)$.

Linear Second Order ODE

Solve: The initial value problem

$$y'' - y = 0, \quad y(0) = 1, \quad y'(0) = 3. \quad (2)$$

Step 1: Observe that $p(x) = 0$ and $q(x) = -1$ are continuous on $(-\infty, \infty)$.

Step 2: Verify $y_1 = e^x$ and $y_2 = e^{-x}$ are solutions of $y'' - y = 0$.

Step 3: Verify that if c_1 and c_2 are arbitrary constants, $y = c_1 e^x + c_2 e^{-x}$ is a solution of (2) on $(-\infty, \infty)$.

$$\begin{aligned} y'' - y &= (c_1 e^x + c_2 e^{-x}) - (c_1 e^x + c_2 e^{-x}) \\ &= c_1(e^x - e^x) + c_2(e^{-x} - e^{-x}) = 0 \end{aligned}$$

Linear Second Order ODE

Solve: The initial value problem

$$y'' - y = 0, \quad y(0) = 1, \quad y'(0) = 3. \quad (2)$$

Step 1: Observe that $p(x) = 0$ and $q(x) = -1$ are continuous on $(-\infty, \infty)$.

Step 2: Verify $y_1 = e^x$ and $y_2 = e^{-x}$ are solutions of $y'' - y = 0$.

Step 3: Verify that if c_1 and c_2 are arbitrary constants, $y = c_1 e^x + c_2 e^{-x}$ is a solution of (2) on $(-\infty, \infty)$.

$$\begin{aligned} y'' - y &= (c_1 e^x + c_2 e^{-x}) - (c_1 e^x + c_2 e^{-x}) \\ &= c_1(e^x - e^x) + c_2(e^{-x} - e^{-x}) = 0 \end{aligned}$$

Therefore $y = c_1 e^x + c_2 e^{-x}$ is a solution of (2) on $(-\infty, \infty)$.

Since $y = c_1 e^x + c_2 e^{-x}$ is a solution of ODE $y'' - y = 0$ on $(-\infty, \infty)$.

Setting $y(0) = 1$ and $y'(0) = 3$, we get

$$c_1 + c_2 = 1$$

$$c_1 - c_2 = 3.$$

Therefore $y = 2e^x - e^{-x}$ is the unique solution on $(-\infty, \infty)$.

Subsection 1

Fundamental solutions : Linearly independent solution

Theorem: Consider the homogeneous ODE:

$$y'' + p(x)y' + q(x)y = 0. \quad (3)$$

If y_1 and y_2 are solutions of (3) on (a, b) , then any linear combination

$$y = c_1y_1 + c_2y_2 \quad (4)$$

is also a solution of (3) on (a, b) .

Theorem: Consider the homogeneous ODE:

$$y'' + p(x)y' + q(x)y = 0. \quad (3)$$

If y_1 and y_2 are solutions of (3) on (a, b) , then any linear combination

$$y = c_1y_1 + c_2y_2 \quad (4)$$

is also a solution of (3) on (a, b) .

Definition: If every solution of (3) on (a, b) can be written as a linear combination of y_1 and y_2 as in (4), we say that $\{y_1, y_2\}$ is a **fundamental set of solutions of (3) on (a, b)** .

Linear Second Order ODE

Theorem: Consider the homogeneous ODE:

$$y'' + p(x)y' + q(x)y = 0. \quad (3)$$

If y_1 and y_2 are solutions of (3) on (a, b) , then any linear combination

$$y = c_1y_1 + c_2y_2 \quad (4)$$

is also a solution of (3) on (a, b) .

Definition: If every solution of (3) on (a, b) can be written as a linear combination of y_1 and y_2 as in (4), we say that $\{y_1, y_2\}$ is a **fundamental set of solutions of (3) on (a, b)** .

Moreover: We say that (4) is **general solution of (3) on (a, b)** .

Second Order Linear Homogeneous ODE

Theorem: Suppose p and q are continuous on (a, b) . Then a set $\{y_1, y_2\}$ of solutions of

$$y'' + p(x)y' + q(x)y = 0 \tag{5}$$

on (a, b) is a fundamental set if and only if $\{y_1, y_2\}$ is linearly independent on (a, b) .

Revision: Linearly independent

$V = C^\infty(a, b)$: Vector space of real valued differentiable functions of any order.

Revision: Linearly independent

$V = C^\infty(a, b)$: Vector space of real valued differentiable functions of any order.

Question. Let $f_1(x)$ and $f_2(x)$ are elements of V . How to show that $f_1(x)$ and $f_2(x)$ are linearly independent in V ?

Revision: Linearly independent

$V = C^\infty(a, b)$: Vector space of real valued differentiable functions of any order.

Question. Let $f_1(x)$ and $f_2(x)$ are elements of V . How to show that $f_1(x)$ and $f_2(x)$ are linearly independent in V ?

By definition if $c_1 f_1 + c_2 f_2 = 0$ for all $x \in (a, b)$, then c_1 and c_2 must be zero.

Revision: Linearly independent

$V = C^\infty(a, b)$: Vector space of real valued differentiable functions of any order.

Question. Let $f_1(x)$ and $f_2(x)$ are elements of V . How to show that $f_1(x)$ and $f_2(x)$ are linearly independent in V ?

By definition if $c_1 f_1 + c_2 f_2 = 0$ for all $x \in (a, b)$, then c_1 and c_2 must be zero.

Why are we discussing this Linear Algebra problem in Differential Equations course?

Section 2

Second order ODE with Constant coefficients

Second Order Linear Homogeneous ODE with constant coeff.

If $a, b, c \in \mathbb{R}$ with $a \neq 0$, then

$$ay'' + by' + cy = 0 \tag{6}$$

is called a constant coefficient 2nd order homogeneous ODE.

Second Order Linear Homogeneous ODE with constant coeff.

If $a, b, c \in \mathbb{R}$ with $a \neq 0$, then

$$ay'' + by' + cy = 0 \tag{6}$$

is called a constant coefficient 2nd order homogeneous ODE.

Let us look for a solution of the type e^{mx} , where m is a constant. Then,

$$\begin{aligned} am^2 e^{mx} + bme^{mx} + ce^{mx} &= 0, \\ e^{mx} (am^2 + bm + c) &= 0. \end{aligned}$$

Second Order Linear Homogeneous ODE with constant coeff.

If $a, b, c \in \mathbb{R}$ with $a \neq 0$, then

$$ay'' + by' + cy = 0 \tag{6}$$

is called a constant coefficient 2nd order homogeneous ODE.

Let us look for a solution of the type e^{mx} , where m is a constant. Then,

$$\begin{aligned} am^2 e^{mx} + bme^{mx} + ce^{mx} &= 0, \\ e^{mx} (am^2 + bm + c) &= 0. \end{aligned}$$

Since $e^{mx} \neq 0$ for all $x \in \mathbb{R}$ and for any constant m , we get that

$$am^2 + bm + c = 0.$$

Second Order Linear Homogeneous ODE with constant coeff.

The quadratic polynomial

$$p(m) = am^2 + bm + c = 0.$$

is the characteristic polynomial of (6), and $p(m) = 0$ is the characteristic equation.

Second Order Linear Homogeneous ODE with constant coeff.

The quadratic polynomial

$$p(m) = am^2 + bm + c = 0.$$

is the characteristic polynomial of (6), and $p(m) = 0$ is the characteristic equation.

Therefore, e^{mx} is a solution of (6) if and only if $p(m) = 0$.

The roots of the characteristic equation are given by $m = \frac{-b \pm \sqrt{b^2 - 4ac}}{2a}$.

Second Order Linear Homogeneous ODE with constant coeff.

The quadratic polynomial

$$p(m) = am^2 + bm + c = 0.$$

is the characteristic polynomial of (6), and $p(m) = 0$ is the characteristic equation.

Therefore, e^{mx} is a solution of (6) if and only if $p(m) = 0$.

The roots of the characteristic equation are given by $m = \frac{-b \pm \sqrt{b^2 - 4ac}}{2a}$.

Case 1: When $b^2 - 4ac > 0$, the characteristic equation $p(m) = 0$ has two distinct real roots.

Second Order Linear Homogeneous ODE with constant coeff.

The quadratic polynomial

$$p(m) = am^2 + bm + c = 0.$$

is the characteristic polynomial of (6), and $p(m) = 0$ is the characteristic equation.

Therefore, e^{mx} is a solution of (6) if and only if $p(m) = 0$.

The roots of the characteristic equation are given by $m = \frac{-b \pm \sqrt{b^2 - 4ac}}{2a}$.

Case 1: When $b^2 - 4ac > 0$, the characteristic equation $p(m) = 0$ has two distinct real roots.

Case 2: When $b^2 - 4ac = 0$, the characteristic equation $p(m) = 0$ has repeated real roots.

Second Order Linear Homogeneous ODE with constant coeff.

The quadratic polynomial

$$p(m) = am^2 + bm + c = 0.$$

is the characteristic polynomial of (6), and $p(m) = 0$ is the characteristic equation.

Therefore, e^{mx} is a solution of (6) if and only if $p(m) = 0$.

The roots of the characteristic equation are given by $m = \frac{-b \pm \sqrt{b^2 - 4ac}}{2a}$.

Case 1: When $b^2 - 4ac > 0$, the characteristic equation $p(m) = 0$ has two distinct real roots.

Case 2: When $b^2 - 4ac = 0$, the characteristic equation $p(m) = 0$ has repeated real roots.

Case 3: When $b^2 - 4ac < 0$, the characteristic equation $p(m) = 0$ has two distinct complex roots which are conjugates.

Subsection 1

Distinct real roots

Example: Find general solution of

$$y'' + 6y' + 5y = 0 \tag{7}$$

Example: Find general solution of

$$y'' + 6y' + 5y = 0 \tag{7}$$

The characteristic equation is

$$p(m) = m^2 + 6m + 5 = (m + 1)(m + 5).$$

Distinct real roots case

Example: Find general solution of

$$y'' + 6y' + 5y = 0 \tag{7}$$

The characteristic equation is

$$p(m) = m^2 + 6m + 5 = (m + 1)(m + 5).$$

Thus roots of characteristic equation are -1 and -5 .

Example: Find general solution of

$$y'' + 6y' + 5y = 0 \quad (7)$$

The characteristic equation is

$$p(m) = m^2 + 6m + 5 = (m + 1)(m + 5).$$

Thus roots of characteristic equation are -1 and -5 .

Thus $y_1 = e^{-x}$ and $y_2 = e^{-5x}$ are solutions of (7).

Example: Find general solution of

$$y'' + 6y' + 5y = 0 \tag{7}$$

The characteristic equation is

$$p(m) = m^2 + 6m + 5 = (m + 1)(m + 5).$$

Thus roots of characteristic equation are -1 and -5 .

Thus $y_1 = e^{-x}$ and $y_2 = e^{-5x}$ are solutions of (7).

Note that any linear combination of the above two is also a solution

$$y = c_1 e^{-x} + c_2 e^{-5x}.$$

Example: Solve IVP

$$y'' + 6y' + 5y = 0, \text{ where } y(0) = 3, y'(0) = 1. \quad (8)$$

Example: Solve IVP

$$y'' + 6y' + 5y = 0, \text{ where } y(0) = 3, y'(0) = 1. \quad (8)$$

If we solve for c_1 and c_2 using the initial conditions, we get $c_1 = 4$ and $c_2 = -1$.

Example: Solve IVP

$$y'' + 6y' + 5y = 0, \text{ where } y(0) = 3, y'(0) = 1. \quad (8)$$

If we solve for c_1 and c_2 using the initial conditions, we get $c_1 = 4$ and $c_2 = -1$.

Thus the solution to IVP is

$$y = 4e^{-x} - e^{-5x}.$$

Subsection 2

Repeated real roots

A repeated real root case

Example: Find the general solution of

$$y'' + 6y' + 9y = 0. \quad (9)$$

A repeated real root case

Example: Find the general solution of

$$y'' + 6y' + 9y = 0. \quad (9)$$

The characteristic equation is

$$p(m) = m^2 + 6m + 9 = (m + 3)^2.$$

A repeated real root case

Example: Find the general solution of

$$y'' + 6y' + 9y = 0. \quad (9)$$

The characteristic equation is

$$p(m) = m^2 + 6m + 9 = (m + 3)^2.$$

The characteristic equation has repeated roots $-3, -3$. Hence $y_1 = e^{-3x}$ is one solution.

For other solution,

A repeated real root case

Example: Find the general solution of

$$y'' + 6y' + 9y = 0. \quad (9)$$

The characteristic equation is

$$p(m) = m^2 + 6m + 9 = (m + 3)^2.$$

The characteristic equation has repeated roots $-3, -3$. Hence $y_1 = e^{-3x}$ is one solution.

For other solution, let us look for a solution of the type $y_2 = ue^{-3x}$. So,

A repeated real root case

Example: Find the general solution of

$$y'' + 6y' + 9y = 0. \quad (9)$$

The characteristic equation is

$$p(m) = m^2 + 6m + 9 = (m + 3)^2.$$

The characteristic equation has repeated roots $-3, -3$. Hence $y_1 = e^{-3x}$ is one solution.

For other solution, let us look for a solution of the type $y_2 = ue^{-3x}$. So,

$$\begin{aligned} y'' + 6y' + 9y &= e^{-3x} [(u'' - 6u' + 9u) + 6(u' - 3u) + 9u] \\ &= u'' e^{-3x} = 0. \end{aligned}$$

A repeated real root case

Example: Find the general solution of

$$y'' + 6y' + 9y = 0. \quad (9)$$

The characteristic equation is

$$p(m) = m^2 + 6m + 9 = (m + 3)^2.$$

The characteristic equation has repeated roots $-3, -3$. Hence $y_1 = e^{-3x}$ is one solution.

For other solution, let us look for a solution of the type $y_2 = ue^{-3x}$. So,

$$\begin{aligned} y'' + 6y' + 9y &= e^{-3x} [(u'' - 6u' + 9u) + 6(u' - 3u) + 9u] \\ &= u'' e^{-3x} = 0. \end{aligned}$$

This shows that $u(x) = c_1x + c_2$, where c_1 and c_2 are constants. Therefore

$$y = e^{-3x}(c_1 + c_2x) \quad (10)$$

is a solution of (9).

A repeated real root case

Example: Solve the initial value problem

$$y'' + 6y' + 9y = 0, \quad y(0) = 3, \quad y'(0) = -1. \quad (11)$$

A repeated real root case

Example: Solve the initial value problem

$$y'' + 6y' + 9y = 0, \quad y(0) = 3, \quad y'(0) = -1. \quad (11)$$

Recall

$$y = e^{-3x}(c_1 + c_2x) \quad (12)$$

is a solution of (27). Using initial conditions, get $c_1 = 3$ and $c_2 = 10$.

Thus solution of IVP is

$$y = e^{-3x}(3 + 10x). \quad (13)$$

Subsection 3

Complex conjugate roots

Two distinct complex conjugate roots case

Example: Find the general solution of

$$y'' + 4y' + 13y = 0. \tag{14}$$

Two distinct complex conjugate roots case

Example: Find the general solution of

$$y'' + 4y' + 13y = 0. \quad (14)$$

The characteristic polynomial of (14) is

$$p(m) = m^2 + 4m + 13 = m^2 + 4m + 4 + 9 = (m + 2)^2 + 9.$$

Two distinct complex conjugate roots case

Example: Find the general solution of

$$y'' + 4y' + 13y = 0. \quad (14)$$

The characteristic polynomial of (14) is

$$p(m) = m^2 + 4m + 13 = m^2 + 4m + 4 + 9 = (m + 2)^2 + 9.$$

The roots of the characteristic equation are $m_1 = -2 + 3i$ and $m_2 = -2 - 3i$.

Two distinct complex conjugate roots case

Example: Find the general solution of

$$y'' + 4y' + 13y = 0. \quad (14)$$

The characteristic polynomial of (14) is

$$p(m) = m^2 + 4m + 13 = m^2 + 4m + 4 + 9 = (m + 2)^2 + 9.$$

The roots of the characteristic equation are $m_1 = -2 + 3i$ and $m_2 = -2 - 3i$.

It is reasonable to expect that $e^{(-2+3i)x}$ and $e^{(-2-3i)x}$ are solutions of (14).

Two distinct complex conjugate roots case

Example: Find the general solution of

$$y'' + 4y' + 13y = 0. \quad (14)$$

The characteristic polynomial of (14) is

$$p(m) = m^2 + 4m + 13 = m^2 + 4m + 4 + 9 = (m + 2)^2 + 9.$$

The roots of the characteristic equation are $m_1 = -2 + 3i$ and $m_2 = -2 - 3i$.

It is reasonable to expect that $e^{(-2+3i)x}$ and $e^{(-2-3i)x}$ are solutions of (14).

In fact that is true. But they are complex valued solutions

Two distinct complex conjugate roots case

Example: Find the general solution of

$$y'' + 4y' + 13y = 0. \quad (14)$$

The characteristic polynomial of (14) is

$$p(m) = m^2 + 4m + 13 = m^2 + 4m + 4 + 9 = (m + 2)^2 + 9.$$

The roots of the characteristic equation are $m_1 = -2 + 3i$ and $m_2 = -2 - 3i$.

It is reasonable to expect that $e^{(-2+3i)x}$ and $e^{(-2-3i)x}$ are solutions of (14).

In fact that is true. But they are complex valued solutions and we want real solutions.

Two distinct complex conjugate roots case

Let us write

$$y_1 = e^{(-2+3i)x} = e^{-2x} e^{3ix} \quad \text{and} \quad y_2 = e^{(-2-3i)x} = e^{-2x} e^{-3ix}.$$

Two distinct complex conjugate roots case

Let us write

$$y_1 = e^{(-2+3i)x} = e^{-2x} e^{3ix} \quad \text{and} \quad y_2 = e^{(-2-3i)x} = e^{-2x} e^{-3ix}.$$

Assume that linear combination of y_1 and y_2 can be written as $y = ue^{-2x}$, where u depends upon x . Let us find u .

Two distinct complex conjugate roots case

Let us write

$$y_1 = e^{(-2+3i)x} = e^{-2x} e^{3ix} \quad \text{and} \quad y_2 = e^{(-2-3i)x} = e^{-2x} e^{-3ix}.$$

Assume that linear combination of y_1 and y_2 can be written as $y = ue^{-2x}$, where u depends upon x . Let us find u .

Since $y = ue^{-2x}$ then

$$y' = u'e^{-2x} - 2ue^{-2x} \quad \text{and} \quad y'' = u''e^{-2x} - 4u'e^{-2x} + 4ue^{-2x}.$$

Now from ODE, we get

$$\begin{aligned} y'' + 4y' + 13y &= e^{-2x} [(u'' - 4u' + 4u) + 4(u' - 2u) + 13u] \\ &= e^{-2x} [u'' - (4 - 4)u' + (4 - 8 + 13)u] = e^{-2x} (u'' + 9u). \end{aligned}$$

Therefore $y = ue^{-2x}$ is a solution of (14) if and only if

$$u'' + 9u = 0.$$

Two distinct complex conjugate roots case

The general solution of the equation $u'' + 9u = 0$ is

$$u = c_1 \cos 3x + c_2 \sin 3x.$$

Check that $\cos 3x$ and $\sin 3x$ are solutions of ODE $u'' + 9u = 0$, and hence their linear combinations.

Therefore any function of the form

$$y = e^{-2x}(c_1 \cos 3x + c_2 \sin 3x) \tag{15}$$

is a solution of (14).

Two distinct complex conjugate roots case

Example: Solve the initial value problem

$$y'' + 4y' + 13y = 0, \quad y(0) = 2, \quad y'(0) = -3. \quad (16)$$

Two distinct complex conjugate roots case

Example: Solve the initial value problem

$$y'' + 4y' + 13y = 0, \quad y(0) = 2, \quad y'(0) = -3. \quad (16)$$

Using initial conditions, get $c_1 = 2$ and $c_2 = \frac{1}{3}$ and hence the solution is

$$y = e^{-2x} \left(2 \cos 3x + \frac{1}{3} \sin 3x \right).$$

Theorem: Let $p(m) = am^2 + bm + c$ be the characteristic polynomial of

$$ay'' + by' + cy = 0. \quad (17)$$

Then:

(a) If $p(m) = 0$ has distinct real roots m_1 and m_2 , then the general solution is

$$y = c_1 e^{m_1 x} + c_2 e^{m_2 x}.$$

Theorem: Let $p(m) = am^2 + bm + c$ be the characteristic polynomial of

$$ay'' + by' + cy = 0. \quad (17)$$

Then:

(a) If $p(m) = 0$ has distinct real roots m_1 and m_2 , then the general solution is

$$y = c_1 e^{m_1 x} + c_2 e^{m_2 x}.$$

(b) If $p(m) = 0$ has a repeated root m_1 , then the general solution is

$$y = e^{m_1 x} (c_1 + c_2 x).$$

Theorem: Let $p(m) = am^2 + bm + c$ be the characteristic polynomial of

$$ay'' + by' + cy = 0. \quad (17)$$

Then:

(a) If $p(m) = 0$ has distinct real roots m_1 and m_2 , then the general solution is

$$y = c_1 e^{m_1 x} + c_2 e^{m_2 x}.$$

(b) If $p(m) = 0$ has a repeated root m_1 , then the general solution is

$$y = e^{m_1 x} (c_1 + c_2 x).$$

(c) If $p(m) = 0$ has complex conjugate roots $m_1 = \lambda + i\omega$ and $m_2 = \lambda - i\omega$ (where $\omega > 0$), then the general solution is

$$y = e^{\lambda x} (c_1 \cos \omega x + c_2 \sin \omega x).$$

How to prove linearly independent!!!

How to prove linearly independent!!!

- ▶ $e^{m_1 x}$ and $e^{m_2 x}$;
- ▶ e^{mx} and xe^{mx} , or
- ▶ $e^{\lambda x} \cos \omega x$ and $e^{\lambda x} \sin \omega x$.

Let $p(m) = am^2 + bm + c$ be the characteristic polynomial of

$$ay'' + by' + cy = 0. \tag{18}$$

Then $p(m) = 0$ can have at most two distinct roots.

Let $p(m) = am^2 + bm + c$ be the characteristic polynomial of

$$ay'' + by' + cy = 0. \quad (18)$$

Then $p(m) = 0$ can have at most two distinct roots.

Observe that solution to ODE (18) will have linear combination of the following form

► $e^{m_1 x}$ and $e^{m_2 x}$;

Let $p(m) = am^2 + bm + c$ be the characteristic polynomial of

$$ay'' + by' + cy = 0. \quad (18)$$

Then $p(m) = 0$ can have at most two distinct roots.

Observe that solution to ODE (18) will have linear combination of the following form

- ▶ e^{m_1x} and e^{m_2x} ;
- ▶ e^{mx} and xe^{mx} , or

Let $p(m) = am^2 + bm + c$ be the characteristic polynomial of

$$ay'' + by' + cy = 0. \quad (18)$$

Then $p(m) = 0$ can have at most two distinct roots.

Observe that solution to ODE (18) will have linear combination of the following form

- ▶ e^{m_1x} and e^{m_2x} ;
- ▶ e^{mx} and xe^{mx} , or
- ▶ $e^{\lambda x} \cos \omega x$ and $e^{\lambda x} \sin \omega x$.

Let $p(m) = am^2 + bm + c$ be the characteristic polynomial of

$$ay'' + by' + cy = 0. \quad (18)$$

Then $p(m) = 0$ can have at most two distinct roots.

Observe that solution to ODE (18) will have linear combination of the following form

- ▶ e^{m_1x} and e^{m_2x} ;
- ▶ e^{mx} and xe^{mx} , or
- ▶ $e^{\lambda x} \cos \omega x$ and $e^{\lambda x} \sin \omega x$.

Something to think about!!!

Suppose $p(m) = a_0m^\ell + a_1m^{\ell-1} + \cdots + a_{m-1}m + a_m = 0$ be the characteristic polynomial of

$$a_0y^{(\ell)} + a_1y^{(\ell-1)} + \cdots + a_{m-1}y' + a_my = 0 \quad (19)$$

Recall: Second Order Linear ODE

Theorem: Suppose $p(x)$ and $q(x)$ are continuous function on an open interval (a, b) , let x_0 be any point in (a, b) . Then the initial value problem

$$y'' + p(x)y' + q(x)y = 0, \quad y(x_0) = w_0, \quad y'(x_0) = w_1 \quad (20)$$

has a unique solution on (a, b) .

Recall: Second Order Linear ODE

Theorem: Suppose $p(x)$ and $q(x)$ are continuous function on an open interval (a, b) , let x_0 be any point in (a, b) . Then the initial value problem

$$y'' + p(x)y' + q(x)y = 0, \quad y(x_0) = w_0, \quad y'(x_0) = w_1 \quad (20)$$

has a unique solution on (a, b) .

Note that the solution to ODE (20) is of the form

$$y(x) = c_1 y_1 + c_2 y_2.$$

Recall: Second Order Linear ODE

Theorem: Suppose $p(x)$ and $q(x)$ are continuous function on an open interval (a, b) , let x_0 be any point in (a, b) . Then the initial value problem

$$y'' + p(x)y' + q(x)y = 0, \quad y(x_0) = w_0, \quad y'(x_0) = w_1 \quad (20)$$

has a unique solution on (a, b) .

Note that the solution to ODE (20) is of the form

$$y(x) = c_1 y_1 + c_2 y_2.$$

Since ODE (20) has a unique solution, we must have unique c_1 and c_2 . Can we also give expression for c_1 and c_2 ?

Recall: Second Order Linear ODE

Theorem: Suppose $p(x)$ and $q(x)$ are continuous function on an open interval (a, b) , let x_0 be any point in (a, b) . Then the initial value problem

$$y'' + p(x)y' + q(x)y = 0, \quad y(x_0) = w_0, \quad y'(x_0) = w_1 \quad (20)$$

has a unique solution on (a, b) .

Note that the solution to ODE (20) is of the form

$$y(x) = c_1 y_1 + c_2 y_2.$$

Since ODE (20) has a unique solution, we must have unique c_1 and c_2 . Can we also give expression for c_1 and c_2 ?

The system of equation

$$\begin{aligned} c_1 y_1(x_0) + c_2 y_2(x_0) &= w_0 \\ c_1 y_1'(x_0) + c_2 y_2'(x_0) &= w_1 \end{aligned} \quad (21)$$

has a solution (c_1, c_2) for every choice of (w_0, w_1) .

Second Order Linear ODE

Theorem: Suppose $p(x)$ and $q(x)$ are continuous on (a, b) , let y_1 and y_2 be solutions of

$$y'' + p(x)y' + q(x)y = 0 \quad (22)$$

on (a, b) , and define

$$W = y_1 y_2' - y_1' y_2. \quad (23)$$

Let x_0 be any point in (a, b) . Then

Second Order Linear ODE

Theorem: Suppose $p(x)$ and $q(x)$ are continuous on (a, b) , let y_1 and y_2 be solutions of

$$y'' + p(x)y' + q(x)y = 0 \quad (22)$$

on (a, b) , and define

$$W = y_1 y_2' - y_1' y_2. \quad (23)$$

Let x_0 be any point in (a, b) . Then

$$W(x) = W(x_0)e^{-\int_{x_0}^x p(t) dt}, \quad a < x < b. \quad (24)$$

Therefore either W has no zeros in (a, b) or $W \equiv 0$ on (a, b) .

Second Order Linear ODE

Theorem: Suppose $p(x)$ and $q(x)$ are continuous on (a, b) , let y_1 and y_2 be solutions of

$$y'' + p(x)y' + q(x)y = 0 \quad (22)$$

on (a, b) , and define

$$W = y_1 y_2' - y_1' y_2. \quad (23)$$

Let x_0 be any point in (a, b) . Then

$$W(x) = W(x_0)e^{-\int_{x_0}^x p(t) dt}, \quad a < x < b. \quad (24)$$

Therefore either W has no zeros in (a, b) or $W \equiv 0$ on (a, b) .

Proof Given $W = y_1 y_2' - y_1' y_2$. On differentiating (23), we get

$$W' = y_1' y_2' + y_1 y_2'' - y_1' y_2' - y_1'' y_2 = y_1 y_2'' - y_1'' y_2.$$

Second Order Linear ODE

Theorem: Suppose $p(x)$ and $q(x)$ are continuous on (a, b) , let y_1 and y_2 be solutions of

$$y'' + p(x)y' + q(x)y = 0 \quad (22)$$

on (a, b) , and define

$$W = y_1 y_2' - y_1' y_2. \quad (23)$$

Let x_0 be any point in (a, b) . Then

$$W(x) = W(x_0)e^{-\int_{x_0}^x p(t) dt}, \quad a < x < b. \quad (24)$$

Therefore either W has no zeros in (a, b) or $W \equiv 0$ on (a, b) .

Proof Given $W = y_1 y_2' - y_1' y_2$. On differentiating (23), we get

$$W' = y_1' y_2' + y_1 y_2'' - y_1'' y_2 - y_1' y_2' = y_1 y_2'' - y_1'' y_2.$$

Also notice that

$$y_1'' = -p y_1' - q y_1 \quad \text{and} \quad y_2'' = -p y_2' - q y_2.$$

Why are we proving this theorem?

Why are we proving this theorem?

Proof (continued...) Simplify the expression of W' and get

$$\begin{aligned}W' &= -y_1(py'_2 + qy_2) + y_2(py'_1 + qy_1) \\&= -p(y_1y'_2 - y_2y'_1) - q(y_1y_2 - y_2y_1) \\&= -p(y_1y'_2 - y_2y'_1) = -pW.\end{aligned}$$

Why are we proving this theorem?

Proof (continued...) Simplify the expression of W' and get

$$\begin{aligned}W' &= -y_1(py_2' + qy_2) + y_2(py_1' + qy_1) \\&= -p(y_1y_2' - y_2y_1') - q(y_1y_2 - y_2y_1) \\&= -p(y_1y_2' - y_2y_1') = -pW.\end{aligned}$$

Therefore $W' + p(x)W = 0$;

Thus

$$W(x) = ce^{-\int_{x_0}^x p(t) dt}, \quad a < x < b.$$

But $c = W(x_0)$, and hence

$$W(x) = W(x_0)e^{-\int_{x_0}^x p(t) dt}, \quad a < x < b.$$

Why are we proving this theorem?

Proof (continued...) Simplify the expression of W' and get

$$\begin{aligned}W' &= -y_1(py_2' + qy_2) + y_2(py_1' + qy_1) \\&= -p(y_1y_2' - y_2y_1') - q(y_1y_2 - y_2y_1) \\&= -p(y_1y_2' - y_2y_1') = -pW.\end{aligned}$$

Therefore $W' + p(x)W = 0$;

Thus

$$W(x) = ce^{-\int_{x_0}^x p(t) dt}, \quad a < x < b.$$

But $c = W(x_0)$, and hence

$$W(x) = W(x_0)e^{-\int_{x_0}^x p(t) dt}, \quad a < x < b.$$

Observe that $e^{-\int_{x_0}^x p(t) dt} \neq 0$ for any $a < x < b$.

Why are we proving this theorem?

Proof (continued...) Simplify the expression of W' and get

$$\begin{aligned}W' &= -y_1(py_2' + qy_2) + y_2(py_1' + qy_1) \\&= -p(y_1y_2' - y_2y_1') - q(y_1y_2 - y_2y_1) \\&= -p(y_1y_2' - y_2y_1') = -pW.\end{aligned}$$

Therefore $W' + p(x)W = 0$;

Thus

$$W(x) = ce^{-\int_{x_0}^x p(t) dt}, \quad a < x < b.$$

But $c = W(x_0)$, and hence

$$W(x) = W(x_0)e^{-\int_{x_0}^x p(t) dt}, \quad a < x < b.$$

Observe that $e^{-\int_{x_0}^x p(t) dt} \neq 0$ for any $a < x < b$.

Therefore either W has no zeros in (a, b) or $W \equiv 0$ on (a, b) .

Second Order Linear ODE

Theorem: Suppose $p(x)$ and $q(x)$ are continuous on (a, b) , let y_1 and y_2 be solutions of

$$y'' + p(x)y' + q(x)y = 0 \quad (25)$$

on (a, b) , and let $W = y_1y_2' - y_1'y_2$.

Second Order Linear ODE

Theorem: Suppose $p(x)$ and $q(x)$ are continuous on (a, b) , let y_1 and y_2 be solutions of

$$y'' + p(x)y' + q(x)y = 0 \quad (25)$$

on (a, b) , and let $W = y_1y_2' - y_1'y_2$.

Then y_1 and y_2 are linearly independent on (a, b) if and only if W has no zeros on (a, b) .

Second Order Linear ODE

Theorem: Suppose $p(x)$ and $q(x)$ are continuous on (a, b) , let y_1 and y_2 be solutions of

$$y'' + p(x)y' + q(x)y = 0 \quad (25)$$

on (a, b) , and let $W = y_1y_2' - y_1'y_2$.

Then y_1 and y_2 are linearly independent on (a, b) if and only if W has no zeros on (a, b) .

Proof We omit the proof in this course.

Second Order Linear ODE

Theorem: Suppose $p(x)$ and $q(x)$ are continuous on (a, b) , let y_1 and y_2 be solutions of

$$y'' + p(x)y' + q(x)y = 0 \quad (25)$$

on (a, b) , and let $W = y_1y_2' - y_1'y_2$.

Then y_1 and y_2 are linearly independent on (a, b) if and only if W has no zeros on (a, b) .

Proof We omit the proof in this course.

Observations We have two expressions for W , namely

► $W(x) = y_1y_2' - y_1'y_2,$

► $W(x) = W(x_0)e^{-\int_{x_0}^x p(t) dt}, \quad a < x < b.,$

and

Second Order Linear ODE

Theorem: Suppose $p(x)$ and $q(x)$ are continuous on (a, b) , let y_1 and y_2 be solutions of

$$y'' + p(x)y' + q(x)y = 0 \quad (25)$$

on (a, b) , and let $W = y_1y_2' - y_1'y_2$.

Then y_1 and y_2 are linearly independent on (a, b) if and only if W has no zeros on (a, b) .

Proof We omit the proof in this course.

Observations We have two expressions for W , namely

- ▶ $W(x) = y_1y_2' - y_1'y_2$,
- ▶ $W(x) = W(x_0)e^{-\int_{x_0}^x p(t) dt}$, $a < x < b$,

and

- ▶ W decides linearly independence of y_1 and y_2 (important!).

Subsection 4

Wronskian

The Wronskian and Abel's Formula

Definition: The Wronskian of y_1 and y_2 is

$$W = \begin{vmatrix} y_1 & y_2 \\ y_1' & y_2' \end{vmatrix} = y_1 y_2' - y_1' y_2.$$

The Wronskian and Abel's Formula

Definition: The Wronskian of y_1 and y_2 is

$$W = \begin{vmatrix} y_1 & y_2 \\ y_1' & y_2' \end{vmatrix} = y_1 y_2' - y_1' y_2.$$

Abel's formula

$$W(x) = W(x_0) e^{-\int_{x_0}^x p(t) dt}, \quad a < x < b.$$

Compute the Wronskian and verify Abel's formula

Example: Solve IVP

$$y'' + 6y' + 5y = 0, \text{ where } y(0) = 3, y'(0) = 1. \quad (26)$$

Compute the Wronskian and verify Abel's formula

Example: Solve IVP

$$y'' + 6y' + 5y = 0, \text{ where } y(0) = 3, y'(0) = 1. \quad (26)$$

The solution to IVP is $y = 4e^{-x} - e^{-5x}$.

Compute the Wronskian and verify Abel's formula

Example: Solve IVP

$$y'' + 6y' + 5y = 0, \text{ where } y(0) = 3, y'(0) = 1. \quad (26)$$

The solution to IVP is $y = 4e^{-x} - e^{-5x}$.

The Wronskian of e^{-x} and e^{-5x} is

$$W = \begin{vmatrix} e^{-x} & e^{-5x} \\ -e^{-x} & -5e^{-5x} \end{vmatrix} = -4e^{-6x} \neq 0 \quad \text{for any } x \in \mathbb{R}.$$

Compute the Wronskian and verify Abel's formula

Example: Solve IVP

$$y'' + 6y' + 5y = 0, \text{ where } y(0) = 3, y'(0) = 1. \quad (26)$$

The solution to IVP is $y = 4e^{-x} - e^{-5x}$.

The Wronskian of e^{-x} and e^{-5x} is

$$W = \begin{vmatrix} e^{-x} & e^{-5x} \\ -e^{-x} & -5e^{-5x} \end{vmatrix} = -4e^{-6x} \neq 0 \quad \text{for any } x \in \mathbb{R}.$$

Abel's formula

$$W(x) = W(0)e^{-\int_0^x p(t) dt}.$$

Compute the Wronskian and verify Abel's formula

Example: Solve IVP

$$y'' + 6y' + 5y = 0, \text{ where } y(0) = 3, y'(0) = 1. \quad (26)$$

The solution to IVP is $y = 4e^{-x} - e^{-5x}$.

The Wronskian of e^{-x} and e^{-5x} is

$$W = \begin{vmatrix} e^{-x} & e^{-5x} \\ -e^{-x} & -5e^{-5x} \end{vmatrix} = -4e^{-6x} \neq 0 \quad \text{for any } x \in \mathbb{R}.$$

Abel's formula

$$W(x) = W(0)e^{-\int_0^x p(t) dt}.$$

Since $p(t) = 6$ and $W(0) = -4$, we get $W(x) = -4e^{-6x}$.

Compute the Wronskian and verify Abel's formula

Example: Solve the initial value problem

$$y'' + 6y' + 9y = 0, \quad y(0) = 3, \quad y'(0) = -1. \quad (27)$$

Compute the Wronskian and verify Abel's formula

Example: Solve the initial value problem

$$y'' + 6y' + 9y = 0, \quad y(0) = 3, \quad y'(0) = -1. \quad (27)$$

Recall, solution of IVP is $y = e^{-3x}(3 + 10x)$.

Compute the Wronskian and verify Abel's formula

Example: Solve the initial value problem

$$y'' + 6y' + 9y = 0, \quad y(0) = 3, \quad y'(0) = -1. \quad (27)$$

Recall, solution of IVP is $y = e^{-3x}(3 + 10x)$.

The Wronskian of $y_1 = e^{-3x}$ and $y_2 = xe^{-3x}$ is

$$W = \begin{vmatrix} y_1 & y_2 \\ y_1' & y_2' \end{vmatrix} = e^{-6x} \neq 0 \text{ for any } x \in \mathbb{R}.$$

Compute the Wronskian and verify Abel's formula

Example: Solve the initial value problem

$$y'' + 6y' + 9y = 0, \quad y(0) = 3, \quad y'(0) = -1. \quad (27)$$

Recall, solution of IVP is $y = e^{-3x}(3 + 10x)$.

The Wronskian of $y_1 = e^{-3x}$ and $y_2 = xe^{-3x}$ is

$$W = \begin{vmatrix} y_1 & y_2 \\ y_1' & y_2' \end{vmatrix} = e^{-6x} \neq 0 \text{ for any } x \in \mathbb{R}.$$

Abel's formula

$$W(x) = W(x_0)e^{-\int_{x_0}^x p(t) dt}, \quad a < x < b.$$

Compute the Wronskian and verify Abel's formula

Example: Solve the initial value problem

$$y'' + 4y' + 13y = 0, \quad y(0) = 2, \quad y'(0) = -3. \quad (28)$$

Recall that the solution to ODE is $y = e^{-2x}(c_1 \cos 3x + c_2 \sin 3x)$.

Compute the Wronskian and verify Abel's formula

Example: Solve the initial value problem

$$y'' + 4y' + 13y = 0, \quad y(0) = 2, \quad y'(0) = -3. \quad (28)$$

Recall that the solution to ODE is $y = e^{-2x}(c_1 \cos 3x + c_2 \sin 3x)$.

The Wronskian of $y_1 = e^{-2x} \cos 3x$ and $y_2 = e^{-2x} \sin 3x$ is

$$W = \begin{vmatrix} y_1 & y_2 \\ y_1' & y_2' \end{vmatrix} = ?$$

Compute the Wronskian and verify Abel's formula

Example: Solve the initial value problem

$$y'' + 4y' + 13y = 0, \quad y(0) = 2, \quad y'(0) = -3. \quad (28)$$

Recall that the solution to ODE is $y = e^{-2x}(c_1 \cos 3x + c_2 \sin 3x)$.

The Wronskian of $y_1 = e^{-2x} \cos 3x$ and $y_2 = e^{-2x} \sin 3x$ is

$$W = \begin{vmatrix} y_1 & y_2 \\ y_1' & y_2' \end{vmatrix} = ?$$

Abel's formula

$$W(x) = W(x_0)e^{-\int_{x_0}^x p(t) dt}, \quad a < x < b.$$

Section 3

Non-homogeneous second order linear ODE

Non-homogeneous second order linear ODE

Consider the nonhomogeneous 2nd order linear equation

$$y'' + p(x)y' + q(x)y = f(x), \quad (29)$$

where $f(x) \neq 0$ identically.

Non-homogeneous second order linear ODE

Consider the nonhomogeneous 2nd order linear equation

$$y'' + p(x)y' + q(x)y = f(x), \quad (29)$$

where $f(x) \neq 0$ identically.

Theorem: Suppose $p(x)$, $q(x)$ and $f(x)$ are continuous functions on an open interval (a, b) , let x_0 be any point in (a, b) . Then the initial value problem

$$y'' + p(x)y' + q(x)y = f(x), \quad y(x_0) = w_0, \quad y'(x_0) = w_1$$

has a unique solution on (a, b) .

Non-homogeneous second order linear ODE

Consider the nonhomogeneous 2nd order linear equation

$$y'' + p(x)y' + q(x)y = f(x), \quad (29)$$

where $f(x) \neq 0$ identically.

Theorem: Suppose $p(x)$, $q(x)$ and $f(x)$ are continuous functions on an open interval (a, b) , let x_0 be any point in (a, b) . Then the initial value problem

$$y'' + p(x)y' + q(x)y = f(x), \quad y(x_0) = w_0, \quad y'(x_0) = w_1$$

has a unique solution on (a, b) .

We know how to solve the associated homogeneous equation, initial value problem

$$y'' + p(x)y' + q(x)y = 0 \quad y(x_0) = w_0, \quad y'(x_0) = w_1$$

Non-homogeneous second order linear ODE

Set-up: The homogeneous IVP

$$y'' + p(x)y' + q(x)y = 0$$

on (a, b) is called complementary equation.

Non-homogeneous second order linear ODE

Set-up: The homogeneous IVP

$$y'' + p(x)y' + q(x)y = 0$$

on (a, b) is called complementary equation.

Particular solution: Any solution of non-homogeneous IVP

$$y'' + p(x)y' + q(x)y = f(x)$$

on (a, b) is called particular solution.

Non-homogeneous second order linear ODE

Set-up: The homogeneous IVP

$$y'' + p(x)y' + q(x)y = 0$$

on (a, b) is called complementary equation.

Particular solution: Any solution of non-homogeneous IVP

$$y'' + p(x)y' + q(x)y = f(x)$$

on (a, b) is called particular solution.

Solve the initial value problem

$$y'' + y = 1, \quad y(0) = 2, \quad y'(0) = 7.$$

Solve the initial value problem

$$y'' + y = 1, \quad y(0) = 2, \quad y'(0) = 7. \quad (30)$$

Particular solution of non-homogeneous second order linear ODE

Solve the initial value problem

$$y'' + y = 1, \quad y(0) = 2, \quad y'(0) = 7. \quad (30)$$

The characteristic equation is

$$p(m) = m^2 + 1 = 0.$$

Particular solution of non-homogeneous second order linear ODE

Solve the initial value problem

$$y'' + y = 1, \quad y(0) = 2, \quad y'(0) = 7. \quad (30)$$

The characteristic equation is

$$p(m) = m^2 + 1 = 0.$$

Thus $y_1 = \cos x$ and $y_2 = \sin x$ form a fundamental set of solutions of the complementary equation.

Particular solution of non-homogeneous second order linear ODE

Solve the initial value problem

$$y'' + y = 1, \quad y(0) = 2, \quad y'(0) = 7. \quad (30)$$

The characteristic equation is

$$p(m) = m^2 + 1 = 0.$$

Thus $y_1 = \cos x$ and $y_2 = \sin x$ form a fundamental set of solutions of the complementary equation.

The general solution of (30) is

$$y = y_p + \cos x + 7 \sin x. \quad (31)$$

Any guess for y_p .

Particular solution of non-homogeneous second order linear ODE

Solve the initial value problem

$$y'' + y = 1, \quad y(0) = 2, \quad y'(0) = 7. \quad (30)$$

The characteristic equation is

$$p(m) = m^2 + 1 = 0.$$

Thus $y_1 = \cos x$ and $y_2 = \sin x$ form a fundamental set of solutions of the complementary equation.

The general solution of (30) is

$$y = y_p + \cos x + 7 \sin x. \quad (31)$$

Any guess for y_p . take $y_p = 1$.