Dilated Singer sets and the excess in Erdős's Sidon problem: a curve-counting theorem, forbidden-structure laws, a resonance theorem, exact Galois autocorrelations, and a reduction

Vichien Fugsukjit
Independent researcher, Bangkok
(designed and carried out with AI assistance: Anthropic Claude)

Version 5, November 2026 (corrective version; Version 4: 30 October 2026; Version 3: 21 October 2026; Version 2: 19 October 2026; Version 1: 16 October 2026)

Abstract

Let $h(N)$ be the maximal size of a Sidon set in $\{1,\dots,N\}$. Erdős asked whether $h(N) = N^{1/2} + O(1)$ (Guy C9); the weaker $h(N) = N^{1/2} + O_\varepsilon(N^\varepsilon)$ is Erdős problem #30. Both questions remain open, and we do not settle them. We study dilates of Singer perfect difference sets S ($|S| = q+1$, modulus $m = q^2+q+1$) and the number Z of points of a random dilate in a window of length Cq. *Proved:* (i) if the factorial moments of Z converge to the Poisson ones for every fixed order and every C, then $h(N) - N^{1/2}$ is unbounded (Proposition 2; experts regard this reduction as folklore, and we give a written proof); (ii) for a ratio ρ, the number $N(\rho)$ of triples of S with $c - a = \rho(b - a)$ is the number of $\mathbb{F}_q$-points of a twisted form of a curve that does not depend on q, so $N(\rho)/E - 1 = O_\rho(q^{-1/2})$ for 17 ratio orbits certified by exact computation (Theorem 1); (iii) for prime m, $\sum_\rho N(\rho)^2 = \sum_{\lambda,\mu}(|S \cap (\lambda S + \mu)|)_3$; (iv) $3k \mid N(\rho)$ for $q = p^k$, and a resonance theorem for cube $q = r^3$: $N(\rho) = r(q+1-e)$, $e \in \{0,1\}$, for the six ratios of the S_3-orbit of $r + r^{-1}$ (m prime); (v) for prime $q \geq 5$ and every unit λ, the per-dilation third moment $\sum_\mu(|S \cap (\lambda S + \mu)|)_3$ equals $P(\lambda) - (q-2)^2$, where $P(\lambda)$ counts solutions of $c\,y^\lambda + d(1-y)^\lambda = 1$ ($y \in \mathbb{F}_{q^3}$, $c, d \in \mathbb{F}_q^\times$); it is 0 at $\lambda = 2, -1$ (the classical conic fact), $(q-2)(q+1)$ at $\lambda = 3$ ($q \equiv 2 \pmod 3$), and $(q-2)(q+1) + b_q$ at $\lambda = -2$, where b_q is the q-th coefficient of the CM newform $\eta(2\tau)^3\eta(6\tau)^3$ (Theorem 3; one step uses an evaluation of a finite-field hypergeometric value due to Ono, in the form of Barman and Kalita). *Observed* (registrations are self-attested, see Section 14): to first order the k-th factorial moment of Z falls short of the random value by $\binom{k}{3}P_3 + \binom{k}{4}P_4$, where P_3, P_4 are the window probabilities of 3-term progressions and parallelograms ($k = 3,\dots,6$ at $C = 2$ on 129 primes; $k = 7$ at $C = 2$ and $k = 3,\dots,8$ at $C = 4$ on 64 of them); and $\sum_\rho(N(\rho) - E)^2$ is close to the order of the multiplier group times $\sum_\rho E$ for prime q and non-cube prime powers, while most of the excess at cube q is explained by (iv). *What remains:* a negative answer would follow from Poisson convergence of all fixed-order factorial moments. For order 3 this reduces to a variance bound, which by (iii) and (v) is a statement about the average of $P(\lambda)$ over all λ; for order ≥ 4 we have no such reduction.

1 Background

A set $A \subset \mathbb{Z}$ is Sidon if all sums $a + b$ ($a \leq b$) are distinct. Erdős and Turán [4] showed $h(N) \leq N^{1/2} + O(N^{1/4})$; Singer's construction [26] with the Baker–Harman–Pintz prime-gap theorem [1]

gives $h(N) \geq N^{1/2} - O(N^{0.2625})$; the constant in the $N^{1/4}$ term has since been improved several times (Balogh–Füredi–Roy [2] and later work listed at [5]). Erdős offered USD 500 for deciding whether $h(N) = N^{1/2} + O(1)$ [7, 9]. The site [5] lists the weaker question $h(N) = N^{1/2} + O_\varepsilon(N^\varepsilon)$ as problem #30 (USD 1000), with the $O(1)$ form as a remark; the evidence below concerns the $O(1)$ question only. A Sidon set A modulo m with $|A| = k$, cut at its largest cyclic gap G, gives a Sidon set of integers in an interval of length $m - G$; for Bose sets [3] ($m = q^2 - 1$, $k = q$) this gives $k - \sqrt{N} \approx G/(2q)$. Martin and O'Bryant [9] note that the question would be answered (excess unbounded) if Bose sets had gaps that are not $O(q)$. Dilating A by a unit t preserves the Sidon property and moves the gaps; the use of dilates was suggested in B. Green's comment on [6] (see also [15]; earlier computer searches of dilates and cuts: [27, 11]), and Shakan [10] proved that every $A \subset \mathbb{F}_p$ with $|A| > 1$ has a dilate with a gap of at least $2p/|A| - 2$. Throughout, "registered" and "blind" mean that a plan file and its SHA-256 hash were recorded locally before the computation; these records are self-attested (there is no third-party timestamp), and later amendments to a plan are listed in the release.

2 Dilate gaps of Bose sets

For every prime q in the ranges below we built Bose(q) $\subset \mathbb{Z}/(q^2-1)$, checked the Sidon property, and computed $G^*(q) = \max_t G(tA)$ over one multiplier per class $\{\pm t, \pm tq\}$. Positive control: $q = 829$ gives excess 10.2797, matching Zhang's bound $K(p_{145}) > 10.279$ ($p_{145} = 829$) [27].

- $101 \leq q \leq 2003$ (279 primes): a $\log q$ law fitted on $q \leq 1000$, $g^* \approx -6.92 + 3.63 \ln q$, gives held-out ($q > 1000$) RMSE 1.22, against 4.25 for a constant fitted on the same data. (Saturating bounded models were not tested.)
- Blind test (law registered before computing): 48 primes $2011 \leq q \leq 5881$ (every 10th prime in $(2003, 6000]$): RMSE 1.52 (log) against 7.34 (constant), mean signed error -0.57. The largest q (5881) gives a Sidon set of size 5881 in [1, 34,453,078] with excess 11.33; the largest excess in this set is 13.18 ($q = 5039$).
- Null model: random q-subsets of $\mathbb{Z}/(q^2-1)$ show the same growth (Bose slightly lower at 15 of 20 primes), so the $\log q$ growth is random-set behaviour; counts of 3-term relations $e_1L_1 + e_2L_2 + e_3L_3 \equiv 0$ among the discrete logarithms are at the random level (mean per-q ratio 0.983 over 11 primes $101 \leq q \leq 383$; percentile bootstrap over the primes, 95% interval $[0.96, 1.00]$).

3 The moment route for Singer sets

Let S be a Singer set [26] ($|S| = n = q + 1$, $m = q^2 + q + 1$), $\ell = Cq$, and $Z_{t,s} = \#\{a \in S : (ta + s) \bmod m \in [0, \ell)\}$ for units t and shifts s. Let F_k be the average of the falling factorial $(Z)_k$ and $H_k = (n)_k(\ell)_k/(m)_k$ its value for a uniformly random n-subset; $R_k = F_k/H_k - 1$.

Proposition 1. *For every unit t, $\mathrm{Var}_s\, Z_{t,s} = q\ell(m-\ell)/m^2$ and $R_2 = 0$. If m is prime, averaging over all units,*

$$R_3 = \sum_{\rho \neq 0,1} \Big(\frac{N(\rho)}{E} - 1\Big) \frac{A(\rho)}{\sum_\sigma A(\sigma)}, \qquad N(\rho) = \#\{(a,b,c) \in S^3 \text{ distinct} : c - a \equiv \rho(b-a)\},\ E = \frac{(n)_3}{m-2},$$

where $A(\rho)$ is the average over $u \in (\mathbb{Z}/m)^\times$ of the number of windows $[s, s+\ell)$ containing $\{0, u, u\rho\}$, and $\sum_\sigma A(\sigma) = \ell(\ell-1)(\ell-2)/(m-1)$.

(The first part holds because every nonzero residue is exactly one difference of tS and $(n)_2 = m-1$; for the second, m prime lets every triple be written as (a,b,c) with $c-a=\rho(b-a)$, and averaging over all units u sends it to every pair $(u,u\rho)$.) Since a perfect difference set has $N(-1)=N(2)=N(1/2)=0$ and $A(\rho)\approx \ell^2/(2m)$ for these ρ, the three forbidden ratios contribute $-3/(2Cq)(1+O(1/q))$ (within 0.2% for $q\ge 293$, 1.2% at $q=41$).

- $R_k \sim c_k(C)/q$ (moments estimated from 4000 sampled multipliers): registered blind test for $k=4$ on 11 new primes $409\le q\le 601$ ($C=2$: RMSE 0.0028 for c/q against 0.0238 for a constant; $C=4$: 0.0013 against 0.0112); the same is seen descriptively for $k=3,5,6$, with $|c_k|$ growing quickly in k.

- Exact over all units: registered tests on 18 primes $293\le q\le 1373$ (m prime) and, from a sealed data set opened under a frozen plan, 17 primes $1559\le q\le 2969$: the mean of $C\,qR_3$ is between -1.45 and -1.41, resp. between -1.47 and -1.46, inside the registered bands around $-3/2$ for all $C\in\{2,3,4,5,8\}$. The first test could not separate $-3/2$ from an earlier empirical value -1.17; the second band ($[-1.59,-1.31]$) excludes it.

- The RMS deviation D_{mid} of $N(\rho)/E$ over ratios of height 3–6: the registered decay test on $293\le q\le 1373$ alone *failed* (slope -0.13 ± 0.17, outside the band $[-0.65,-0.35]$; few independent values per q). A second test, frozen before opening the sealed data but pooling them with the already seen first set, gives slope -0.504 ± 0.098 over $293\le q\le 2969$ (the sealed set alone: -1.28 ± 0.39, i.e. noisy); $\sqrt{q}\,D_{\mathrm{mid}}$ lies between 0.49 and 1.84. Over all ratios the exponent is -0.50 in both sets, with $\sqrt{q}\,D_{\mathrm{all}}\approx 1.73$ for Singer sets against about 1.0 for random sets, consistent with counting noise inflated by the size-3 orbits of the multiplier q. These data are consistent with, but do not establish, a $q^{-1/2}$ law.

Sections 4–8 and 13 (new in Version 2; Sections 9–11 are new in Version 3) turn the heuristic of Version 1 into a theorem for each fixed ratio, give exact identities, explain the constants c_k, and state the reduction precisely.

4 The reduction

Proposition 2. *Fix $C>0$ and let $\ell=\lfloor Cq\rfloor$. Suppose that for every fixed $k\ge 1$, $R_k=R_k(q,C)\to 0$ as $q\to\infty$ through primes. Then for all large q some dilate of S has an empty arc of length ℓ, and $\limsup_{N\to\infty}(h(N)-N^{1/2})\ge (C+1)/2$. In particular, if this holds for every C, then $h(N)-N^{1/2}$ is unbounded.*

Proof. Choose (t,s) uniformly. Since $Z\le n$, inclusion–exclusion gives $\Pr(Z=0)=\sum_{k\ge 0}(-1)^k F_k/k!$, and the Bonferroni inequalities give $\Pr(Z=0)\ge \sum_{k=0}^{2K+1}(-1)^k F_k/k!$ for every K. Since $H_k=(n)_k(\ell)_k/(m)_k\to C^k$, the hypothesis gives $F_k\to C^k$ for each fixed k. Choose K with $\left|\sum_{k\le 2K+1}(-1)^k C^k/k!-e^{-C}\right|<e^{-C}/2$; then $\Pr(Z=0)>0$ for all large q, so some dilate tS misses a window of length ℓ. Rotating, tS lies in an interval of length $m-\ell$; a Sidon set modulo m is a Sidon set of integers, so we get $q+1$ elements in $[1,N]$ with $N=m-\ell=q^2+(1-C)q+O(1)$ and $q+1-N^{1/2}=(C+1)/2+O(1/q)$. □

The cutting step is classical (Section 1). The reduction of unbounded excess to the convergence of fixed-order factorial moments of window counts is regarded as folklore by experts (private communication); we did not find a written proof and include one here.

5 Universal curves: a theorem for each fixed ratio

Write $S = \{i : g^i \in L\}$ with g a generator of $\mathbb{F}_{q^3}^\times$ and L a 2-dimensional $\mathbb{F}_q$-subspace. For $\rho = u/v$ put $(\alpha, \beta, \gamma) = (u - v, v, -u)$, so $\alpha + \beta + \gamma = 0$, and let

$$C_{u,v}: \qquad s^\alpha r^\beta t^\gamma = 1, \qquad (s-1)^\alpha (r-1)^\beta (t-1)^\gamma = 1.$$

Theorem 1. *(a) For q prime, $N(\rho)$ equals the number of $\mathbb{F}_q$-points of a twisted form of $C_{u,v}$ off its diagonal $s = r = t$, up to $O_\rho(1)$; the twist is Frobenius composed with the order-3 map $s \mapsto 1 - 1/s$. (b) For $\rho \in \{-1, 2, 1/2\}$, $C_{u,v}$ is the diagonal, so $N(\rho) = 0$. (c) If $C_{u,v}$ minus the diagonal is absolutely irreducible over $\overline{\mathbb{Q}}$, then $N(\rho)/E - 1 = O_\rho(q^{-1/2})$ for all primes $q > q_0(\rho)$. (d) The hypothesis of (c) holds for every ratio of height ≤ 4 other than $-1, 2, 1/2$ (five orbits) and for the orbits (under $\rho \mapsto 1 - \rho, 1/\rho$) of $-5, -5/2, -5/4, -6, -6/5, -7, -7/2, -7/6, -8, -8/3, -8/5, -8/7$: 17 orbits in all. With (b), 18 of the 22 orbits of height ≤ 8 are settled.*

Proof sketch. (a) With $x = g^a$, $y = g^b$, $z = g^c$ the condition is $x^{u-v} z^v y^{-u} \in \mathbb{F}_q^\times$. Parametrise $P(L) \cong \mathbb{P}^1(\mathbb{F}_q)$ by $x = e + sf$; the three conjugate linear forms $e^{q^i} + sf^{q^i}$ have distinct roots, and a Möbius change of variable over $\mathbb{F}_{q^3}$ sending them to $\infty, 0, 1$ turns $w \in \mathbb{F}_q^\times$, i.e. $w = w^q = w^{q^2}$, into the two equations of $C_{u,v}$ (the constants cancel because $\alpha + \beta + \gamma = 0$); Frobenius becomes the stated twist. (b) For exponents $(-2, 1, 1)$ the equations give $r + t = 2s$ and $rt = s^2$. (c) Absolute irreducibility over $\overline{\mathbb{Q}}$ persists modulo all large primes; apply the Weil bound for singular curves [12] to a p-independent projective model; excluded points are $O_\rho(1)$ and $E = q + O(1)$. (d) Eliminating one variable by a resultant gives a plane curve G; we exhibit smooth points of G over number fields whose degrees have $\gcd 1$ (or one quadratic or cubic field over which G stays irreducible). A geometric component through a smooth K-point is defined over a subfield of K, so the number of components divides every such degree and equals 1. Birationality of the projection is checked by subresultants. Exact arithmetic in SymPy; certificates and code are released. A separate AI review checked (a)–(c) and the certificate logic, and confirmed (a) by exact counts for $q = 5, 7, 11, 13$. □

Numerically, the genus of $C_{u,v}$ appears to grow at least quadratically in the height (for example $(n-3)(n-4)/2$ for exponents $(-n, 1, n-1)$, $n \le 6$, by Riemann–Hurwitz with numerical monodromy), so Theorem 1 alone does not give $R_3 \to 0$; see Section 13. Four further orbits $(-5/3, -7/3, -7/4, -7/5)$ exceeded our 10-hour certification budget; none failed.

6 Exact identities

Let $I_\lambda(\mu) = |S \cap (\lambda S + \mu)|$ and let m be prime. Two ordered triples of S with the same ratio determine a unique affine map $x \mapsto \lambda x + \mu$ taking one to the other, hence

$$\sum_{\rho \ne 0,1} N(\rho)^2 = \sum_{\lambda \in (\mathbb{Z}/m)^\times} \sum_{\mu} \big(I_\lambda(\mu)\big)_3,$$

which we also checked exactly for $q = 5, 17, 41, 59$. For prime q each inner sum $\sum_\mu (I_\lambda(\mu))_3$ is an explicit point count over $\mathbb{F}_{q^3}$ (Lemma 1 in Section 12). Since every nonzero residue is a unique difference, $\sum_\mu (I_\lambda(\mu))_2 = m - 1$ for every λ: second moments are exactly Poisson. For the multipliers $\lambda \in \{1, q, q^2\}$, λS is a translate of S and one μ gives $I = n$, contributing exactly $3(n)_3$. Under $x \mapsto x^\lambda$, a dilate by λ whose base-q digits have sum d maps the line L into a curve of degree d, so $I_\lambda \le d$ by Bézout unless L lies on that curve (this does not happen

for $d = 2$ and odd q); for $d = 2$ (for example $\lambda = -1$, a conic) $(I_\lambda)_3 = 0$. Numerically, the non-multiplier λ with $\sum_\mu (I_\lambda(\mu))_3 = 0$ are exactly the nine elements $\{-1, 2, 1/2\} \cdot \{1, q, q^2\}$ for $q = 41, 59, 167, 173, 293, 383$ (all λ computed); the arc and oval facts behind -1 and 2 are classical [13, 14].

7 Higher moments: the forbidden-structure law (blind tests)

A Sidon set contains no 3-term progression and no non-trivial parallelogram $x + y = z + w$. Let $P_3(\ell)$, resp. $P_4(\ell)$, be the probability that an ordered distinct triple of positions in $[0, \ell)$ contains a 3-term progression, resp. that an ordered distinct quadruple is a parallelogram; $P_3 \sim 3/(2\ell)$ and $P_4 \sim 2/\ell$. The first-order prediction is

$$R_k \approx -\Big[\tbinom{k}{3} P_3(\ell) + \tbinom{k}{4} P_4(\ell)\Big] \sim -\frac{\frac{3}{2}\binom{k}{3} + 2\binom{k}{4}}{Cq}.$$

Two registered blind tests (blind for $k \geq 4$ in the first, for all k in the second) used plans frozen with a recorded SHA-256 before computing (self-attested; two amendments made before any computation are listed in the release), 3000 sampled unit classes per q, and the inverse-variance pooled ratio measured/predicted. On 65 primes $607 \leq q \leq 1499$ with $C = 2$, $k = 3, \ldots, 6$ gave $1.02, 1.06, 1.06, 1.05$ (all 95% intervals contain 1). On the other 64 primes of that range, $C = 2$ gave $0.97, 0.97, 0.97, 0.97, 1.01$ for $k = 3, \ldots, 7$ (the $k = 8$ cell is void because its null control failed), and $C = 4$ gave $0.95, 0.98, 1.00, 1.00, 1.00, 1.01$ for $k = 3, \ldots, 8$. Random sets were within 3 standard errors of 0 in every other cell. So no new forbidden structure appears up to 8 points, and $c_k \sim k^4$ explains the fast growth noted in Version 1.

8 Variance law and prime powers

Conjecture 1. *$\sum_\rho (N(\rho) - E)^2 = |G| \sum_\rho E\,(1 + o(1))$, where $|G|$ is the order of the multiplier group ($|G| = 3$ for prime q, $|G| = 3k$ for $q = p^k$), the sums run over ρ with ρ and $\rho - 1$ units, and E is the mean of $N(\rho)$ over these ρ (equal to $(n)_3/(m-2)$ when m is prime).*

By Proposition 4 (Section 9), $3k \mid N(\rho)$ for every prime power $q = p^k$. For prime q with m prime, the ratio has mean 2.998 (s.d. 0.008, range $[2.979, 3.019]$) over 37 primes $293 \leq q \leq 2969$. For prime powers (Singer sets built in $\mathbb{F}_{p^{3k}}$; registered test) the ratio divided by $3k$ is given in Table 1. The per-λ errors $\sum_\mu (I_\lambda(\mu))_3$ minus the random value have mean about 0; excluding the nine null λ they are close to Gaussian, with standard deviation about $4.2\,q$ ($q = 383$: $4.21q$, kurtosis 3.01).

Proposition 3 (conditional)**.** *If $\sum_\rho (N(\rho) - E)^2 \ll q^3$, then $R_3 = O\big(\sqrt{\log q/q}\big)$.*

Proof sketch. By the Proposition of Section 3, $R_3 = \sum_\rho (N/E - 1)(w(\rho) - \bar{w})$ with $w = A/\sum A$, because $\sum_\rho N(\rho) = (n)_3$ makes the uniform part vanish. Cauchy–Schwarz gives $|R_3| \leq E^{-1}\big(\sum_\rho (N - E)^2\big)^{1/2} \|w - \bar{w}\|_2$, and lattice-point counting for $\{(x, y) : y \equiv \rho x\}$ in $(-\ell, \ell)^2$ gives $\|w - \bar{w}\|_2 \ll \sqrt{\log q}/q$. □

9 Orbits of the multiplier group

Proposition 4. *Let $q = p^k$ and let $N(\rho)$ count triples $(a, b, c) \in S^3$ with $b - a$ a unit and $c - a = \rho(b - a)$ (for prime m this is the count of Section 3). Then $3k \mid N(\rho)$ for every ρ.*

q	p^k	$3k$	good ρ	ratio	ratio/$3k$
121	11^2	6	2975	6.13	1.022
125	5^3	9	14059	9.91	1.101
128	2^7	21	11725	20.30	0.967
169	13^2	6	9145	6.53	1.089
243	3^5	15	50149	14.62	0.974
256	2^8	24	13145	24.35	1.014
289	17^2	6	16775	6.32	1.053
343	7^3	9	37135	10.62	1.180
361	19^2	6	30625	6.12	1.020
512	2^9	27	262655	28.04	1.039
529	23^2	6	55055	5.75	0.959
625	5^4	12	86855	12.05	1.004
729	3^6	18	449225	18.76	1.042
841	29^2	6	192335	6.00	1.000
961	31^2	6	195755	6.00	0.999
1024	2^{10}	30	245105	29.96	0.999
1331	11^3	9	1772891	9.48	1.053
1369	37^2	6	386425	6.07	1.012
1681	41^2	6	937945	6.00	1.001
1849	43^2	6	947903	5.98	0.996
2048	2^{11}	33	2997385	33.22	1.007
2187	3^7	21	4048957	21.03	1.001
2197	13^3	9	1609667	9.79	1.088

Table 1: Variance ratio $\sum(N-E)^2/\sum E$ over ρ with $\rho, \rho-1$ units, for Singer sets with $q = p^k$, $k \geq 2$ (registered test: within 10% of $3k$ for at least 80% of $q \geq 121$; here 21 of 23, with $q = 125$ (1.1013) a borderline miss under the rule $|\text{ratio}/3k - 1| \leq 0.10$; the two misses and most of the excess at cube q are accounted for by the resonance of Section 10 (Table 2); for $q < 121$, outside the registered range, the ratio/$3k$ is far from 1, e.g. 0.20 at $q = 9$, 0.69 at $q = 49$, 0.83 at $q = 81$; later runs: all 11 later non-cube q $(2209, 2401, 2809, 3125, 3481, 3721, 4489, 6561, 8192, 14641, 16384, 16807)$ give ratio/$3k \in [0.9980, 1.0026]$; the later cube $q = 4096, 4913, 6859$ give $1.022, 1.038, 1.064$, see Table 2).

Proof. By the multiplier theorem, some translate of S is fixed by multiplication by p [16, 17]. N is translation invariant, so assume $pS = S$. We have $p^{3k} - 1 = (q-1)m$. If $p^j \equiv 1 \pmod{m}$ with $0 < j < 3k$, then $m \mid p^d - 1$ for $d = \gcd(j, 3k) \leq 2k$, which is impossible because $0 < p^d - 1 < m$. So p has order $3k$ modulo m. The map $(a, b, c) \mapsto (pa, pb, pc)$ preserves the set of triples. A triple fixed by p^j, $0 < j < 3k$, would give $(p^j - 1)(b - a) = 0$ with $b - a$ a unit, which is impossible. So the action is free and every orbit has $3k$ elements. □

This is elementary and probably known. The order statement and the freeness step are machine-checked in Lean 4 with Mathlib (files `C9Order.lean`, `C9Orbit.lean`; standard axioms only). The multiplier theorem itself is not formalised. So Conjecture 1 says that $M = N/3k$ is Poisson-like. This holds for non-cube q (Table 1) but fails for cube q; the next two sections explain most of this failure.

10 A resonance theorem for cube q

Let $q = r^3$, where r is a prime power. Let $F = \mathbb{F}_q$ and $K = \mathbb{F}_{q^3} = \mathbb{F}_{r^9}$. Write $\tau(u) = u^r$ and $\varphi = \tau^3$ (the q-Frobenius). Let $L = \ker \mathrm{Tr}_{K/F}$, so that $S = \{i : g^i \in L\}$. Since τ commutes with φ, we have $\tau(L) = L$. Moreover φ is F-linear on L with $\varphi^3 = 1$. Since $r^9 = q^3 \equiv 1$ and $r^3 = q \not\equiv 1 \pmod m$, r has order 9 modulo m. Put $\rho_1 = r + r^{-1} \bmod m$.

Theorem 2. *Let $x \in P(L)$ be a point of the Singer line.*

- *If $\varphi x \neq x$ (projectively), then there are exactly r points $y \in P(L) \setminus \{x\}$ with $z/x = (y/x)^{\rho_1}$ for some $z \in P(L)$.*
- *If $\varphi x = x$, there are none.*

The number e of φ-fixed points of $P(L)$ is 0, 1 or 2 according as $r \equiv 2$, 0 or 1 (mod 3). Consequently, if m is prime, $N(\rho) = r\,(q + 1 - e)$ for each of the six ratios ρ in the S_3-orbit of ρ_1; this orbit contains $\rho_0 = r^4 + r^{-4} = (r + 1)r^4$. If $r \equiv 1 \pmod 3$ then $3 \mid m$, so for prime m we have $e \in \{0, 1\}$, and $e = 1$ exactly when $p = 3$.

Proof. (i) *Reduction.* Put $u = y/x$. Then $y \in P(L)$ iff $\mathrm{Tr}(xu) = 0$. If x is not φ-fixed, then $\psi = \varphi(x)/x \notin F$ and $\mathrm{Tr}(x\psi) = \mathrm{Tr}(\varphi x) = 0$, so $\{u : \mathrm{Tr}(xu) = 0\} = F + F\psi$. In $K^\times/F^\times$ the exponent ρ_1 acts as $u \mapsto \tau(u)\tau^{-1}(u)$, so we need $z = x\,\tau(u)\,\tau^{-1}(u) \in F^\times L$. That is, $\mathrm{Tr}(x\,\tau(u)\,\tau^{-1}(u)) = 0$. Take $u = 1 + s\psi$ with $s \in F$; on F we have $\tau s = s^r$ and $\tau^{-1}s = s^{r^2}$. Using $\mathrm{Tr}(x) = 0$, the condition reads $As^r + Bs^{r^2} + Cs^{r+r^2} = 0$, where $A = \mathrm{Tr}(x\tau\psi)$, $B = \mathrm{Tr}(x\tau^{-1}\psi)$ and $C = \mathrm{Tr}(x\,\tau\psi\,\tau^{-1}\psi)$ lie in F. Remove $s = 0$ (that is, $y = x$) and put $Z = s^{-r}$, with $s = \infty$ corresponding to $Z = 0$. This gives the affine $\mathbb{F}_r$-linear equation

$$AZ^r + BZ + C = 0, \qquad Z \in F.$$

(ii) *Two identities.* Write $\xi_j = \tau^j(x)$ for j mod 9, with $\xi_j + \xi_{j+3} + \xi_{j+6} = 0$. Then A, B, C are rational functions of $\xi_0, \ldots, \xi_5$ with integer coefficients and monomial denominators, and τ acts as the index shift. Let D be the Dickson matrix of $f(Z) = AZ^r + BZ$. Exact computation in Sage gives

$$N_{F/\mathbb{F}_r}(A) + N_{F/\mathbb{F}_r}(B) = 0 \quad \text{and} \quad \mathrm{adj}(D)\,(C, C^r, C^{r^2})^T = 0.$$

After clearing the (nonzero) denominators, both are identities between polynomials with integer coefficients that hold over $\mathbb{Q}$. They therefore hold in every characteristic. They were also rechecked directly over $\mathbb{F}_2, \mathbb{F}_3, \mathbb{F}_5, \mathbb{F}_7$. (iii) *Non-degeneracy.* The numerator of A factors as $(\xi_1\xi_3 - \xi_0\xi_4)(\xi_1^2 + \xi_1\xi_4 + \xi_4^2)$, and the denominator is $-\xi_1\xi_4\xi_7 \neq 0$. The second factor vanishes iff ψ is a root of $t^2 + t + 1$. The first vanishes iff $\tau x = cx$ with $c \in F$, which gives $\varphi x = N_{F/\mathbb{F}_r}(c)\,x$. In both cases x is φ-fixed. So $A \neq 0$, hence $N(B) = -N(A) \neq 0$ and $\mathrm{rank}\, D = 2$. (iv) *Counting.* $N(-B/A) = N(-1)\,N(B)/N(A) = (-1)^{1+r+r^2} \cdot (-1) = 1$, so the kernel of f has exactly r elements and $|\mathrm{Im}\, f| = r^2$. Since $\mathrm{adj}(D)\,D = \det(D)\,I = 0$, each row (w_0, w_1, w_2) of $\mathrm{adj}(D)$ annihilates (b, b^r, b^{r^2}) for every $b \in \mathrm{Im}\, f$. A nonzero row gives a nonzero linearised polynomial of degree at most r^2. Its zero set therefore has at most r^2 elements and contains $\mathrm{Im}\, f$, so it equals $\mathrm{Im}\, f$. By (ii), $-C \in \mathrm{Im}\, f$, so there are exactly r solutions Z, that is, r partners y. (v) *Fixed points.* The φ-fixed points of $P(L)$ are the eigenlines of the F-linear map $\varphi|_L$, whose eigenvalues are cube roots of unity in F. The eigenvalue 1 needs $x \in F$, and then $\mathrm{Tr}(x) = 3x$ vanishes only if $p = 3$.

- If $r \equiv 2 \pmod 3$, then F has no primitive cube root of unity, so $e = 0$.
- If $p = 3$, then $\varphi|_L$ is unipotent with fixed line $F \cdot 1$, so $e = 1$.

- If $r \equiv 1 \pmod 3$, then $\varphi|_L$ has the two eigenvalues ω, ω^2 ($\omega \in F$ a primitive cube root of unity), so $e = 2$.

For a fixed point:

- If $\varphi x = \omega x$, then K splits into φ-eigenlines, which τ preserves, so $\tau x = cx$ and $\tau^{-1}x = c'x$ with $c, c' \in F$, and $\{u : \mathrm{Tr}(xu) = 0\} = F + Fx$. For $u = \beta_0 + \beta x$ the condition becomes $3cc'\beta^{r+r^2}x^3 = 0$, i.e. $\beta = 0$, i.e. $y = x$.

- If $p = 3$, then $L = F + Fv$ with $\varphi v = v + \alpha$, $\alpha \in F^\times$. For $y = \beta_0 + \beta v$, the condition reduces to $\mathrm{Tr}(v^2)\,\beta^{r+r^2} = 0$ with $\mathrm{Tr}(v^2) = -\alpha^2 \neq 0$, so again $y = x$.

Summing over x gives $N(\rho_1)$ when m is prime, because then every nonzero difference is a unit. The maps $(a, b, c) \mapsto (b, a, c)$ and $(a, b, c) \mapsto (a, c, b)$ realise $\rho \mapsto 1 - \rho$ and $\rho \mapsto 1/\rho$ and preserve the unit conditions, so N is constant on the S_3-orbit. □

The theorem was also checked by independent brute force for $r \leq 13$; for $q = 8, 27, 512, 1331$ it was checked on all six ratios. A separate AI review accepted it after the fixes included here.

Algebraic form. The six ratios of the orbit are roots modulo m of $x^3 - 3x + 1$ and $x^3 - 3x^2 + 1$. For prime m they are all of the roots; for composite m these cubics can have more roots, by the Chinese remainder theorem. The first is the minimal polynomial of $2\cos(2\pi/9) = \zeta_9 + \zeta_9^{-1}$, with $\zeta_9 \mapsto r$. Its root set is invariant under $x \mapsto 1/(1 - x)$, the inverse of the order-3 map $s \mapsto 1 - 1/s$ of Theorem 1(a).

11 The resonance and the cube excess

For the six resonant ratios $M(\rho) \approx r\mu$. If all other ratios behaved as in Conjecture 1, then $\mathrm{Var}\, M/\mathbb{E}\, M$ would exceed 1 by about

$$\Delta = 6(r-1)^2\mu/n_{\mathrm{good}}, \qquad n_{\mathrm{good}} = \prod_{\ell^e \| m} \ell^{e-1}(\ell - 2), \quad \mu = \frac{(q+1)q(q-1)}{3k\, n_{\mathrm{good}}} \prod_{\ell \mid m} \frac{(\ell-1)(\ell-2)}{\ell^2}.$$

This is a heuristic, not a theorem; it decays like $q^{-1/3}$.

Registration.

- The value of Δ was registered before computation only for $q = 12167$.

- The statements "the six largest $M(\rho)$ are roots of the cubics" and "$M/\mu \in [0.8r, 1.2r]$" were registered before computation for $q = 4913, 6859$. For $q = 15625$ the registration was "the cubic orbit is among the twelve largest, with $M/\mu \in [20, 30]$".

- All other entries of Table 2 are retrospective.

The plan file and its SHA-256 ledger are released. Registration times are self-attested.

For $k = 3$ and $q \geq 1331$, the formula gives 94–100% of the observed excess. In the other rows of Table 2 it gives 54–100% (lowest at $q = 512$). Two earlier registered readings failed at $q = 6859$: a $q^{-1/2}$ decay band, and a $p \equiv 1 \pmod 3$ effect. Registered tests at $q = 24389$ and $q = 32768$ are pending. For $k = 6, 9, 12$ a second, weaker orbit appears, with $M/\mu = 2.77$, 3.91, 4.00, 2.98 at $q = 729, 4096, 15625, 19683$. Its ratios are not roots of integer polynomials of degree ≤ 4 with coefficients bounded by 4, and they are not of the form $p^a \pm p^b \pm p^c$. We have not identified this orbit. At $q = 19683$ it contributes only about 10% of the residual 0.0015; most of the residual comes from several hundred ratios with $M/\mu \approx 2$. Weaker levels with $M/\mu \approx 2$ also occur for $k = 3$ ($q = 4913, 12167$) and for the non-cube control $q = 16384$. They are also unexplained.

q	k	Δ formula	observed	top-6 M/μ (r)	status
125	3	0.089	0.101	4.91 (5)	retrospective
343	3	0.143	0.180	7.19 (7)	retrospective
512	9	0.021	0.039	8.03 (8)	retrospective
729	6	0.032	0.042	8.86 (9)	retrospective
1331	3	0.0500	0.0534	11.0 (11)	retrospective (local computation)
2197	3	0.0873	0.0877	– (13)	retrospective
4096	12	0.020	0.022	15.93 (16)	retrospective
4913	3	0.0368	0.0377	17.01 (17)	top-6, M/μ registered
6859	3	0.0631	0.0635	18.94 (19)	top-6, M/μ registered; Δ retrospective
12167	3	0.0281	0.0281	22.99 (23)	**Δ registered (blind)**; top-6, M/μ retrospective
15625	6	0.0261	0.0261	24.95 (25)	top-6, M/μ registered; Δ not registered
19683	9	0.0077	0.0092	27.01 (27)	exploratory ($p = 3$)

Table 2: The resonance and the cube excess. Wherever the six largest $M(\rho)$ were recorded, they are roots of the two cubics and $M/\mu \approx r$. Non-cube controls ($q = 8192, 14641, 16384$) give ratio/$3k \in [0.9997, 1.0024]$. For composite m, M uses the count restricted to unit differences (for example $N(\rho_1) = 576$ at $q = 125$, against $r(q + 1 - e) = 630$ for all triples in Theorem 2), so $M/\mu \approx r$ is approximate partly for this reason. Rows $q = 4096, 4913, 6859$ were computed on Kaggle with `run_nb_top12.py` (all triples) and rows $q \geq 12167$ with the orbit-reduced program `run_orb.py`; both are released with Version 5, with a regression test of `run_orb.py` against an independent brute-force count and a per-row provenance table.

12 Galois autocorrelation of Jacobi sums: exact values

Let $K = \mathbb{F}_{q^3}$ (q an odd prime; Theorem 3 needs $q \geq 5$), $m = q^2 + q + 1$, and let $L : K^\times \to \mathbb{Z}/m$ be the discrete logarithm modulo m; its kernel is $\mathbb{F}_q^\times$. Let ψ be a character of $K^\times$ of order m (trivial on $\mathbb{F}_q^\times$) and put $J[a, b] = \sum_{y \in K \setminus \{0,1\}} \psi^a(1 - y)\psi^b(y)$, a Jacobi sum. For a unit l modulo m define the *Galois autocorrelation*

$$C(l) = \sum_{a \neq 0} \sum_{b} J[a, b]\, \overline{J[la, lb]},$$

i.e. the correlation of the Jacobi sums with their conjugates under $\zeta_m \mapsto \zeta_m^l$. By Lemma 1 below, these correlations are the per-dilation third moments of Section 6 in another form. We also use the elementary count

$$P(l) = \#\{(y, c, d) :\ y \in K \setminus \{0, 1\},\ c, d \in \mathbb{F}_q^\times,\ c\, y^l + d\, (1 - y)^l = 1\}$$

(equivalently, pairs y, y' with $L(1 - y') = l\, L(1 - y)$ and $L(y') = l\, L(y)$).

Proposition 5. *$C(l) = m^2 P(l) - ((q - 1)m - 1)^2 - (m - 1)$ for every unit l, and $P(l) = P(l^{-1})$.*

Proof. By orthogonality, $\sum_{a,b} J[a, b]\overline{J[la, lb]} = m^2 \#\{(y, y') : L(1 - y) = lL(1 - y'),\ L(y) = lL(y')\} = m^2 P(l^{-1})$, by the pair description of P; counting the same set of pairs with the other coordinate as the free variable gives $m^2 P(l)$, hence $P(l^{-1}) = P(l)$. (The power y^l depends on the integer representative of l only up to a factor in $\mathbb{F}_q^\times$, which is absorbed into (c, d).) The row $a = 0$ equals $\sum_b J[0, b]\overline{J[0, lb]}$ with $J[0, 0] = q^3 - 2 = (q - 1)m - 1$ and $J[0, b] = -1$ for $b \neq 0$. □

Lemma 1. *For every unit l modulo m, $P(l) - (q - 2)^2 = \sum_\mu \big(I_l(\mu)\big)_3$, where $I_l(\mu) = |S \cap (lS + \mu)|$ as in Section 6.*

Proof. Identify $\mathbb{Z}/m$ with the points of $\mathrm{PG}(2,q) = K^\times/\mathbb{F}_q^\times$ by the discrete logarithm; S is a line, translation by μ is multiplication by g^μ, and dilation by l is $x \mapsto x^l$. Because l is a unit modulo m, this map is a bijection of the points, so it maps distinct points to distinct points (this is the only use of the unit condition, and it is essential: for non-units the identity fails). Since the Singer group acts regularly on lines, $\sum_\mu (I_l(\mu))_3$ is the number T_S of ordered triples of distinct points of S whose l-th powers are collinear. Multiplication by g^ν maps lines to lines and $(g^\nu x)^l = g^{l\nu}x^l$, so $T_{S'} = T_S$ for every line S'. Counting the triples of distinct collinear points with collinear l-th powers in two ways (by their line, or by their first point, which multiplication moves to 1 freely) gives $m\,T_S = m\,T_1$, where T_1 is the number of ordered pairs (x_2, x_3) of points such that $1, x_2, x_3$ are distinct and collinear and $1, x_2^l, x_3^l$ are collinear. The map $y \mapsto ([y], [1-y])$ is a bijection from $K \setminus \mathbb{F}_q$ onto the pairs (x_2, x_3) with $1, x_2, x_3$ distinct and collinear: given such a pair, $1 = ay_0 + bz_0$ for representatives y_0, z_0 and unique $a, b \in \mathbb{F}_q^\times$, and $y = ay_0$ is the unique representative of x_2 with $1 - y$ representing x_3. Collinearity of the three distinct points $1, y^l, (1-y)^l$ means that $c\,y^l + d(1-y)^l = 1$ for exactly one $(c,d) \in (\mathbb{F}_q^\times)^2$. Hence the sum counts the solutions of $P(l)$ with $y \notin \mathbb{F}_q$. For $y \in \mathbb{F}_q \setminus \{0,1\}$ there are exactly $q-2$ admissible pairs (c,d), which gives the term $(q-2)^2$. □

The lemma was found by an AI referee and checked numerically in the release (`v5_lemma_I3.py`, $5 \le q \le 19$, ten values of l each, with the controls $l = 1$, where both sides equal $q^3 - q$, and $l = -1, 2$, where both sides vanish). In particular $P(l) \ge (q-2)^2$ always. For $l = 3$ and $l = -2$ the equation is a polynomial over $\mathbb{F}_q$ of degree ≤ 4 in y, and its roots in $K \setminus \mathbb{F}_q$ come from irreducible cubic factors; hence $P(l) = (q-2)^2 + 3I(l)$, where $I(l)$ is the number of $(c,d) \in (\mathbb{F}_q^\times)^2$ for which the polynomial has an irreducible cubic factor (I is not the triple count $N(\rho)$ of the earlier sections).

Theorem 3. *Let $q \ge 5$ be prime. Define $b_q = 0$ if $q \equiv 2 \pmod 3$ and $b_q = 4x^2 - 2q$ if $q = x^2 + 3y^2$ ($q \equiv 1 \pmod 3$). (b_q is the q-th coefficient of the weight-3 newform $\eta(2\tau)^3\eta(6\tau)^3$ of level 12, which has CM by $\mathbb{Q}(\sqrt{-3})$ [8]; this is the standard Hecke-character description of a CM form, and we checked it for all primes $5 \le q \le 199$. It is not used below.)*

- *(a) $P(2) = P(-1) = (q-2)^2$, the least possible value. By Lemma 1 this is equivalent to the conic fact $(I_\lambda)_3 = 0$ of Section 6 [13, 14]; we include it for completeness.*
- *(b) If $q \equiv 2 \pmod 3$, then $P(3) = (q-2)(2q-1)$.*
- *(c) $I(-2) = \big((q-2)(q+1) + b_q\big)/3$, hence $P(-2) = (q-2)(2q-1) + b_q$. In particular $P(3) = P(-2)$ when $q \equiv 2 \pmod 3$.*

Equivalently (Lemma 1), $\sum_\mu (I_3(\mu))_3 = (q-2)(q+1)$ for $q \equiv 2 \pmod 3$ and $\sum_\mu (I_{-2}(\mu))_3 = (q-2)(q+1) + b_q$. (The unit condition: $l = \pm 3$ is a unit modulo m only for $q \equiv 2 \pmod 3$.)

Corollary. If m is prime, then $\sum_{\rho \ne 0,1} N(\rho)^2 = \sum_\lambda \big(P(\lambda) - (q-2)^2\big)$, the sum over all units λ (Section 6 and Lemma 1). So Conjecture 1, which is a sufficient condition for the order-3 case of the hypothesis of Proposition 2, is a statement about the average of $P(\lambda)$ over all λ. Theorem 3 evaluates four of the roughly q^2 terms.

Proof of (a). For $l = 2$ and $l = -1$ (after clearing denominators) the equation is a nonzero polynomial of degree ≤ 2 over $\mathbb{F}_q$. Its roots lie in $\mathbb{F}_{q^2} \cap K = \mathbb{F}_q$, so only the $(q-2)^2$ solutions with $y \in \mathbb{F}_q$ remain. □

Proof of (b), sketch. If $c = d$ the polynomial has degree 2 and, as in (a), has no root outside $\mathbb{F}_q$. For $c \neq d$ write $c\,y^3 + d(1-y)^3 - 1$ as $(c-d)\,g(y)$ with $g(y) = y^3 + b(y^2 - y) + e$. The map $(c,d) \mapsto (b,e)$ is a bijection onto the pairs with $b \notin \{0,-3\}$ and $b \neq 3e$. On the excluded lines, g has a root in $\mathbb{F}_q$, since cubing is bijective on $\mathbb{F}_q$. In the family $x^3 + b(x^2 - x) + e$, two distinct roots determine the third, except when the two roots sum to 1, which needs $a^2 - a + 1 = 0$; this has no solution because -3 is a non-square. Counting triple roots (2), double roots $(q-3)$, ordered pairs of roots and root incidences gives $(q^2 - 4q + 7)/6$ split, $(q^2+1)/2$ one-root and $(q-2)(q+1)/3$ irreducible members. Hence $I(3) = (q-2)(q+1)/3$ and $P(3) = (q-2)^2 + 3I(3)$. □

Proof of (c), outline. The full proof is in the release (file `N2_PROOF.md`); we list the steps. Let ϕ be the quadratic character of $\mathbb{F}_q$. (1) For distinct roots $y_1, y_2 \in \mathbb{F}_q \setminus \{0,1\}$ the two root conditions are linear in (c,d), with determinant $(y_2 - y_1)(y_1 + y_2 - 2y_1y_2)$, $c = y_1^2 y_2^2 (y_2 - y_1)(2 - y_1 - y_2)/\det$ and $d = (1-y_1)^2(1-y_2)^2(y_2^2 - y_1^2)/\det$. Put $u = y_1 + y_2$ and $v = y_1 y_2$. (2) The number of admissible unordered pairs is $P_2 = (q-3)(q-5)/2$. (3) The residual quadratic of the quartic has discriminant $u(u-2)(2v-u+2)/(2v-u)$. Summing its quadratic character over admissible pairs gives $S_2 = (T - 4q + 14)/2$ with $T = \sum_{x,z} \phi\big((x^2-1)(z^2-1)(x^2+1-2z)\big)$. The excluded lines contribute $2q-4$, $2q-6$, -3 and 0. (4) A change of variables gives $T = 1 + \phi(-1)G(4)$, where $G(\lambda) = \sum_{x,y} \phi\big(xy(1-x)(1-y)(x-\lambda y)\big)$, and $q^2\,{}_3F_2(4) = \phi(-1)G(4)$ for Greene's ${}_3F_2$ with quadratic characters. (5) The repeated-root types have counts $E_1 = (q+1-z_0)/2$ (type $(1)^2(2)$), $E_2 = (q-5-z_0)/2$ (type $(1)(1)(1)^2$) and $E_3 = z_0/2$ (type $(1)^2(1)^2$), where $z_0 = 1 + \phi(-3)$. A double root a forces $c = a^3$ and $d = -(a-1)^3$. (6) Counting roots, pairs and character-weighted pairs by factorisation type gives $I(-2) = (q-2)^2 - E_1 - \frac{4}{3}P_2 + \frac{2}{3}S_2 - \frac{1}{3}E_2 - \frac{4}{3}E_3$. (7) The value $q^2\,{}_3F_2(4) = q + b_q$ follows from [18, Thm. 1.7] (Greene's ${}_3F_2$ [21], character $S = \phi$, allowed there since ϕ has order 2); the authors of [18] state that $S = \phi$ gives a result of Ono [25]. For $q \equiv 2 \pmod 3$ the value is $1/q$. For $q \equiv 1 \pmod 3$ the theorem gives $[(\phi\chi_3 \text{ over } \chi_3) + (\phi\chi_3^2 \text{ over } \chi_3^2)]^2 - 1/q$; here the Hasse–Davenport relation [22] gives $J(\phi\chi_3, \bar{\chi}_3) = \chi_3(2)\overline{J(\chi_3,\chi_3)}$, and the cubic character of 2 [22] makes the real part $\pm x$, so $q^2\,{}_3F_2(4) = 4x^2 - q = q + b_q$. Substituting (2)–(7) into (6) gives (c). □

Verification. Each step was checked by exact computation:

- steps (1)–(6) on all primes $5 \le q \le 31$;
- step (7): the Jacobi-sum part on all primes $q \equiv 1 \pmod 3$, $7 \le q \le 97$ (including the choice of associate), the value $1/q$ for $q \equiv 2 \pmod 3$ on all primes $5 \le q \le 97$;
- the final formula for $I(-2)$ on all primes $5 \le q \le 53$, by an independent brute-force count;
- the identity $b_q = a_q(\eta(2\tau)^3\eta(6\tau)^3)$ on all primes $5 \le q \le 199$ (numerical only).

The symbolic identities were checked with SymPy. A separate AI instance reviewed steps (1)–(6) with its own code and found no error. Step (7) for $q \equiv 1 \pmod 3$ was written after that review; it was checked numerically and by the AI referee reports prepared for Version 5, which corrected one sentence of the released proof (the choice of associate). None of these is a human expert review.

Geometric explanation (heuristic, not used in the proof). In the twisted form, the level sets of $\mathrm{Tr}(\alpha)\mathrm{Tr}(\alpha^{-1})$ are the members of the pencil $(x+y)(y+z)(z+x) = k\,xyz$. This pencil is the extremal rational elliptic surface with singular fibres I_6, I_3, I_2, I_1, i.e. the universal curve for $\Gamma_1(6)$. The fibre types were computed exactly. We expect the double cover branched at the I_3 and I_1 fibres to be the singular K3 surface with fibres $3I_6 + 3I_2$ and transcendental lattice $\mathrm{diag}(2,6)$ [20],

which has complex multiplication by $\mathbb{Q}(\sqrt{-3})$, but we have not written down the isomorphism. This would explain why the error term is a CM coefficient.

Observations.

- $P(4) = P(-2)$, i.e. $\sum_\mu (I_4)_3 = \sum_\mu (I_{-2})_3$, for all 20 primes $5 \le q \le 79$ tested. Besides $P(ql) = P(l)$ (Frobenius) and $P(l^{-1}) = P(l)$, a search over Möbius maps with integer coefficients in $[-3, 3]$ found no further symmetry of P ($q = 11, \dots, 23$), so this equality is not explained by such a map.

- Among the rationals a/b with $|a|, |b| \le 12$ that are units modulo m for every tested q ($11 \le q \le 41$; this excludes all a/b involving 3 or 7), the only classes of equal P beyond $\{l, l^{-1}\}$ are $\{2, -1, \frac{1}{2}\}$ and $\{-2, -\frac{1}{2}, 4, \frac{1}{4}\}$. For $q \equiv 2 \pmod 3$ the second class also contains 3 and $\frac{1}{3}$, by Theorem 3(b),(c).

- For other small heights, $(P(l) - (q-2)^2 - q^2)/q$ fluctuates more as the height grows: over the primes $47 \le q \le 79$ (only 4 to 8 values of q for each l) its standard deviation is about 1 at $l = -2$ and about 4 at $l = 7$, but it is not monotone in the height, so this is a rough observation (smaller q include multiplier coincidences such as $l = q$, which inflate it). This is consistent with cohomology that grows with the height, so these exact formulas do not by themselves give the averaged bound of Section 13.

- $N(\rho)$ is an average of root counts of the trinomials $c\,y^\rho + y - 1$. Kelley proved, under GRH, that the root counts of random-coefficient trinomials over prime fields are Poisson, and conjectured the same for a fixed prime [23, 24, 19]. The Poisson-type laws of Sections 7–8 are an extension-field analogue in a different regime.

13 What remains

By Proposition 2, a negative answer to the $O(1)$ question would follow from Poisson convergence of all fixed-order factorial moments of Z, that is, $R_k \to 0$ for each fixed k and each C. This is a full Poisson limit law for the window counts, not a single remaining estimate. For $k = 3$ it reduces to a variance bound: Conjecture 1, or any bound $o(q^4/\log q)$ on the variance, and by the Corollary of Section 12 this is a statement about the average of $P(\lambda)$ over all units λ. For $k \ge 4$ we have no comparable reduction. In Fourier terms the missing input for $k = 3$ is cancellation, on average over the dilation parameter, in correlations of Jacobi sums over $\mathbb{F}_{q^3}$, beyond the square-root (Weil–Deligne) bound for single sums. We do not know how to prove it. Section 12 evaluates these correlations exactly at a few small dilations. Three cheap routes to the averaged bound failed in our tests: the exact formulas grow in complexity with the height of the dilation; among small heights (numerators and denominators ≤ 12, avoiding 3 and 7) we found no further coincidences of P beyond those explained by Theorem 3 and one unexplained class; and maximum bounds for roots of trinomials [19] are far too weak for an L^2 statement.

14 Limitations, literature and disclosure

Theorem 1 covers fixed ratios only (prime q, or large characteristic). The higher-moment and variance laws are numerical. Theorem 2 is proved; two polynomial identities and one factorisation in its proof were verified by exact computer algebra (Sage 10.7; the scripts and their SHA-256 hashes are released) rather than by hand. The excess formula of Section 11 is a heuristic, not a

theorem. For Version 3, a search (web, zbMATH Open, three full texts; 20 October 2026) found no prior statement of Theorem 2. Proposition 4 is elementary and probably known. Proposition 2 is conditional on moment convergence, which is not proved. A full-text literature review and a zbMATH Open search (19 October 2026) found the forbidden ratios, the arc and oval facts [13, 14] and the dilate strategy ([6], Shakan [10]). They did not find the curve count, the two laws, the identity of Section 6 or the reduction of Section 4. Guy's C9 entry and MathSciNet were not checked. Computations and drafting were done with AI assistance (Anthropic Claude). The statements labelled as proved were checked by separate AI instances and by exact computation, but not yet by a human expert. For Version 4 (searches on 30 October 2026: zbMATH Open, web, GitHub), we found no prior statement of Theorem 3(b),(c) or of Proposition 5; Theorem 3(a) is, by Lemma 1, equivalent to the classical conic fact [13, 14]. Also not checked for Version 4: the original paper of Jungnickel and Vedder, the full text of the paper of Ono, and a 2005 essay on Singer sets (A. Chmelnitzki), whose server was unreachable. Theorem 3(c) relies on [18, Thm. 1.7] (statement read; a preprint; we did not confirm a journal version) and on standard facts from [22]. All registrations in this paper are self-attested (Section 1).

Changes in Version 5. This version corrects Version 4 after two AI referee reports (mathematics; reproducibility). They found no error in the statements or proofs of the paper; one sentence of the released proof file was wrong and is corrected (item 4). (1) Lemma 1: $P(l) - (q-2)^2$ is the per-dilation third moment of Section 6, so Theorem 3(a) is a known fact and (b), (c) are exact third moments; we added the Corollary. (2) The summary of what remains no longer speaks of a single remaining step that we isolate; the reduction of Section 4 is described as folklore. (3) The abstract gives the exact prime sets of the higher-moment tests (129 primes for $k \le 6$ at $C = 2$, 64 primes otherwise) and states that registrations are self-attested. (4) Released: the programs behind the rows $q \ge 4096$ of Table 2 (`run_nb_top12.py`; `run_orb.py`, which produced the blind test at $q = 12167$) with a regression test and a per-row provenance table, and the raw output for $q = 16807$; a script for Lemma 1; a corrected sentence in the released proof of Theorem 3(c) (step 7: the associate congruent to 1 modulo 2 is $\overline{\chi_3(2)}J(\chi_3, \chi_3)$); a claim-to-script table. (5) Smaller fixes: the case $c = d$ in the proof of (b), the wording of the proof of Proposition 5, hedged geometric explanation and height observation, two numbers (54%, $q = 16807$), the borderline row $q = 125$ of Table 1, the date of Version 4, the disclosure that step (7) was written after the first review, and the note on composite m in Table 2. Still open: a notation table, and exact theorem numbers for [25] and [22].

Data and code

Version 1 release: `https://github.com/vichienFF/ramanujan-explorer/tree/main/erdos_sidon`. Version 2 additions (certificates, blind-test plans with SHA-256 ledger, Kaggle outputs): `https://github.com/vichienFF/ramanujan-explorer/tree/main/erdos_sidon/v2`. Version 3 additions (Lean files, Sage scripts for Theorem 2, resonance blind tests): `https://github.com/vichienFF/ramanujan-explorer/tree/main/erdos_sidon/v3`. Version 4 additions (proof of Theorem 3, verification scripts, independent review): `https://github.com/vichienFF/ramanujan-explorer/tree/main/erdos_sidon/v4`. Version 5 additions (`run_orb.py` with regression test, Lemma 1 script, corrected proof file, claim-to-script table, referee reports): `https://github.com/vichienFF/ramanujan-explorer/tree/main/erdos_sidon/v5`.

References

[1] R. C. Baker, G. Harman, J. Pintz, The difference between consecutive primes II, Proc. London Math. Soc. (3) 83 (2001) 532–562.

[2] J. Balogh, Z. Füredi, S. Roy, An upper bound on the size of Sidon sets, Amer. Math. Monthly 130(5) (2023) 437–445.

[3] R. C. Bose, An affine analogue of Singer's theorem, J. Indian Math. Soc. (N.S.) 6 (1942) 1–15.

[4] P. Erdős, P. Turán, On a problem of Sidon in additive number theory, J. London Math. Soc. 16 (1941) 212–215.

[5] T. Bloom, Erdős problems, problem #30, `https://www.erdosproblems.com/30`.

[6] W. T. Gowers, What are dense Sidon subsets of $\{1, 2, \ldots, n\}$ like?, blog post and comments (2012), `https://gowers.wordpress.com/2012/07/13/what-are-dense-sidon-subsets-of-12-n-like/`.

[7] R. K. Guy, Unsolved Problems in Number Theory, 3rd ed., Springer 2004, problem C9.

[8] The LMFDB Collaboration, The L-functions and modular forms database, newform 12.3.c.a, `https://www.lmfdb.org/ModularForm/GL2/Q/holomorphic/12/3/c/a/`.

[9] G. Martin, K. O'Bryant, Constructions of generalized Sidon sets, J. Combin. Theory Ser. A 113 (2006) 591–607.

[10] G. Shakan, A large gap in a dilate of a set, arXiv:2004.14828 (2020).

[11] T. Rokicki, G. Dogon, Golomb ruler and Sidon set tables, `https://cube20.org/golomb/`.

[12] Y. Aubry, M. Perret, A Weil theorem for singular curves, in: R. Pellikaan, M. Perret, S. G. Vlăduţ (eds.), Arithmetic, Geometry and Coding Theory (Luminy 1993), de Gruyter, Berlin 1996, 1–7.

[13] D. Jungnickel, K. Vedder, On the geometry of planar difference sets, European J. Combin. 5 (1984) 143–148.

[14] G. Pickert, Differenzmengen und Ovale, Discrete Math. 73 (1989) 165–179.

[15] B. Green, 100 open problems, Problems 31–32 (manuscript, updated 2026).

[16] M. Hall, Jr., Cyclic projective planes, Duke Math. J. 14 (1947) 1079–1090.

[17] R. L. McFarland, B. F. Rice, Translates and multipliers of abelian difference sets, Proc. Amer. Math. Soc. 68 (1978) 375–379.

[18] R. Barman, G. Kalita, Certain values of Gaussian hypergeometric series and a family of algebraic curves, arXiv:1208.0495 (2012).

[19] J. Bi, Q. Cheng, J. M. Rojas, Sub-linear root detection, and new hardness results, for sparse polynomials over finite fields, Proc. ISSAC 2013, 61–68.

[20] K. Hulek, M. Schütt, Arithmetic of singular Enriques surfaces, Algebra Number Theory 6 (2012) 195–230; arXiv:1002.1598, Example 3.1.

[21] J. Greene, Hypergeometric functions over finite fields, Trans. Amer. Math. Soc. 301 (1987) 77–101.

[22] K. Ireland, M. Rosen, A Classical Introduction to Modern Number Theory, 2nd ed., Springer 1990, Chapters 8, 9 and 11.

[23] Z. Kelley, The number of roots of trinomials over prime fields, Texas A&M REU report and slides (2015).

[24] Z. Kelley, S. Owen, Estimating the number of roots of trinomials over finite fields, arXiv:1510.01758 (2015).

[25] K. Ono, Values of Gaussian hypergeometric series, Trans. Amer. Math. Soc. 350 (1998) 1205–1223.

[26] J. Singer, A theorem in finite projective geometry and some applications to number theory, Trans. Amer. Math. Soc. 43 (1938) 377–385.

[27] Z. Zhang, Finding finite B_2-sequences with larger $m - a_m^{1/2}$, Math. Comp. 63 (1994) 403–414.