

Submonoid Membership in n -dimensional lamplighter groups

Ruiwen Dong

Magdalen College, University of Oxford

December 2024



MAGDALEN COLLEGE
OXFORD



European Research Council
Established by the European Commission

Membership problems

Let G be a group (infinite, finitely generated, with solvable Word Problem).
Consider the following **membership** problems:

Definition (Subgroup Membership)

Input: Elements $g_1, g_2, \dots, g_n, g \in G$.

Question: Is g in the subgroup $\langle g_1, g_2, \dots, g_n \rangle_{\text{grp}}$?

Let G be an group (infinite, finitely generated, with solvable Word Problem).
Consider the following **membership** problems:

Definition (Subgroup Membership)

Input: Elements $g_1, g_2, \dots, g_n, g \in G$.

Question: Is g in the subgroup $\langle g_1, g_2, \dots, g_n \rangle_{\text{grp}}$?

Definition (Submonoid Membership)

Input: Elements $g_1, g_2, \dots, g_n, g \in G$.

Question: Is g in the submonoid $\langle g_1, g_2, \dots, g_n \rangle_{\text{mnd}}$?

Let G be an group (infinite, finitely generated, with solvable Word Problem).
Consider the following **membership** problems:

Definition (Subgroup Membership)

Input: Elements $g_1, g_2, \dots, g_n, g \in G$.

Question: Is g in the subgroup $\langle g_1, g_2, \dots, g_n \rangle_{\text{grp}}$?

Definition (Submonoid Membership)

Input: Elements $g_1, g_2, \dots, g_n, g \in G$.

Question: Is g in the submonoid $\langle g_1, g_2, \dots, g_n \rangle_{\text{mnd}}$?

Definition (Rational Subset Membership)

Input: A **rational subset** $S \subseteq G$ and an element $g \in G$.

Question: Is g in the rational subset S ?

Here, $S \subseteq G$ is called a **rational subset** if there is an alphabet Σ , a monoid homomorphism $\varphi: \Sigma^* \rightarrow G$, and a regular language $L \subseteq \Sigma^*$, such that $\varphi(L) = S$.

Example: $S = \{g_1\}^* \{g_2\}^* = \{g_1^n g_2^m \mid n, m \in \mathbb{N}\} \subseteq G$.

Obviously, Subgroup Mshp. \leq Submonoid Mshp. \leq Rational Subset Mshp.

Theorem (Benois 1969, Grunschlag 1999)

*If G is an **abelian** group or a **free** group, then Subgroup Membership, Submonoid Membership and Rational Subset Membership are decidable in G .*

Obviously, Subgroup Mshp. \leq Submonoid Mshp. \leq Rational Subset Mshp.

Theorem (Benois 1969, Grunschlag 1999)

*If G is an **abelian** group or a **free** group, then Subgroup Membership, Submonoid Membership and Rational Subset Membership are decidable in G .*

Theorem (Malcev 1958, Romanovskii 1974, Roman'kov 2022)

*If G is a **nilpotent** group or a **metabelian** group (i.e. $[G, G]$ is abelian), then Subgroup Membership is decidable. But there are nilpotent groups (such as $H_3(\mathbb{Z})^{10000}$) and metabelian groups where Submonoid Membership and Rational Subset Membership are undecidable.*

Obviously, Subgroup Mshp. \leq Submonoid Mshp. \leq Rational Subset Mshp.

Theorem (Benois 1969, Grunschlag 1999)

*If G is an **abelian** group or a **free** group, then Subgroup Membership, Submonoid Membership and Rational Subset Membership are decidable in G .*

Theorem (Malcev 1958, Romanovskii 1974, Roman'kov 2022)

*If G is a **nilpotent** group or a **metabelian** group (i.e. $[G, G]$ is abelian), then Subgroup Membership is decidable. But there are nilpotent groups (such as $H_3(\mathbb{Z})^{10000}$) and metabelian groups where Submonoid Membership and Rational Subset Membership are undecidable.*

Theorem (Shafrir 2018, Bodart 2024)

There exists a group G (such as the two-dimensional lamplighter group $(\mathbb{Z}/2\mathbb{Z}) \wr \mathbb{Z}^2$), where Submonoid Membership is decidable but Rational Subset Membership is undecidable.

Hence, Subgroup Mshp. \leq Submonoid Mshp. \leq Rational Subset Mshp.

Theorem (folklore)

*Let G be a group and $\tilde{G} \leq G$ be a finite index subgroup. Then **Subgroup Membership** is decidable in G if and only if it is decidable in \tilde{G} .*

Idea: if H is a f.g. subgroup of G , then $H \cap \tilde{G}$ is a f.g. subgroup of \tilde{G} .

Theorem (folklore)

*Let G be a group and $\tilde{G} \leq G$ be a finite index subgroup. Then **Subgroup Membership** is decidable in G if and only if it is decidable in \tilde{G} .*

Idea: if H is a f.g. subgroup of G , then $H \cap \tilde{G}$ is a f.g. subgroup of \tilde{G} .

Theorem (Grunschlag 1999)

*Let G be a group and $\tilde{G} \leq G$ be a finite index subgroup. Then **Rational Subset Membership** is decidable in G if and only if it is decidable in \tilde{G} .*

Idea: if S is a rational subset of G , then $S \cap \tilde{G}$ is a rational subset of \tilde{G} .

Theorem (folklore)

Let G be a group and $\tilde{G} \leq G$ be a finite index subgroup. Then **Subgroup Membership** is decidable in G if and only if it is decidable in \tilde{G} .

Idea: if H is a f.g. subgroup of G , then $H \cap \tilde{G}$ is a f.g. subgroup of \tilde{G} .

Theorem (Grunschlag 1999)

Let G be a group and $\tilde{G} \leq G$ be a finite index subgroup. Then **Rational Subset Membership** is decidable in G if and only if it is decidable in \tilde{G} .

Idea: if S is a rational subset of G , then $S \cap \tilde{G}$ is a rational subset of \tilde{G} .

Problem

What about **Submonoid Membership**?

Difficulty: if M is a f.g. submonoid of G , then $M \cap \tilde{G}$ can be an **infinitely generated** submonoid of \tilde{G} .

Theorem (folklore)

Let G be a group and $\tilde{G} \leq G$ be a finite index subgroup. Then **Subgroup Membership** is decidable in G if and only if it is decidable in \tilde{G} .

Idea: if H is a f.g. subgroup of G , then $H \cap \tilde{G}$ is a f.g. subgroup of \tilde{G} .

Theorem (Grunschlag 1999)

Let G be a group and $\tilde{G} \leq G$ be a finite index subgroup. Then **Rational Subset Membership** is decidable in G if and only if it is decidable in \tilde{G} .

Idea: if S is a rational subset of G , then $S \cap \tilde{G}$ is a rational subset of \tilde{G} .

Problem

What about **Submonoid Membership**?

Difficulty: if M is a f.g. submonoid of G , then $M \cap \tilde{G}$ can be an **infinitely generated** submonoid of \tilde{G} .

Theorem (D. 2024 + Shafrir 2024)

There exists a group G and a finite index subgroup \tilde{G} , such that Submonoid Membership is decidable in \tilde{G} but undecidable in G .

Definition (n -dimensional lamplighter group)

The n -dimensional lamplighter group $(\mathbb{Z}/2\mathbb{Z}) \wr \mathbb{Z}^n$ is defined as a matrix group

$$\left\{ \begin{pmatrix} X_1^{z_1} X_2^{z_2} \cdots X_n^{z_n} & f \\ 0 & 1 \end{pmatrix} \mid z_1, \dots, z_n \in \mathbb{Z}, \underbrace{f \in \mathbb{F}_2[X_1^{\pm}, \dots, X_n^{\pm}]}_{\text{Laurent polynomial over } \mathbb{F}_2} \right\}.$$

Definition (n -dimensional lamplighter group)

The n -dimensional lamplighter group $(\mathbb{Z}/2\mathbb{Z}) \wr \mathbb{Z}^n$ is defined as a matrix group

$$\left\{ \begin{pmatrix} X_1^{z_1} X_2^{z_2} \cdots X_n^{z_n} & f \\ 0 & 1 \end{pmatrix} \mid z_1, \dots, z_n \in \mathbb{Z}, \underbrace{f \in \mathbb{F}_2[X_1^{\pm}, \dots, X_n^{\pm}]}_{\text{Laurent polynomial over } \mathbb{F}_2} \right\}.$$

Take for example $n = 2$. Each element in $(\mathbb{Z}/2\mathbb{Z}) \wr \mathbb{Z}^2$ can be drawn as a \mathbb{Z}^2 -grid of lamps, each on or off, plus a person in some position.

$$\begin{pmatrix} X_1^1 X_2^1 & X_1^2 + X_2^2 \\ 0 & 1 \end{pmatrix}$$


Definition (n -dimensional lamplighter group)

The n -dimensional lamplighter group $(\mathbb{Z}/2\mathbb{Z}) \wr \mathbb{Z}^n$ is defined as a matrix group

$$\left\{ \begin{pmatrix} X_1^{z_1} X_2^{z_2} \cdots X_n^{z_n} & f \\ 0 & 1 \end{pmatrix} \mid z_1, \dots, z_n \in \mathbb{Z}, \underbrace{f \in \mathbb{F}_2[X_1^{\pm}, \dots, X_n^{\pm}]}_{\text{Laurent polynomial over } \mathbb{F}_2} \right\}.$$

Take for example $n = 2$. Each element in $(\mathbb{Z}/2\mathbb{Z}) \wr \mathbb{Z}^2$ can be drawn as a \mathbb{Z}^2 -grid of lamps, each on or off, plus a person in some position.

$$\begin{pmatrix} X_1^1 X_2^1 & X_1^2 + X_2^2 \\ 0 & 1 \end{pmatrix}$$




Definition (n -dimensional lamplighter group)

The n -dimensional lamplighter group $(\mathbb{Z}/2\mathbb{Z}) \wr \mathbb{Z}^n$ is defined as a matrix group

$$\left\{ \begin{pmatrix} X_1^{z_1} X_2^{z_2} \cdots X_n^{z_n} & f \\ 0 & 1 \end{pmatrix} \mid z_1, \dots, z_n \in \mathbb{Z}, \underbrace{f \in \mathbb{F}_2[X_1^{\pm}, \dots, X_n^{\pm}]}_{\text{Laurent polynomial over } \mathbb{F}_2} \right\}.$$

Take for example $n = 2$. Each element in $(\mathbb{Z}/2\mathbb{Z}) \wr \mathbb{Z}^2$ can be drawn as a \mathbb{Z}^2 -grid of lamps, each on or off, plus a person in some position.

$$\begin{pmatrix} X_1^1 X_2^1 & X_1^2 + X_2^2 \\ 0 & 1 \end{pmatrix}$$




Definition (n -dimensional lamplighter group)

The n -dimensional lamplighter group $(\mathbb{Z}/2\mathbb{Z}) \wr \mathbb{Z}^n$ is defined as a matrix group

$$\left\{ \begin{pmatrix} X_1^{z_1} X_2^{z_2} \cdots X_n^{z_n} & f \\ 0 & 1 \end{pmatrix} \mid z_1, \dots, z_n \in \mathbb{Z}, \underbrace{f \in \mathbb{F}_2[X_1^{\pm}, \dots, X_n^{\pm}]}_{\text{Laurent polynomial over } \mathbb{F}_2} \right\}.$$

Take for example $n = 2$. Each element in $(\mathbb{Z}/2\mathbb{Z}) \wr \mathbb{Z}^2$ can be drawn as a \mathbb{Z}^2 -grid of lamps, each on or off, plus a person in some position.

$$\begin{pmatrix} X_1^1 X_2^1 & X_1^2 + X_2^2 \\ 0 & 1 \end{pmatrix}$$




Definition (n -dimensional lamplighter group)

The n -dimensional lamplighter group $(\mathbb{Z}/2\mathbb{Z}) \wr \mathbb{Z}^n$ is defined as a matrix group

$$\left\{ \begin{pmatrix} X_1^{z_1} X_2^{z_2} \cdots X_n^{z_n} & f \\ 0 & 1 \end{pmatrix} \mid z_1, \dots, z_n \in \mathbb{Z}, \underbrace{f \in \mathbb{F}_2[X_1^{\pm}, \dots, X_n^{\pm}]}_{\text{Laurent polynomial over } \mathbb{F}_2} \right\}.$$

Take for example $n = 2$. Each element in $(\mathbb{Z}/2\mathbb{Z}) \wr \mathbb{Z}^2$ can be drawn as a \mathbb{Z}^2 -grid of lamps, each on or off, plus a person in some position.

$$\begin{pmatrix} X_1^1 X_2^1 & X_1^2 + X_2^2 \\ 0 & 1 \end{pmatrix}$$



$$\begin{pmatrix} X_1^1 X_2^1 & X_1^2 + X_2^2 \\ 0 & 1 \end{pmatrix}$$




Illustration: multiplication

$$\begin{pmatrix} X_1^1 X_2^1 & X_1^2 + X_2^2 \\ 0 & 1 \end{pmatrix} \times \begin{pmatrix} X_1^1 & 1 \\ 0 & 1 \end{pmatrix}$$








Illustration: multiplication

$$\begin{pmatrix} X_1^1 X_2^1 & X_1^2 + X_2^2 \\ 0 & 1 \end{pmatrix} \times \begin{pmatrix} X_1^1 & 1 \\ 0 & 1 \end{pmatrix} = \begin{pmatrix} X_1^2 X_2 & X_1^2 + X_1 X_2 + X_2^2 \\ 0 & 1 \end{pmatrix}$$



Definition (semidirect products)

Let \mathcal{Y} be a finitely presented $\mathbb{F}_2[X_1^\pm, \dots, X_n^\pm]$ -module. One can define a *semidirect product* $\mathcal{Y} \rtimes \mathbb{Z}^n$ as a matrix group

$$\left\{ \begin{pmatrix} X_1^{z_1} & X_2^{z_2} & \dots & X_n^{z_n} & y \\ & 0 & & & 1 \end{pmatrix} \mid z_1, \dots, z_n \in \mathbb{Z}, y \in \mathcal{Y} \right\}.$$

Definition (semidirect products)

Let \mathcal{Y} be a finitely presented $\mathbb{F}_2[X_1^\pm, \dots, X_n^\pm]$ -module. One can define a *semidirect product* $\mathcal{Y} \rtimes \mathbb{Z}^n$ as a matrix group

$$\left\{ \begin{pmatrix} X_1^{z_1} X_2^{z_2} \cdots X_n^{z_n} & y \\ 0 & 1 \end{pmatrix} \mid z_1, \dots, z_n \in \mathbb{Z}, y \in \mathcal{Y} \right\}.$$

Example:

One can consider $\mathcal{Y} = \mathbb{F}_2[X_1^\pm]$ as an $\mathbb{F}_2[X_1^\pm, X_2^\pm]$ -module. Then $\mathcal{Y} \rtimes \mathbb{Z}^2$ is simply the direct product of $(\mathbb{Z}/2\mathbb{Z}) \wr \mathbb{Z}$ with \mathbb{Z} .

Definition (semidirect products)

Let \mathcal{Y} be a finitely presented $\mathbb{F}_2[X_1^\pm, \dots, X_n^\pm]$ -module. One can define a *semidirect product* $\mathcal{Y} \rtimes \mathbb{Z}^n$ as a matrix group


$$\left\{ \begin{pmatrix} X_1^{z_1} X_2^{z_2} \cdots X_n^{z_n} & y \\ 0 & 1 \end{pmatrix} \mid z_1, \dots, z_n \in \mathbb{Z}, y \in \mathcal{Y} \right\}.$$

Example:

One can consider $\mathcal{Y} = \mathbb{F}_2[X_1^\pm]$ as an $\mathbb{F}_2[X_1^\pm, X_2^\pm]$ -module. Then $\mathcal{Y} \rtimes \mathbb{Z}^2$ is simply the direct product of $(\mathbb{Z}/2\mathbb{Z}) \wr \mathbb{Z}$ with \mathbb{Z} .

Since $\mathcal{Y} = \mathbb{F}_2[X_1^\pm] = \mathbb{F}_2[X_1^\pm, X_2^\pm]/(X_2 - 1)$, the group $\mathcal{Y} \rtimes \mathbb{Z}^2$ can be seen as a quotient of $(\mathbb{Z}/2\mathbb{Z}) \wr \mathbb{Z}^2$ by “wiring” all lamps in the same column together.

$$\begin{pmatrix} X_1^1 X_2^1 & 1 + X_1^2 \\ 0 & 1 \end{pmatrix}$$



Theorem (Lohrey, Steinberg, and Zetsche 2015)

Rational Subset Membership in $(\mathbb{Z}/2\mathbb{Z}) \wr \mathbb{Z}^2$ is undecidable even for a fixed rational subset.

Theorem (Lohrey, Steinberg, and Zetsche 2015)

Rational Subset Membership in $(\mathbb{Z}/2\mathbb{Z}) \wr \mathbb{Z}^2$ is undecidable even for a fixed rational subset. (Because it can encode tiling problems on a plane.)

Theorem (Lohrey, Steinberg, and Zetsche 2015)

Rational Subset Membership in $(\mathbb{Z}/2\mathbb{Z}) \wr \mathbb{Z}^2$ is undecidable even for a fixed rational subset. (Because it can encode tiling problems on a plane.)

Theorem (Shafrir 2024)

Let G be a group. Rational Subset Membership (for a fixed rational subset) in G reduces to Submonoid Membership in $G \times H$ for some virtually abelian H .

In particular, for LSZ'15, H can be taken as a finite extension of \mathbb{Z}^5 .

Theorem (Lohrey, Steinberg, and Zetsche 2015)

Rational Subset Membership in $(\mathbb{Z}/2\mathbb{Z}) \wr \mathbb{Z}^2$ is undecidable even for a fixed rational subset. (Because it can encode tiling problems on a plane.)

Theorem (Shafrir 2024)

Let G be a group. Rational Subset Membership (for a fixed rational subset) in G reduces to Submonoid Membership in $G \times H$ for some virtually abelian H .

In particular, for LSZ'15, H can be taken as a finite extension of \mathbb{Z}^5 .

Corollary

The direct product $\left((\mathbb{Z}/2\mathbb{Z}) \wr \mathbb{Z}^2\right) \times \mathbb{Z}^5$ has a finite extension with undecidable Submonoid Membership.

Constructing the counter-example: old and new results

Theorem (Lohrey, Steinberg, and Zetsche 2015)

Rational Subset Membership in $(\mathbb{Z}/2\mathbb{Z}) \wr \mathbb{Z}^2$ is undecidable even for a fixed rational subset. (Because it can encode tiling problems on a plane.)

Theorem (Shafrir 2024)

Let G be a group. Rational Subset Membership (for a fixed rational subset) in G reduces to Submonoid Membership in $G \times H$ for some virtually abelian H .

In particular, for LSZ'15, H can be taken as a finite extension of \mathbb{Z}^5 .

Corollary

The direct product $\left((\mathbb{Z}/2\mathbb{Z}) \wr \mathbb{Z}^2\right) \times \mathbb{Z}^5$ has a finite extension with undecidable Submonoid Membership.

Theorem (D. 2024)

Let \mathcal{Y} be a finitely presented $\mathbb{F}_2[X_1^\pm, \dots, X_n^\pm]$ -module. Then the semidirect product $\mathcal{Y} \rtimes \mathbb{Z}^n$ has decidable Submonoid Membership.

But $\left((\mathbb{Z}/2\mathbb{Z}) \wr \mathbb{Z}^2\right) \times \mathbb{Z}^5 = \mathbb{F}_2[X_1^\pm, X_2^\pm] \rtimes \mathbb{Z}^7$!!!

Theorem (D. 2024)

Let \mathcal{Y} be a finitely presented $\mathbb{F}_2[X_1^\pm, \dots, X_n^\pm]$ -module. Then the semidirect product $\mathcal{Y} \rtimes \mathbb{Z}^n$ has decidable Submonoid Membership.

Theorem (D. 2024)

Let \mathcal{Y} be a finitely presented $\mathbb{F}_2[X_1^\pm, \dots, X_n^\pm]$ -module. Then the semidirect product $\mathcal{Y} \rtimes \mathbb{Z}^n$ has decidable Submonoid Membership.

How do submonoids of $\mathcal{Y} \rtimes \mathbb{Z}^n$ look like? Example: $n = 2$, consider the submonoid generated by

$$g_1 = \begin{pmatrix} X_1^1 & ? \\ 0 & 1 \end{pmatrix}, g_2 = \begin{pmatrix} X_1^{-1} & ? \\ 0 & 1 \end{pmatrix}, g_3 = \begin{pmatrix} X_2^1 & ? \\ 0 & 1 \end{pmatrix}.$$

Theorem (D. 2024)

Let \mathcal{Y} be a finitely presented $\mathbb{F}_2[X_1^\pm, \dots, X_n^\pm]$ -module. Then the semidirect product $\mathcal{Y} \rtimes \mathbb{Z}^n$ has decidable Submonoid Membership.

How do submonoids of $\mathcal{Y} \rtimes \mathbb{Z}^n$ look like? Example: $n = 2$, consider the submonoid generated by

$$g_1 = \begin{pmatrix} X_1^1 & ? \\ 0 & 1 \end{pmatrix}, g_2 = \begin{pmatrix} X_1^{-1} & ? \\ 0 & 1 \end{pmatrix}, g_3 = \begin{pmatrix} X_2^1 & ? \\ 0 & 1 \end{pmatrix}.$$

We want to decide whether $g = \begin{pmatrix} X_1^1 X_2^2 & ? \\ 0 & 1 \end{pmatrix}$ is in the monoid $\langle g_1, g_2, g_3 \rangle_{\text{mnd}}$.


Theorem (D. 2024)

Let \mathcal{Y} be a finitely presented $\mathbb{F}_2[X_1^\pm, \dots, X_n^\pm]$ -module. Then the semidirect product $\mathcal{Y} \rtimes \mathbb{Z}^n$ has decidable Submonoid Membership.

How do submonoids of $\mathcal{Y} \rtimes \mathbb{Z}^n$ look like? Example: $n = 2$, consider the submonoid generated by

$$g_1 = \begin{pmatrix} X_1^1 & ? \\ 0 & 1 \end{pmatrix}, g_2 = \begin{pmatrix} X_1^{-1} & ? \\ 0 & 1 \end{pmatrix}, g_3 = \begin{pmatrix} X_2^1 & ? \\ 0 & 1 \end{pmatrix}.$$

We want to decide whether $g = \begin{pmatrix} X_1^1 X_2^2 & ? \\ 0 & 1 \end{pmatrix}$ is in the monoid $\langle g_1, g_2, g_3 \rangle_{\text{mnd}}$.




Theorem (D. 2024)

Let \mathcal{Y} be a finitely presented $\mathbb{F}_2[X_1^\pm, \dots, X_n^\pm]$ -module. Then the semidirect product $\mathcal{Y} \rtimes \mathbb{Z}^n$ has decidable Submonoid Membership.

How do submonoids of $\mathcal{Y} \rtimes \mathbb{Z}^n$ look like? Example: $n = 2$, consider the submonoid generated by

$$g_1 = \begin{pmatrix} X_1^1 & ? \\ 0 & 1 \end{pmatrix}, g_2 = \begin{pmatrix} X_1^{-1} & ? \\ 0 & 1 \end{pmatrix}, g_3 = \begin{pmatrix} X_2^1 & ? \\ 0 & 1 \end{pmatrix}.$$

We want to decide whether $g = \begin{pmatrix} X_1^1 X_2^2 & ? \\ 0 & 1 \end{pmatrix}$ is in the monoid $\langle g_1, g_2, g_3 \rangle_{\text{mnd}}$.




Theorem (D. 2024)

Let \mathcal{Y} be a finitely presented $\mathbb{F}_2[X_1^\pm, \dots, X_n^\pm]$ -module. Then the semidirect product $\mathcal{Y} \rtimes \mathbb{Z}^n$ has decidable Submonoid Membership.

How do submonoids of $\mathcal{Y} \rtimes \mathbb{Z}^n$ look like? Example: $n = 2$, consider the submonoid generated by

$$g_1 = \begin{pmatrix} X_1^1 & ? \\ 0 & 1 \end{pmatrix}, g_2 = \begin{pmatrix} X_1^{-1} & ? \\ 0 & 1 \end{pmatrix}, g_3 = \begin{pmatrix} X_2^1 & ? \\ 0 & 1 \end{pmatrix}.$$

We want to decide whether $g = \begin{pmatrix} X_1^1 X_2^2 & ? \\ 0 & 1 \end{pmatrix}$ is in the monoid $\langle g_1, g_2, g_3 \rangle_{\text{mnd}}$.




Theorem (D. 2024)

Let \mathcal{Y} be a finitely presented $\mathbb{F}_2[X_1^\pm, \dots, X_n^\pm]$ -module. Then the semidirect product $\mathcal{Y} \rtimes \mathbb{Z}^n$ has decidable Submonoid Membership.

How do submonoids of $\mathcal{Y} \rtimes \mathbb{Z}^n$ look like? Example: $n = 2$, consider the submonoid generated by

$$g_1 = \begin{pmatrix} X_1^1 & ? \\ 0 & 1 \end{pmatrix}, g_2 = \begin{pmatrix} X_1^{-1} & ? \\ 0 & 1 \end{pmatrix}, g_3 = \begin{pmatrix} X_2^1 & ? \\ 0 & 1 \end{pmatrix}.$$

We want to decide whether $g = \begin{pmatrix} X_1^1 X_2^2 & ? \\ 0 & 1 \end{pmatrix}$ is in the monoid $\langle g_1, g_2, g_3 \rangle_{\text{mnd}}$.




Theorem (D. 2024)

Let \mathcal{Y} be a finitely presented $\mathbb{F}_2[X_1^\pm, \dots, X_n^\pm]$ -module. Then the semidirect product $\mathcal{Y} \rtimes \mathbb{Z}^n$ has decidable Submonoid Membership.

How do submonoids of $\mathcal{Y} \rtimes \mathbb{Z}^n$ look like? Example: $n = 2$, consider the submonoid generated by

$$g_1 = \begin{pmatrix} X_1^1 & ? \\ 0 & 1 \end{pmatrix}, g_2 = \begin{pmatrix} X_1^{-1} & ? \\ 0 & 1 \end{pmatrix}, g_3 = \begin{pmatrix} X_2^1 & ? \\ 0 & 1 \end{pmatrix}.$$

We want to decide whether $g = \begin{pmatrix} X_1^1 X_2^2 & ? \\ 0 & 1 \end{pmatrix}$ is in the monoid $\langle g_1, g_2, g_3 \rangle_{\text{mnd}}$.




Theorem (D. 2024)

Let \mathcal{Y} be a finitely presented $\mathbb{F}_2[X_1^\pm, \dots, X_n^\pm]$ -module. Then the semidirect product $\mathcal{Y} \rtimes \mathbb{Z}^n$ has decidable Submonoid Membership.

How do submonoids of $\mathcal{Y} \rtimes \mathbb{Z}^n$ look like? Example: $n = 2$, consider the submonoid generated by

$$g_1 = \begin{pmatrix} X_1^1 & ? \\ 0 & 1 \end{pmatrix}, g_2 = \begin{pmatrix} X_1^{-1} & ? \\ 0 & 1 \end{pmatrix}, g_3 = \begin{pmatrix} X_2^1 & ? \\ 0 & 1 \end{pmatrix}.$$

We want to decide whether $g = \begin{pmatrix} X_1^1 X_2^2 & ? \\ 0 & 1 \end{pmatrix}$ is in the monoid $\langle g_1, g_2, g_3 \rangle_{\text{mnd}}$.




Theorem (D. 2024)

Let \mathcal{Y} be a finitely presented $\mathbb{F}_2[X_1^\pm, \dots, X_n^\pm]$ -module. Then the semidirect product $\mathcal{Y} \rtimes \mathbb{Z}^n$ has decidable Submonoid Membership.

How do submonoids of $\mathcal{Y} \rtimes \mathbb{Z}^n$ look like? Example: $n = 2$, consider the submonoid generated by

$$g_1 = \begin{pmatrix} X_1^1 & ? \\ 0 & 1 \end{pmatrix}, g_2 = \begin{pmatrix} X_1^{-1} & ? \\ 0 & 1 \end{pmatrix}, g_3 = \begin{pmatrix} X_2^1 & ? \\ 0 & 1 \end{pmatrix}.$$

We want to decide whether $g = \begin{pmatrix} X_1^1 X_2^2 & ? \\ 0 & 1 \end{pmatrix}$ is in the monoid $\langle g_1, g_2, g_3 \rangle_{\text{mnd}}$.




Theorem (D. 2024)

Let \mathcal{Y} be a finitely presented $\mathbb{F}_2[X_1^\pm, \dots, X_n^\pm]$ -module. Then the semidirect product $\mathcal{Y} \rtimes \mathbb{Z}^n$ has decidable Submonoid Membership.

How do submonoids of $\mathcal{Y} \rtimes \mathbb{Z}^n$ look like? Example: $n = 2$, consider the submonoid generated by

$$g_1 = \begin{pmatrix} X_1^1 & ? \\ 0 & 1 \end{pmatrix}, g_2 = \begin{pmatrix} X_1^{-1} & ? \\ 0 & 1 \end{pmatrix}, g_3 = \begin{pmatrix} X_2^1 & ? \\ 0 & 1 \end{pmatrix}.$$

We want to decide whether $g = \begin{pmatrix} X_1^1 X_2^2 & ? \\ 0 & 1 \end{pmatrix}$ is in the monoid $\langle g_1, g_2, g_3 \rangle_{\text{mnd}}$.




Theorem (D. 2024)

Let \mathcal{Y} be a finitely presented $\mathbb{F}_2[X_1^\pm, \dots, X_n^\pm]$ -module. Then the semidirect product $\mathcal{Y} \rtimes \mathbb{Z}^n$ has decidable Submonoid Membership.

How do submonoids of $\mathcal{Y} \rtimes \mathbb{Z}^n$ look like? Example: $n = 2$, consider the submonoid generated by

$$g_1 = \begin{pmatrix} X_1^1 & ? \\ 0 & 1 \end{pmatrix}, g_2 = \begin{pmatrix} X_1^{-1} & ? \\ 0 & 1 \end{pmatrix}, g_3 = \begin{pmatrix} X_2^1 & ? \\ 0 & 1 \end{pmatrix}.$$

We want to decide whether $g = \begin{pmatrix} X_1^1 X_2^2 & ? \\ 0 & 1 \end{pmatrix}$ is in the monoid $\langle g_1, g_2, g_3 \rangle_{\text{mnd}}$.




Theorem (D. 2024)

Let \mathcal{Y} be a finitely presented $\mathbb{F}_2[X_1^\pm, \dots, X_n^\pm]$ -module. Then the semidirect product $\mathcal{Y} \rtimes \mathbb{Z}^n$ has decidable Submonoid Membership.

How do submonoids of $\mathcal{Y} \rtimes \mathbb{Z}^n$ look like? Example: $n = 2$, consider the submonoid generated by

$$g_1 = \begin{pmatrix} X_1^1 & ? \\ 0 & 1 \end{pmatrix}, g_2 = \begin{pmatrix} X_1^{-1} & ? \\ 0 & 1 \end{pmatrix}, g_3 = \begin{pmatrix} X_2^1 & ? \\ 0 & 1 \end{pmatrix}.$$

We want to decide whether $g = \begin{pmatrix} X_1^1 X_2^2 & ? \\ 0 & 1 \end{pmatrix}$ is in the monoid $\langle g_1, g_2, g_3 \rangle_{\text{mnd}}$.




Theorem (D. 2024)

Let \mathcal{Y} be a finitely presented $\mathbb{F}_2[X_1^\pm, \dots, X_n^\pm]$ -module. Then the semidirect product $\mathcal{Y} \rtimes \mathbb{Z}^n$ has decidable Submonoid Membership.

How do submonoids of $\mathcal{Y} \rtimes \mathbb{Z}^n$ look like? Example: $n = 2$, consider the submonoid generated by

$$g_1 = \begin{pmatrix} X_1^1 & ? \\ 0 & 1 \end{pmatrix}, g_2 = \begin{pmatrix} X_1^{-1} & ? \\ 0 & 1 \end{pmatrix}, g_3 = \begin{pmatrix} X_2^1 & ? \\ 0 & 1 \end{pmatrix}.$$

We want to decide whether $g = \begin{pmatrix} X_1^1 X_2^2 & ? \\ 0 & 1 \end{pmatrix}$ is in the monoid $\langle g_1, g_2, g_3 \rangle_{\text{mnd}}$.




Theorem (D. 2024)

Let \mathcal{Y} be a finitely presented $\mathbb{F}_2[X_1^\pm, \dots, X_n^\pm]$ -module. Then the semidirect product $\mathcal{Y} \rtimes \mathbb{Z}^n$ has decidable Submonoid Membership.

How do submonoids of $\mathcal{Y} \rtimes \mathbb{Z}^n$ look like? Example: $n = 2$, consider the submonoid generated by

$$g_1 = \begin{pmatrix} X_1^1 & ? \\ 0 & 1 \end{pmatrix}, g_2 = \begin{pmatrix} X_1^{-1} & ? \\ 0 & 1 \end{pmatrix}, g_3 = \begin{pmatrix} X_2^1 & ? \\ 0 & 1 \end{pmatrix}.$$

We want to decide whether $g = \begin{pmatrix} X_1^1 X_2^2 & ? \\ 0 & 1 \end{pmatrix}$ is in the monoid $\langle g_1, g_2, g_3 \rangle_{\text{mnd}}$.




Theorem (D. 2024)

Let \mathcal{Y} be a finitely presented $\mathbb{F}_2[X_1^\pm, \dots, X_n^\pm]$ -module. Then the semidirect product $\mathcal{Y} \rtimes \mathbb{Z}^n$ has decidable Submonoid Membership.

How do submonoids of $\mathcal{Y} \rtimes \mathbb{Z}^n$ look like? Example: $n = 2$, consider the submonoid generated by

$$g_1 = \begin{pmatrix} X_1^1 & ? \\ 0 & 1 \end{pmatrix}, g_2 = \begin{pmatrix} X_1^{-1} & ? \\ 0 & 1 \end{pmatrix}, g_3 = \begin{pmatrix} X_2^1 & ? \\ 0 & 1 \end{pmatrix}.$$

We want to decide whether $g = \begin{pmatrix} X_1^1 X_2^2 & ? \\ 0 & 1 \end{pmatrix}$ is in the monoid $\langle g_1, g_2, g_3 \rangle_{\text{mnd}}$.




Theorem (D. 2024)

Let \mathcal{Y} be a finitely presented $\mathbb{F}_2[X_1^\pm, \dots, X_n^\pm]$ -module. Then the semidirect product $\mathcal{Y} \rtimes \mathbb{Z}^n$ has decidable Submonoid Membership.

How do submonoids of $\mathcal{Y} \rtimes \mathbb{Z}^n$ look like? Example: $n = 2$, consider the submonoid generated by

$$g_1 = \begin{pmatrix} X_1^1 & ? \\ 0 & 1 \end{pmatrix}, g_2 = \begin{pmatrix} X_1^{-1} & ? \\ 0 & 1 \end{pmatrix}, g_3 = \begin{pmatrix} X_2^1 & ? \\ 0 & 1 \end{pmatrix}.$$

We want to decide whether $g = \begin{pmatrix} X_1^1 X_2^2 & ? \\ 0 & 1 \end{pmatrix}$ is in the monoid $\langle g_1, g_2, g_3 \rangle_{\text{mnd}}$.




Theorem (D. 2024)

Let \mathcal{Y} be a finitely presented $\mathbb{F}_2[X_1^\pm, \dots, X_n^\pm]$ -module. Then the semidirect product $\mathcal{Y} \rtimes \mathbb{Z}^n$ has decidable Submonoid Membership.

How do submonoids of $\mathcal{Y} \rtimes \mathbb{Z}^n$ look like? Example: $n = 2$, consider the submonoid generated by

$$g_1 = \begin{pmatrix} X_1^1 & ? \\ 0 & 1 \end{pmatrix}, g_2 = \begin{pmatrix} X_1^{-1} & ? \\ 0 & 1 \end{pmatrix}, g_3 = \begin{pmatrix} X_2^1 & ? \\ 0 & 1 \end{pmatrix}.$$

We want to decide whether $g = \begin{pmatrix} X_1^1 X_2^2 & ? \\ 0 & 1 \end{pmatrix}$ is in the monoid $\langle g_1, g_2, g_3 \rangle_{\text{mnd}}$.




Theorem (D. 2024)

Let \mathcal{Y} be a finitely presented $\mathbb{F}_2[X_1^\pm, \dots, X_n^\pm]$ -module. Then the semidirect product $\mathcal{Y} \rtimes \mathbb{Z}^n$ has decidable Submonoid Membership.

How do submonoids of $\mathcal{Y} \rtimes \mathbb{Z}^n$ look like? Example: $n = 2$, consider the submonoid generated by

$$g_1 = \begin{pmatrix} X_1^1 & ? \\ 0 & 1 \end{pmatrix}, g_2 = \begin{pmatrix} X_1^{-1} & ? \\ 0 & 1 \end{pmatrix}, g_3 = \begin{pmatrix} X_2^1 & ? \\ 0 & 1 \end{pmatrix}.$$

We want to decide whether $g = \begin{pmatrix} X_1^1 X_2^2 & ? \\ 0 & 1 \end{pmatrix}$ is in the monoid $\langle g_1, g_2, g_3 \rangle_{\text{mnd}}$.




Theorem (D. 2024)

Let \mathcal{Y} be a finitely presented $\mathbb{F}_2[X_1^\pm, \dots, X_n^\pm]$ -module. Then the semidirect product $\mathcal{Y} \rtimes \mathbb{Z}^n$ has decidable Submonoid Membership.

How do submonoids of $\mathcal{Y} \rtimes \mathbb{Z}^n$ look like? Example: $n = 2$, consider the submonoid generated by

$$g_1 = \begin{pmatrix} X_1^1 & ? \\ 0 & 1 \end{pmatrix}, g_2 = \begin{pmatrix} X_1^{-1} & ? \\ 0 & 1 \end{pmatrix}, g_3 = \begin{pmatrix} X_2^1 & ? \\ 0 & 1 \end{pmatrix}.$$

We want to decide whether $g = \begin{pmatrix} X_1^1 X_2^2 & ? \\ 0 & 1 \end{pmatrix}$ is in the monoid $\langle g_1, g_2, g_3 \rangle_{\text{mnd}}$.




Theorem (D. 2024)

Let \mathcal{Y} be a finitely presented $\mathbb{F}_2[X_1^\pm, \dots, X_n^\pm]$ -module. Then the semidirect product $\mathcal{Y} \rtimes \mathbb{Z}^n$ has decidable Submonoid Membership.

How do submonoids of $\mathcal{Y} \rtimes \mathbb{Z}^n$ look like? Example: $n = 2$, consider the submonoid generated by

$$g_1 = \begin{pmatrix} X_1^1 & ? \\ 0 & 1 \end{pmatrix}, g_2 = \begin{pmatrix} X_1^{-1} & ? \\ 0 & 1 \end{pmatrix}, g_3 = \begin{pmatrix} X_2^1 & ? \\ 0 & 1 \end{pmatrix}.$$

We want to decide whether $g = \begin{pmatrix} X_1^1 X_2^2 & ? \\ 0 & 1 \end{pmatrix}$ is in the monoid $\langle g_1, g_2, g_3 \rangle_{\text{mnd}}$.




Theorem (D. 2024)

Let \mathcal{Y} be a finitely presented $\mathbb{F}_2[X_1^\pm, \dots, X_n^\pm]$ -module. Then the semidirect product $\mathcal{Y} \rtimes \mathbb{Z}^n$ has decidable Submonoid Membership.

How do submonoids of $\mathcal{Y} \rtimes \mathbb{Z}^n$ look like? Example: $n = 2$, consider the submonoid generated by

$$g_1 = \begin{pmatrix} X_1^1 & ? \\ 0 & 1 \end{pmatrix}, g_2 = \begin{pmatrix} X_1^{-1} & ? \\ 0 & 1 \end{pmatrix}, g_3 = \begin{pmatrix} X_2^1 & ? \\ 0 & 1 \end{pmatrix}.$$

We want to decide whether $g = \begin{pmatrix} X_1^1 X_2^2 & ? \\ 0 & 1 \end{pmatrix}$ is in the monoid $\langle g_1, g_2, g_3 \rangle_{\text{mnd}}$.



Satisfy: $X_1^{z_1} \cdot m_1 + X_1^{z_2} \cdot m_2 = m_0$ for some $m_1, m_2, m_0 \in \mathcal{Y}/\mathcal{Y}'$.

The quotient by some submodule \mathcal{Y}' accounts for “zigzags”.

Theorem (D. 2024)

Let \mathcal{Y} be a finitely presented $\mathbb{F}_2[X_1^\pm, \dots, X_n^\pm]$ -module. Then the semidirect product $\mathcal{Y} \rtimes \mathbb{Z}^n$ has decidable Submonoid Membership.

The proof has two steps. **Step 1** is rather elementary:

Proposition (From Submonoid Membership to S-unit equations)

*Submonoid Membership in $\mathcal{Y} \rtimes \mathbb{Z}^n$ reduces to solving **S-unit equations** in some finitely presented $\mathbb{F}_2[X_1^\pm, \dots, X_n^\pm]$ -module $M = \mathcal{Y}/\mathcal{Y}'$.*

Proof of decidability result: from submonoids to S-unit equations

Theorem (D. 2024)

Let \mathcal{Y} be a finitely presented $\mathbb{F}_2[X_1^\pm, \dots, X_n^\pm]$ -module. Then the semidirect product $\mathcal{Y} \rtimes \mathbb{Z}^n$ has decidable Submonoid Membership.

The proof has two steps. **Step 1** is rather elementary:

Proposition (From Submonoid Membership to S-unit equations)

Submonoid Membership in $\mathcal{Y} \rtimes \mathbb{Z}^n$ reduces to solving **S-unit equations** in some finitely presented $\mathbb{F}_2[X_1^\pm, \dots, X_n^\pm]$ -module $M = \mathcal{Y}/\mathcal{Y}'$.

Definition (S-unit equations in modules)

Input: a finitely presented $\mathbb{F}_2[X_1^\pm, \dots, X_n^\pm]$ -module M , and elements $m_1, \dots, m_k, m_0 \in M$.

Question: Does the following equation have solutions $z_{11}, z_{12}, \dots, z_{kn} \in \mathbb{Z}$?

$$X_1^{z_{11}} X_2^{z_{12}} \dots X_n^{z_{1n}} \cdot m_1 + \dots + X_1^{z_{k1}} X_2^{z_{k2}} \dots X_n^{z_{kn}} \cdot m_k = m_0.$$

For example, if $M = \mathbb{F}_2[X^\pm, Y^\pm]/(X + Y + 1)$, then we are asking whether

$$X + Y + 1 \mid X^{z_{11}} Y^{z_{12}} m_1 + \dots + X^{z_{k1}} Y^{z_{k2}} m_k - m_0$$

has integer solutions $z_{11}, z_{12}, \dots, z_{k1}, z_{k2}$.

Proposition (Solution of S-unit equations)

*The solutions $(z_{11}, z_{12}, \dots, z_{kn}) \in \mathbb{Z}^{kn}$ of an **S-unit equation***

$$X_1^{z_{11}} X_2^{z_{12}} \dots X_n^{z_{1n}} \cdot m_1 + \dots + X_1^{z_{k1}} X_2^{z_{k2}} \dots X_n^{z_{kn}} \cdot m_k = m_0.$$

over a finitely presented $\mathbb{F}_2[X_1^\pm, \dots, X_n^\pm]$ -module M , form an effectively 2-automatic subset of \mathbb{Z}^{kn} .

Proposition (Solution of S-unit equations)

*The solutions $(z_{11}, z_{12}, \dots, z_{kn}) \in \mathbb{Z}^{kn}$ of an **S-unit equation***

$$X_1^{z_{11}} X_2^{z_{12}} \dots X_n^{z_{1n}} \cdot m_1 + \dots + X_1^{z_{k1}} X_2^{z_{k2}} \dots X_n^{z_{kn}} \cdot m_k = m_0.$$

over a finitely presented $\mathbb{F}_2[X_1^\pm, \dots, X_n^\pm]$ -module M , form an effectively 2-automatic subset of \mathbb{Z}^{kn} .

Example

Let $M = \mathbb{F}_2[X^\pm, Y^\pm]/(X + Y + 1)$. The following equation in M

$$X^{z_{11}} Y^{z_{12}} + X^{z_{21}} Y^{z_{22}} = 1$$

is equivalent to the following equation in $\mathbb{F}_2(X)$:

$$X^{z_{11}}(X + 1)^{z_{12}} + X^{z_{21}}(X + 1)^{z_{22}} = 1.$$

Proposition (Solution of S-unit equations)

*The solutions $(z_{11}, z_{12}, \dots, z_{kn}) \in \mathbb{Z}^{kn}$ of an **S-unit equation***

$$X_1^{z_{11}} X_2^{z_{12}} \dots X_n^{z_{1n}} \cdot m_1 + \dots + X_1^{z_{k1}} X_2^{z_{k2}} \dots X_n^{z_{kn}} \cdot m_k = m_0.$$

over a finitely presented $\mathbb{F}_2[X_1^\pm, \dots, X_n^\pm]$ -module M , form an effectively 2-automatic subset of \mathbb{Z}^{kn} .

Example

Let $M = \mathbb{F}_2[X^\pm, Y^\pm]/(X + Y + 1)$. The following equation in M

$$X^{z_{11}} Y^{z_{12}} + X^{z_{21}} Y^{z_{22}} = 1$$

is equivalent to the following equation in $\mathbb{F}_2(X)$:

$$X^{z_{11}}(X + 1)^{z_{12}} + X^{z_{21}}(X + 1)^{z_{22}} = 1.$$

The solution set is

$$(z_{11}, z_{12}, z_{21}, z_{22}) \in \{(2^k, 0, 0, 2^k) \mid k \in \mathbb{N}\} \cup \{(0, 2^k, 2^k, 0) \mid k \in \mathbb{N}\}.$$

Proposition (Solution of S-unit equations)

The solutions $(z_{11}, z_{12}, \dots, z_{kn}) \in \mathbb{Z}^{kn}$ of an **S-unit equation**

$$X_1^{z_{11}} X_2^{z_{12}} \dots X_n^{z_{1n}} \cdot m_1 + \dots + X_1^{z_{k1}} X_2^{z_{k2}} \dots X_n^{z_{kn}} \cdot m_k = m_0.$$

over a finitely presented $\mathbb{F}_2[X_1^\pm, \dots, X_n^\pm]$ -module M , form an effectively 2-automatic subset of \mathbb{Z}^{kn} .

Example

Let $M = \mathbb{F}_2[X^\pm, Y^\pm]/(X + Y + 1)$. The following equation in M

$$X^{z_{11}} Y^{z_{12}} + X^{z_{21}} Y^{z_{22}} = 1$$

is equivalent to the following equation in $\mathbb{F}_2(X)$:

$$X^{z_{11}}(X + 1)^{z_{12}} + X^{z_{21}}(X + 1)^{z_{22}} = 1.$$

The solution set is

$$(z_{11}, z_{12}, z_{21}, z_{22}) \in \{(2^k, 0, 0, 2^k) \mid k \in \mathbb{N}\} \cup \{(0, 2^k, 2^k, 0) \mid k \in \mathbb{N}\}.$$

Hint: $(X + 1)^{2^k} = X^{2^k} + 1$, so $(X + 1)^{2^k} + X^{2^k} = 1$.

Proof of 2-automaticity: Derkson's idea

Let's give the proof idea of step 2.


Instead of an equation with kn integer variables, let's consider a similar equation with one integer variable z .

For example, we want to solve in $M = \mathbb{F}_2[X^\pm, Y^\pm]/(X + Y + 1)$:

$$Y^z + X^z = 1.$$

That is, we need to solve in $\mathbb{F}_2(X)$:

$$(X + 1)^z + X^z = 1.$$


$$(X + 1)^z + X^z = 1$$

Proof of 2-automaticity: Derkson's idea

Let's give the proof idea of step 2.


Instead of an equation with kn integer variables, let's consider a similar equation with one integer variable z .

For example, we want to solve in $M = \mathbb{F}_2[X^\pm, Y^\pm]/(X + Y + 1)$:

$$Y^z + X^z = 1.$$

That is, we need to solve in $\mathbb{F}_2(X)$:

$$(X + 1)^z + X^z = 1.$$



Proof of 2-automaticity: Derkson's idea

Let's give the proof idea of step 2.


Instead of an equation with kn integer variables, let's consider a similar equation with one integer variable z .

For example, we want to solve in $M = \mathbb{F}_2[X^\pm, Y^\pm]/(X + Y + 1)$:

$$Y^z + X^z = 1.$$

That is, we need to solve in $\mathbb{F}_2(X)$:

$$(X + 1)^z + X^z = 1.$$



Proof of 2-automaticity: Derkson's idea

Let's give the proof idea of step 2.


Instead of an equation with kn integer variables, let's consider a similar equation with one integer variable z .

For example, we want to solve in $M = \mathbb{F}_2[X^\pm, Y^\pm]/(X + Y + 1)$:

$$Y^z + X^z = 1.$$

That is, we need to solve in $\mathbb{F}_2(X)$:

$$(X + 1)^z + X^z = 1.$$



Proof of 2-automaticity: Derkson's idea

Let's give the proof idea of step 2.


Instead of an equation with kn integer variables, let's consider a similar equation with one integer variable z .

For example, we want to solve in $M = \mathbb{F}_2[X^\pm, Y^\pm]/(X + Y + 1)$:

$$Y^z + X^z = 1.$$

That is, we need to solve in $\mathbb{F}_2(X)$:

$$(X + 1)^z + X^z = 1.$$



Proof of 2-automaticity: Derkson's idea

Let's give the proof idea of step 2.

Instead of an equation with kn integer variables, let's consider a similar equation with one integer variable z .

For example, we want to solve in $M = \mathbb{F}_2[X^\pm, Y^\pm]/(X + Y + 1)$:

$$Y^z + X^z = 1.$$

That is, we need to solve in $\mathbb{F}_2(X)$:

$$(X + 1)^z + X^z = 1.$$

The diagram illustrates the reduction of the equation $(X + 1)^z + X^z = 1$ to a simpler form. The equation is enclosed in an oval. Above the oval, the substitution $z = 2z'$ is shown with a curved arrow pointing to the z in the equation. Below the oval, the substitution $z = 2z' + 1$ is shown with a straight arrow pointing away from the oval.

Proof of 2-automaticity: Derkson's idea

Let's give the proof idea of step 2.


Instead of an equation with kn integer variables, let's consider a similar equation with one integer variable z .

For example, we want to solve in $M = \mathbb{F}_2[X^\pm, Y^\pm]/(X + Y + 1)$:

$$Y^z + X^z = 1.$$

That is, we need to solve in $\mathbb{F}_2(X)$:

$$(X + 1)^z + X^z = 1.$$



Proof of 2-automaticity: Derkson's idea

Let's give the proof idea of step 2.


Instead of an equation with kn integer variables, let's consider a similar equation with one integer variable z .

For example, we want to solve in $M = \mathbb{F}_2[X^\pm, Y^\pm]/(X + Y + 1)$:

$$Y^z + X^z = 1.$$

That is, we need to solve in $\mathbb{F}_2(X)$:

$$(X + 1)^z + X^z = 1.$$



Proof of 2-automaticity: Derkson's idea

Let's give the proof idea of step 2.


Instead of an equation with kn integer variables, let's consider a similar equation with one integer variable z .

For example, we want to solve in $M = \mathbb{F}_2[X^\pm, Y^\pm]/(X + Y + 1)$:

$$Y^z + X^z = 1.$$

That is, we need to solve in $\mathbb{F}_2(X)$:

$$(X + 1)^z + X^z = 1.$$



From this automaton, we see directly that $z = 2^k, k \in \mathbb{N}$.

Proposition (Solution of S-unit equations)

*The solutions $(z_{11}, z_{12}, \dots, z_{kn}) \in \mathbb{Z}^{kn}$ of an **S-unit equation***

$$X_1^{z_{11}} X_2^{z_{12}} \dots X_n^{z_{1n}} \cdot m_1 + \dots + X_1^{z_{k1}} X_2^{z_{k2}} \dots X_n^{z_{kn}} \cdot m_k = m_0.$$

over a finitely presented $\mathbb{F}_2[X_1^\pm, \dots, X_n^\pm]$ -module M , form an effectively 2-automatic subset of \mathbb{Z}^{kn} .

Proposition (Solution of S-unit equations)

*The solutions $(z_{11}, z_{12}, \dots, z_{kn}) \in \mathbb{Z}^{kn}$ of an **S-unit equation***

$$X_1^{z_{11}} X_2^{z_{12}} \dots X_n^{z_{1n}} \cdot m_1 + \dots + X_1^{z_{k1}} X_2^{z_{k2}} \dots X_n^{z_{kn}} \cdot m_k = m_0.$$

over a finitely presented $\mathbb{F}_2[X_1^\pm, \dots, X_n^\pm]$ -module M , form an effectively 2-automatic subset of \mathbb{Z}^{kn} .

We can construct an automaton that recognizes the binary expansion of $(z_{11}, z_{12}, \dots, z_{kn})$ by guessing the parity of each z_{ij} each time.

Proposition (Solution of S-unit equations)

*The solutions $(z_{11}, z_{12}, \dots, z_{kn}) \in \mathbb{Z}^{kn}$ of an **S-unit equation***

$$X_1^{z_{11}} X_2^{z_{12}} \dots X_n^{z_{1n}} \cdot m_1 + \dots + X_1^{z_{k1}} X_2^{z_{k2}} \dots X_n^{z_{kn}} \cdot m_k = m_0.$$

over a finitely presented $\mathbb{F}_2[X_1^\pm, \dots, X_n^\pm]$ -module M , form an effectively 2-automatic subset of \mathbb{Z}^{kn} .

We can construct an automaton that recognizes the binary expansion of $(z_{11}, z_{12}, \dots, z_{kn})$ by guessing the parity of each z_{ij} each time.

Some extra difficulty: the module M is not as nice as a field $\mathbb{F}_2(X)$, so can't use " $(X + 1)^2 = X^2 + 1$ ".

Solution: use primary decomposition and Noether Normalization to reduce to case of fields.

Summary: Submonoid Membership \longrightarrow S-unit equations \longrightarrow decidable

Theorem (D. 2024)

Let \mathcal{Y} be a finitely presented $\mathbb{F}_2[X_1^\pm, \dots, X_n^\pm]$ -module. Then the semidirect product $\mathcal{Y} \rtimes \mathbb{Z}^n$ has decidable Submonoid Membership.

Summary: Submonoid Membership \longrightarrow S-unit equations \longrightarrow decidable

Theorem (D. 2024)

Let \mathcal{Y} be a finitely presented $\mathbb{F}_p[X_1^\pm, \dots, X_n^\pm]$ -module. Then the semidirect product $\mathcal{Y} \rtimes \mathbb{Z}^n$ has decidable Submonoid Membership.

(In fact we can replace 2 by any prime p)

Summary: Submonoid Membership \longrightarrow S-unit equations \longrightarrow decidable

Theorem (D. 2024)

Let \mathcal{Y} be a finitely presented $\mathbb{F}_p[X_1^\pm, \dots, X_n^\pm]$ -module. Then the semidirect product $\mathcal{Y} \rtimes \mathbb{Z}^n$ has decidable Submonoid Membership.

(In fact we can replace 2 by any prime p)

But can we replace the finite field \mathbb{F}_p by the ring $\mathbb{Z}/k\mathbb{Z}$ for any integer k ?

Summary: Submonoid Membership \longrightarrow S-unit equations \longrightarrow decidable

Theorem (D. 2024)

Let \mathcal{Y} be a finitely presented $\mathbb{F}_p[X_1^\pm, \dots, X_n^\pm]$ -module. Then the semidirect product $\mathcal{Y} \rtimes \mathbb{Z}^n$ has decidable Submonoid Membership.

(In fact we can replace 2 by any prime p)

But can we replace the finite field \mathbb{F}_p by the ring $\mathbb{Z}/k\mathbb{Z}$ for any integer k ?

Conjecture

Let $k \in \mathbb{N}$ and \mathcal{Y} be a finitely presented $(\mathbb{Z}/k\mathbb{Z})[X_1^\pm, \dots, X_n^\pm]$ -module. Then the semidirect product $\mathcal{Y} \rtimes \mathbb{Z}^n$ has decidable Submonoid Membership.

Summary: Submonoid Membership \longrightarrow S-unit equations \longrightarrow decidable

Theorem (D. 2024)

Let \mathcal{Y} be a finitely presented $\mathbb{F}_p[X_1^\pm, \dots, X_n^\pm]$ -module. Then the semidirect product $\mathcal{Y} \rtimes \mathbb{Z}^n$ has decidable Submonoid Membership.

(In fact we can replace 2 by any prime p)

But can we replace the finite field \mathbb{F}_p by the ring $\mathbb{Z}/k\mathbb{Z}$ for any integer k ?

Conjecture

Let $k \in \mathbb{N}$ and \mathcal{Y} be a finitely presented $(\mathbb{Z}/k\mathbb{Z})[X_1^\pm, \dots, X_n^\pm]$ -module. Then the semidirect product $\mathcal{Y} \rtimes \mathbb{Z}^n$ has decidable Submonoid Membership. Moreover, let G be a metabelian group with torsion commutator subgroup (i.e. $[G, G]$ is torsion abelian). Then Submonoid Membership is decidable in G .

Summary: Submonoid Membership \longrightarrow S-unit equations \longrightarrow decidable

Theorem (D. 2024)

Let \mathcal{Y} be a finitely presented $\mathbb{F}_p[X_1^\pm, \dots, X_n^\pm]$ -module. Then the semidirect product $\mathcal{Y} \rtimes \mathbb{Z}^n$ has decidable Submonoid Membership.

(In fact we can replace 2 by any prime p)

But can we replace the finite field \mathbb{F}_p by the ring $\mathbb{Z}/k\mathbb{Z}$ for any integer k ?

Conjecture

Let $k \in \mathbb{N}$ and \mathcal{Y} be a finitely presented $(\mathbb{Z}/k\mathbb{Z})[X_1^\pm, \dots, X_n^\pm]$ -module. Then the semidirect product $\mathcal{Y} \rtimes \mathbb{Z}^n$ has decidable Submonoid Membership. Moreover, let G be a metabelian group with torsion commutator subgroup (i.e. $[G, G]$ is torsion abelian). Then Submonoid Membership is decidable in G .

However we cannot replace the finite field \mathbb{F}_p by \mathbb{Z} :

Theorem (Lohrey, Steinberg, and Zetsche 2015)

Submonoid Membership in $\mathbb{Z} \wr \mathbb{Z} = \mathbb{Z}[X^\pm] \rtimes \mathbb{Z}$ is undecidable. (Because it can encode the halting problem for Minsky machines.)