

# Submonoid Membership in $n$ -dimensional lamplighter groups and $S$ -unit equations

Ruiwen Dong<sup>1</sup>

Magdalen College, University of Oxford

ICALP 2025

---

<sup>1</sup>partially supported by ERC Advanced Grant 101097307.

## **Submonoid Membership** in $n$ -dimensional lamplighter groups and $S$ -unit equations

## Membership problems

Let  $G$  be an infinite group (e.g. a matrix group).

## Membership problems

Let  $G$  be an infinite group (e.g. a matrix group).

### Definition (Subgroup Membership Problem)

**Input:** A finite set  $S \subset G$ , an element  $t \in G$ .

**Question:** Does  $t$  belong to the group  $\langle S \rangle_{\text{group}}$ ?

## Membership problems

Let  $G$  be an infinite group (e.g. a matrix group).

### Definition (Subgroup Membership Problem)

**Input:** A finite set  $S \subset G$ , an element  $t \in G$ .

**Question:** Does  $t$  belong to the group  $\langle S \rangle_{\text{group}}$ ?

### Definition (Submonoid Membership Problem)

**Input:** A finite set  $S \subset G$ , an element  $t \in G$ .

**Question:** Does  $t$  belong to the monoid  $\langle S \rangle_{\text{monoid}}$ ?

## Membership problems

Let  $G$  be an infinite group (e.g. a matrix group).

### Definition (Subgroup Membership Problem)

**Input:** A finite set  $S \subset G$ , an element  $t \in G$ .

**Question:** Does  $t$  belong to the group  $\langle S \rangle_{\text{group}}$ ?

### Definition (Submonoid Membership Problem)

**Input:** A finite set  $S \subset G$ , an element  $t \in G$ .

**Question:** Does  $t$  belong to the monoid  $\langle S \rangle_{\text{monoid}}$ ?

### Definition (Rational Subset Membership Problem)

**Input:** A rational subset  $R \subseteq G$  and an element  $t \in G$ .

**Question:** Does  $t$  belong to the set  $R$ ?


rational subset = set defined by a rational expression in  $G$   
= set recognized by a finite state automaton over  $G$

## Relation between the three membership problems

**Example of a rational subset:**  $\{g_1\}^* \{g_2\}^* = \{g_1^n g_2^m \mid n, m \in \mathbb{N}\}.$

## Relation between the three membership problems


**Example of a rational subset:**  $\{g_1\}^* \{g_2\}^* = \{g_1^n g_2^m \mid n, m \in \mathbb{N}\}$ .





## Relation between the three membership problems


**Example of a rational subset:**  $\{g_1\}^* \{g_2\}^* = \{g_1^n g_2^m \mid n, m \in \mathbb{N}\}$ .



**Fin. gen. submonoids are rational subsets:**

## Relation between the three membership problems


**Example of a rational subset:**  $\{g_1\}^* \{g_2\}^* = \{g_1^n g_2^m \mid n, m \in \mathbb{N}\}$ .




**Fin. gen. submonoids are rational subsets:**  $\langle g_1, g_2 \rangle_{\text{monoid}} = \{g_1, g_2\}^*$ .

## Relation between the three membership problems

**Example of a rational subset:**  $\{g_1\}^* \{g_2\}^* = \{g_1^n g_2^m \mid n, m \in \mathbb{N}\}$ .




**Fin. gen. submonoids are rational subsets:**  $\langle g_1, g_2 \rangle_{\text{monoid}} = \{g_1, g_2\}^*$ .




## Relation between the three membership problems

**Example of a rational subset:**  $\{g_1\}^* \{g_2\}^* = \{g_1^n g_2^m \mid n, m \in \mathbb{N}\}$ .




**Fin. gen. submonoids are rational subsets:**  $\langle g_1, g_2 \rangle_{\text{monoid}} = \{g_1, g_2\}^*$ .




**Fin. gen. subgroups are submonoids:**

## Relation between the three membership problems

**Example of a rational subset:**  $\{g_1\}^* \{g_2\}^* = \{g_1^n g_2^m \mid n, m \in \mathbb{N}\}$ .




**Fin. gen. submonoids are rational subsets:**  $\langle g_1, g_2 \rangle_{\text{monoid}} = \{g_1, g_2\}^*$ .




**Fin. gen. subgroups are submonoids:**  $\langle g_1, g_2 \rangle_{\text{group}} = \langle g_1, g_2, g_1^{-1}, g_2^{-1} \rangle_{\text{monoid}}$

## Relation between the three membership problems


**Example of a rational subset:**  $\{g_1\}^* \{g_2\}^* = \{g_1^n g_2^m \mid n, m \in \mathbb{N}\}$ .



**Fin. gen. submonoids are rational subsets:**  $\langle g_1, g_2 \rangle_{\text{monoid}} = \{g_1, g_2\}^*$ .




**Fin. gen. subgroups are submonoids:**  $\langle g_1, g_2 \rangle_{\text{group}} = \langle g_1, g_2, g_1^{-1}, g_2^{-1} \rangle_{\text{monoid}}$




## Relation between the three membership problems


**Example of a rational subset:**  $\{g_1\}^* \{g_2\}^* = \{g_1^n g_2^m \mid n, m \in \mathbb{N}\}$ .



**Fin. gen. submonoids are rational subsets:**  $\langle g_1, g_2 \rangle_{\text{monoid}} = \{g_1, g_2\}^*$ .




**Fin. gen. subgroups are submonoids:**  $\langle g_1, g_2 \rangle_{\text{group}} = \langle g_1, g_2, g_1^{-1}, g_2^{-1} \rangle_{\text{monoid}}$




Subgroup Membership  $<$  Submonoid Membership  $<$  Rational Subset Mshp.

## Relation between the three membership problems


**Example of a rational subset:**  $\{g_1\}^* \{g_2\}^* = \{g_1^n g_2^m \mid n, m \in \mathbb{N}\}$ .



**Fin. gen. submonoids are rational subsets:**  $\langle g_1, g_2 \rangle_{\text{monoid}} = \{g_1, g_2\}^*$ .



**Fin. gen. subgroups are submonoids:**  $\langle g_1, g_2 \rangle_{\text{group}} = \langle g_1, g_2, g_1^{-1}, g_2^{-1} \rangle_{\text{monoid}}$



Subgroup Membership < Submonoid Membership < Rational Subset Mshp.


Decidability depends on the group  $G$ .

- All three problems are **decidable** in  $\mathbb{Z}^d$  and the free group  $F_2$ . (Benois '69)




# Relation between the three membership problems


**Example of a rational subset:**  $\{g_1\}^* \{g_2\}^* = \{g_1^n g_2^m \mid n, m \in \mathbb{N}\}$ .



**Fin. gen. submonoids are rational subsets:**  $\langle g_1, g_2 \rangle_{\text{monoid}} = \{g_1, g_2\}^*$ .



**Fin. gen. subgroups are submonoids:**  $\langle g_1, g_2 \rangle_{\text{group}} = \langle g_1, g_2, g_1^{-1}, g_2^{-1} \rangle_{\text{monoid}}$



Subgroup Membership < Submonoid Membership < Rational Subset Mshp.

Decidability depends on the group  $G$ .

- ▶ All three problems are **decidable** in  $\mathbb{Z}^d$  and the free group  $F_2$ . (Benois '69)
- ▶ All three problems are **undecidable** in  $F_2 \times F_2$ . (Mikhailova '66)

## Is decidability preserved in finite index subgroups?

$$G \geq_{\text{finite index}} H, \quad (G = H \cup g_1 H \cup \cdots \cup g_k H)$$

# Is decidability preserved in finite index subgroups?

$$G \geq_{\text{finite index}} H, \quad (G = H \cup g_1 H \cup \dots \cup g_k H)$$

$\uparrow$   
decidable Subgroup Membership

# Is decidability preserved in finite index subgroups?

$$\begin{array}{ccc} G & \geq_{\text{finite index}} & H, & (G = H \cup g_1 H \cup \dots \cup g_k H) \\ \uparrow & & \uparrow & \\ ? & \text{decidable} & \text{Subgroup Membership} & \end{array}$$

# Is decidability preserved in finite index subgroups?

$$\begin{array}{ccc} G & \geq_{\text{finite index}} & H, & (G = H \cup g_1 H \cup \dots \cup g_k H) \\ \uparrow & & \uparrow & \\ ? & \text{decidable} & \text{Subgroup Membership} & \end{array}$$

Theorem (folklore, Gilman 1987?, Stallings 1983?)

Let  $H$  be a finite index subgroup of  $G$ . Then **Subgroup Membership** is *decidable* in  $G$  if and only if it is *decidable* in  $H$ .

# Is decidability preserved in finite index subgroups?

$$\begin{array}{ccc} G & \geq_{\text{finite index}} & H, & (G = H \cup g_1 H \cup \dots \cup g_k H) \\ \uparrow & & \uparrow & \\ ? & \text{decidable} & \text{Subgroup Membership} & \end{array}$$

Theorem (folklore, Gilman 1987?, Stallings 1983?)

Let  $H$  be a finite index subgroup of  $G$ . Then **Subgroup Membership** is *decidable* in  $G$  if and only if it is *decidable* in  $H$ .

Theorem (Grunschlag 1999)

Let  $H$  be a finite index subgroup of  $G$ . Then **Rational Subset Membership** is *decidable* in  $G$  if and only if it is *decidable* in  $H$ .

# Is decidability preserved in finite index subgroups?

$$\begin{array}{ccc} G & \geq_{\text{finite index}} & H, & (G = H \cup g_1 H \cup \dots \cup g_k H) \\ \uparrow & & \uparrow & \\ ? & \text{decidable} & \text{Subgroup Membership} & \end{array}$$

Theorem (folklore, Gilman 1987?, Stallings 1983?)

Let  $H$  be a finite index subgroup of  $G$ . Then **Subgroup Membership** is *decidable* in  $G$  if and only if it is *decidable* in  $H$ .

## Open Problem

Is decidability of **Submonoid Membership** stable under taking finite index subgroups?

Theorem (Grunschlag 1999)

Let  $H$  be a finite index subgroup of  $G$ . Then **Rational Subset Membership** is *decidable* in  $G$  if and only if it is *decidable* in  $H$ .

Theorem (D. 2025, this talk)

*Decidability of **Submonoid Membership** is not stable under taking finite index subgroup.*



Theorem (D. 2025, this talk)

*Decidability of **Submonoid Membership** is not stable under taking finite index subgroup.*

i.e. we construct

$$\begin{array}{ccc}
 G & \geq_{\text{finite index}} & H \\
 \uparrow & & \uparrow \\
 \text{undecidable} & & \text{decidable}
 \end{array}
 \qquad
 (G = H \cup g_1 H \cup \dots \cup g_k H)$$

Theorem (D. 2025, this talk)

*Decidability of **Submonoid Membership** is not stable under taking finite index subgroup.*

i.e. we construct

$$\begin{array}{ccc}
 G & \geq_{\text{finite index}} & H \\
 \uparrow & & \uparrow \\
 \text{undecidable} & & \text{decidable}
 \end{array}
 \quad (G = H \cup g_1 H \cup \dots \cup g_k H)$$

the title of this talk:

**Submonoid Membership in n-dimensional  
lamplighter groups and S-unit equations**

## Definition: 2-dimensional lamplighter groups

Let  $\mathbb{F}_2$  be the finite field  $\{0, 1\}$  ( $\approx \{\text{off}, \text{on}\}$  for lamps).

## Definition: 2-dimensional lamplighter groups

Let  $\mathbb{F}_2$  be the finite field  $\{0, 1\}$  ( $\approx \{\text{off}, \text{on}\}$  for lamps).

### Definition

The *2-dimensional lamplighter group* is defined as a matrix group

$$\left\{ \begin{pmatrix} X_1^{z_1} X_2^{z_2} & f \\ 0 & 1 \end{pmatrix} \mid z_1, z_2 \in \mathbb{Z}, \underbrace{f \in \mathbb{F}_2[X_1^{\pm}, X_2^{\pm}]}_{\text{Laurent polynomial over } \mathbb{F}_2} \right\}.$$

Each element is a  $\mathbb{Z}^2$ -grid of lamps, each on  $\circ$  or off  $\bullet$ , along with a person  $\text{⚡}$ .

## Definition: 2-dimensional lamplighter groups

Let  $\mathbb{F}_2$  be the finite field  $\{0, 1\}$  ( $\approx \{\text{off}, \text{on}\}$  for lamps).

### Definition

The *2-dimensional lamplighter group* is defined as a matrix group

$$\left\{ \begin{pmatrix} X_1^{z_1} X_2^{z_2} & f \\ 0 & 1 \end{pmatrix} \mid z_1, z_2 \in \mathbb{Z}, \underbrace{f \in \mathbb{F}_2[X_1^{\pm}, X_2^{\pm}]}_{\text{Laurent polynomial over } \mathbb{F}_2} \right\}.$$

Each element is a  $\mathbb{Z}^2$ -grid of lamps, each on  $\circ$  or off  $\bullet$ , along with a person  $\text{⚡}$ .

$$\begin{pmatrix} X_1 X_2 & X_1^2 + X_2^2 \\ 0 & 1 \end{pmatrix}$$

## Definition: 2-dimensional lamplighter groups

Let  $\mathbb{F}_2$  be the finite field  $\{0, 1\}$  ( $\approx \{\text{off}, \text{on}\}$  for lamps).


### Definition

The *2-dimensional lamplighter group* is defined as a matrix group

$$\left\{ \begin{pmatrix} X_1^{z_1} X_2^{z_2} & f \\ 0 & 1 \end{pmatrix} \mid z_1, z_2 \in \mathbb{Z}, \underbrace{f \in \mathbb{F}_2[X_1^{\pm}, X_2^{\pm}]}_{\text{Laurent polynomial over } \mathbb{F}_2} \right\}.$$

Each element is a  $\mathbb{Z}^2$ -grid of lamps, each on  $\circ$  or off  $\bullet$ , along with a person  $\text{⚡}$ .

$$\begin{pmatrix} X_1 X_2 & X_1^2 + X_2^2 \\ 0 & 1 \end{pmatrix}$$



## Definition: 2-dimensional lamplighter groups

Let  $\mathbb{F}_2$  be the finite field  $\{0, 1\}$  ( $\approx$  {off, on} for lamps).


### Definition

The 2-dimensional lamplighter group is defined as a matrix group

$$\left\{ \begin{pmatrix} X_1^{z_1} X_2^{z_2} & f \\ 0 & 1 \end{pmatrix} \mid z_1, z_2 \in \mathbb{Z}, \underbrace{f \in \mathbb{F}_2[X_1^{\pm}, X_2^{\pm}]}_{\text{Laurent polynomial over } \mathbb{F}_2} \right\}.$$

Each element is a  $\mathbb{Z}^2$ -grid of lamps, each on  $\circ$  or off  $\bullet$ , along with a person  $\text{⚡}$ .

$$\begin{pmatrix} X_1 X_2 & X_1^2 + X_2^2 \\ 0 & 1 \end{pmatrix}$$



## Definition: 2-dimensional lamplighter groups

Let  $\mathbb{F}_2$  be the finite field  $\{0, 1\}$  ( $\approx \{\text{off}, \text{on}\}$  for lamps).


### Definition

The *2-dimensional lamplighter group* is defined as a matrix group

$$\left\{ \begin{pmatrix} X_1^{z_1} X_2^{z_2} & f \\ 0 & 1 \end{pmatrix} \mid z_1, z_2 \in \mathbb{Z}, \underbrace{f \in \mathbb{F}_2[X_1^{\pm}, X_2^{\pm}]}_{\text{Laurent polynomial over } \mathbb{F}_2} \right\}.$$

Each element is a  $\mathbb{Z}^2$ -grid of lamps, each on  $\circ$  or off  $\bullet$ , along with a person  $\text{⚡}$ .

$$\begin{pmatrix} X_1 X_2 & X_1^2 + X_2^2 \\ 0 & 1 \end{pmatrix}$$





## Definition: 2-dimensional lamplighter groups

Let  $\mathbb{F}_2$  be the finite field  $\{0, 1\}$  ( $\approx \{\text{off}, \text{on}\}$  for lamps).


### Definition

The *2-dimensional lamplighter group* is defined as a matrix group

$$\left\{ \begin{pmatrix} X_1^{z_1} X_2^{z_2} & f \\ 0 & 1 \end{pmatrix} \mid z_1, z_2 \in \mathbb{Z}, \underbrace{f \in \mathbb{F}_2[X_1^{\pm}, X_2^{\pm}]}_{\text{Laurent polynomial over } \mathbb{F}_2} \right\}.$$


Each element is a  $\mathbb{Z}^2$ -grid of lamps, each on  $\circ$  or off  $\bullet$ , along with a person  $\text{⚡}$ .

$$\begin{pmatrix} X_1 X_2 & X_1^2 + X_2^2 \\ 0 & 1 \end{pmatrix}$$





# Multiplication in the 2-dimensional lamplighter group

$$\begin{pmatrix} X_1 X_2 & X_1^2 + X_2^2 \\ 1 & 1 \end{pmatrix}$$




# Multiplication in the 2-dimensional lamplighter group

$$\begin{pmatrix} X_1 X_2 & X_1^2 + X_2^2 \\ 1 & 1 \end{pmatrix} \times \begin{pmatrix} X_1 & 1 \\ 1 & 1 \end{pmatrix}$$



# Multiplication in the 2-dimensional lamplighter group


$$\begin{pmatrix} X_1 X_2 & X_1^2 + X_2^2 \\ 1 & 1 \end{pmatrix} \times \begin{pmatrix} X_1 & 1 \\ 1 & 1 \end{pmatrix}$$



align the **origin** of 2nd graph to the **person** in 1st graph,

## Multiplication in the 2-dimensional lamplighter group


$$\begin{pmatrix} X_1 X_2 & X_1^2 + X_2^2 \\ 1 & 1 \end{pmatrix} \times \begin{pmatrix} X_1 & 1 \\ 1 & 1 \end{pmatrix}$$



align the **origin** of 2nd graph to the **person** in 1st graph,  
move the 2nd graph onto the 1st graph,

## Multiplication in the 2-dimensional lamplighter group



$$\begin{pmatrix} X_1 X_2 & X_1^2 + X_2^2 \\ 1 & 1 \end{pmatrix} \times \begin{pmatrix} X_1 & 1 \\ 1 & 1 \end{pmatrix}$$



align the **origin** of 2nd graph to the **person** in 1st graph,  
move the 2nd graph onto the 1st graph,  
and do pointwise addition on all lamps.

## Multiplication in the 2-dimensional lamplighter group

$$= \begin{pmatrix} X_1^2 X_2 & X_1^2 + X_1 X_2 + X_2^2 \\ 1 & 1 \end{pmatrix} \quad \begin{pmatrix} X_1 & 1 \\ 1 & 1 \end{pmatrix}$$



align the **origin** of 2nd graph to the **person** in 1st graph,  
move the 2nd graph onto the 1st graph,  
and do pointwise addition on all lamps.

**$n$ -dimensional lamplighter group:** replace 2-dimension grid with  $n$ -dimension grid.



**$n$ -dimensional lamplighter group:** replace 2-dimension grid with  $n$ -dimension grid.

Theorem (Lohrey, Steinberg and Zetsche 2015)

**Submonoid Membership** *is decidable in the 1-dimensional lamplighter group.*

Theorem (Shafrir 2018)

**Submonoid Membership** *is decidable in the 2-dimensional lamplighter group.*

Proof is specific to each dimension.

**$n$ -dimensional lamplighter group:** replace 2-dimension grid with  $n$ -dimension grid.

Theorem (Lohrey, Steinberg and Zetsche 2015)

**Submonoid Membership** *is decidable in the 1-dimensional lamplighter group.*

Theorem (Shafrir 2018)

**Submonoid Membership** *is decidable in the 2-dimensional lamplighter group.*

Proof is specific to each dimension.

Theorem (This paper)

**Submonoid Membership** *is decidable in the  $n$ -dimensional lamplighter group for all  $n$ .*

**$n$ -dimensional lamplighter group:** replace 2-dimension grid with  $n$ -dimension grid.

Theorem (Lohrey, Steinberg and Zetsche 2015)

**Submonoid Membership** *is decidable in the 1-dimensional lamplighter group.*

Theorem (Shafrir 2018)

**Submonoid Membership** *is decidable in the 2-dimensional lamplighter group.*


Proof is specific to each dimension.

Theorem (This paper)

**Submonoid Membership** *is decidable in the  $n$ -dimensional lamplighter group and its quotients for all  $n$ .*


# Quotients of the lamplighter group

$$\left\{ \begin{pmatrix} X_1^{z_1} X_2^{z_2} & f \\ 0 & 1 \end{pmatrix} \mid z_1, z_2 \in \mathbb{Z}, \underbrace{f \in \mathbb{F}_2[X_1^{\pm}, X_2^{\pm}]}_{\text{Laurent polynomial over } \mathbb{F}_2} \right\}.$$



# Quotients of the lamplighter group


$$\left\{ \begin{pmatrix} X_1^{z_1} X_2^{z_2} & f \\ 0 & 1 \end{pmatrix} \mid z_1, z_2 \in \mathbb{Z}, \underbrace{f \in \mathbb{F}_2[X_1^\pm, X_2^\pm]}_{\text{Laurent polynomial over } \mathbb{F}_2} \right\}.$$



Modify the 2-dimensional lamplighter group by "wiring" all lamps in the same column together.

# Quotients of the lamplighter group


$$\left\{ \begin{pmatrix} X_1^{z_1} X_2^{z_2} & f \\ 0 & 1 \end{pmatrix} \mid z_1, z_2 \in \mathbb{Z}, \underbrace{f \in \mathbb{F}_2[X_1^{\pm}, X_2^{\pm}]}_{\text{Laurent polynomial over } \mathbb{F}_2} \right\}.$$



Modify the 2-dimensional lamplighter group by "wiring" all lamps in the same column together.

# Quotients of the lamplighter group


$$\left\{ \begin{pmatrix} X_1^{z_1} X_2^{z_2} & f \\ 0 & 1 \end{pmatrix} \mid z_1, z_2 \in \mathbb{Z}, \underbrace{f \in \mathbb{F}_2[X_1^{\pm}, X_2^{\pm}] / \langle X_2 - 1 \rangle}_{\substack{\text{Laurent polynomial over } \mathbb{F}_2, \\ \text{quotiented by the ideal } \langle X_2 - 1 \rangle}} \right\}.$$



Modify the 2-dimensional lamplighter group by “wiring” all lamps in the same column together. ( $\iff$  “quotienting”  $\mathbb{F}_2[X_1^{\pm}, X_2^{\pm}]$  by the ideal  $\langle X_2 - 1 \rangle$ )

# Quotients of the lamplighter group

$$\left\{ \begin{pmatrix} X_1^{z_1} X_2^{z_2} & f \\ 0 & 1 \end{pmatrix} \mid z_1, z_2 \in \mathbb{Z}, \underbrace{f \in \mathbb{F}_2[X_1^{\pm}, X_2^{\pm}] / \langle X_2 - 1 \rangle}_{\substack{\text{Laurent polynomial over } \mathbb{F}_2, \\ \text{quotiented by the ideal } \langle X_2 - 1 \rangle}} \right\}.$$



Modify the 2-dimensional lamplighter group by “wiring” all lamps in the same column together. ( $\iff$  “quotienting”  $\mathbb{F}_2[X_1^{\pm}, X_2^{\pm}]$  by the ideal  $\langle X_2 - 1 \rangle$ )  
different “wiring” patterns  $\approx$  quotient by different ideals



# The counterexample

Theorem (D. 2025)

**Submonoid Membership** *is decidable in the  $n$ -dimensional lamplighter group and its quotients for all  $n$ .*

# The counterexample

Theorem (D. 2025)

**Submonoid Membership** is decidable in the  $n$ -dimensional lamplighter group and its quotients for all  $n$ .

Let

$$H := (\text{2-dim. lamplighter group}) \times \mathbb{Z}^5,$$

# The counterexample

Theorem (D. 2025)

**Submonoid Membership** is decidable in the  $n$ -dimensional lamplighter group and its quotients for all  $n$ .

Let

$$H := (\text{2-dim. lamplighter group}) \times \mathbb{Z}^5,$$

Theorem (Lohrey, Steinberg and Zetsche 2015)

**Rational Subset Membership** in the 2-dimensional lamplighter group is undecidable.

# The counterexample

Theorem (D. 2025)

**Submonoid Membership** is decidable in the  $n$ -dimensional lamplighter group and its quotients for all  $n$ .

Let

$$H := (\text{2-dim. lamplighter group}) \times \mathbb{Z}^5,$$

Theorem (Lohrey, Steinberg and Zetsche 2015)

**Rational Subset Membership** in the 2-dimensional lamplighter group is undecidable. (Because it can encode the 2-dimensional tiling problem.)

# The counterexample

Theorem (D. 2025)

**Submonoid Membership** is decidable in the  $n$ -dimensional lamplighter group and its quotients for all  $n$ .

Let

$$\begin{array}{c} G \geq_{\text{finite index}} H := (2\text{-dim. lamplighter group}) \times \mathbb{Z}^5, \\ \uparrow \\ \text{undecidable} \end{array}$$

Theorem (Lohrey, Steinberg and Zetsche 2015)

**Rational Subset Membership** in the 2-dimensional lamplighter group is undecidable. (Because it can encode the 2-dimensional tiling problem.)

Corollary (Shafrir 2024)

$H$  has a finite index overgroup  $G$  with undecidable **Submonoid Membership**.

# The counterexample

Theorem (D. 2025)

**Submonoid Membership** is decidable in the  $n$ -dimensional lamplighter group and its quotients for all  $n$ .

Let

$$\begin{array}{ccc} G & \geq_{\text{finite index}} & H := \underbrace{(2\text{-dim. lamplighter group})}_{\text{quotient of 7-dim. lamplighter group}} \times \mathbb{Z}^5, \\ \uparrow & & \\ \text{undecidable} & & \end{array}$$

Theorem (Lohrey, Steinberg and Zetsche 2015)

**Rational Subset Membership** in the 2-dimensional lamplighter group is undecidable. (Because it can encode the 2-dimensional tiling problem.)

Corollary (Shafrir 2024)

$H$  has a finite index overgroup  $G$  with undecidable **Submonoid Membership**.

# The counterexample

Theorem (D. 2025)

**Submonoid Membership** is decidable in the  $n$ -dimensional lamplighter group and its quotients for all  $n$ .

Let

$$\begin{array}{ccc} G & \geq_{\text{finite index}} & H := \underbrace{(2\text{-dim. lamplighter group}) \times \mathbb{Z}^5}_{\text{quotient of 7-dim. lamplighter group}} \\ \uparrow & & \uparrow \\ \text{undecidable} & & \text{decidable} \end{array}$$

Theorem (Lohrey, Steinberg and Zetsche 2015)

**Rational Subset Membership** in the 2-dimensional lamplighter group is undecidable. (Because it can encode the 2-dimensional tiling problem.)

Corollary (Shafrir 2024)

$H$  has a finite index overgroup  $G$  with undecidable **Submonoid Membership**.

## Deciding Submonoid Membership in lamplighter groups



## Submonoid Membership in n-dimensional lamplighter groups and **S-unit equations**

## Submonoid Membership in n-dimensional lamplighter groups and **S-unit equations**

An **S-unit equation** is a linear equation

$$x_1 m_1 + \cdots + x_K m_K = m_0,$$

where we look for solutions  $x_1, \dots, x_K$  in a multiplicative subgroup of a field  $\mathbb{K}$ .

## Submonoid Membership in n-dimensional lamplighter groups and **S-unit equations**

An **S-unit equation** is a linear equation

$$x_1 m_1 + \cdots + x_K m_K = m_0,$$

where we look for solutions  $x_1, \dots, x_K$  in a multiplicative subgroup of a field  $\mathbb{K}$ .

**Example:** solve  $x_1 + 3x_2 = 6$ , where  $x_1, x_2 \in \mathbb{Q}$  are of the form  $2^n 3^m$ .

## Submonoid Membership in n-dimensional lamplighter groups and **S-unit equations**

An **S-unit equation** is a linear equation

$$x_1 m_1 + \cdots + x_K m_K = m_0,$$

where we look for solutions  $x_1, \dots, x_K$  in a multiplicative subgroup of a field  $\mathbb{K}$ .

**Example:** solve  $x_1 + 3x_2 = 6$ , where  $x_1, x_2 \in \mathbb{Q}$  are of the form  $2^n 3^m$ .

Proposition (D. 2025)

*Submonoid Membership in any lamplighter group and its quotient reduces to solving S-unit equations in fields of characteristic 2. (such as  $\mathbb{F}_2(X)$ )*

# Submonoid Membership in n-dimensional lamplighter groups and **S-unit equations**

An **S-unit equation** is a linear equation

$$x_1 m_1 + \cdots + x_K m_K = m_0,$$

where we look for solutions  $x_1, \dots, x_K$  in a multiplicative subgroup of a field  $\mathbb{K}$ .

**Example:** solve  $x_1 + 3x_2 = 6$ , where  $x_1, x_2 \in \mathbb{Q}$  are of the form  $2^n 3^m$ .

**Proposition (D. 2025)**

*Submonoid Membership in any lamplighter group and its quotient reduces to solving S-unit equations in fields of characteristic 2. (such as  $\mathbb{F}_2(X)$ )*

**Theorem (Adamczewski and Bell 2012, Derksen and Masser 2012)**

*The solution set of an S-unit equation over a field of characteristic  $p > 0$  is effectively  $p$ -automatic.*