

Relationship between solar wind, D_{st} , and plasmasphere trough mass density on one-hour time scales

Victoir Veibell*

R.S. Weigel†

March 25, 2016

Abstract

This paper compares various magnetosphere conditions around the onset of geomagnetic events, defined as decreases in D_{st} below a threshold value or ρ_{eq} above a threshold value.

[More detail](#)

1 Introduction

The plasma trough is a transition region between the high density and low temperature plasmasphere and the lower density and higher energy magnetosphere [Mayr and Volland, 1968]. Inside of the plasmopause, the density is primarily due to ionospheric photoionization; outside of the plasmopause, the density has contributions from the co-rotating plasmasphere and convecting magnetosphere plasmas.

The dependence of the plasmopause location on geomagnetic activity was observed as early as ?. There are well-validated models of the plasmopause boundary, as a function of L , MLT , and geomagnetic activity (Lemaire [1998], Moldwin et al. [2002], O’Brien and Moldwin [2003b]). Models also exist for the plasmasphere density (Gallagher et al. [1988], Lemaire [1998], Takahashi et al. [2006], and Takahashi et al. [2010]), but the dependence of the plasma trough on geomagnetic and solar wind conditions has not been extensively studied until the past decade.

Takahashi et al. [2006] estimated magnetospheric mass density using measurements from the CRRES satellite during a 73-day period in 1991 (observations were made when CRRES was in an elliptical, near equatorial orbit and primarily in the plasma trough region). They found that the average ion mass, $M = \rho/n_e$, had some correlation with geomagnetic activity, with more negative hourly D_{st} values cor-

responding to higher M and 1.5- and 3-day averages of K_p corresponding to higher M (3-hour averages of K_p had little visual correlation with M). The mass density estimates were obtained from CRRES observations in the MLT range of 12:00 and 18:00 with most observations having L between 5 and 7 R_E .

Denton et al. [2006] found a weak relationship between D_{st} and K_p and the field line distribution of ρ using observations of the first three toroidal Alfvén harmonic frequencies for L in the range of 6-8 R_E . A more pronounced peak in the distribution along a field line near the equator was found for lower D_{st} and larger K_p . The equatorial value of ρ appeared identical for the bins of D_{st} (-142 to -31 nT and -31 to 37 nT) and K_p (1.5 to 3.4 and 3.4 to 5.9) considered.

Takahashi et al. [2010] developed a mass density dataset using measurements from the Space Environment Monitor instruments on the Geostationary Operational Environmental Satellites (GOES) satellites from 1980 through 1992, with most measurements in the range of $L = 6.8 \pm 0.2 R_E$. The mass density ρ was estimated using the Alfvén wave velocity relationship, $V_A = B/\sqrt{\mu_0 \rho}$, **Connect to harmonic toroidal frequency** a magnetic field model, and a numerical solution to a wave equation with an ionospheric boundary and the assumption of a zero resistance ionosphere. The equatorial mass density, ρ_{eq} , was derived from the estimated mass density using a power law dependence on the geocentric distance to the field line of the observation, R ,

*vveibell@gmu.edu

†rweigel@gmu.edu

$\rho = \rho_{eq} L / (R/R_E)$. They found a high correlation (~ 0.93) between 27-day averages of F10.7 and 27-day medians of ρ_{eq} . [2008]].

Takahashi et al. [2010] noted that downward spikes in the D_{st} index coincided with significant changes in ρ_{eq} for L near $6.8 R_E$. For five storms, two had ρ_{eq} spikes after the D_{st} drop, two had ρ_{eq} spikes before the drop, and one showed little change in ρ_{eq} . A key result was that when daily-averaged measurements were considered, an enhancement in ρ_{eq} appeared on the same day as minimum D_{st} .

Yao et al. [2008] studied the relationship between D_{st} and ion number density for low-energy O^+ in different regions (ring current and plasma sheet) using the TEAMS and ESA instruments on the low-altitude and high-inclination orbiting FAST satellite and found that the average N_{O^+} across the sampled L-shells (2-14) had a strong correlation (0.88). Although the correlation in the plasma trough was not calculated, the data presented show that the variations in N_{O^+} tended to appear near D_{st} minimum (± 2 hours).

The results in the papers discussed indicate that (1) the mass density in the plasma trough region is best correlated with F10.7 and (2) there exists a correlation between mass density and the geomagnetic activity indices K_p and D_{st} . In this work we consider the dependence of mass density estimates in the Takahashi et al. [2010] and attempt to identify and statistically characterize the relationship between geomagnetic activity with mass density in the plasma trough region. In addition, we attempt to identify solar wind processes may drive changes in mass density.

There are several issues that are addressed in detail: (1) the plasma density measurements are sparse, (2) The plasma trough density has a high correlation with F10.7, and geomagnetic and solar wind parameters also correlate with F10.7, (3) the dependence of mass density on the north/south component of the solar wind magnetic field is expected to be nonlinear. For northward IMF, corresponding to low solar wind driving of the magnetosphere, the plasmasphere can expand past geosynchronous orbit where trough estimates of mass density are made resulting in an increase in observed density that are not due to solar wind driving; for southward IMF, events studies have shown an increase in mass density associated with strong southward IMF and enhanced magnetospheric activity [Takahashi et al. [2010]; Yao et al.

Issue (1) is addressed by considering in detail the number of observations that are used in epoch averages around the time of two types of events, decreases in D_{st} below a threshold, and increases in ρ_{eq} above a threshold. Issue (2) is addressed using various methods to remove or isolate the F10.7 dependences. Issue (3) is addressed by considering epochs and linear correlations independent of the direction of the IMF along with separation of the epoch averages based on the northward and southward IMF.

In addition, we consider the influence of time averaging by comparing results using daily means and medians and hourly means and medians of solar wind and magnetospheric parameters.

–

Gallagher et al. [1988] used measurements from the Retarding Ion Mass Spectrometer onboard the Dynamics Explorer 1 satellite to model plasmasphere density using a relationship dependent on L shell, altitude, geomagnetic latitude, and observations grouped by a weighted average of K_p . The developed statistical model showed significant improvements when a weighted average of K_p over three days was used to group the observations used in the model.

Discuss plasma trough relevance.

Hamilton et al. [1988] analyzed a strong geomagnetic storm and compared energy measurements of the ring current to D_{st} , along with measurements of ring current compositions before, during, and after the storm. They found D_{st} to be correlated with the inner ring current energy density during and after storm onset, an inward movement of energy density, and a rapid D_{st} recovery from the quick loss of heavier ions.

Loto'aniu et al. [1999] used measurements from the Plasma Wave Experiment instrument on the CRRES satellite and compared in-situ measured plasma mass density to that inferred by ground-based ULF wave measurements from the CANOPUS array. By extending the ground footprints to the plasmatrough, they found agreement between the two methods for certain plasmapause conditions. **Mention specifics? Worth including paper?**

2 Data Preparation and Overview

The parameters ρ_{eq} and $F_{10.7}$ used in this work are from the dataset of Denton [2007], which was used in Takahashi et al. [2010]; data are available from 1980

through 1991 from GOES 2,3,5, 6 and 7. All other parameters used are from Kondrashov et al. [2014] from 1972 through 2013, which are on a 1-hour time grid. ρ_{eq} is the inferred equatorial mass density based on the 3rd harmonic toroidal frequency of magnetic field measurements as described in Takahashi et al. [2010]. The smallest cadence for ρ_{eq} values is 10 minutes. To compute an hourly average over the same interval for which the solar wind parameters were averaged, the median of all ρ_{eq} values in a given hour window was used. Fill (NaN) values were used for hours in which no measurements were available. In cases where ρ_{eq} was available from multiple GOES satellites at the same time, the value from GOES 6 was selected. In the dataset, measurements from GOES 6 had the longest time span of coverage. An overview of the data used is shown in Figure 1.

In this work, events are defined to occur when D_{st} or ρ_{eq} crosses a threshold value, as indicated by horizontal dashed lines in their respective panels of Figure 1. **need to put line in ρ_{eq} plot.**

Figure 2 shows the long-term trends of $\log(\rho_{eq})$ from GOES 6 and $F_{10.7}$ computed using the median values in 27-day non-overlapping windows. A scatter plot of these two lines is shown in the lower panel of Figure 2. The linear correlation for GOES 6 using all measurements is found to be 0.94, which is the same value documented by Takahashi et al. [2010] who used measurements from all satellites in the time interval of 1980 through 1991 using the constraint $0600 \leq MLT \leq 1200$, $K_{p3d} \geq 1.0$ and $D_{st} \geq -50$ nT. Our 27-day correlations for GOES 2, 5, and 7 are 0.81, 0.78, and 0.87, and the time range of coverage of measurements from these satellites are 1980-01-01/1983-05-16, 1983-01-01/1987-02-28, and 1983-05-27/1991-08-29 respectively.

3 Epoch Average Results

Two types of events are considered. The first is a drop in the D_{st} index Cite Kyoto below the threshold of -50 nT. These types of events are considered in order to determine if there is a well-defined ρ_{eq} dependence on geomagnetic activity/storms as indicated by D_{st} index. It is expected that the plasmasphere will exhibit two responses near these threshold crossings. First is the movement of the plasmasphere earthward and the second is a possible increase in density due to magneto-

spheric processes **cite.**

The second type of event is an increase in ρ_{eq} above a threshold of 20 amu/cm³. It is expected that these increases will typically occur after period of quiet geomagnetic activity that follow a geomagnetic storm. In this case, the plasmopause boundary may move outward past the measurement location and the increase will be due to measurements being made inside the plasmopause. A second possible reason for the increase could be due to the above-mentioned increase in due to magnetospheric processes.

3.1 D_{st} Events

Our first analysis uses data only from the time interval 1989-1991, which corresponds with the interval used in the analysis in Takahashi et al. [2010]. In this interval, there were 329 D_{st} events and 70 ρ_{eq} events. For both types of events, the hour of the threshold crossing defines the zero epoch time, and for each event, 4·24 hours were considered before each event and 8·24 hours after each event. To compute the epoch averages on a daily time scale, the median value of all available measurements for all events centered on a window of ± 12 hours of the epoch day was computed, and these averaging windows were shifted in increments of 24 hours. Similar results are obtained if we first reduce each event time series to have a 1-day cadence by computing medians in 1-day bins and then compute the medians across events. Error bars for each parameter are the standard deviation of the values used in computing the median divided by the square root of the number of values used.

The stack plot in the upper panel of Figure 3 shows the epoch averages for these events. The minimum D_{st} median is -48 nT. Consistent with Takahashi et al. [2010], D_{st} events correspond to elevated ρ_{eq} , although the magnitude of increase observed here is 1.5 amu/cm³ (from 18 to 19.5 amu/cm³) instead of the increase of ~ 10 amu/cm³ found in Takahashi et al. [2010]. The vertical green bars in the ρ_{eq} plot show the number of ρ_{eq} values that were used to compute the medians and error bars. For the non- ρ_{eq} medians, the number of measurements used in computing the medians is typically equal to the number of events, which is much larger than the number of ρ_{eq} values as we used all available measurements

instead of restricting to only values where a non-fill ρ_{eq} value existed.

In the lower panel of Figure 3 it is shown that the trends seen in the analysis of 1989-1991 also hold for the entire span of measurements from GOES 6, which spans 1983-1991 - a small elevation on the day of the event relative to the day before and day after, with median averages after the day of the event being slightly lower than that before.

To determine if the magnitude of the density medians on a given day were statistically different than another day, we used a two-population t-test. For the interval of 1989-1991, corresponding to the top panel of 3, the null hypothesis that any of the medians from days -1, 0, and 1 are different from each other is rejected with a maximum p value of ??.

Days p -1 and 0 -1 and 1 0 and 1

Caption: Results of test on different between means of ρ_{eq} shown in the top panel of ?? on days near or on the day of a D_{st} event. A difference in the means can be rejected with a probability of false rejection of p .

Days p -1 and 0 -1 and 1 0 and 1

Caption: Results of test on different between means of ρ_{eq} shown in the bottom panel of ?? on days near or on the day of a D_{st} event.

Both of the results in 3 indicate that there is not a statistically significant change in medians of ρ_{eq} on the day before and day after a threshold crossing in D_{st} .

Because the dynamics of the magnetosphere take place on a time scale of less than one day, we have also considered hourly averages as shown in the top panel of 4. A total of 668 events were found in the interval of May 1983 through August 1991 with an average duration of 9 hours and a median duration of 3 hours. The error bars were computed in the same way as 3. In this plot, the overall trend is similar to that found in the bottom panel of 4 - ρ_{eq} is slightly elevated at the time of the threshold crossing of D_{st} with respect to the values one day before and after, but the elevation is small and in general not statistically significant. At

this timescale no obvious change in mass density at hourly timescale is visible. This points to an issue with only looking at long-timescale trends between density and D_{st} , and allows for the possibility that other factors are influencing the long term correlation because there is no obvious connection on a short timescale. One possibility is that, as suggested in Takahashi et al. [2010], $F_{10.7}$ plays a significant role in driving long term density values which biases the long-term correlation of density and D_{st} . ===== Results of test on different between means of ρ_{eq} shown in the bottom panel of ?? on days near or on the day of a D_{st} event.

It was found that ρ_{eq} enhancements do not clearly correspond to a signature in D_{st} , but that there is a trend for such enhancements to occur when B_z in the IMF is positive. This is consistent with Takahashi et al. [2010] who note that very large increases in ρ_{eq} often corresponded to movement of the plasmasphere to beyond geosynchronous distances.

This method finds 669 periods of D_{st} crossing beneath the selected threshold between May 1983 and August 1991 with an average duration of 9 hours and a median duration of 3 hours Make sure I defined "duration". Figure 4 shows the average values of all events over a window of 24 hours before onset and 48 hours after. Figure 4a shows events selected by looking for event onsets where D_{st} crossed a threshold of -40 nT. The dashed red lines indicate plus and minus one standard deviation of values from all events that were used to compute the median. The final plot in the stack shows both ρ_{eq} and a bar plot of how many valid data points went into the ρ_{eq} average. Since ρ_{eq} comes from a sparser dataset, it has less valid points contributing to the averages than the other parameters in the stack.

Figure 4 shows a definite spike in the Z component of the magnetic field, as well as the defined drop in D_{st} , but no obvious change in mass density at an hourly timescale. This points to an issue with only looking at long-timescale trends between density and D_{st} , and allows for the possibility that other factors are influencing the long term correlation since there's no obvious connection on a short timescale. One possibility is that, as suggested

in Takahashi et al. [2010], $F_{10.7}$ plays a significant role in driving long term density values which biases the long term correlation of density and D_{st} .

Two types of events are considered. The first is a drop in D_{st} below the threshold of -40 nT. The second is an increase in ρ_{eq} above 40 amu/cm³.

Our first analysis uses data only from the time interval 1989-1991. In this interval, there were 329 D_{st} events and 70 ρ_{eq} events. For both types of events, the hour of the threshold crossing defines the zero epoch time, and for each event $8 \cdot 24$ hours were kept after each event and $4 \cdot 24$ hours before each event. To compute the epoch averages on a daily time scale, the median value of all measurements for all events centered window of ± 12 hours was computed, and these windows were shifted in increments of 24 hours. (Similar results are obtained if we first reduce each event time series to have a 1-day cadence by computing medians in 1-day bins and then compute the medians across events.) Error bars for each parameter were computed using the standard deviation of values used in computing the median divided by the square root of the number of values used.

In the upper panel of 3, D_{st} events are shown. The minimum D_{st} median is -48 nT. Consistent with Takahashi et al. [2010], D_{st} events correspond to elevated ρ_{eq} , although the magnitude of increase observed here is on the order of ~ 2 amu/cm³ instead of ~ 10 amu/cm³ found in Takahashi et al. [2010]. The vertical green bars in the ρ_{eq} plot show the number of ρ_{eq} values that were used to compute its median and error bars. Note that for D_{st} , the number of measurements used in computing the medians is much larger as we use all available D_{st} measurements instead of restricting to only values where a non-fill ρ_{eq} value existed.

Both of the results in 3 indicate that there is not a statistically significant change in ρ_{eq} on the day before and day after a threshold crossing in D_{st} .

Because the dynamics of the magnetosphere

take place on a time scale of less than one day, we have also considered hourly averages as shown in the top panel of 4. A total of 668 events were found in the interval of May 1983 through August 1991 with an average duration of 9 hours and a median duration of 3 hours. The error bars were computed in the same way as 3. In this plot, the overall trend is similar to that found in the bottom panel of 4 - ρ_{eq} is slightly elevated at the time of the threshold crossing of D_{st} with respect to the values one day before and after, but the elevation is small and in general not statistically significant.

The bottom panel of 4 contains the events shown in the top panel under the constraint that D_{st} remained below -50 nT for at least twelve hours after the event onset time.

and the event study of Yao et al. [2008] which showed ...

3.2 ρ_{eq} Events

An alternative type of event can be defined that corresponds to large increases in ρ . In finding ρ_{eq} events, all fill values were replaced with linearly interpolated values. Not including ρ_{eq} events?

Figure 5 shows the epoch time series for events in which an increase of ρ_{eq} above 20 amu/cm² defines the start of the event. Here we see the expected result that such events tend to be associated with positive B_z^{IMF} . To determine if the sign of B_z^{IMF} influences the ρ epoch time series, we separated the events by the average of B_z^{IMF} in a time window of 4 hours.

Figure (a) shows the The epoch curves for $B_z^{IMF} \geq 0$ (257) events and $B_z^{IMF} \leq 0$ (139) have no statistically significant difference.

4 F10.7 dependence

In an effort to analyze the dependence of ρ_{eq} on $F_{10.7}$, a few tests were performed. Takahashi et al. [2010] mention a strong correlation between the two. The long term correlation could be a bias for D_{st} 's effects, so a linear model was created, recreating mass density purely from $F_{10.7}$ in the form of $\rho_{eq}(t) = a \cdot F_{10.7}(t)$. This re-created ρ_{eq} shows around a

45% correlation with the actual ρ_{eq} , suggesting a strong influence, while doing the same procedure with D_{st} shows only a 20-25% correlation.

Figure 7 takes the events and bins them by the value of $F_{10.7}$ at event onset, then looks at both ρ_{eq} and B_z . Error bars calculated in the same manner as before are added to the highest and lowest bins. A significant difference is seen in ρ_{eq} between events taking place during periods of high $F_{10.7}$ and low $F_{10.7}$, while a much less significant, but still noticeable, difference is seen in B_z . Also of interest is how ρ_{eq} only sharply reacts to D_{st} events during periods of high $F_{10.7}$. The positive relationship seen in Takahashi et al. [2010] could be due to the short time period looked at, and that it coincides with periods of higher $F_{10.7}$.

Looking at GOES 2,5,6, and 7 shows that this post-onset D_{st} spike happens in almost every case for periods of high $F_{10.7}$. Creating a linear model directly relating variables (and combinations of variables) to ρ_{eq} also shows that $F_{10.7}$ most often has the highest correlation with ρ_{eq} . A sample of these correlations is shown in Table ?? . The correlations were made by training a one-input linear model of 5-hour post-onset medians of each variable (or variables) against the median of ρ_{eq} at, and four hours after, event onset. This was done by randomly sampling half of the data, generating the model, and testing against the other half, doing this process 100 times and then taking the median correlations.

	GOES 2	GOES 5	GOES 6	GOES 7
DoY	-0.08 ± 0.08	$+0.14 \pm 0.13$	-0.06 ± 0.06	$+0.09 \pm 0.10$
MLT	-0.10 ± 0.21	-0.07 ± 0.12	$+0.01 \pm 0.23$	-0.06 ± 0.05
B_z	$+0.16 \pm 0.21$	-0.13 ± 0.15	$+0.08 \pm 0.14$	-0.07 ± 0.06
V_{sw}	-0.04 ± 0.10	$+0.27 \pm 0.09$	$+0.06 \pm 0.11$	$+0.06 \pm 0.06$
D_{st}	$+0.26 \pm 0.17$	$+0.66 \pm 0.08$	$+0.06 \pm 0.13$	$+0.23 \pm 0.14$
ρ_{sw}	$+0.35 \pm 0.24$	$+0.63 \pm 0.31$	$+0.12 \pm 0.19$	$+0.36 \pm 0.17$
$F_{10.7}$	$+0.43 \pm 0.08$	$+0.12 \pm 0.12$	$+0.51 \pm 0.06$	$+0.40 \pm 0.06$
$B_z + V_{sw}$	$+0.11 \pm 0.17$	$+0.20 \pm 0.17$	$+0.12 \pm 0.10$	-0.12 ± 0.06
$D_{st} + F_{10.7}$	$+0.44 \pm 0.09$	$+0.71 \pm 0.08$	$+0.54 \pm 0.07$	$+0.47 \pm 0.06$
<i>All</i>	-0.03 ± 0.19	$+0.34 \pm 0.27$	$+0.61 \pm 0.11$	$+0.49 \pm 0.12$

Table 1: Table of linear model test correlations showing the median of 100 random samples. Each sample trained on half of the data (via randomly selected rows of the least squares matrix) and tested on the other half

5 Appendix

5.1 Bias

While attempting to reproduce Figure 11 from Takahashi et al. [2010], an ambiguity in data-handling was found. It is unknown how they got from hourly ρ_{eq} to daily medians, whether in one step or two steps, and whether the hourly medians are an hour ahead of each hour grid point, or a median of points a half hour to either side, so attempts were made to reproduce all possibilities and compare in order to find a data-handling method with the least bias. These attempts are shown in Figure ?? . This also shows the effect of only considering events with a minimum before noon. All show a similar spike in ρ_{eq} at the time of D_{st} minimum, though none quite reproduce the exact values seen in Takahashi et al. [2010]. Figure ?? shows where the median falls for all of the found events during that period, as a sanity check for why the prior work would get median values nearing 30 amu/cm³. Both authors of this paper conducted independent trials to confirm that the data and analysis don't quite yield the results of Figure 11 in Takahashi et al. [2010] with the processes described in that paper, but the same general trend is still seen.

To check for potential bias in the data as used by our analysis, we checked how the data availability varied with hour, shown in Figure ?? . We also looked at how storm conditions affected data availability. Both a two sample t-test for difference in means and a Wilcoxon rank sum test showed significance at the 1% level that D_{st} during periods with available data was of a different distribution than D_{st} during periods missing data. When testing for other significant D_{st} differences, the Wilcoxon test was significant while the t-test was not in pre-noon vs post-noon distributions, as well as $\rho_{eq} > 40$ vs $\rho_{eq} < 40$ distributions indicating perhaps an increase in variability without an increase in mean.

While the GOES Satellites tend to keep an average geostationary distance of around $L = 6.8R_E$ shown in Takahashi et al. [2010], the actual plasmopause location varies significantly, as shown in O'Brien and Moldwin [2003a]. This means that during periods of large D_{st} the plasmopause may be far from the point of measurement of ρ_{eq} creating a discord in the

correlation. Gallagher et al. [2000] provides a model for ρ_{eq} at a range of L-shells and a brief discussion of the elemental contributions.

References

- R. Denton. Database of Input Parameters for Tsyganenko Magnetic Field Models, 2007. URL <http://www.dartmouth.edu/~rdenton/magpar/index.html>.
- R. E. Denton, K. Takahashi, I. A. Galkin, P. A. Nsumei, X. Huang, B. W. Reinisch, R. R. Anderson, M. K. Sleeper, and W. J. Hughes. Distribution of density along magnetospheric field lines. *Journal of Geophysical Research (Space Physics)*, 111:A04213, April 2006. doi:10.1029/2005JA011414.
- D. L. Gallagher, P. D. Craven, and R. H. Comfort. An empirical model of the earth's plasmasphere. *Advances in Space Research*, 8:15–21, 1988. doi:10.1016/0273-1177(88)90258-X.
- D. L. Gallagher, P. D. Craven, and R. H. Comfort. Global core plasma model. *Journal of Geophysical Research*, 105:18819, August 2000. doi:10.1029/1999JA000241.
- D. C. Hamilton, G. Gloeckler, F. M. Ipavich, B. Wilken, and W. Stuedemann. Ring current development during the great geomagnetic storm of February 1986. *J. Geophys. Res.*, 93:14343–14355, December 1988. doi:10.1029/JA093iA12p14343.
- D. Kondrashov, R. Denton, Y. Y. Shprits, and H. J. Singer. Reconstruction of gaps in the past history of solar wind parameters. *Geophysical Research Letters*, 41:2702–2707, April 2014. doi:10.1002/2014GL059741.
- J Lemaire. *The Earth's Plasmasphere*. Cambridge University Press, Cambridge, U.K. New York, 1998. ISBN 9780521430913.
- T. M. Loto'aniu, C. L. Waters, B. J. Fraser, and J. C. Samson. Plasma mass density in the plasmatrough: Comparison using ULF waves and CRRES. *Geophys. Res. Lett.*, 26:3277–3280, 1999. doi:10.1029/1999GL003641.
- H. G. Mayr and H. Volland. Model of magnetospheric temperature distribution. *J. Geophys. Res.*, 73:4851–4858, August 1968. doi:10.1029/JA073i015p04851.
- M. B. Moldwin, L. Downward, H. K. Rassoul, R. Amin, and R. R. Anderson. A new model of the location of the plasmopause: CRRES results. *Journal of Geophysical Research (Space Physics)*, 107:1339, November 2002. doi:10.1029/2001JA009211.
- T. P. O'Brien and M. B. Moldwin. Empirical plasmopause models from magnetic indices. *Geophysical Research Letters*, 30:1152, February 2003a. doi:10.1029/2002GL016007.
- T. P. O'Brien and M. B. Moldwin. Empirical plasmopause models from magnetic indices. *Geophys. Res. Lett.*, 30:1152, February 2003b. doi:10.1029/2002GL016007.
- K. Takahashi, R. E. Denton, R. R. Anderson, and W. J. Hughes. Mass density inferred from toroidal wave frequencies and its comparison to electron density. *Journal of Geophysical Research (Space Physics)*, 111:A01201, January 2006. doi:10.1029/2005JA011286.

- K. Takahashi, R. E. Denton, and H. J. Singer. Solar cycle variation of geosynchronous plasma mass density derived from the frequency of standing Alfvén waves. *Journal of Geophysical Research (Space Physics)*, 115:A07207, July 2010. doi:[10.1029/2009JA015243](https://doi.org/10.1029/2009JA015243).
- Y. Yao, K. Seki, Y. Miyoshi, J. P. McFadden, E. J. Lund, and C. W. Carlson. Effect of solar wind variation on low-energy O⁺ populations in the magnetosphere during geomagnetic storms: FAST observations. *Journal of Geophysical Research (Space Physics)*, 113, 2008. doi:[10.1029/2007JA012681](https://doi.org/10.1029/2007JA012681).

6 Figures




Figure 1: Overview of data used in this study. The top four panels show parameters from [Kondrashov et al. \[2014\]](#) and the bottom panel contains ρ_{eq} based on GOES 6 measurements from [Denton \[2007\]](#) after interpolation and averaging described in the text. Dashed horizontal lines in the D_{st} and ρ_{eq} panels indicate sample event cutoff thresholds of $D_{st} = -50$ nT and $\rho_{eq} = 40$ amu/cm³




Figure 2: Top: 27-day medians of $F_{10.7}$ and ρ_{eq} from GOES 6. Bottom: Correlation between $\log(\rho_{eq})$ and $F_{10.7}$ using medians in hour, day, and 27-day non-overlapping; compare to [Takahashi et al. \[2010\]](#) Fig. 14.




Figure 3: D_{st} events from GOES 6 using daily medians. Top: Events in the interval 1989-1991; compare to [Takahashi et al. \[2010\]](#) Fig. 11. Bottom: Events in the interval 1983-1991.




Figure 4: D_{st} events from GOES 6 using hourly medians. Top: Events in the interval 1983-1991; compare to bottom panel DailyAveragedDstEvents. Bottom: Same as Top except for constraint that D_{st} stayed below -50 nT for at least 12 hours after crossing below -50 nT.




Figure 5: $\rho_{eq} > 20$ amu/cm³ events from GOES 6 using hourly medians. As described in the text, there is no statistically significant dependence of the ρ_{eq} plot on B_z in the time interval of ± 2 hours around the event.




Figure 6: Binning ρ_{eq} events by B_z value (a) at onset and four hours before, and (b) at onset and four hours after.




Figure 7: Binning events by $F_{10.7}$ value at onset for ρ_{eq} and, in the middle panel, B_z . Lower panel shows number of valid points in each bin

7 Appendix




Figure 8: (a) Same as Figure 4(a) except with constraint that D_{st} crossed below -80 nT




Figure 9: GOES Satellite 6 and 7




Figure 10: GOES Satellite 2 and 5




Figure 11: Top: GOES 6 from 89-91 (compare to Takahashi 2010). Bottom: $D_{st} < -80$ nT.




Figure 12: GOES 6 from 83-87 and 87-91




Figure 13: $D_{st} < -30$ nT events, ρ_{eq} binned by $F_{10.7}$. Lower panel: valid hourly points




Figure 14: $\rho_{eq} > 40$ amu/cm³ events, ρ_{eq} binned by $F_{10.7}$




Figure 15: Top: $AE > 400$ nT events, ρ_{eq} binned by $F_{10.7}$ Bottom: averages of $AE > 400$ events




Figure 16: Top: $AE > 400$ nT events. Bottom: $D_{st} < -50$ nT events. Both ρ_{eq} binned by D_{st} .




Figure 17: Top: ρ_{eq} binned by $F_{10.7}$ for randomly selected times. Bottom: averages of randomly selected times




Figure 18: Top: B_z binned by D_{st} for $D_{st} < -50$ nT events. Bottom: B_z binned by D_{st} for $AE > 400$ nT events




Figure 19: Median ρ_{eq} vs magnetic local time and universal time




Figure 20: Median D_{st} vs universal time, and median AE vs universal time