

NUMERICAL METHODS FOR FLUID DYNAMICS

EMMANUEL DORMY (DMA-ENS)


emmanuel.dormy@ens.fr

CFD

```
lv=np.size(vec) # lv = n+1;
D1=np.zeros([lv,lv])
D2=np.zeros([lv,lv])
D3=np.zeros([lv,lv])
D4=np.zeros([lv,lv])
D1[:,1]=D0[:,0]
D1[:,2]=4*D0[:,1]
D2[:,2]=4*D0[:,0]
for i in range(n-2): # We compute Tj, which is in column j+1
    j=i+2
    jj=j+1
    D1[:,j+1]=2*jj*D0[:,j]+jj*D1[:,j-1]/(jj-2);
    D2[:,j+1]=2*jj*D1[:,j]+jj*D2[:,j-1]/(jj-2);
    D3[:,j+1]=2*jj*D2[:,j]+jj*D3[:,j-1]/(jj-2);
    D4[:,j+1]=2*jj*D3[:,j]+jj*D4[:,j-1]/(jj-2);
return D0,D1,D2,D4

au=np.linalg.solve(D0,u)
du_Cheb=D1.dot(au)
err_Cheb[k]=np.max(abs(du-du_Cheb))
```

CFD



CFD

```

for i in range(NT):
    # Laplacian(psi) = -w
    psi_hat = -w_hat/k2_fixed

    # Velocity from streamfunction
    u = spec_phys(ky*psi_hat,NX,NY,NXSp,NYSp)
    v = spec_phys(-kx*psi_hat,NX,NY,NXSp,NYSp)

    # Gradients of vorticity
    w_x = spec_phys(kx*w_hat,NX,NY,NXSp,NYSp)
    w_y = spec_phys(ky*w_hat,NX,NY,NXSp,NYSp)

    # Non-linear term
    NL      = u*w_x + v*w_y
    NL_hat = phys_spec(NL,NX,NY,NXSp,NYSp)


    # Time stepping (Crank-Nicholson)
    w_hat = (1/dt+k2/(2*Re))*w_hat
    w_hat = w_hat-NL_hat

    w_hat = w_hat/(1/dt-k2/(2*Re))

```


CFD

Vortices interactions



Leweke et al (2016)

CFD



Hydrodynamic turbulence



Leonardo da Vinci

c1510

Hydrodynamic turbulence

Kolmogorov K41

$$[\hat{E}(k)] = L^3 T^{-2}.$$

$$[k] = L^{-1} \quad \text{et} \quad [\varepsilon] = L^2 T^{-3}.$$

$$\hat{E}(k) \propto \varepsilon^{2/3} k^{-5/3}.$$

Hydrodynamic turbulence

Kolmogorov K41

$$\ell_{\text{diss}} \sim \left(\nu^3 / \varepsilon \right)^{1/4},$$

$$\ell_{\text{diss}} \sim \text{Re}^{-3/4}$$

$$N \sim \text{Re}^{9/4}$$

Hydrodynamic turbulence

The Lorenz system

$$\dot{X} = \sigma(Y - X)$$

$$\dot{Y} = X(\rho - Z) - Y$$

$$\dot{Z} = \beta Z + XY$$

CFD

There are two different stability concepts:


- **Zero stability:** If $t > 0$ is held constant, do the values $u(t)$ remain bounded as $\Delta t \rightarrow 0$?
(i.e. vanishing time-step for fixed time)
- **Absolute stability:** If Δt is kept fixed, do the computed values $u(t)$ remain bounded as $t \rightarrow \infty$?
(i.e. finite time-step, infinite time)

What's the relevance
of numerical simulations?

- The attractor is robust
 - Shadowing lemma in dynamical systems

CFD

To explore further:



CFD

7. Compressibility and waves

CFD

Waves are a common feature in many fluid dynamic applications:

- oceanography/geology
- acoustics
- instabilities



CFD

Waves are a common feature in many fluid dynamic applications:



- oceanography/geology
- **acoustics**
- instabilities



CFD

Waves are a common feature in many fluid dynamic applications:

- oceanography/geology
- acoustics
- **instabilities**



CFD

7. Compressibility and waves

7.1 Staggering

CFD

Linear acoustics

Start with compressible Euler equations

$$\frac{\partial \rho}{\partial t} + \nabla \cdot (\rho \mathbf{u}) = 0$$

$$\rho \frac{D\mathbf{u}}{Dt} = -\nabla p + \eta \left(\nabla^2 \mathbf{u} + \frac{1}{3} \nabla (\nabla \cdot \mathbf{u}) \right) + \mathbf{F}$$

Homentropic fluid $p = p(\rho)$

CFD

Linear acoustics

Inertial effects are negligible; leads to linearized Euler equations

$$\frac{1}{\rho_0} \frac{\partial \rho'}{\partial t} + \nabla \cdot \mathbf{u} = q$$

$$\rho_0 \frac{\partial \mathbf{u}}{\partial t} + \nabla p' = \mathbf{F}$$

Eliminate velocity field by cross-differentiation

CFD

Linear acoustics

$$\frac{\partial^2 \rho'}{\partial t^2} - \nabla^2 p' = \rho_0 \frac{\partial q}{\partial t} - \nabla \cdot \mathbf{F}$$

Density and pressure are related via the EOS

$$p_0 + p' = p(\rho_0 + \rho') = p(\rho_0) + \left(\frac{\partial p}{\partial \rho} \right)_0 \rho' + \dots$$

c_0^2 →

CFD

Linear acoustics

Finally, we get an inhomogeneous wave equation for the pressure

$$\left(\frac{1}{c_0^2} \frac{\partial^2}{\partial t^2} - \nabla^2 \right) p = \boxed{\rho_0 \frac{\partial q}{\partial t} - \nabla \cdot \mathbf{F}}$$

↑
Source term

CFD


$$\frac{1}{\rho_0} \frac{\partial \rho'}{\partial t} + \nabla \cdot \mathbf{u} = q$$

$$\rho_0 \frac{\partial \mathbf{u}}{\partial t} + c_0^2 \nabla \rho' = \mathbf{F}$$

CFD

$$\frac{1}{\rho_0} \frac{\partial \rho'}{\partial t} + \nabla \cdot \mathbf{u} = q$$


$$\rho_0 \frac{\partial \mathbf{u}}{\partial t} + c_0^2 \nabla \rho' = \mathbf{F}$$




CFD

$$\frac{1}{\rho_0} \frac{\partial \rho'}{\partial t} + \nabla \cdot \mathbf{u} = q$$

$$\rho_0 \frac{\partial \mathbf{u}}{\partial t} + c_0^2 \nabla \rho' = \mathbf{F}$$



CFD



CFD

$$\frac{1}{\rho_0} \frac{\partial \rho'}{\partial t} + \nabla \cdot \mathbf{u} = q$$

$$\rho_0 \frac{\partial \mathbf{u}}{\partial t} + c_0^2 \nabla \rho' = \mathbf{F}$$

$$\left(\frac{1}{c_0^2} - \Delta \right) p = \rho_0 \frac{\partial q}{\partial t} - \nabla \cdot \mathbf{F}$$

$$\frac{1}{c_0^2} \frac{p_j^{n+1} - 2p_j^n + p_j^{n-1}}{\Delta t^2} = \frac{p_{j+1}^n - 2p_j^n + p_{j-1}^n}{\Delta x^2}$$

CFD

$$\frac{1}{c_0^2} \frac{p_j^{n+1} - 2p_j^n + p_j^{n-1}}{\Delta t^2} = \frac{p_{j+1}^n - 2p_j^n + p_{j-1}^n}{\Delta x^2}$$

This scheme is second-order in time and second-order in space.

CFD

Von Neuman stability analysis

$$\frac{1}{c_0^2} \frac{p_j^{n+1} - 2p_j^n + p_j^{n-1}}{\Delta t^2} = \frac{p_{j+1}^n - 2p_j^n + p_{j-1}^n}{\Delta x^2},$$

$$p_j^{n+1} = 2p_j^n - p_j^{n-1} - \frac{c_0^2 \Delta t^2}{\Delta x^2} (p_{j+1}^n - 2p_j^n + p_{j-1}^n)$$

CFD

Von Neuman stability analysis

$$\frac{\xi_j^{n+1} - 2\xi_j^n + \xi_j^{n-1}}{\Delta t^2} = c^2 \xi^n \frac{e^{ik\Delta x} - 2 + e^{-ik\Delta x}}{\Delta x^2}$$

$$\xi^2 - 2 [2 + \lambda^2(2 \cos k\Delta x - 2)] \xi + 1 = 0$$

$$\lambda = c\Delta t/\Delta x$$

CFD

$$\xi^2 - 2 [2 + \lambda^2(2 \cos k\Delta x - 2)] \xi + 1 = 0$$

Both roots have to be inside the unit disk for stability.

CFL restriction $\lambda \leq 1$

$\lambda > 1$: $\pi/\Delta x$ waves become unstable first

CFD

von Neumann stability analysis

In two dimensions, the CFL restriction extends to

$$\lambda \leq \frac{1}{\sqrt{2}}$$


In three dimensions, the CFL restriction extends to

$$\lambda \leq \frac{1}{\sqrt{3}}$$

CFD

Application


Propagation of a pressure pulse with reflecting boundaries



CFD

Application


Propagation of a pressure pulse with reflecting boundaries



CFD

Application


Propagation of a pressure pulse with reflecting boundaries



CFD

Application


Propagation of a pressure pulse with reflecting boundaries



CFD

Application

Propagation of a pressure pulse with reflecting boundaries



CFD

Grid anisotropy

Different waves travel at different speeds on the grid, although continuous wave equation is nondispersive

$$c_p = \omega/k \quad \xi = e^{i\omega\Delta t}$$

$$c_p(k_x, k_y) = \left| \frac{\ln (\xi(k_x, k_y)/|\xi(k_x, k_y)|)}{i \sqrt{k_x^2 + k_y^2} \Delta t} \right|$$

CFD

With:


$$\xi^2(k_x, k_y) + B(k_x, k_y) \xi(k_x, k_y) + 1 = 0$$

$$B(k_x, k_y) = -2 (1 + \lambda^2 (\cos(k_x \Delta) + \cos(k_y \Delta) - 2))$$

CFD

Grid anisotropy

$$B(k_x, k_y) = -2 (1 + \lambda^2 (\cos(k_x \Delta) + \cos(k_y \Delta) - 2))$$



strongly anisotropic wave propagation

CFD

Grid anisotropy

$$\begin{aligned}
 p_{i,j}^{n+1} = & -p_{i,j}^{n-1} + \lambda^2 a (p_{i,j+1}^n + p_{i,j-1}^n + p_{i+1,j}^n + p_{i-1,j}^n) \\
 & + \lambda^2 b (p_{i+1,j+1}^n + p_{i+1,j-1}^n + p_{i-1,j+1}^n + p_{i-1,j-1}^n) \\
 & + \lambda^2 c p_{i,j}^n
 \end{aligned}$$



$$\begin{aligned}
 B(k_x, k_y) = & -2\lambda^2 (a(\cos(k_x \Delta) + \cos(k_y \Delta))) \\
 & +(1 - a) \cos(k_x \Delta) \cos(k_y \Delta) - 1 - a) - 2
 \end{aligned}$$

CFD

Grid anisotropy

$$\begin{aligned}
 B(k_x, k_y) = & -2\lambda^2 (a(\cos(k_x \Delta) + \cos(k_y \Delta))) \\
 & +(1 - a) \cos(k_x \Delta) \cos(k_y \Delta) - 1 - a) - 2
 \end{aligned}$$

$$a = 0.62$$



more isotropic wave propagation

CFD


7. Compressibility and waves

7.2 Water waves





CFD



CFD

$$\omega^2 = gk \tanh kH .$$

Shallow water

$$\omega^2 = gHk^2 , \quad c = \omega/k = \pm \sqrt{gH} .$$

Deep water

$$\omega = \pm \sqrt{gk} , \quad c = \omega/k = \pm \sqrt{g/k} .$$

CFD

$$\rho \frac{Du}{Dt} = \rho \frac{\partial u}{\partial t} + \rho u \frac{\partial u}{\partial x} + \rho w \frac{\partial u}{\partial z} = - \frac{\partial p}{\partial x},$$

$$\rho \frac{Dw}{Dt} = \rho \frac{\partial w}{\partial t} + \rho u \frac{\partial w}{\partial x} + \rho w \frac{\partial w}{\partial z} = - \frac{\partial p}{\partial z} - \rho g.$$

$$\frac{\partial u}{\partial x} + \frac{\partial w}{\partial z} = 0.$$

$$\frac{dh}{dt} = w \mid_{z=h} \quad \text{and} \quad p \mid_{z=h} = p_0.$$

CFD

Hydrostatic basic state

$$u = w = 0, \quad h = H, \quad p = p_0 + \rho g (H - z),$$

$$\rho \frac{Du}{Dt} = \rho \frac{\partial u}{\partial t} + \rho u \frac{\partial u}{\partial x} + \rho w \frac{\partial u}{\partial z} = - \frac{\partial p}{\partial x},$$

$$\rho \frac{Dw}{Dt} = \rho \frac{\partial w}{\partial t} + \rho u \frac{\partial w}{\partial x} + \rho w \frac{\partial w}{\partial z} = - \frac{\partial p}{\partial z} - \rho g.$$

$$\frac{\partial u}{\partial x} + \frac{\partial w}{\partial z} = 0.$$

CFD

Hydrostatic basic state

$$u = w = 0, \quad h = H, \quad p = p_0 + \rho g (H - z),$$

Perturbation

$$\rho \frac{\partial u'}{\partial t} = -\frac{\partial p'}{\partial x}, \quad \rho \frac{\partial w'}{\partial t} = -\frac{\partial p'}{\partial z},$$

$$\frac{\partial u'}{\partial x} + \frac{\partial w'}{\partial z} = 0.$$

CFD

Hydrostatic basic state

$$u = w = 0, \quad h = H, \quad p = p_0 + \rho g (H - z),$$

$$\rho \frac{\partial u'}{\partial t} = -\frac{\partial p'}{\partial x}, \quad \rho \frac{\partial w'}{\partial t} = -\frac{\partial p'}{\partial z},$$

$$\frac{\partial u'}{\partial x} + \frac{\partial w'}{\partial z} = 0.$$

$$\frac{\partial^2 p'}{\partial x^2} + \frac{\partial^2 p'}{\partial z^2} = 0.$$

$$\mathsf{CFD}$$

$$p'(x,z,t)=\hat p(z,t)\,{\mathrm e}^{ikx}\,+c.c.\,,$$

$$\frac{\partial^2 \hat{p}}{\partial z^2}-k^2\,\hat{p}=0\,,$$

$$\hat p(z,t)=p_1(t)\,{\mathrm e}^{kz}+p_2(t)\,{\mathrm e}^{-kz}\,.$$

$$\mathsf{CFD}$$

$$\hat p(z,t)=p_1(t)\,{\mathrm e}^{kz}+p_2(t)\,{\mathrm e}^{-kz}\,.$$

$$\frac{\partial \hat{p}}{\partial z}(0,t)=0\,.$$

$$P=p_1/2=p_2/2$$

$$\mathbf{CFD}$$

$$p'(x,z,t)=P(t)\,\cosh kz\,{\rm e}^{ikx}+c.c.\,.$$

$$u'(x,z,t)=-i(k/\rho)\,Q(t)\,\cosh kz\,{\rm e}^{ikx}+c.c.\,,$$

$$w'(x,z,t)=-(k/\rho)\,Q(t)\,\sinh kz\,{\rm e}^{ikx}+c.c.\,,$$

$${\mathrm d} Q/{\mathrm d} t = P \qquad \qquad p' = \rho g h'$$

$$h'(x,t)=\frac{1}{g\rho}P(t)\cosh kH\,{\rm e}^{ikx}+c.c.\,,$$

$$\frac{\partial p'}{\partial t}(x,H,t)=g\rho w'(x,H,t)\,.$$

$$\left(\frac{\partial^2 \psi}{\partial x^2} + \frac{\partial^2 \psi}{\partial y^2}\right)_{\rm max} = 0$$

$$\mathbf{CFD}$$

$$\frac{{\mathrm d} P}{{\mathrm d} t}\,\cosh kH=-gk\,Q(t)\,\sinh kH\,,$$

$$\frac{{\mathrm d}^2 Q}{{\mathrm d} t^2}+gk\,Q(t)\,\tanh kH=0\,.$$

$$\omega^2=gk\tanh kH\,.$$

CFD

$$\omega^2 = gk \tanh kH .$$


Shallow water

$$\omega^2 = gHk^2 , \quad c = \omega/k = \pm\sqrt{gH} .$$

Deep water

$$\omega = \pm\sqrt{gk} , \quad c = \omega/k = \pm\sqrt{g/k} .$$

CFD



©2011, Dan Russell

CFD



CFD



CFD

7. Compressibility and waves

7.3 Conservation laws

and Finite Volumes


CFD

Finite Volumes:

$$u_j^n = \frac{1}{\Delta x} \int_{x_j - 1/2}^{x_j + 1/2} u(x, t) dx$$

CFD

Finite Volumes:



CFD

Finite Volumes:

$$u_j^{n+1} - u_j^n = -\frac{\Delta t}{\Delta x} \left(f_{j+1/2}^{n+1/2} - f_{j-1/2}^{n+1/2} \right)$$

$$f_{j+1/2}^{n+1/2} = \frac{1}{\Delta t} \int_{t^n}^{t^{n+1}} f(u(x_{j+1/2}, t)) dt$$

CFD



Suhas V. Patankar
(1941)

series in computational
methods in mechanics
and thermal sciences

Numerical
Heat
Transfer
and Fluid
Flow

Suhas V.
Patankar



Suhas V. Patankar
(1941)

CFD

series in computational
methods in mechanics
and thermal sciences

Numerical
Heat
Transfer
and Fluid
Flow

Suhas V.
Patankar

“much like preparing mashed potatoes from dehydrated potato powder. After all, textbook derivations of differential equations always start from the conservation principle applied to a small control volume.”

CFD

Upwind:

$$f_{j+1/2}^{up} = \begin{cases} c u_j & \text{if } c > 0 \\ c u_{j+1} & \text{if } c < 0 \end{cases}$$

Lax-Wendroff:

$$f_{j+1/2}^{LW} = \frac{1}{2}c(u_j + u_{j+1}) - \frac{\Delta t}{2\Delta x}c^2(u_{j+1} - u_j)$$

CFD

Finite Volumes in 2D:

$$\begin{aligned} \psi_{i,j}^{n+1} = & \psi_{i,j}^n - \frac{\Delta t}{\Delta x} (F_{i+1/2,j} - F_{i-1/2,j}) \\ & - \frac{\Delta t}{\Delta y} (G_{i,j+1/2} - G_{i,j-1/2}) \end{aligned}$$

CFD


Non-linear conservation laws

$$\frac{\partial u}{\partial t} = -\nabla \cdot \mathbf{F}(u)$$

$$\frac{\partial u}{\partial t} = -\frac{\partial f(u)}{\partial x}$$

CFD


$$f_{j+1/2}^{n+1/2} = \frac{1}{\Delta t} \int_{t^n}^{t^{n+1}} f(u(x_{j+1/2}, t) dt$$



CFD

Riemann problem:

$$u(x, 0) = \begin{cases} u_L & \text{if } x < 0 \\ u_R & \text{if } x > 0 \end{cases}$$
$$u_{j+1/2}^*(x/t) = RP [u_j^n, u_{j+1}^n]$$



CFD

Riemann problem:

$$u(x, 0) = \begin{cases} u_L & \text{if } x < 0 \\ u_R & \text{if } x > 0 \end{cases}$$
$$u_{j+1/2}^*(x/t) = RP [u_j^n, u_{j+1}^n]$$

$$f_{j+1/2}^{n+1/2} = f(u_{j+1/2}^*(0))$$

CFD

The Burgers equation

$$\frac{\partial u}{\partial t} = -\frac{\partial f(u)}{\partial x}$$

$$f(u) = \frac{u^2}{2}$$

CFD

The Burgers equation

$$f(u) = \frac{u^2}{2}$$

$$\frac{\partial u}{\partial t} = -\frac{\partial}{\partial x} \left(\frac{u^2}{2} \right)$$

CFD


The Burgers equation

$$f(u) = \frac{u^2}{2}$$

$$\frac{\partial u}{\partial t} = -\frac{\partial}{\partial x} \left(\frac{u^2}{2} \right)$$

$$\frac{\partial u}{\partial t} = -u \frac{\partial u}{\partial x}$$

CFD



CFD

The Burgers equation


$$\frac{\partial u}{\partial t} = -\frac{\partial}{\partial x} \left(\frac{u^2}{2} \right)$$

$$\frac{\partial u}{\partial t} = -\frac{\partial}{\partial x} \left(\frac{u^2}{2} \right) + \varepsilon \frac{\partial^2 u}{\partial x^2}$$

KdV: $\frac{\partial u}{\partial t} = -\frac{\partial}{\partial x} \left(\frac{u^2}{2} \right) - \alpha \frac{\partial^3 u}{\partial x^3}$

CFD

$u_L > u_R$ **Shock**



CFD

$u_L > u_R$ **Shock**


$$S = \frac{u_L + u_R}{2}$$

$$u\left(\frac{x}{t}\right) = \begin{cases} u_L & \text{if } S \geq x/t \\ u_R & \text{if } S < x/t \end{cases}$$

$$\begin{cases} u_L & \text{if } S > 0 \\ u_R & \text{if } S < 0 \end{cases}$$

CFD

$u_L \leq u_R$ **Rarefaction**



CFD

$u_L \leq u_R$ **Rarefaction**

$$u\left(\frac{x}{t}\right) = \begin{cases} u_L & \text{if } x/t \leq u_L \\ x/t & \text{if } u_L < x/t < u_R \\ u_R & \text{if } u_R \leq x/t \end{cases}$$

$$\begin{cases} u_L & \text{if } 0 \leq u_L \\ 0 & \text{if } u_L < 0 < u_R \\ u_R & \text{if } u_R \leq 0 \end{cases}$$

CFD

Final Projects


CFD



Kelvin wake

CFD

Final Projects



Vortices interactions

CFD


Final Projects



Thermal convection

CFD

Final Projects



Thermo-haline convection



CFD

Final Projects




Avalanche model



CFD

Final Projects





Propulsion



CFD

Final Projects



Tennis/Football lift, Magnus effect

