

NUMERICAL METHODS FOR FLUID DYNAMICS



EMMANUEL DORMY (ENS)
dormy@dma.ens.fr

CFD

9. Flows in
Complex Domains


CFD

Most industrial applications of computational fluid dynamics take place in complex geometries.




CFD

Most industrial applications of computational fluid dynamics take place in complex geometries.




CFD

Most industrial applications of computational fluid dynamics take place in complex geometries.



CFD


Most industrial applications of computational fluid dynamics take place in complex geometries.



CFD

Two different approaches:

- **Unstructured grids**




CFD

Two different approaches:

- **Domain cutting**

cut region from a Cartesian grid



CFD

Variational form

$$\begin{cases} \Delta u = -f & \text{in } \Omega, \\ u = 0 & \text{on } \partial\Omega. \end{cases}$$

CFD

Variational form

$$\begin{cases} \Delta u = -f & \text{in } \Omega, \\ u = 0 & \text{on } \partial\Omega. \end{cases}$$

$$H^1(\Omega) = \left\{ w \in L^2(\Omega), \frac{\partial w}{\partial x}, \frac{\partial w}{\partial y} \in L^2(\Omega) \right\}.$$

CFD

Variational form

Find $u \in H_0^1(\Omega)$

such that $\int_{\Omega} \Delta u w \, d\mathbf{x} = - \int_{\Omega} f w \, d\mathbf{x},$

$\forall w \in H_0^1(\Omega)$

CFD

Variational form

Find $u \in H_0^1(\Omega)$

such that $\int_{\Omega} \nabla u \cdot \nabla w \, d\mathbf{x} = \int_{\Omega} f w \, d\mathbf{x},$

$\forall w \in H_0^1(\Omega)$

CFD

A simple 1D periodic example

$$\frac{\partial^2 u}{\partial x^2} = -f$$

CFD

A simple 1D periodic example

$$\frac{\partial^2 u}{\partial x^2} = -f$$

$$\int_{\Omega} \frac{\partial^2 u}{\partial x^2} w \, dx = - \int_{\Omega} f w \, dx \quad \forall w \in H^1(\Omega)$$

CFD

A simple 1D periodic example

$$\frac{\partial^2 u}{\partial x^2} = -f$$

$$\int_{\Omega} \frac{\partial^2 u}{\partial x^2} w \, dx = - \int_{\Omega} f w \, dx \quad \forall w \in H^1(\Omega)$$

$$\int_{\Omega} \frac{\partial u}{\partial x} \frac{\partial w}{\partial x} \, dx = \int_{\Omega} f w \, dx \quad \forall w \in H^1(\Omega)$$

CFD

Finite Element « shape functions »

$$\phi_i(x) = \begin{cases} 0 & \text{if } x \leq x_{i-1}, \\ \frac{x - x_{i-1}}{x_i - x_{i-1}} & \text{if } x_{i-1} < x \leq x_i, \\ \frac{x_{i+1} - x}{x_{i+1} - x_i} & \text{if } x_i < x \leq x_{i+1}, \\ 0 & \text{if } x_{i+1} < x. \end{cases}$$

CFD

Finite Element « shape functions »

$$u(x) = \sum_{i=1}^N u_i \phi_i(x),$$

$$f(x) = \sum_{i=1}^N f_i \phi_i(x),$$

$$w(x) = \sum_{i=1}^N w_i \phi_i(x).$$

CFD

Discrete problem

$$\int_{\Omega} \frac{\partial u}{\partial x} \frac{\partial w}{\partial x} \, dx = \int_{\Omega} f w \, dx \quad \forall w \in H^1(\Omega)$$

$$\int_{\Omega} \sum_i u_i \frac{\partial \phi_i}{\partial x} \frac{\partial \phi_j}{\partial x} \, dx = \int_{\Omega} \sum_i f_i \phi_i \phi_j \, dx, \quad \forall j \in [1, N].$$

CFD

$$\int_{\Omega} \sum_i u_i \frac{\partial \phi_i}{\partial x} \frac{\partial \phi_j}{\partial x} \, \mathrm{d}x = \int_{\Omega} \sum_i f_i \phi_i \phi_j \, \mathrm{d}x \,, \qquad \forall j \in [1,\, N] \,.$$

CFD

$$\int_{\Omega} \sum_i u_i \frac{\partial \phi_i}{\partial x} \frac{\partial \phi_j}{\partial x} \, \mathrm{d}x = \int_{\Omega} \sum_i f_i \phi_i \phi_j \, \mathrm{d}x \,, \qquad \forall j \in [1,\, N] \,.$$

$$\int_{\Omega} \frac{\partial \phi_i}{\partial x} \frac{\partial \phi_j}{\partial x} \, \mathrm{d}x = \left\{ \begin{array}{ll} 2/h & \text{if } i=j \,, \\ -1/h & \text{if } |i-j|=1 \,, \\ 0 & \text{otherwise}\,. \end{array} \right.$$

CFD

$$\int_{\Omega} \sum_i u_i \frac{\partial \phi_i}{\partial x} \frac{\partial \phi_j}{\partial x} \, dx = \int_{\Omega} \sum_i f_i \phi_i \phi_j \, dx , \quad \forall j \in [1, N] .$$

$$\int_{\Omega} \frac{\partial \phi_i}{\partial x} \frac{\partial \phi_j}{\partial x} \, dx = \begin{cases} \quad 2/h & \text{if } i = j , \\ -1/h & \text{if } |i - j| = 1 , \\ \quad 0 & \text{otherwise .} \end{cases}$$


$$\int_{\Omega} \phi_i \phi_j \, dx = \begin{cases} \quad 2h/3 & \text{if } i = j , \\ h/6 & \text{if } |i - j| = 1 , \\ \quad 0 & \text{otherwise .} \end{cases}$$

CFD

$$\frac{1}{h} \begin{pmatrix} 2 & -1 & & & -1 \\ -1 & 2 & -1 & & \\ & \ddots & \ddots & \ddots & \\ & & \ddots & \ddots & \ddots \\ & & & \ddots & \ddots & -1 \\ -1 & & & & -1 & 2 \end{pmatrix} \begin{pmatrix} u_1 \\ \vdots \\ \vdots \\ \vdots \\ \vdots \\ u_N \end{pmatrix} = h \begin{pmatrix} 2/3 & 1/6 & & & 1/6 \\ 1/6 & 2/3 & 1/6 & & \\ & \ddots & \ddots & \ddots & \\ & & \ddots & \ddots & \ddots \\ & & & \ddots & \ddots & 1/6 \\ 1/6 & & & & 1/6 & 2/3 \end{pmatrix} \begin{pmatrix} f_1 \\ \vdots \\ \vdots \\ \vdots \\ \vdots \\ f_N \end{pmatrix}$$


CFD

Shape functions




CFD

Shape functions in 2D, P1




CFD

Shape functions in 2D, P1




CFD

Shape functions in 2D, P2




CFD

Mesh generation



CFD

Mesh generation




Voronoi



Georgi Fedoseevich
Voronoi
(1868-1908)

CFD

Mesh generation




Boris Delaunay
(1890-1980)

Delaunay - Voronoï

CFD


Mesh generation



Delaunay - Voronoï

CFD

Integration



CFD

Velocity Pressure

$$\int_{\Omega} [\nabla \mathbf{u} : \nabla \mathbf{w}](\mathbf{x}) \, d\mathbf{x} + \int_{\Omega} [\mathbf{w} \cdot \nabla p](\mathbf{x}) \, d\mathbf{x} = 0 \quad \forall \mathbf{w} \in [H_0^1(\Omega)]^2 ,$$
$$\int_{\Omega} [q \, \nabla \cdot \mathbf{u}](\mathbf{x}) \, d\mathbf{x} = 0 \quad \forall q \in L^2(\Omega) .$$

CFD

Velocity Pressure

$$\begin{aligned} \int_{\Omega} [\nabla \mathbf{u} : \nabla \mathbf{w}](\mathbf{x}) \, d\mathbf{x} + \int_{\Omega} [\mathbf{w} \cdot \nabla p](\mathbf{x}) \, d\mathbf{x} &= 0 \quad \forall \mathbf{w} \in [H_0^1(\Omega)]^2, \\ \int_{\Omega} [q \, \nabla \cdot \mathbf{u}](\mathbf{x}) \, d\mathbf{x} &= 0 \quad \forall q \in L^2(\Omega). \end{aligned}$$

$$\begin{aligned} \int_{\Omega_h} [\nabla \mathbf{u}_h : \nabla \mathbf{w}_h](\mathbf{x}) \, d\mathbf{x} + \int_{\Omega_h} [\mathbf{w}_h \cdot \nabla p_h](\mathbf{x}) \, d\mathbf{x} &= 0 \quad \forall \mathbf{w}_h \in V_{0h}, \\ \int_{\Omega_h} [q_h \, \nabla \cdot \mathbf{u}_h](\mathbf{x}) \, d\mathbf{x} &= 0 \quad \forall q_h \in Q_h. \end{aligned}$$

CFD

Ladyzenskaya-Brezzi-Babuska

$$\inf_{q_h \in Q_h} \sup_{\mathbf{u}_h \in V_{0h}} \frac{1}{\|\mathbf{u}_h\|_{V_{0h}} \|q_h\|_{Q_h}} \int_{\Omega_h} [q_h \, \nabla \cdot \mathbf{u}_h](\mathbf{x}) \, d\mathbf{x} \geq C > 0.$$

CFD

Ladyzenskaya-Brezzi-Babuska (LBB or Inf-Sup condition)

$$\inf_{q_h \in Q_h} \sup_{\mathbf{u}_h \in V_{0h}} \frac{1}{\|\mathbf{u}_h\|_{V_{0h}} \|q_h\|_{Q_h}} \int_{\Omega_h} [q_h \nabla \cdot \mathbf{u}_h](\mathbf{x}) d\mathbf{x} \geq C > 0.$$


CFD

Freefem++

<https://freefem.org>


CFD

To explore further:




FINITE ELEMENT ANALYSIS

FROM CONCEPTS TO APPLICATIONS · DAVID S. BURNETT



CFD

Adaptative stencils



CFD

Adaptive stencils


Very intuitive idea: adjust the finite-difference discretization near the boundary and include the boundary values directly.

- Shortley-Weller technique
- ghost-point reflection

CFD

Adaptive stencils

Shortley-Weller technique



$$u_W \frac{2}{\Delta x(\Delta_E + \Delta x)} + u_N \frac{1}{\Delta y^2} + u_S \frac{1}{\Delta y^2} -$$


$$u_C \left(\frac{2}{\Delta_E \Delta x} + \frac{2}{\Delta y^2} \right) = f_C - U \frac{2}{\Delta_E (\Delta_E + \Delta x)}$$

CFD

Adaptive stencils

ghost-point reflection

reflection along the boundary normal



bilinear interpolation from neighboring points to mapped point

ghost value = +/- interpolated mapped value

CFD


Adaptive stencils

Advantage: ease of implementation

Disadvantage: uniform stencil structure is compromised;
convergence issues; algorithmic issues;

CFD

Penalisation



CFD

Penalisation

$$\nabla \cdot \mathbf{u} = 0$$

$$\frac{\partial \mathbf{u}}{\partial t} + (\mathbf{u} \cdot \nabla) \mathbf{u} = -\nabla p + \frac{1}{Re} \Delta \mathbf{u} + \mathbf{f}$$

CFD

Penalisation

