


Turbulence Equations and Conserved Quantities



Alexandros ALEXAKIS

alexakis@phys.ens.fr

Dep. Physique ENS Ulm

The Incompressible Navier Stokes equations



Claude-Louis Navier
1785-1836



Sir George Stokes
1819-1903



Isaac Newton
1642-1727



Leonhard Euler
1707-1783

$$\partial_t \mathbf{u} + \mathbf{u} \cdot \nabla \mathbf{u} = -\nabla P + \nu \nabla^2 \mathbf{u} + \mathbf{f}$$

Incompressibility condition

$$\nabla \cdot \mathbf{u} = 0$$

$$\nu = \mu / \rho \equiv \text{kinematic viscosity}$$

Navier Stokes related equations



Claude-Louis Navier
1785-1836



Sir George Stokes
1819-1903



Isaac Newton
1642-1727



Leonhard Euler
1707-1783

Plus advection of a passive scalar

$$\partial_t \mathbf{u} + \mathbf{u} \cdot \nabla \mathbf{u} = -\nabla P + \nu \nabla^2 \mathbf{u} + \mathbf{f}$$

$$\partial_t \phi + \mathbf{u} \cdot \nabla \phi = \kappa \nabla^2 \phi + S$$

In a rotating reference frame

$$\partial_t \mathbf{u} + \mathbf{u} \cdot \nabla \mathbf{u} + 2\boldsymbol{\Omega} \times \mathbf{u} = -\nabla P + \nu \nabla^2 \mathbf{u} + \mathbf{f}$$

In uniformly Stratified medium

$$\partial_t \mathbf{u} + \mathbf{u} \cdot \nabla \mathbf{u} = -\nabla P + \nu \nabla^2 \mathbf{u} - \alpha g \boldsymbol{\theta} + \mathbf{f}$$

$$\partial_t \theta + \mathbf{u} \cdot \nabla \theta = \kappa \nabla^2 \theta + \mathbf{e}_z S u_z$$

Breaking down the Navier Stokes equations



Claude-Louis Navier
1785-1836



Sir George Stokes
1819-1903

Navier Stokes equations

$$\partial_t \mathbf{u} + \mathbf{u} \cdot \nabla \mathbf{u} = -\nabla P + \nu \nabla^2 \mathbf{u} + \mathbf{f}$$

$$\nabla \cdot \mathbf{u} = 0$$

- time derivative
- **Non-linear terms**: advection and pressure
- **Linear terms**: viscous terms responsible for energy dissipation
- **Non-homogeneous terms**: Forcing - energy injection
- **Divergence-free constraint**

Breaking down the Navier Stokes equations



Leonhard Euler
1707-1783

Euler Equations

$$\partial_t \mathbf{u} + \mathbf{u} \cdot \nabla \mathbf{u} = -\nabla P$$

$$\nabla \cdot \mathbf{u} = 0$$

- time derivative
- **Non-linear terms**: advection and pressure
- **Linear terms**: viscous terms responsible for energy dissipation
- **Non-homogeneous terms**: Forcing - energy injection
- **Divergence-free constraint**

Breaking down the Navier Stokes equations



Sir George Stokes
1819-1903

Stokes Equations

$$\partial_t \mathbf{u} = -\nabla \pi + \nu \nabla^2 \mathbf{u} + \mathbf{f}$$

$$\nabla \cdot \mathbf{u} = 0$$

- time derivative
- **Non-linear terms**: advection and pressure
- **Linear terms**: viscous terms responsible for energy dissipation
- **Non-homogeneous terms**: Forcing - energy injection
- **Divergence-free constraint**

The Navier-Stokes

$$\partial_t \mathbf{u} + \mathbf{u} \cdot \nabla \mathbf{u} = -\nabla P + \nu \nabla^2 \mathbf{u} + \mathbf{f}$$

$$\nabla \cdot \mathbf{u} = 0$$

The pressure

$$\partial_t \mathbf{u} + \mathbf{u} \cdot \nabla \mathbf{u} = -\nabla P + \nu \nabla^2 \mathbf{u} + \mathbf{f}$$

Take the divergence ($\nabla \cdot$) of the Navier Stokes equation

$$\partial_t (\nabla \cdot \mathbf{u}) + \nabla \cdot (\mathbf{u} \cdot \nabla \mathbf{u}) = -\nabla^2 P + \nu \nabla^2 (\nabla \cdot \mathbf{u}) + \nabla \cdot \mathbf{f}$$

by incompressibility

$$\cancel{\partial_t (\nabla \cdot \mathbf{u})} + \nabla \cdot (\mathbf{u} \cdot \nabla \mathbf{u}) = -\nabla^2 P + \cancel{\nu \nabla^2 (\nabla \cdot \mathbf{u})} + \cancel{\nabla \cdot \mathbf{f}}$$

$$\boxed{\nabla^2 P = -\nabla \cdot (\mathbf{u} \cdot \nabla \mathbf{u})}$$

To find the pressure we have to invert a laplacian

Alternative forms of the Navier Stokes equations

$$\partial_t \mathbf{u} + \mathbf{u} \cdot \nabla \mathbf{u} = -\nabla P + \nu \nabla^2 \mathbf{u} + \mathbf{f}$$

using the vector identity

$$\mathbf{A} \times (\nabla \times \mathbf{A}) = -\mathbf{A} \cdot \nabla \mathbf{A} + \frac{1}{2} \nabla (\mathbf{A} \cdot \mathbf{A})$$

$$\partial_t \mathbf{u} = \mathbf{u} \times \mathbf{w} - \nabla P' + \nu \nabla^2 \mathbf{u} + \mathbf{f}$$

where $\mathbf{w} = \nabla \times \mathbf{u}$ and $P' = P + \frac{1}{2} |\mathbf{u}|^2$

vorticity equation

$$\partial_t \mathbf{w} = \nabla \times (\mathbf{u} \times \mathbf{w}) + \nu \nabla^2 \mathbf{w} + \nabla \times \mathbf{f}$$


where $\nabla \times \mathbf{w} = \nabla \times \nabla \times \mathbf{u} = \nabla (\nabla \cdot \mathbf{u}) - \nabla^2 \mathbf{u} = -\nabla^2 \mathbf{u}$

Alternative forms of the Navier Stokes equations


vorticity equation

$$\partial_t \mathbf{w} = \nabla \times (\mathbf{u} \times \mathbf{w}) + \nu \nabla^2 \mathbf{w} + \nabla \times \mathbf{f}$$

Vorticity field lines move with the flow




Domain \mathcal{D}



For 99,99% of the course $L_x = L_y = L_z = L$.

In many cases we want $L \rightarrow \infty$

Boundary Conditions on $\partial\mathcal{D}$



- No-slip

$$\mathbf{u} = 0$$


- Free slip

$$\mathbf{u} \cdot \mathbf{e}_n = 0 \quad \& \quad \mathbf{w} \times \mathbf{e}_n = 0$$

- Periodic

$$\mathbf{u}(x, y, z, t) = \mathbf{u}(x + n_x L_x, y + n_y L_y, z + n_z L_z, t)$$

Fourier Space (Finite box)



$$\mathbf{u}(\mathbf{x}, t) = \sum \tilde{\mathbf{u}}_{\mathbf{k}}(t) e^{i\mathbf{k} \cdot \mathbf{x}}, \quad \tilde{\mathbf{u}}_{\mathbf{k}}(t) = \langle \mathbf{u} e^{-i\mathbf{k} \cdot \mathbf{x}} \rangle$$

where

$$\langle f(\mathbf{x}) \rangle = \frac{1}{L_x L_y L_z} \int f(\mathbf{x}) dV$$

and

$$\mathbf{k} = \left(\frac{2\pi n_x}{L_x}, \frac{2\pi n_y}{L_y}, \frac{2\pi n_z}{L_z} \right), \quad \text{with } n_x, n_y, n_z \in \mathbb{Z}$$

Fourier Space (finite and infinite space)

$$\tilde{\mathbf{u}}(\mathbf{k}, t) = \frac{1}{L^3} \int \mathbf{u} e^{-i\mathbf{k} \cdot \mathbf{x}} dV,$$

$$\tilde{\mathbf{u}}(\mathbf{k}, t) = \frac{1}{(2\pi)^3} \int \mathbf{u} e^{-i\mathbf{k} \cdot \mathbf{x}} dV$$

$$\mathbf{u}(\mathbf{x}, t) = \sum_{\mathbf{k} \in \mathbb{N}^3} \tilde{\mathbf{u}}(\mathbf{k}, t) e^{i\mathbf{k} \cdot \mathbf{x}},$$

$$\mathbf{u}(\mathbf{x}, t) = \int_{\mathbf{k} \in \mathbb{R}^3} \tilde{\mathbf{u}}(\mathbf{k}, t) e^{i\mathbf{k} \cdot \mathbf{x}} dk^3$$

Energy Spectrum:

- Finite Box

$$E(k) = \frac{L}{2} \sum_{k \leq |\mathbf{k}| < k+1} |\tilde{\mathbf{u}}_{\mathbf{k}}|^2$$

- Infinite space

$$E(k) = \frac{1}{2} \int |\tilde{\mathbf{u}}|^2 \delta(k - |\mathbf{k}|) dk^3$$

Navier Stokes in Fourier Space

$$\left\langle (\partial_t \mathbf{u} + \mathbf{u} \cdot \nabla \mathbf{u}) e^{-i\mathbf{k} \cdot \mathbf{x}} \right\rangle = \left\langle (-\nabla P + \nu \nabla^2 \mathbf{u} + \mathbf{f}) e^{-i\mathbf{k} \cdot \mathbf{x}} \right\rangle$$

$$\partial_t \tilde{\mathbf{u}}_{\mathbf{k}} + \left\langle (\mathbf{u} \cdot \nabla \mathbf{u}) e^{-i\mathbf{k} \cdot \mathbf{x}} \right\rangle = -\nabla \tilde{P}_{\mathbf{k}} - \nu |\mathbf{k}|^2 \tilde{\mathbf{u}}_{\mathbf{k}} + \tilde{\mathbf{f}}_{\mathbf{k}}$$

Navier Stokes in Fourier Space

$$\left\langle (\partial_t \mathbf{u} + \mathbf{u} \cdot \nabla \mathbf{u}) e^{-i\mathbf{k} \cdot \mathbf{x}} \right\rangle = \left\langle (-\nabla P + \nu \nabla^2 \mathbf{u} + \mathbf{f}) e^{-i\mathbf{k} \cdot \mathbf{x}} \right\rangle$$

$$\partial_t \tilde{\mathbf{u}}_{\mathbf{k}} + \left\langle (\mathbf{u} \cdot \nabla \mathbf{u}) e^{-i\mathbf{k} \cdot \mathbf{x}} \right\rangle = -\nabla \tilde{P}_{\mathbf{k}} - \nu |\mathbf{k}|^2 \tilde{\mathbf{u}}_{\mathbf{k}} + \tilde{\mathbf{f}}_{\mathbf{k}}$$

$$\begin{aligned} \left\langle (\mathbf{u} \cdot \nabla \mathbf{u}) e^{-i\mathbf{k} \cdot \mathbf{x}} \right\rangle &= \left\langle \left(\left[\sum_{\mathbf{q}} \tilde{\mathbf{u}}_{\mathbf{q}} e^{i\mathbf{q} \cdot \mathbf{x}} \right] \cdot \nabla \left[\sum_{\mathbf{p}} \tilde{\mathbf{u}}_{\mathbf{p}} e^{i\mathbf{p} \cdot \mathbf{x}} \right] \right) e^{-i\mathbf{k} \cdot \mathbf{x}} \right\rangle \\ &= \left\langle \left(\sum_{\mathbf{p}, \mathbf{q}} \tilde{\mathbf{u}}_{\mathbf{q}} \cdot (i\mathbf{p}) \tilde{\mathbf{u}}_{\mathbf{p}} \right) e^{i(\mathbf{q} + \mathbf{p} - \mathbf{k}) \cdot \mathbf{x}} \right\rangle \\ &= \sum_{\mathbf{p} + \mathbf{q} = \mathbf{k}} i\mathbf{p} \cdot \tilde{\mathbf{u}}_{\mathbf{q}} \tilde{\mathbf{u}}_{\mathbf{p}} \end{aligned}$$

Navier Stokes in Fourier Space: Pressure term

$$\left\langle e^{-i\mathbf{k}\cdot\mathbf{x}} \nabla \cdot (\mathbf{u} \cdot \nabla \mathbf{u}) \right\rangle = \left\langle -(\nabla^2 P) e^{-i\mathbf{k}\cdot\mathbf{x}} \right\rangle$$

$$i\mathbf{k} \cdot \sum_{\mathbf{p}+\mathbf{q}=\mathbf{k}} i(\mathbf{p} \cdot \tilde{\mathbf{u}}_{\mathbf{q}}) \tilde{\mathbf{u}}_{\mathbf{p}} = |\mathbf{k}|^2 \tilde{P}_{\mathbf{k}}$$

$$\tilde{P}_{\mathbf{k}} = - \sum_{\mathbf{p}+\mathbf{q}=\mathbf{k}} \frac{(\mathbf{k} \cdot \tilde{\mathbf{u}}_{\mathbf{p}})(\mathbf{p} \cdot \tilde{\mathbf{u}}_{\mathbf{q}})}{|\mathbf{k}|^2}$$

Navier Stokes:

$$\partial_t \tilde{\mathbf{u}}_{\mathbf{k}} = - \sum_{\mathbf{p}+\mathbf{q}=\mathbf{k}} i \left((\mathbf{p} \cdot \tilde{\mathbf{u}}_{\mathbf{q}}) \tilde{\mathbf{u}}_{\mathbf{p}} - \mathbf{k} \frac{(\mathbf{k} \cdot \tilde{\mathbf{u}}_{\mathbf{p}})(\mathbf{p} \cdot \tilde{\mathbf{u}}_{\mathbf{q}})}{|\mathbf{k}|^2} \right) - \nu |\mathbf{k}|^2 \tilde{\mathbf{u}}_{\mathbf{k}} + \tilde{\mathbf{f}}_{\mathbf{k}}$$

and

$$\mathbf{k} \cdot \mathbf{u}_{\mathbf{k}} = 0$$

Volume averaged quantities

$$\langle f(\mathbf{x}) \rangle = \frac{1}{L_x L_y L_z} \int f(\mathbf{x}) dV$$

Identities

$$\langle \nabla \cdot \mathbf{a} \rangle = \frac{1}{V} \int_{\partial \mathcal{D}} (\mathbf{a} \cdot \mathbf{n}) d\mathcal{S},$$

$$\langle \nabla \times \mathbf{a} \rangle = \frac{1}{V} \iiint_{\partial \mathcal{A}} \nabla \times \mathbf{a} d\mathcal{A} dz = \frac{1}{V} \int \oint_{\partial \mathcal{C}} \mathbf{a} \cdot d\ell dz,$$

In periodic domains this implies

$$\langle \nabla \cdot \mathbf{a} \rangle = 0$$

$$\langle \nabla \times \mathbf{a} \rangle = 0$$

Volume averaged quantities

in periodic domains we thus have

- $\langle \partial_i f \rangle = 0$
- $\langle (\nabla f)g \rangle = \langle \nabla(fg) \rangle - \langle f(\nabla g) \rangle = -\langle f(\nabla g) \rangle$
- $\langle (\nabla^2 f)g \rangle = \langle \nabla(g\nabla f) \rangle - \langle (\nabla f)(\nabla g) \rangle = -\langle (\nabla f)(\nabla g) \rangle$
- $\langle \mathbf{a} \cdot \nabla \mathbf{b} \rangle = \langle \nabla_i(a_i b_j) \rangle - \langle \mathbf{b} \nabla \cdot \mathbf{a} \rangle = -\langle \mathbf{b} \nabla \cdot \mathbf{a} \rangle$
 $= 0 \quad \text{if} \quad \nabla \cdot \mathbf{a} = 0$
- $\langle \mathbf{a} \cdot \nabla \times \mathbf{b} \rangle = \langle \nabla \cdot (\mathbf{b} \times \mathbf{a}) \rangle + \langle \mathbf{b} \cdot \nabla \times \mathbf{a} \rangle = \langle \mathbf{b} \cdot \nabla \times \mathbf{a} \rangle$
- $\langle \mathbf{a} \cdot \nabla^2 \mathbf{b} \rangle = -\langle \mathbf{a} \cdot \nabla \times \nabla \times \mathbf{b} \rangle = -\langle (\nabla \times \mathbf{b}) \cdot (\nabla \times \mathbf{a}) \rangle$

Conservation laws: Momentum

$$\partial_t \mathbf{u} + \mathbf{u} \cdot \nabla \mathbf{u} = -\nabla P + \nu \nabla^2 \mathbf{u} + \mathbf{f}$$

$$\partial_t \mathbf{u} + \mathbf{u} \cdot \nabla \mathbf{u} = -\nabla P + \nu \nabla^2 \mathbf{u} + \mathbf{f}$$

$$\partial_t \langle \mathbf{u} \rangle + \langle \mathbf{u} \cdot \nabla \mathbf{u} \rangle = -\langle \nabla P \rangle + \nu \langle \nabla^2 \mathbf{u} \rangle + \langle \mathbf{f} \rangle$$

$$\partial_t \langle \mathbf{u} \rangle + \langle \nabla_i (u_i u_j) \rangle - \langle \mathbf{u} \nabla \cdot \mathbf{u} \rangle = -\langle \nabla P \rangle + \nu \langle \nabla^2 \mathbf{u} \rangle + \langle \mathbf{f} \rangle$$

$$\partial_t \langle \mathbf{u} \rangle + \cancel{\langle \nabla_i (u_i u_j) \rangle} - \cancel{\langle \mathbf{u} \nabla \cdot \mathbf{u} \rangle} = -\cancel{\langle \nabla P \rangle} + \nu \cancel{\langle \nabla^2 \mathbf{u} \rangle} + \cancel{\langle \mathbf{f} \rangle}$$

$$\partial_t \langle \mathbf{u} \rangle = 0$$

(here $\langle \mathbf{f} \rangle = 0$ is assumed)

Conservation laws: Energy

$$\partial_t \mathbf{u} + \mathbf{u} \cdot \nabla \mathbf{u} = -\nabla P + \nu \nabla^2 \mathbf{u} + \mathbf{f}$$

$$\langle \mathbf{u} \cdot \partial_t \mathbf{u} \rangle + \langle \mathbf{u} \cdot (\mathbf{u} \cdot \nabla \mathbf{u}) \rangle = -\langle \mathbf{u} \cdot \nabla P \rangle + \nu \langle \mathbf{u} \cdot \nabla^2 \mathbf{u} \rangle + \langle \mathbf{u} \cdot \mathbf{f} \rangle$$

$$\frac{1}{2} \frac{d}{dt} \langle |\mathbf{u}|^2 \rangle + \frac{1}{2} \langle (\mathbf{u} \cdot \nabla |\mathbf{u}|^2) \rangle = -\langle \mathbf{u} \cdot \nabla P \rangle + \nu \langle \mathbf{u} \cdot \nabla^2 \mathbf{u} \rangle + \langle \mathbf{u} \cdot \mathbf{f} \rangle$$

$$\begin{aligned} \frac{1}{2} \frac{d}{dt} \langle |\mathbf{u}|^2 \rangle + \frac{1}{2} \langle (\nabla_i u_i |\mathbf{u}|^2) \rangle &= -\langle \nabla \cdot (\mathbf{u} P) \rangle + \langle P \nabla \cdot \mathbf{u} \rangle \\ &\quad + \nu \langle \nabla_i (u_j \nabla_i u_j) \rangle - \nu \langle \nabla_j u_i \nabla_j u_i \rangle \\ &\quad + \langle \mathbf{u} \cdot \mathbf{f} \rangle \end{aligned}$$

Conservation laws: Energy

$$\partial_t \mathbf{u} + \mathbf{u} \cdot \nabla \mathbf{u} = -\nabla P + \nu \nabla^2 \mathbf{u} + \mathbf{f}$$

$$\langle \mathbf{u} \cdot \partial_t \mathbf{u} \rangle + \langle \mathbf{u} \cdot (\mathbf{u} \cdot \nabla \mathbf{u}) \rangle = -\langle \mathbf{u} \cdot \nabla P \rangle + \nu \langle \mathbf{u} \cdot \nabla^2 \mathbf{u} \rangle + \langle \mathbf{u} \cdot \mathbf{f} \rangle$$

$$\frac{1}{2} \frac{d}{dt} \langle |\mathbf{u}|^2 \rangle + \frac{1}{2} \langle (\mathbf{u} \cdot \nabla |\mathbf{u}|^2) \rangle = -\langle \mathbf{u} \cdot \nabla P \rangle + \nu \langle \mathbf{u} \cdot \nabla^2 \mathbf{u} \rangle + \langle \mathbf{u} \cdot \mathbf{f} \rangle$$

$$\begin{aligned} \frac{1}{2} \frac{d}{dt} \langle |\mathbf{u}|^2 \rangle + \frac{1}{2} \langle \cancel{(\nabla_i u_i |\mathbf{u}|^2)} \rangle &= -\langle \cancel{\nabla \cdot (\mathbf{u} P)} \rangle + \langle P \cancel{\nabla \cdot \mathbf{u}} \rangle \\ &\quad + \nu \langle \cancel{\nabla_i (u_j \nabla_i u_j)} \rangle - \nu \langle \nabla_j u_i \nabla_j u_i \rangle \\ &\quad + \langle \mathbf{u} \cdot \mathbf{f} \rangle \end{aligned}$$

$$\frac{1}{2} \frac{d}{dt} \langle |\mathbf{u}|^2 \rangle = -\nu \langle \nabla_j u_i \nabla_j u_i \rangle + \langle \mathbf{u} \cdot \mathbf{f} \rangle$$

$$\boxed{\frac{d}{dt} \mathcal{E} = -\epsilon + \mathcal{I}}$$

Energy Dissipation rate ϵ

Let $\mathcal{S}_{ij} = (\partial_i u_j + \partial_j u_i)/2$ and $\Omega_{ij} = (\partial_i u_j - \partial_j u_i)/2$

The local energy dissipation rate is then given by

$$2\nu \mathcal{S}_{i,j} \mathcal{S}_{i,j}$$

In **periodic** domains we have

$$\begin{aligned}\epsilon &= 2\nu \langle (\partial_i u_j + \partial_j u_i)(\partial_i u_j + \partial_j u_i) \rangle / 4 \\ &= \nu \langle (\partial_i u_j)^2 + 2\partial_j u_i \partial_i u_j + (\partial_j u_i)^2 \rangle / 2 \\ &= \nu \langle (\partial_i u_j)^2 \rangle + \nu \langle \partial_j (u_i \partial_i u_j) \rangle \\ &= \nu \langle (\partial_i u_j)^2 \rangle + \cancel{\nu \langle \partial_j (u_i \partial_i u_j) \rangle} \\ &= \boxed{\nu \langle (\partial_i u_j)^2 \rangle} \\ &= \nu \langle (\partial_i u_j)^2 - 2\partial_j u_i \partial_i u_j + (\partial_j u_i)^2 \rangle / 2 \\ &= \boxed{2\nu \langle \Omega_{i,j} \Omega_{i,j} \rangle = \nu \langle |\mathbf{w}|^2 \rangle} \quad (1)\end{aligned}$$

Conservation laws: Helicity

$$\mathcal{H} = \frac{1}{2} \langle \mathbf{u} \cdot \mathbf{w} \rangle$$

$$\partial_t \mathbf{u} = \mathbf{u} \times \mathbf{w} - \nabla P + \nu \nabla^2 \mathbf{u} + \mathbf{f}$$

$$\partial_t \mathbf{w} = \nabla \times (\mathbf{u} \times \mathbf{w}) + \nu \nabla^2 \mathbf{w} + \nabla \times \mathbf{f}$$

$$\begin{aligned} \frac{d}{dt} \langle \mathbf{u} \cdot \mathbf{w} \rangle &= \langle \mathbf{u} \cdot \partial_t \mathbf{w} \rangle + \langle \mathbf{w} \partial_t \mathbf{u} \rangle \\ &= \langle \mathbf{u} \cdot \nabla \times (\mathbf{u} \times \mathbf{w}) \rangle + \nu \langle \mathbf{u} \cdot \nabla^2 \mathbf{w} \rangle + \langle \mathbf{u} \cdot \nabla \times \mathbf{f} \rangle + \\ &\quad \langle \mathbf{w} \cdot (\mathbf{u} \times \mathbf{w}) \rangle + \langle \mathbf{w} \cdot \nabla P \rangle + \langle \mathbf{w} \cdot \nabla^2 \mathbf{u} \rangle + \langle \mathbf{w} \cdot \mathbf{f} \rangle \end{aligned}$$

Conservation laws: Helicity

$$\mathcal{H} = \frac{1}{2} \langle \mathbf{u} \cdot \mathbf{w} \rangle$$

$$\partial_t \mathbf{u} = \mathbf{u} \times \mathbf{w} - \nabla P + \nu \nabla^2 \mathbf{u} + \mathbf{f}$$

$$\partial_t \mathbf{w} = \nabla \times (\mathbf{u} \times \mathbf{w}) + \nu \nabla^2 \mathbf{w} + \nabla \times \mathbf{f}$$


$$\begin{aligned} \frac{d}{dt} \langle \mathbf{u} \cdot \mathbf{w} \rangle &= \langle \mathbf{u} \cdot \partial_t \mathbf{w} \rangle + \langle \mathbf{w} \partial_t \mathbf{u} \rangle \\ &= \cancel{\langle \mathbf{u} \cdot \nabla \times (\mathbf{u} \times \mathbf{w}) \rangle} + \nu \langle \mathbf{u} \cdot \nabla^2 \mathbf{w} \rangle + \langle \mathbf{u} \cdot \nabla \times \mathbf{f} \rangle + \\ &\quad \cancel{\langle \mathbf{w} \cdot (\mathbf{u} \times \mathbf{w}) \rangle} + \cancel{\langle \mathbf{w} \cdot \nabla P \rangle} + \nu \langle \mathbf{w} \cdot \nabla^2 \mathbf{u} \rangle + \langle \mathbf{w} \cdot \mathbf{f} \rangle \end{aligned}$$

$$\frac{1}{2} \frac{d}{dt} \langle \mathbf{u} \cdot \mathbf{w} \rangle = -\nu \langle \mathbf{w} \cdot \nabla \times \mathbf{w} \rangle + \langle \mathbf{w} \cdot \mathbf{f} \rangle$$

$$\frac{d}{dt} \mathcal{H} = -\epsilon_{\mathcal{H}} + \mathcal{I}_{\mathcal{H}}$$

Conservation laws: Helicity

$$\mathcal{H} = \frac{1}{2} \langle \mathbf{u} \cdot \mathbf{w} \rangle$$



Consider two infinitesimal vorticity tubes

$$\begin{aligned}\mathcal{H} &= \frac{1}{2} \int_{V_1} \mathbf{u} \cdot \mathbf{w} dx^3 + \frac{1}{2} \int_{V_2} \mathbf{u} \cdot \mathbf{w} dx^3 \\ &= \frac{1}{2} \oint_{C_\infty} \int_{S_1} \mathbf{u} \cdot \mathbf{w} ds dl + \frac{1}{2} \oint_{C_\epsilon} \int_{S_2} \mathbf{u} \cdot \mathbf{w} ds dl \\ &= \frac{1}{2} \Phi_1 \oint_{C_\infty} \mathbf{u} \cdot d\ell + \frac{1}{2} \Phi_2 \oint_{C_\epsilon} \mathbf{u} \cdot d\ell \\ &= \frac{1}{2} \Phi_1 \oint_{A_\infty} \mathbf{w} \cdot d\mathbf{s} + \frac{1}{2} \Phi_2 \oint_{A_\epsilon} \mathbf{w} \cdot d\mathbf{s} \\ &= \Phi_1 \Phi_2\end{aligned}$$



Thank you
for your attention!