

Weighted Chairman Assignment and Flow-Time Scheduling

Victor Reis

joint work with

Siyue Liu

to appear in ITCS 2026

Microsoft Research

University of Washington Theory Seminar

November 18, 2025

Warmup I: discrepancy of two permutations

1 2 3 4 5 6 7

6 7 4 5 2 3 1

Goal: Color numbers red and blue so that $|\text{\#red} - \text{\#blue}| \leq 1$ in all $2n$ prefixes


Warmup I: discrepancy of two permutations

1 2 3 4 5 6 7 8

6 7 4 5 2 3 1 8


Goal: Color numbers red and blue so that $|\text{\#red} - \text{\#blue}| \leq 1$ in all $2n$ prefixes

Warmup I: discrepancy of two permutations




Goal: Color numbers red and blue so that $|\text{\#red} - \text{\#blue}| \leq 1$ in all $2n$ prefixes


Warmup I: discrepancy of two permutations




Goal: Color numbers red and blue so that $|\text{\#red} - \text{\#blue}| \leq 1$ in all $2n$ prefixes




Warmup I: discrepancy of two permutations




Goal: Color numbers red and blue so that $|\text{\#red} - \text{\#blue}| \leq 1$ in all $2n$ prefixes




Warmup I: discrepancy of two permutations




Goal: Color numbers red and blue so that $|\#red - \#blue| \leq 1$ in all $2n$ prefixes



Warmup I: discrepancy of two permutations




Goal: Color numbers red and blue so that $|\#red - \#blue| \leq 1$ in all $2n$ prefixes



Open problem


Given *weights* $w \in [0, 1]^n$ can we get $|w(\text{red}) - w(\text{blue})| \leq O(1)$ in all $2n$ prefixes?

Warmup II: edge orientation




Goal: Orient edges so that $|\#in - \#out| \leq 1$ in all $2|E|$ prefixes

Warmup II: edge orientation




Goal: Orient edges so that $|\#in - \#out| \leq 1$ in all $2|E|$ prefixes

Warmup II: edge orientation




Goal: Orient edges so that $|\#in - \#out| \leq 1$ in all $2|E|$ prefixes

Warmup II: edge orientation




Goal: Orient edges so that $|\#in - \#out| \leq 1$ in all $2|E|$ prefixes

Warmup II: edge orientation




Goal: Orient edges so that $|\#in - \#out| \leq 1$ in all $2|E|$ prefixes

Warmup II: edge orientation




Goal: Orient edges so that $|\#in - \#out| \leq 1$ in all $2|E|$ prefixes

Warmup II: edge orientation




Goal: Orient edges so that $|\#in - \#out| \leq 1$ in all $2|E|$ prefixes

Warmup II: edge orientation




Goal: Orient edges so that $|\#in - \#out| \leq 1$ in all $2|E|$ prefixes

Warmup II: edge orientation




Goal: Orient edges so that $|\#in - \#out| \leq 1$ in all $2|E|$ prefixes

Warmup II: edge orientation




Goal: Orient edges so that $|\#in - \#out| \leq 1$ in all $2|E|$ prefixes

Warmup II: edge orientation




Goal: Orient edges so that $|\#in - \#out| \leq 1$ in all $2|E|$ prefixes

Warmup II: edge orientation




Goal: Orient edges so that $|\#in - \#out| \leq 1$ in all $2|E|$ prefixes

Warmup II: edge orientation




Goal: Orient edges so that $|\#in - \#out| \leq 1$ in all $2|E|$ prefixes

Warmup II: edge orientation




Goal: Orient edges so that $|\text{\#in} - \text{\#out}| \leq 1$ in all $2|E|$ prefixes

Warmup II: edge orientation




Goal: Orient edges so that $|\#in - \#out| \leq 1$ in all $2|E|$ prefixes

Warmup II: edge orientation




Goal: Orient edges so that $|\#in - \#out| \leq 1$ in all $2|E|$ prefixes

Warmup II: edge orientation



Goal: Orient edges so that $|\#in - \#out| \leq 1$ in all $2|E|$ prefixes

Warmup II: edge orientation




Goal: Orient edges so that $|\#in - \#out| \leq 1$ in all $2|E|$ prefixes

Open problem

Given *weights* $w \in [0, 1]^E$ can we get $|w(in) - w(out)| \leq O(1)$ in all $2|E|$ prefixes?


Single-source unsplittable flow (SSUF)

(splittable) flow




Single-source unsplittable flow (SSUF)

(splittable) flow




unsplittable flow




Single-source unsplittable flow (SSUF)

(splittable) flow





unsplittable flow



Theorem (Dinitz, Garg, Goemans, 1999)

Given directed $D = (V, E)$ and flow $x \in \mathbb{R}_{\geq 0}^E$, there exists an *unsplittable* flow $y \in \mathbb{R}_{\geq 0}^E$ such that $y(e) \leq x(e) + d_{\max} \forall e \in E$.

Single-source unsplittable flow (SSUF)





Theorem (Dinitz, Garg, Goemans, 1999)

Given directed $D = (V, E)$ and flow $x \in \mathbb{R}_{\geq 0}^E$, there exists an *unsplittable* flow $y \in \mathbb{R}_{\geq 0}^E$ such that $y(e) \leq x(e) + d_{\max} \forall e \in E$.

Theorem (Morell, Skutella, 2020)

Given directed $D = (V, E)$ and flow $x \in \mathbb{R}_{\geq 0}^E$, there exists an *unsplittable* flow $y \in \mathbb{R}_{\geq 0}^E$ such that $y(e) \geq x(e) - d_{\max} \forall e \in E$.

Single-source unsplittable flow (SSUF)



Theorem (Dinitz, Garg, Goemans, 1999)

Given directed $D = (V, E)$ and flow $x \in \mathbb{R}_{\geq 0}^E$, there exists an *unsplittable* flow $y \in \mathbb{R}_{\geq 0}^E$ such that $y(e) \leq x(e) + d_{\max} \forall e \in E$.

Theorem (Morell, Skutella, 2020)

Given directed $D = (V, E)$ and flow $x \in \mathbb{R}_{\geq 0}^E$, there exists an *unsplittable* flow $y \in \mathbb{R}_{\geq 0}^E$ such that $y(e) \geq x(e) - d_{\max} \forall e \in E$.

Conjecture (Morell, Skutella, 2020)

Given directed $D = (V, E)$ and flow $x \in \mathbb{R}_{\geq 0}^E$, there exists an *unsplittable* flow $y \in \mathbb{R}_{\geq 0}^E$ such that $|x(e) - y(e)| \leq d_{\max} \forall e \in E$.

Morell-Skutella conjecture

Conjecture (Morell, Skutella, 2020)

Given directed $D = (V, E)$ and flow $x \in \mathbb{R}_{\geq 0}^E$, there exists an *unsplittable* flow $y \in \mathbb{R}_{\geq 0}^E$ such that $|x(e) - y(e)| \leq d_{\max} \forall e \in E$.

Morell-Skutella conjecture

Conjecture (Morell, Skutella, 2020)

Given directed $D = (V, E)$ and flow $x \in \mathbb{R}_{\geq 0}^E$, there exists an *unsplittable* flow $y \in \mathbb{R}_{\geq 0}^E$ such that $|x(e) - y(e)| \leq d_{\max} \forall e \in E$.

- Known for bipartite digraphs [Lenstra, Shmoys, Tardos, 1990]

Morell-Skutella conjecture

Conjecture (Morell, Skutella, 2020)

Given directed $D = (V, E)$ and flow $x \in \mathbb{R}_{\geq 0}^E$, there exists an *unsplittable* flow $y \in \mathbb{R}_{\geq 0}^E$ such that $|x(e) - y(e)| \leq d_{\max} \forall e \in E$.

- ▶ Known for bipartite digraphs [Lenstra, Shmoys, Tardos, 1990]
- ▶ Known for series-parallel digraphs [Almoghrahi, Skutella, Warode, 2024]

Morell-Skutella conjecture

Conjecture (Morell, Skutella, 2020)

Given directed $D = (V, E)$ and flow $x \in \mathbb{R}_{\geq 0}^E$, there exists an *unsplittable* flow $y \in \mathbb{R}_{\geq 0}^E$ such that $|x(e) - y(e)| \leq d_{\max} \forall e \in E$.

- ▶ Known for bipartite digraphs [Lenstra, Shmoys, Tardos, 1990]
- ▶ Known for series-parallel digraphs [Almoghrahi, Skutella, Warode, 2024]
- ▶ Known for planar digraphs [Traub, Koch, Zenklusen, 2024]

Morell-Skutella conjecture

Conjecture (Morell, Skutella, 2020)

Given directed $D = (V, E)$ and flow $x \in \mathbb{R}_{\geq 0}^E$, there exists an *unsplittable* flow $y \in \mathbb{R}_{\geq 0}^E$ such that $|x(e) - y(e)| \leq d_{\max} \forall e \in E$.

- ▶ Known for bipartite digraphs [Lenstra, Shmoys, Tardos, 1990]
- ▶ Known for series-parallel digraphs [Almoghrahi, Skutella, Warode, 2024]
- ▶ Known for planar digraphs [Traub, Koch, Zenklusen, 2024]
- ▶ Known when $d_{i+1}/d_i \in \mathbb{N}$ for each $i < n$ [Morell, Skutella, 2020]

Morell-Skutella conjecture

Conjecture (Morell, Skutella, 2020)

Given directed $D = (V, E)$ and flow $x \in \mathbb{R}_{\geq 0}^E$, there exists an *unsplittable* flow $y \in \mathbb{R}_{\geq 0}^E$ such that $|x(e) - y(e)| \leq d_{\max} \forall e \in E$.

- ▶ Known for bipartite digraphs [Lenstra, Shmoys, Tardos, 1990]
- ▶ Known for series-parallel digraphs [Almoghrahi, Skutella, Warode, 2024]
- ▶ Known for planar digraphs [Traub, Koch, Zenklusen, 2024]
- ▶ Known when $d_{i+1}/d_i \in \mathbb{N}$ for each $i < n$ [Morell, Skutella, 2020]
- ▶ Known to imply a *cost-augmented* one-sided bound conjectured by Goemans [Swamy, Traub, Koch, Zenklusen, 2025]

Application in scheduling: makespan minimization

- ▶ Jobs J with processing times d_j , machines M , bipartite graph $D = (M \cup J, E)$

Application in scheduling: makespan minimization

- ▶ Jobs J with processing times d_j , machines M , bipartite graph $D = (M \cup J, E)$
- ▶ Objective: minimize the time to complete all jobs

Application in scheduling: makespan minimization

- ▶ Jobs J with processing times d_j , machines M , bipartite graph $D = (M \cup J, E)$
- ▶ Objective: minimize the time to complete all jobs

min T


$$\text{s.t. } \sum_{i \in M} x_{ij} = 1, \quad \forall j \in J,$$

$$\sum_{j \in J} d_j x_{ij} \leq T, \quad \forall i \in M,$$

$$x \geq 0,$$

$$x_{ij} = 0, \quad \forall (i, j) \notin E.$$

(splittable) flow



Application in scheduling: makespan minimization

- Jobs J with processing times d_j , machines M , bipartite graph $D = (M \cup J, E)$
- Objective: minimize the time to complete all jobs


min T

$$\text{s.t. } \sum_{i \in M} x_{ij} = 1, \quad \forall j \in J,$$

$$\sum_{j \in J} d_j x_{ij} \leq T, \quad \forall i \in M,$$

$$x \geq 0,$$

$$x_{ij} = 0, \quad \forall (i, j) \notin E.$$



$$\sum_{j \in J} d_j y_{ij} \leq \sum_{j \in J} d_j x_{ij} + d_{\max} \leq OPT + OPT = 2OPT$$

Application in scheduling II: flow-time minimization

- ▶ Jobs J with processing times d_j , machines M , bipartite graph $D = (M \cup J, E)$

Application in scheduling II: flow-time minimization

- ▶ Jobs J with processing times d_j , machines M , bipartite graph $D = (M \cup J, E)$
- ▶ Jobs ordered by release time $r_1 \leq r_2 \leq \dots \leq r_n$

Application in scheduling II: flow-time minimization

- ▶ Jobs J with processing times d_j , machines M , bipartite graph $D = (M \cup J, E)$
- ▶ Jobs ordered by release time $r_1 \leq r_2 \leq \dots \leq r_n$
- ▶ Minimize the max *flow-time* $\max_{j \in J} (C_j - r_j)$ where C_j is the completion time

Application in scheduling II: flow-time minimization

- ▶ Jobs J with processing times d_j , machines M , bipartite graph $D = (M \cup J, E)$
- ▶ Jobs ordered by release time $r_1 \leq r_2 \leq \dots \leq r_n$
- ▶ Minimize the max *flow-time* $\max_{j \in J} (C_j - r_j)$ where C_j is the completion time

$$\min T$$

$$\text{s.t. } \sum_{i \in M} x_{ij} = 1, \quad \forall j \in J,$$

$$\sum_{j=a}^b d_j x_{ij} \leq (r_b - r_a) + T, \quad \forall i \in M, 1 \leq a \leq b \leq n,$$

$$x \geq 0, x_{ij} = 0 \quad \forall (i, j) \notin E.$$

Application in scheduling II: flow-time minimization

- ▶ Jobs J with processing times d_j , machines M , bipartite graph $D = (M \cup J, E)$
- ▶ Jobs ordered by release time $r_1 \leq r_2 \leq \dots \leq r_n$
- ▶ Minimize the max *flow-time* $\max_{j \in J} (C_j - r_j)$ where C_j is the completion time

$$\min T$$

$$\text{s.t. } \sum_{i \in M} x_{ij} = 1, \quad \forall j \in J,$$

$$\sum_{j=a}^b d_j x_{ij} \leq (r_b - r_a) + T, \quad \forall i \in M, 1 \leq a \leq b \leq n,$$

$$x \geq 0, x_{ij} = 0 \quad \forall (i, j) \notin E.$$

- ▶ Why is this an LP relaxation? Suppose we have an integral solution x

Application in scheduling II: flow-time minimization

- ▶ Jobs J with processing times d_j , machines M , bipartite graph $D = (M \cup J, E)$
- ▶ Jobs ordered by release time $r_1 \leq r_2 \leq \dots \leq r_n$
- ▶ Minimize the max *flow-time* $\max_{j \in J} (C_j - r_j)$ where C_j is the completion time

$$\min T$$

$$\text{s.t. } \sum_{i \in M} x_{ij} = 1, \quad \forall j \in J,$$

$$\sum_{j=a}^b d_j x_{ij} \leq (r_b - r_a) + T, \quad \forall i \in M, 1 \leq a \leq b \leq n,$$

$$x \geq 0, x_{ij} = 0 \quad \forall (i, j) \notin E.$$

- ▶ Why is this an LP relaxation? Suppose we have an integral solution x
- ▶ Fix job j with $x_{ij} = 1$ for some i and let t be the last time in $[0, r_j]$ when i idle

Application in scheduling II: flow-time minimization

- ▶ Jobs J with processing times d_j , machines M , bipartite graph $D = (M \cup J, E)$
- ▶ Jobs ordered by release time $r_1 \leq r_2 \leq \dots \leq r_n$
- ▶ Minimize the max *flow-time* $\max_{j \in J} (C_j - r_j)$ where C_j is the completion time

$$\min T$$

$$\text{s.t. } \sum_{i \in M} x_{ij} = 1, \quad \forall j \in J,$$

$$\sum_{j=a}^b d_j x_{ij} \leq (r_b - r_a) + T, \quad \forall i \in M, 1 \leq a \leq b \leq n,$$

$$x \geq 0, x_{ij} = 0 \quad \forall (i, j) \notin E.$$

- ▶ Why is this an LP relaxation? Suppose we have an integral solution x
- ▶ Fix job j with $x_{ij} = 1$ for some i and let t be the last time in $[0, r_j]$ when i idle
- ▶ May assume $t = r_a$ for some job $a \leq j$

Application in scheduling II: flow-time minimization

- ▶ Jobs J with processing times d_j , machines M , bipartite graph $D = (M \cup J, E)$
- ▶ Jobs ordered by release time $r_1 \leq r_2 \leq \dots \leq r_n$
- ▶ Minimize the max *flow-time* $\max_{j \in J} (C_j - r_j)$ where C_j is the completion time

$$\min T$$

$$\text{s.t. } \sum_{i \in M} x_{ij} = 1, \quad \forall j \in J,$$

$$\sum_{j=a}^b d_j x_{ij} \leq (r_b - r_a) + T, \quad \forall i \in M, 1 \leq a \leq b \leq n,$$

$$x \geq 0, x_{ij} = 0 \quad \forall (i, j) \notin E.$$

- ▶ Why is this an LP relaxation? Suppose we have an integral solution x
- ▶ Fix job j with $x_{ij} = 1$ for some i and let t be the last time in $[0, r_j]$ when i idle
- ▶ May assume $t = r_a$ for some job $a \leq j$
- ▶ Machine busy in $[r_a, r_j]$ with total volume assigned $\leq r_j - r_a + T$

Application in scheduling II: flow-time minimization

- ▶ Jobs J with processing times d_j , machines M , bipartite graph $D = (M \cup J, E)$
- ▶ Jobs ordered by release time $r_1 \leq r_2 \leq \dots \leq r_n$
- ▶ Minimize the max *flow-time* $\max_{j \in J} (C_j - r_j)$ where C_j is the completion time

$$\min T$$

$$\text{s.t. } \sum_{i \in M} x_{ij} = 1, \quad \forall j \in J,$$

$$\sum_{j=a}^b d_j x_{ij} \leq (r_b - r_a) + T, \quad \forall i \in M, 1 \leq a \leq b \leq n,$$

$$x \geq 0, x_{ij} = 0 \quad \forall (i, j) \notin E.$$

- ▶ Why is this an LP relaxation? Suppose we have an integral solution x
- ▶ Fix job j with $x_{ij} = 1$ for some i and let t be the last time in $[0, r_j]$ when i idle
- ▶ May assume $t = r_a$ for some job $a \leq j$
- ▶ Machine busy in $[r_a, r_j]$ with total volume assigned $\leq r_j - r_a + T$
- ▶ By time r_j , $\leq (r_j - r_a + T) - |[r_a, r_j]| = T$ volume left, so j finishes by $r_j + T$

Application in scheduling II: flow-time minimization

- Jobs J with processing times d_j , machines M , bipartite graph $D = (M \cup J, E)$
- Jobs ordered by release time $r_1 \leq r_2 \leq \dots \leq r_n$
- Minimize the max *flow-time* $\max_{j \in J} (C_j - r_j)$ where C_j is the completion time


$$\min T$$

$$\text{s.t. } \sum_{i \in M} x_{ij} = 1, \quad \forall j \in J,$$


$$\sum_{j=a}^b d_j x_{ij} \leq (r_b - r_a) + T, \quad \forall i \in M, 1 \leq a \leq b \leq n,$$

$$x \geq 0, x_{ij} = 0 \quad \forall (i, j) \notin E.$$

$D = (M \cup J, A)$



(splittable) flow



Application in scheduling II: flow-time minimization



- Jobs J with processing times d_j , machines M , bipartite graph $D = (M \cup J, E)$
- Jobs ordered by release time $r_1 \leq r_2 \leq \dots \leq r_n$
- Minimize the max *flow-time* $\max_{j \in J} (C_j - r_j)$ where C_j is the completion time

$$\min T$$

$$\text{s.t.} \quad \sum_{i \in M} x_{ij} = 1, \quad \forall j \in J,$$

$$\sum_{j=a}^b d_j x_{ij} \leq (r_b - r_a) + T, \quad \forall i \in M, \quad 1 \leq a \leq b \leq n,$$

$$x \geq 0, x_{ij} = 0 \quad \forall (i, j) \notin E.$$



Application in scheduling II: flow-time minimization

Special case of Morell-Skutella's conjecture

Given bipartite $D = (M \cup J, E)$, and a fractional assignment $x \in [0, 1]^{m \times n}$, there is an assignment $y \in \{0, 1\}^{m \times n}$ such that

$$\left| \sum_{j=1}^t d_j x_{ij} - \sum_{j=1}^t d_j y_{ij} \right| \leq d_{\max}, \quad \forall i \in [m], t \in [n].$$

Application in scheduling II: flow-time minimization

Special case of Morell-Skutella's conjecture

Given bipartite $D = (M \cup J, E)$, and a fractional assignment $x \in [0, 1]^{m \times n}$, there is an assignment $y \in \{0, 1\}^{m \times n}$ such that

$$\left| \sum_{j=1}^t d_j x_{ij} - \sum_{j=1}^t d_j y_{ij} \right| \leq d_{\max}, \quad \forall i \in [m], t \in [n].$$

- Given a feasible LP solution x , can round it to y so that for any $1 \leq a \leq b \leq n$,

$$\begin{aligned} \sum_{j=a}^b d_j y_{ij} &= \sum_{j=1}^b d_j y_{ij} - \sum_{j=1}^{a-1} d_j y_{ij} \leq \left(\sum_{j=1}^b d_j x_{ij} + d_{\max} \right) - \left(\sum_{j=1}^{a-1} d_j x_{ij} - d_{\max} \right) \\ &= \sum_{j=a}^b d_j x_{ij} + 2d_{\max} \\ &\leq (r_b - r_a) + OPT + 2d_{\max} \\ &\leq (r_b - r_a) + 3OPT. \end{aligned}$$

Application in scheduling II: flow-time minimization

Special case of Morell-Skutella's conjecture

Given bipartite $D = (M \cup J, E)$, and a fractional assignment $x \in [0, 1]^{m \times n}$, there is an assignment $y \in \{0, 1\}^{m \times n}$ such that

$$\left| \sum_{j=1}^t d_j x_{ij} - \sum_{j=1}^t d_j y_{ij} \right| \leq d_{\max}, \quad \forall i \in [m], t \in [n].$$

- Given a feasible LP solution x , can round it to y so that for any $1 \leq a \leq b \leq n$,

$$\begin{aligned} \sum_{j=a}^b d_j y_{ij} &= \sum_{j=1}^b d_j y_{ij} - \sum_{j=1}^{a-1} d_j y_{ij} \leq \left(\sum_{j=1}^b d_j x_{ij} + d_{\max} \right) - \left(\sum_{j=1}^{a-1} d_j x_{ij} - d_{\max} \right) \\ &= \sum_{j=a}^b d_j x_{ij} + 2d_{\max} \\ &\leq (r_b - r_a) + OPT + 2d_{\max} \\ &\leq (r_b - r_a) + 3OPT. \end{aligned}$$

- Known up to $O(\sqrt{\log n})$ [Bansal, Svensson, Rohwedder STOC 2022]

Main results

Theorem (Liu, R., 2026)

Given a fractional assignment $x \in [0, 1]^{m \times n}$ and weights $d \in \mathbb{R}_{>0}^n$, there exists an assignment $y \in \{0, 1\}^{m \times n}$ so that for every $i \in [m]$ and $t \in [n]$,

$$\left| \sum_{j=1}^t d_j x_{ij} - \sum_{j=1}^t d_j y_{ij} \right| < \max_{j \in [n]} d_j.$$

Main results

Theorem (Liu, R., 2026)

Given a fractional assignment $x \in [0, 1]^{m \times n}$ and weights $d \in \mathbb{R}_{>0}^n$, there exists an assignment $y \in \{0, 1\}^{m \times n}$ so that for every $i \in [m]$ and $t \in [n]$,

$$\left| \sum_{j=1}^t d_j x_{ij} - \sum_{j=1}^t d_j y_{ij} \right| < \max_{j \in [n]} d_j.$$

- The RHS can be improved to $(1 - \frac{1}{2m-2}) \cdot \max_{j \in [n]} d_j$, which is tight:

	1	2	...	$m-1$
1	$\frac{1}{2m-2}$	$\frac{1}{m-1}$...	$\frac{1}{m-1}$
2	$\frac{1}{2m-2}$	$\frac{1}{m-1}$...	$\frac{1}{m-1}$
\vdots	\vdots	\vdots	\ddots	\vdots
$m-1$	$\frac{1}{2m-2}$	$\frac{1}{m-1}$...	$\frac{1}{m-1}$
m	$\frac{1}{2}$	0	...	0

Application to flow-time scheduling

Corollary (Liu, R., 2026)

A 3-approx. algorithm for maximum flow-time minimization on identical machines.

Application to flow-time scheduling

Corollary (Liu, R., 2026)

A 3-approx. algorithm for maximum flow-time minimization on identical machines.

- ▶ Already known: sort jobs by release times and schedule on least busy machine [Mastrolilli, 2004]

Application to flow-time scheduling

Corollary (Liu, R., 2026)

A 3-approx. algorithm for maximum flow-time minimization on identical machines.

- ▶ Already known: sort jobs by release times and schedule on least busy machine [Mastrolilli, 2004]
- ▶ Suppose now each machine i has a *closing time* b_i

Application to flow-time scheduling

Corollary (Liu, R., 2026)

A 3-approx. algorithm for maximum flow-time minimization on identical machines.

- ▶ Already known: sort jobs by release times and schedule on least busy machine [Mastrolilli, 2004]
- ▶ Suppose now each machine i has a *closing time* b_i
- ▶ Can schedule job j on machine i if $r_j \leq b_i$

$$r = \delta \quad r = 2\delta \quad \dots \quad r = m\delta$$

$b_1 = \delta$	$\frac{1}{m}$		
$b_2 = 2\delta$	$\frac{1}{m}$	$\frac{1}{m-1}$	
\vdots	$\frac{1}{m}$	$\frac{1}{m-1}$	\dots
$b_m = m\delta$	$\frac{1}{m}$	$\frac{1}{m-1}$	\dots

$\frac{1}{m}$	$\frac{1}{m}$	$\frac{1}{m}$	$\frac{1}{m}$
$\frac{1}{m-1}$	$\frac{1}{m-1}$	$\frac{1}{m-1}$	
\dots			
1	1	1	

Application to flow-time scheduling

Corollary (Liu, R., 2026)

A 3-approx. algorithm for maximum flow-time minimization on **identical** machines with closing times.

- ▶ Already known: sort jobs by release times and schedule on least busy machine [Mastrolilli, 2004]
- ▶ Suppose now each machine i has a *closing time* b_i
- ▶ Can schedule job j on machine i if $r_j \leq b_i$

$$r = \delta \quad r = 2\delta \quad \dots \quad r = m\delta$$

$b_1 = \delta$	$\frac{1}{m}$		
$b_2 = 2\delta$	$\frac{1}{m}$	$\frac{1}{m-1}$	
\vdots	$\frac{1}{m}$	$\frac{1}{m-1}$	\dots
$b_m = m\delta$	$\frac{1}{m}$	$\frac{1}{m-1}$	\dots

$\frac{1}{m}$	$\frac{1}{m}$	$\frac{1}{m}$	$\frac{1}{m}$
$\frac{1}{m-1}$	$\frac{1}{m-1}$	$\frac{1}{m-1}$	
\dots			
1			

Algorithm

Given fractional assignment $x \in [0, 1]^{m \times n}$, $d \in (0, 1]^n$, define

$$P_t(i) := \sum_{j=1}^t d_j x_{ij}, \quad N_t(i) := \sum_{j=1}^t d_j y_{ij}, \quad \Delta_t(i) := P_t(i) - N_t(i).$$

Algorithm

Given fractional assignment $x \in [0, 1]^{m \times n}$, $d \in (0, 1]^n$, define

$$P_t(i) := \sum_{j=1}^t d_j x_{ij}, \quad N_t(i) := \sum_{j=1}^t d_j y_{ij}, \quad \Delta_t(i) := P_t(i) - N_t(i).$$

Algorithm 1 EARLIEST DEADLINE ALGORITHM

for $t \in [n]$ **do**

$C(t) \leftarrow \{i \in [m] : P_t(i) > N_{t-1}(i)\}$

 ▷ set of candidates at time t

for $i \in [m]$ **do**

$D_t(i) \leftarrow \min\{T \geq t : P_T(i) \geq N_{t-1}(i) + 1\}$

 ▷ deadline of i at time t

end for

 Choose $s_t \in \arg \min_{i \in C(t)} D_t(i)$

$y_{s_t, t} \leftarrow 1$ and $y_{i, t} \leftarrow 0$ for $i \in [m] \setminus \{s_t\}$

end for

Algorithm

Given fractional assignment $x \in [0, 1]^{m \times n}$, $d \in (0, 1]^n$, define

$$P_t(i) := \sum_{j=1}^t d_j x_{ij}, \quad N_t(i) := \sum_{j=1}^t d_j y_{ij}, \quad \Delta_t(i) := P_t(i) - N_t(i).$$

Algorithm 1 EARLIEST DEADLINE ALGORITHM

```
for  $t \in [n]$  do  
     $C(t) \leftarrow \{i \in [m] : P_t(i) > N_{t-1}(i)\}$  ▷ set of candidates at time  $t$   
    for  $i \in [m]$  do  
         $D_t(i) \leftarrow \min\{T \geq t : P_T(i) \geq N_{t-1}(i) + 1\}$  ▷ deadline of  $i$  at time  $t$   
    end for  
    Choose  $s_t \in \arg \min_{i \in C(t)} D_t(i)$   
     $y_{s_t, t} \leftarrow 1$  and  $y_{i, t} \leftarrow 0$  for  $i \in [m] \setminus \{s_t\}$   
end for
```

Claim 1: The algorithm outputs a valid assignment, i.e. $C(t) \neq \emptyset$ for all $t \in [n]$.

Algorithm

Given fractional assignment $x \in [0, 1]^{m \times n}$, $d \in (0, 1]^n$, define

$$P_t(i) := \sum_{j=1}^t d_j x_{ij}, \quad N_t(i) := \sum_{j=1}^t d_j y_{ij}, \quad \Delta_t(i) := P_t(i) - N_t(i).$$

Algorithm 1 EARLIEST DEADLINE ALGORITHM

for $t \in [n]$ **do**

$C(t) \leftarrow \{i \in [m] : P_t(i) > N_{t-1}(i)\}$ ▷ set of candidates at time t

for $i \in [m]$ **do**

$D_t(i) \leftarrow \min\{T \geq t : P_T(i) \geq N_{t-1}(i) + 1\}$ ▷ deadline of i at time t

end for

Choose $s_t \in \arg \min_{i \in C(t)} D_t(i)$

$y_{s_t, t} \leftarrow 1$ and $y_{i, t} \leftarrow 0$ for $i \in [m] \setminus \{s_t\}$

end for

Claim 1: The algorithm outputs a valid assignment, i.e. $C(t) \neq \emptyset$ for all $t \in [n]$.

$$\sum_{i=1}^m (P_t(i) - N_{t-1}(i)) = \sum_{j=1}^{t-1} d_j \left(\sum_{i=1}^m x_{ij} - \sum_{i=1}^m y_{ij} \right) + d_t \sum_{i=1}^m x_{ij} = d_t > 0.$$

Algorithm

Given fractional assignment $x \in [0, 1]^{m \times n}$, $d \in (0, 1]^n$, define

$$P_t(i) := \sum_{j=1}^t d_j x_{ij}, \quad N_t(i) := \sum_{j=1}^t d_j y_{ij}, \quad \Delta_t(i) := P_t(i) - N_t(i).$$

Algorithm 1 EARLIEST DEADLINE ALGORITHM

for $t \in [n]$ **do**

$C(t) \leftarrow \{i \in [m] : P_t(i) > N_{t-1}(i)\}$

 ▷ set of candidates at time t

for $i \in [m]$ **do**

$D_t(i) \leftarrow \min\{T \geq t : P_T(i) \geq N_{t-1}(i) + 1\}$

 ▷ deadline of i at time t

end for

 Choose $s_t \in \arg \min_{i \in C(t)} D_t(i)$

$y_{s_t, t} \leftarrow 1$ and $y_{i, t} \leftarrow 0$ for $i \in [m] \setminus \{s_t\}$

end for

Claim 1: The algorithm outputs a valid assignment, i.e. $C(t) \neq \emptyset$ for all $t \in [n]$.

Claim 2: $\Delta_t(i) > -1$ for all $i \in [m]$, $t \in [n]$.

Algorithm

Given fractional assignment $x \in [0, 1]^{m \times n}$, $d \in (0, 1]^n$, define

$$P_t(i) := \sum_{j=1}^t d_j x_{ij}, \quad N_t(i) := \sum_{j=1}^t d_j y_{ij}, \quad \Delta_t(i) := P_t(i) - N_t(i).$$

Algorithm 1 EARLIEST DEADLINE ALGORITHM

for $t \in [n]$ **do**

$C(t) \leftarrow \{i \in [m] : P_t(i) > N_{t-1}(i)\}$

 ▷ set of candidates at time t

for $i \in [m]$ **do**

$D_t(i) \leftarrow \min\{T \geq t : P_T(i) \geq N_{t-1}(i) + 1\}$

 ▷ deadline of i at time t

end for

 Choose $s_t \in \arg \min_{i \in C(t)} D_t(i)$

$y_{s_t, t} \leftarrow 1$ and $y_{i, t} \leftarrow 0$ for $i \in [m] \setminus \{s_t\}$

end for

Claim 1: The algorithm outputs a valid assignment, i.e. $C(t) \neq \emptyset$ for all $t \in [n]$.

Claim 2: $\Delta_t(i) > -1$ for all $i \in [m]$, $t \in [n]$.

Claim 3: $\Delta_t(i) < 1$ for all $i \in [m]$, $t \in [n]$.

Lower bound proof

Given fractional assignment $x \in [0, 1]^{m \times n}$, $d \in (0, 1]^n$, define

$$P_t(i) := \sum_{j=1}^t d_j x_{ij}, \quad N_t(i) := \sum_{j=1}^t d_j y_{ij}, \quad \Delta_t(i) := P_t(i) - N_t(i).$$

$$C(t) := \{i \in [m] : P_t(i) > N_{t-1}(i)\}, \quad D_t(i) := \min\{T \geq t : P_T(i) \geq N_{t-1}(i) + 1\}$$

Claim 2

For any $i \in [m]$ and $t \in [n]$, $\Delta_t(i) > -1$.

Lower bound proof

Given fractional assignment $x \in [0, 1]^{m \times n}$, $d \in (0, 1]^n$, define

$$P_t(i) := \sum_{j=1}^t d_j x_{ij}, \quad N_t(i) := \sum_{j=1}^t d_j y_{ij}, \quad \Delta_t(i) := P_t(i) - N_t(i).$$

$$C(t) := \{i \in [m] : P_t(i) > N_{t-1}(i)\}, \quad D_t(i) := \min\{T \geq t : P_T(i) \geq N_{t-1}(i) + 1\}$$

Claim 2

For any $i \in [m]$ and $t \in [n]$, $\Delta_t(i) > -1$.

Proof. If i has not been selected at any time up to t , then $N_t(i) = 0$ and

$$\Delta_t(i) = P_t(i) \geq 0 > -1.$$

Lower bound proof

Given fractional assignment $x \in [0, 1]^{m \times n}$, $d \in (0, 1]^n$, define

$$P_t(i) := \sum_{j=1}^t d_j x_{ij}, \quad N_t(i) := \sum_{j=1}^t d_j y_{ij}, \quad \Delta_t(i) := P_t(i) - N_t(i).$$

$$C(t) := \{i \in [m] : P_t(i) > N_{t-1}(i)\}, \quad D_t(i) := \min\{T \geq t : P_T(i) \geq N_{t-1}(i) + 1\}$$

Claim 2

For any $i \in [m]$ and $t \in [n]$, $\Delta_t(i) > -1$.

Proof. If i has not been selected at any time up to t , then $N_t(i) = 0$ and

$$\Delta_t(i) = P_t(i) \geq 0 > -1.$$

Otherwise, let $j \in [t]$ be the latest time such that $s_j = i$. Then

$$\Delta_t(i) = P_t(i) - N_t(i)$$

Lower bound proof

Given fractional assignment $x \in [0, 1]^{m \times n}$, $d \in (0, 1]^n$, define

$$P_t(i) := \sum_{j=1}^t d_j x_{ij}, \quad N_t(i) := \sum_{j=1}^t d_j y_{ij}, \quad \Delta_t(i) := P_t(i) - N_t(i).$$

$$C(t) := \{i \in [m] : P_t(i) > N_{t-1}(i)\}, \quad D_t(i) := \min\{T \geq t : P_T(i) \geq N_{t-1}(i) + 1\}$$

Claim 2

For any $i \in [m]$ and $t \in [n]$, $\Delta_t(i) > -1$.

Proof. If i has not been selected at any time up to t , then $N_t(i) = 0$ and

$$\Delta_t(i) = P_t(i) \geq 0 > -1.$$

Otherwise, let $j \in [t]$ be the latest time such that $s_j = i$. Then

$$\begin{aligned} \Delta_t(i) &= P_t(i) - N_t(i) \\ &= P_t(i) - N_{j-1}(i) - d_j \end{aligned}$$

Lower bound proof

Given fractional assignment $x \in [0, 1]^{m \times n}$, $d \in (0, 1]^n$, define

$$P_t(i) := \sum_{j=1}^t d_j x_{ij}, \quad N_t(i) := \sum_{j=1}^t d_j y_{ij}, \quad \Delta_t(i) := P_t(i) - N_t(i).$$

$$C(t) := \{i \in [m] : P_t(i) > N_{t-1}(i)\}, \quad D_t(i) := \min\{T \geq t : P_T(i) \geq N_{t-1}(i) + 1\}$$

Claim 2

For any $i \in [m]$ and $t \in [n]$, $\Delta_t(i) > -1$.

Proof. If i has not been selected at any time up to t , then $N_t(i) = 0$ and

$$\Delta_t(i) = P_t(i) \geq 0 > -1.$$

Otherwise, let $j \in [t]$ be the latest time such that $s_j = i$. Then

$$\begin{aligned} \Delta_t(i) &= P_t(i) - N_t(i) \\ &= P_t(i) - N_{j-1}(i) - d_j \\ &\geq P_j(i) - N_{j-1}(i) - d_j \end{aligned}$$

Lower bound proof

Given fractional assignment $x \in [0, 1]^{m \times n}$, $d \in (0, 1]^n$, define

$$P_t(i) := \sum_{j=1}^t d_j x_{ij}, \quad N_t(i) := \sum_{j=1}^t d_j y_{ij}, \quad \Delta_t(i) := P_t(i) - N_t(i).$$

$$C(t) := \{i \in [m] : P_t(i) > N_{t-1}(i)\}, \quad D_t(i) := \min\{T \geq t : P_T(i) \geq N_{t-1}(i) + 1\}$$

Claim 2

For any $i \in [m]$ and $t \in [n]$, $\Delta_t(i) > -1$.

Proof. If i has not been selected at any time up to t , then $N_t(i) = 0$ and

$$\Delta_t(i) = P_t(i) \geq 0 > -1.$$

Otherwise, let $j \in [t]$ be the latest time such that $s_j = i$. Then

$$\begin{aligned} \Delta_t(i) &= P_t(i) - N_t(i) \\ &= P_t(i) - N_{j-1}(i) - d_j \\ &\geq P_j(i) - N_{j-1}(i) - d_j \\ &> -d_j \geq -1. \end{aligned}$$

Upper bound proof I

Given fractional assignment $x \in [0, 1]^{m \times n}$, $d \in (0, 1]^n$, define

$$P_t(i) := \sum_{j=1}^t d_j x_{ij}, \quad N_t(i) := \sum_{j=1}^t d_j y_{ij}, \quad \Delta_t(i) := P_t(i) - N_t(i).$$

$$C(t) := \{i \in [m] : P_t(i) > N_{t-1}(i)\}, \quad D_t(i) := \min\{T \geq t : P_T(i) \geq N_{t-1}(i) + 1\}$$

Claim 3 (Main technical lemma)

For any $i \in [m]$ and $t \in [n]$, $\Delta_t(i) < 1$.

Upper bound proof I

Given fractional assignment $x \in [0, 1]^{m \times n}$, $d \in (0, 1]^n$, define

$$P_t(i) := \sum_{j=1}^t d_j x_{ij}, \quad N_t(i) := \sum_{j=1}^t d_j y_{ij}, \quad \Delta_t(i) := P_t(i) - N_t(i).$$

$$C(t) := \{i \in [m] : P_t(i) > N_{t-1}(i)\}, \quad D_t(i) := \min\{T \geq t : P_T(i) \geq N_{t-1}(i) + 1\}$$

Claim 3 (Main technical lemma)

For any $i \in [m]$ and $t \in [n]$, $\Delta_t(i) < 1$.

Proof. Assuming otherwise $I^+ := \{i \in [m] : \Delta_t(i) \geq 1\} \neq \emptyset$ for some $t \in [n]$.

Upper bound proof I

Given fractional assignment $x \in [0, 1]^{m \times n}$, $d \in (0, 1]^n$, define

$$P_t(i) := \sum_{j=1}^t d_j x_{ij}, \quad N_t(i) := \sum_{j=1}^t d_j y_{ij}, \quad \Delta_t(i) := P_t(i) - N_t(i).$$

$$C(t) := \{i \in [m] : P_t(i) > N_{t-1}(i)\}, \quad D_t(i) := \min\{T \geq t : P_T(i) \geq N_{t-1}(i) + 1\}$$

Claim 3 (Main technical lemma)

For any $i \in [m]$ and $t \in [n]$, $\Delta_t(i) < 1$.

Proof. Assuming otherwise $I^+ := \{i \in [m] : \Delta_t(i) \geq 1\} \neq \emptyset$ for some $t \in [n]$.

$$\sum_{i \in [m]} \Delta_t(i) = 0 \implies \{i \in [m] : \Delta_t(i) < 0\} \neq \emptyset.$$

Upper bound proof I

Given fractional assignment $x \in [0, 1]^{m \times n}$, $d \in (0, 1]^n$, define

$$P_t(i) := \sum_{j=1}^t d_j x_{ij}, \quad N_t(i) := \sum_{j=1}^t d_j y_{ij}, \quad \Delta_t(i) := P_t(i) - N_t(i).$$

$$C(t) := \{i \in [m] : P_t(i) > N_{t-1}(i)\}, \quad D_t(i) := \min\{T \geq t : P_T(i) \geq N_{t-1}(i) + 1\}$$

Claim 3 (Main technical lemma)

For any $i \in [m]$ and $t \in [n]$, $\Delta_t(i) < 1$.

Proof. Assuming otherwise $I^+ := \{i \in [m] : \Delta_t(i) \geq 1\} \neq \emptyset$ for some $t \in [n]$.

$$\sum_{i \in [m]} \Delta_t(i) = 0 \implies \{i \in [m] : \Delta_t(i) < 0\} \neq \emptyset.$$

So $t^- := \max\{j \leq t : \Delta_t(s_j) < 0\}$ is well-defined.

Upper bound proof I

Given fractional assignment $x \in [0, 1]^{m \times n}$, $d \in (0, 1]^n$, define

$$P_t(i) := \sum_{j=1}^t d_j x_{ij}, \quad N_t(i) := \sum_{j=1}^t d_j y_{ij}, \quad \Delta_t(i) := P_t(i) - N_t(i).$$

$$C(t) := \{i \in [m] : P_t(i) > N_{t-1}(i)\}, \quad D_t(i) := \min\{T \geq t : P_T(i) \geq N_{t-1}(i) + 1\}$$

Claim 3 (Main technical lemma)

For any $i \in [m]$ and $t \in [n]$, $\Delta_t(i) < 1$.

Proof. Assuming otherwise $I^+ := \{i \in [m] : \Delta_t(i) \geq 1\} \neq \emptyset$ for some $t \in [n]$.

$$\sum_{i \in [m]} \Delta_t(i) = 0 \implies \{i \in [m] : \Delta_t(i) < 0\} \neq \emptyset.$$

So $t^- := \max\{j \leq t : \Delta_t(s_j) < 0\}$ is well-defined.

We will need to define another time $t^- \leq t' \leq t$ and analyze what happens in $(t', t]$.

Upper bound proof II

$t^- := \max\{j \leq t : \Delta_t(s_j) < 0\}$ is well-defined.

Upper bound proof II

$t^- := \max\{j \leq t : \Delta_t(s_j) < 0\}$ is well-defined.

Claim 3.1.

$t' := \max\{j \leq t : P_t(s_j) < N_{j-1}(s_j) + 1\}$ is well-defined.

Upper bound proof II

$t^- := \max\{j \leq t : \Delta_t(s_j) < 0\}$ is well-defined.

Claim 3.1.

$t' := \max\{j \leq t : P_t(s_j) < N_{j-1}(s_j) + 1\}$ is well-defined.

Proof. It suffices to show that t^- satisfies the inequality:

Upper bound proof II

$t^- := \max\{j \leq t : \Delta_t(s_j) < 0\}$ is well-defined.

Claim 3.1.

$t' := \max\{j \leq t : P_t(s_j) < N_{j-1}(s_j) + 1\}$ is well-defined.

Proof. It suffices to show that t^- satisfies the inequality:

$$P_t(s_{t^-}) < N_t(s_{t^-})$$

Upper bound proof II

$t^- := \max\{j \leq t : \Delta_t(s_j) < 0\}$ is well-defined.

Claim 3.1.

$t' := \max\{j \leq t : P_t(s_j) < N_{j-1}(s_j) + 1\}$ is well-defined.

Proof. It suffices to show that t^- satisfies the inequality:

$$\begin{aligned} P_t(s_{t^-}) &< N_t(s_{t^-}) \\ &= N_{t^-}(s_{t^-}) \end{aligned}$$

Upper bound proof II

$t^- := \max\{j \leq t : \Delta_t(s_j) < 0\}$ is well-defined.

Claim 3.1.

$t' := \max\{j \leq t : P_t(s_j) < N_{j-1}(s_j) + 1\}$ is well-defined.

Proof. It suffices to show that t^- satisfies the inequality:

$$\begin{aligned} P_t(s_{t^-}) &< N_t(s_{t^-}) \\ &= N_{t^-}(s_{t^-}) \\ &= N_{t^- - 1}(s_{t^-}) + d_{t^-} \end{aligned}$$

Upper bound proof II

$t^- := \max\{j \leq t : \Delta_t(s_j) < 0\}$ is well-defined.

Claim 3.1.

$t' := \max\{j \leq t : P_t(s_j) < N_{j-1}(s_j) + 1\}$ is well-defined.

Proof. It suffices to show that t^- satisfies the inequality:

$$\begin{aligned} P_t(s_{t^-}) &< N_t(s_{t^-}) \\ &= N_{t^-}(s_{t^-}) \\ &= N_{t^- - 1}(s_{t^-}) + d_{t^-} \\ &\leq N_{t^- - 1}(s_{t^-}) + 1. \quad \square \end{aligned}$$

Upper bound proof II

$t^- := \max\{j \leq t : \Delta_t(s_j) < 0\}$ is well-defined.

Claim 3.1.

$t' := \max\{j \leq t : P_t(s_j) < N_{j-1}(s_j) + 1\}$ is well-defined.

Proof. It suffices to show that t^- satisfies the inequality:

$$\begin{aligned} P_t(s_{t^-}) &< N_t(s_{t^-}) \\ &= N_{t^-}(s_{t^-}) \\ &= N_{t^- - 1}(s_{t^-}) + d_{t^-} \\ &\leq N_{t^- - 1}(s_{t^-}) + 1. \quad \square \end{aligned}$$

► By maximality, for any $j \in (t', t]$, $P_t(s_j) \geq N_{j-1}(s_j) + 1 \geq N_{t'-1}(s_j) + 1$.

Upper bound proof II

$t^- := \max\{j \leq t : \Delta_t(s_j) < 0\}$ is well-defined.

Claim 3.1.

$t' := \max\{j \leq t : P_t(s_j) < N_{j-1}(s_j) + 1\}$ is well-defined.

Proof. It suffices to show that t^- satisfies the inequality:

$$\begin{aligned} P_t(s_{t^-}) &< N_t(s_{t^-}) \\ &= N_{t^-}(s_{t^-}) \\ &= N_{t^- - 1}(s_{t^-}) + d_{t^-} \\ &\leq N_{t^- - 1}(s_{t^-}) + 1. \quad \square \end{aligned}$$

- ▶ By maximality, for any $j \in (t', t]$, $P_t(s_j) \geq N_{j-1}(s_j) + 1 \geq N_{t'-1}(s_j) + 1$.
- ▶ Define $I := \{i \in [m] : P_t(i) \geq N_{t'-1}(i) + 1\} = \{i \in [m] : D_{t'}(i) \leq t\}$.

Upper bound proof II

$t^- := \max\{j \leq t : \Delta_t(s_j) < 0\}$ is well-defined.

Claim 3.1.

$t' := \max\{j \leq t : P_t(s_j) < N_{j-1}(s_j) + 1\}$ is well-defined.

Proof. It suffices to show that t^- satisfies the inequality:

$$\begin{aligned} P_t(s_{t^-}) &< N_t(s_{t^-}) \\ &= N_{t^-}(s_{t^-}) \\ &= N_{t^- - 1}(s_{t^-}) + d_{t^-} \\ &\leq N_{t^- - 1}(s_{t^-}) + 1. \quad \square \end{aligned}$$

- ▶ By maximality, for any $j \in (t', t]$, $P_t(s_j) \geq N_{j-1}(s_j) + 1 \geq N_{t' - 1}(s_j) + 1$.
- ▶ Define $I := \{i \in [m] : P_t(i) \geq N_{t' - 1}(i) + 1\} = \{i \in [m] : D_{t'}(i) \leq t\}$.
- ▶ $s_{t'} \notin I, s_j \in I \quad \forall j \in (t', t]$.

Upper bound proof II

$t^- := \max\{j \leq t : \Delta_t(s_j) < 0\}$ is well-defined.

Claim 3.1.

$t' := \max\{j \leq t : P_t(s_j) < N_{j-1}(s_j) + 1\}$ is well-defined.

Proof. It suffices to show that t^- satisfies the inequality:

$$\begin{aligned} P_t(s_{t^-}) &< N_t(s_{t^-}) \\ &= N_{t^-}(s_{t^-}) \\ &= N_{t^- - 1}(s_{t^-}) + d_{t^-} \\ &\leq N_{t^- - 1}(s_{t^-}) + 1. \quad \square \end{aligned}$$

- ▶ By maximality, for any $j \in (t', t]$, $P_t(s_j) \geq N_{j-1}(s_j) + 1 \geq N_{t' - 1}(s_j) + 1$.
- ▶ Define $I := \{i \in [m] : P_t(i) \geq N_{t' - 1}(i) + 1\} = \{i \in [m] : D_{t'}(i) \leq t\}$.
- ▶ $s_{t'} \notin I, s_j \in I \quad \forall j \in (t', t]$.
- ▶ $I \neq \emptyset$ since $I^+ \subseteq I$: $P_t(i) \geq N_t(i) + 1 \geq N_{t' - 1}(i) + 1$ for any $i \in I^+$

Upper bound proof (last part!)

- ▶ Define $I := \{i \in [m] : P_t(i) \geq N_{t'-1}(i) + 1\} = \{i \in [m] : D_{t'}(i) \leq t\}$.
- ▶ $s_{t'} \notin I, s_j \in I \quad \forall j \in (t', t], \quad I \neq \emptyset$

Upper bound proof (last part!)

- ▶ Define $I := \{i \in [m] : P_t(i) \geq N_{t'-1}(i) + 1\} = \{i \in [m] : D_{t'}(i) \leq t\}$.
- ▶ $s_{t'} \notin I, s_j \in I \quad \forall j \in (t', t], \quad I \neq \emptyset$

Claim 3.2.

For any $i \in I$ we have $P_t(i) \geq N_t(i)$ and $P_{t'}(i) \leq N_{t'-1}(i)$.

Upper bound proof (last part!)

- ▶ Define $I := \{i \in [m] : P_t(i) \geq N_{t'-1}(i) + 1\} = \{i \in [m] : D_{t'}(i) \leq t\}$.
- ▶ $s_{t'} \notin I, s_j \in I \quad \forall j \in (t', t], \quad I \neq \emptyset$

Claim 3.2.

For any $i \in I$ we have $P_t(i) \geq N_t(i)$ and $P_{t'}(i) \leq N_{t'-1}(i)$.

Contradiction assuming Claim 3.2.:

$$\sum_{j \in (t', t]} d_j < 1 + \sum_{j \in (t', t]} d_j$$

Upper bound proof (last part!)

- ▶ Define $I := \{i \in [m] : P_t(i) \geq N_{t'-1}(i) + 1\} = \{i \in [m] : D_{t'}(i) \leq t\}$.
- ▶ $s_{t'} \notin I, s_j \in I \quad \forall j \in (t', t], \quad I \neq \emptyset$

Claim 3.2.

For any $i \in I$ we have $P_t(i) \geq N_t(i)$ and $P_{t'}(i) \leq N_{t'-1}(i)$.

Contradiction assuming Claim 3.2.:

$$\begin{aligned} \sum_{j \in (t', t]} d_j &< 1 + \sum_{j \in (t', t]} d_j \\ &= 1 + \sum_{i \in I} (N_t(i) - N_{t'-1}(i)) \end{aligned}$$

Upper bound proof (last part!)

- ▶ Define $I := \{i \in [m] : P_t(i) \geq N_{t'-1}(i) + 1\} = \{i \in [m] : D_{t'}(i) \leq t\}$.
- ▶ $s_{t'} \notin I, s_j \in I \quad \forall j \in (t', t], \quad I \neq \emptyset$

Claim 3.2.

For any $i \in I$ we have $P_t(i) \geq N_t(i)$ and $P_{t'}(i) \leq N_{t'-1}(i)$.

Contradiction assuming Claim 3.2.:

$$\begin{aligned} \sum_{j \in (t', t]} d_j &< 1 + \sum_{j \in (t', t]} d_j \\ &= 1 + \sum_{i \in I} (N_t(i) - N_{t'-1}(i)) \\ &\leq \sum_{i \in I} (P_t(i) - N_{t'-1}(i)) \end{aligned}$$

Upper bound proof (last part!)

- ▶ Define $I := \{i \in [m] : P_t(i) \geq N_{t'-1}(i) + 1\} = \{i \in [m] : D_{t'}(i) \leq t\}$.
- ▶ $s_{t'} \notin I, s_j \in I \quad \forall j \in (t', t], \quad I \neq \emptyset$

Claim 3.2.

For any $i \in I$ we have $P_t(i) \geq N_t(i)$ and $P_{t'}(i) \leq N_{t'-1}(i)$.

Contradiction assuming Claim 3.2.:

$$\begin{aligned} \sum_{j \in (t', t]} d_j &< 1 + \sum_{j \in (t', t]} d_j \\ &= 1 + \sum_{i \in I} (N_t(i) - N_{t'-1}(i)) \\ &\leq \sum_{i \in I} (P_t(i) - N_{t'-1}(i)) \\ &\leq \sum_{i \in I} (P_t(i) - P_{t'}(i)) \end{aligned}$$

Upper bound proof (last part!)

- ▶ Define $I := \{i \in [m] : P_t(i) \geq N_{t'-1}(i) + 1\} = \{i \in [m] : D_{t'}(i) \leq t\}$.
- ▶ $s_{t'} \notin I, s_j \in I \quad \forall j \in (t', t], \quad I \neq \emptyset$

Claim 3.2.

For any $i \in I$ we have $P_t(i) \geq N_t(i)$ and $P_{t'}(i) \leq N_{t'-1}(i)$.

Contradiction assuming Claim 3.2.:

$$\begin{aligned} \sum_{j \in (t', t]} d_j &< 1 + \sum_{j \in (t', t]} d_j \\ &= 1 + \sum_{i \in I} (N_t(i) - N_{t'-1}(i)) \\ &\leq \sum_{i \in I} (P_t(i) - N_{t'-1}(i)) \\ &\leq \sum_{i \in I} (P_t(i) - P_{t'}(i)) \\ &\leq \sum_{i \in [m]} (P_t(i) - P_{t'}(i)) \end{aligned}$$

Upper bound proof (last part!)

- ▶ Define $I := \{i \in [m] : P_t(i) \geq N_{t'-1}(i) + 1\} = \{i \in [m] : D_{t'}(i) \leq t\}$.
- ▶ $s_{t'} \notin I, s_j \in I \quad \forall j \in (t', t], \quad I \neq \emptyset$

Claim 3.2.

For any $i \in I$ we have $P_t(i) \geq N_t(i)$ and $P_{t'}(i) \leq N_{t'-1}(i)$.

Contradiction assuming Claim 3.2.:

$$\begin{aligned} \sum_{j \in (t', t]} d_j &< 1 + \sum_{j \in (t', t]} d_j \\ &= 1 + \sum_{i \in I} (N_t(i) - N_{t'-1}(i)) \\ &\leq \sum_{i \in I} (P_t(i) - N_{t'-1}(i)) \\ &\leq \sum_{i \in I} (P_t(i) - P_{t'}(i)) \\ &\leq \sum_{i \in [m]} (P_t(i) - P_{t'}(i)) \\ &= \sum_{j \in (t', t]} d_j. \quad \square \end{aligned}$$

Upper bound proof (last part!)

- ▶ Define $I := \{i \in [m] : P_t(i) \geq N_{t'-1}(i) + 1\} = \{i \in [m] : D_{t'}(i) \leq t\}$.
- ▶ $s_{t'} \notin I, s_j \in I \quad \forall j \in (t', t], \quad I \neq \emptyset$

Claim 3.2.

For any $i \in I$ we have $P_t(i) \geq N_t(i)$ and $P_{t'}(i) \leq N_{t'-1}(i)$.

Proof. Fix $i \in I$. Either i is not chosen in $(t', t]$: $P_t(i) \geq N_{t'-1}(i) + 1 = N_t(i) + 1$.

Upper bound proof (last part!)

- ▶ Define $I := \{i \in [m] : P_t(i) \geq N_{t'-1}(i) + 1\} = \{i \in [m] : D_{t'}(i) \leq t\}$.
- ▶ $s_{t'} \notin I, s_j \in I \quad \forall j \in (t', t], \quad I \neq \emptyset$

Claim 3.2.

For any $i \in I$ we have $P_t(i) \geq N_t(i)$ and $P_{t'}(i) \leq N_{t'-1}(i)$.

Proof. Fix $i \in I$. Either i is not chosen in $(t', t]$: $P_t(i) \geq N_{t'-1}(i) + 1 = N_t(i) + 1$.

OR $s_j = i$ for some $j \in (t', t]$: then $j > t' \geq t^- \implies P_t(i) \geq N_t(i)$.

Upper bound proof (last part!)

- ▶ Define $I := \{i \in [m] : P_t(i) \geq N_{t'-1}(i) + 1\} = \{i \in [m] : D_{t'}(i) \leq t\}$.
- ▶ $s_{t'} \notin I, s_j \in I \quad \forall j \in (t', t], \quad I \neq \emptyset$

Claim 3.2.

For any $i \in I$ we have $P_t(i) \geq N_t(i)$ and $P_{t'}(i) \leq N_{t'-1}(i)$.

Proof. Fix $i \in I$. Either i is not chosen in $(t', t]$: $P_t(i) \geq N_{t'-1}(i) + 1 = N_t(i) + 1$.

OR $s_j = i$ for some $j \in (t', t]$: then $j > t' \geq t^- \implies P_t(i) \geq N_t(i)$.

For the second inequality: $D_{t'}(s_{t'}) > t \implies I \cap C(t') = \emptyset$. □

Other conjectures

Non-uniform weights

Let $d \in \mathbb{R}_{>0}^{m \times n}$. For every fractional assignment $x \in [0, 1]^{m \times n}$, there is an assignment $y \in \{0, 1\}^{m \times n}$ so that for every $i \in [m]$ and $t \in [n]$,

$$\left| \sum_{j \in [t]} d_{ij}(x_{ij} - y_{ij}) \right| \leq \max_{i \in [m], j \in [n]} d_{ij}.$$

Other conjectures

Non-uniform weights

Let $d \in \mathbb{R}_{>0}^{m \times n}$. For every fractional assignment $x \in [0, 1]^{m \times n}$, there is an assignment $y \in \{0, 1\}^{m \times n}$ so that for every $i \in [m]$ and $t \in [n]$,

$$\left| \sum_{j \in [t]} d_{ij}(x_{ij} - y_{ij}) \right| \leq \max_{i \in [m], j \in [n]} d_{ij}.$$

Weighted committee assignment

Let $d \in \mathbb{R}_{>0}^n$. For every $x \in [0, 1]^{m \times n}$ so that $n_j := \sum_{i \in [m]} x_{ij} \in \mathbb{N}$ for all $j \in [n]$, there is $y \in \{0, 1\}^{m \times n}$ so that $\sum_{i \in [m]} y_{ij} = n_j$ for all $j \in [n]$ and

$$\left| \sum_{j \in [t]} d_j(x_{ij} - y_{ij}) \right| \leq \max_{j \in [n]} d_j \quad \forall i \in [m], t \in [n].$$

Other conjectures

Non-uniform weights

Let $d \in \mathbb{R}_{>0}^{m \times n}$. For every fractional assignment $x \in [0, 1]^{m \times n}$, there is an assignment $y \in \{0, 1\}^{m \times n}$ so that for every $i \in [m]$ and $t \in [n]$,

$$\left| \sum_{j \in [t]} d_{ij}(x_{ij} - y_{ij}) \right| \leq \max_{i \in [m], j \in [n]} d_{ij}.$$

Weighted committee assignment

Let $d \in \mathbb{R}_{>0}^n$. For every $x \in [0, 1]^{m \times n}$ so that $n_j := \sum_{i \in [m]} x_{ij} \in \mathbb{N}$ for all $j \in [n]$, there is $y \in \{0, 1\}^{m \times n}$ so that $\sum_{i \in [m]} y_{ij} = n_j$ for all $j \in [n]$ and

$$\left| \sum_{j \in [t]} d_j(x_{ij} - y_{ij}) \right| \leq \max_{j \in [n]} d_j \quad \forall i \in [m], t \in [n].$$

Thanks for your attention!