Linear-size ℓ_1 Sparsifiers

Victor Reis

Joint work with Thomas Rothvoss

Simons Institute Workshop: Beyond Graph Laplacians

November 2, 2026






Graph Sparsification

Theorem (Batson-Spielman-Srivastava '08)


For any graph $G = (V, E)$ one can find weights $s(e) \geqslant 0$ in poly-time with $|\mathrm{supp}(s)| \leqslant O(n/\varepsilon^2)$ so that $|\delta(U)| = (1 \pm \varepsilon) \cdot |s(\delta(U))|$ for every $U \subseteq V$.



Graph Sparsification

Theorem (Batson-Spielman-Srivastava '08)

For any graph $G = (V, E)$ one can find weights $s(e) \geqslant 0$ in poly-time with $|\mathrm{supp}(s)| \leqslant O(n/\varepsilon^2)$ so that $|\delta(U)| = (1 \pm \varepsilon) \cdot |s(\delta(U))|$ for every $U \subseteq V$.




Graph Sparsification

Theorem (Batson-Spielman-Srivastava '08)

For any $A \in \mathbb{R}^{m \times n}$ there is a diagonal matrix $D \in \mathbb{R}^{m \times m}_{\geqslant 0}$ with support size $O(\frac{n}{\varepsilon^2})$ so that

$$(1-\varepsilon)\|Ax\|_2 \leqslant \|DAx\|_2 \leqslant (1+\varepsilon)\|Ax\|_2 \quad \forall x \in \mathbb{R}^n.$$

Matrix D can be found in polytime.



Main result

Theorem (R., Rothvoss'26)

For any matrix $A \in \mathbb{R}^{m \times n}$ and $\varepsilon \in (0, \frac{1}{2}]$, there is a diagonal matrix $D \in \mathbb{R}^{m \times m}_{\geqslant 0}$ with $O(\frac{n}{\varepsilon^2})$ nonzero entries so that

$$(1-\varepsilon)\|Ax\|_1 \leqslant \|DAx\|_1 \leqslant (1+\varepsilon)\|Ax\|_1 \quad \forall x \in \mathbb{R}^n.$$

- Previously known: $O(\frac{n}{\varepsilon^2} \log n)$ [Talagrand '90, Cohen, Peng '14]. See also [Bourgain, Lindenstrauss, Milman '89], [Schechtman '87]

Main result

Theorem (R., Rothvoss'26)

For any matrix $A \in \mathbb{R}^{m \times n}$ and $\varepsilon \in (0, \frac{1}{2}]$, there is a diagonal matrix $D \in \mathbb{R}^{m \times m}_{\geqslant 0}$ with $O(\frac{n}{\varepsilon^2})$ nonzero entries so that


$$(1-\varepsilon)\|Ax\|_1 \leqslant \|DAx\|_1 \leqslant (1+\varepsilon)\|Ax\|_1 \quad \forall x \in \mathbb{R}^n.$$

- Previously known: $O(\frac{n}{\varepsilon^2} \log n)$ [Talagrand '90, Cohen, Peng '14]. See also [Bourgain, Lindenstrauss, Milman '89], [Schechtman '87]
- No polytime algorithm (given A, B, checking whether $\|Ax\|_1 \leqslant \|Bx\|_1 \; \forall x \in \mathbb{R}^n$ is **coNP**-complete)

Zonotopes

Definition

A **zonotope** $Z \subseteq \mathbb{R}^n$ is the projection of a cube.




Zonotopes

Definition

A **zonotope** $Z \subseteq \mathbb{R}^n$ is the projection of a cube.

- We write $Z = \{\sum_{i=1}^{m} y_i A_i \mid y \in [-1,1]^m\}$ where $A \in \mathbb{R}^{m \times n}$




- m is the **number of segments** of Z

Zonotopes

Definition

A **zonotope** $Z \subseteq \mathbb{R}^n$ is the projection of a cube.

- We write $Z = \{\sum_{i=1}^{m} y_i A_i \mid y \in [-1,1]^m\}$ where $A \in \mathbb{R}^{m \times n}$




- m is the **number of segments** of Z
- Example of **zonoids**: B_p^n for $2 \leqslant p \leqslant \infty$

Zonotopes

Definition

A **zonotope** $Z \subseteq \mathbb{R}^n$ is the projection of a cube.

- We write $Z = \{\sum_{i=1}^{m} y_i A_i \mid y \in [-1,1]^m\}$ where $A \in \mathbb{R}^{m \times n}$




- m is the **number of segments** of Z
- Example of **zonoids**: B_p^n for $2 \leqslant p \leqslant \infty$
- NOT a zonoid: B_1^n

Zonotopes

Definition

A **zonotope** $Z \subseteq \mathbb{R}^n$ is the projection of a cube.

- We write $Z = \{\sum_{i=1}^{m} y_i A_i \mid y \in [-1,1]^m\}$ where $A \in \mathbb{R}^{m \times n}$



- m is the **number of segments** of Z
- Example of **zonoids**: B_p^n for $2 \leqslant p \leqslant \infty$
- NOT a zonoid: B_1^n
- **Support function** of zonotope Z:


$$h_Z(x) = \max_{y \in Z} \langle y, x \rangle = \|Ax\|_1$$

Reducing number of segments

Theorem (R., Rothvoss'26)

For any zonotope $Z \subseteq \mathbb{R}^n$ and $\varepsilon \in (0, \frac{1}{2}]$, there is a zonotope $Z' \subseteq \mathbb{R}^n$ with $O(\frac{n}{\varepsilon^2})$ segments so that

$$(1-\varepsilon)Z \subseteq Z' \subseteq (1+\varepsilon)Z.$$




Reducing number of segments

Theorem (R., Rothvoss'26)

For any zonotope $Z \subseteq \mathbb{R}^n$ and $\varepsilon \in (0, \frac{1}{2}]$, there is a zonotope $Z' \subseteq \mathbb{R}^n$ with $O(\frac{n}{\varepsilon^2})$ segments so that


$$(1-\varepsilon)Z \subseteq Z' \subseteq (1+\varepsilon)Z.$$



Discrepancy theory

Theorem

For symmetric convex body $K \subseteq [-1,1]^m$ with $\mathrm{Vol}_m(K) \geqslant c^m$, there is an $x \in s(c)K \cap [-1,1]^m$ with $|\{j \in [m] : |x_j| = 1\}| \geqslant \frac{m}{2}$.




- **Note:** x can be found in poly-time assuming separation oracle for K.

Discrepancy theory

Theorem

For symmetric convex body $K \subseteq [-1,1]^m$ with $\mathrm{Vol}_m(K) \geqslant c^m$, there is an $x \in s(c)K \cap [-1,1]^m$ with $|\{j \in [m] : |x_j| = 1\}| \geqslant \frac{m}{2}$.



- **Note:** x can be found in poly-time assuming separation oracle for K.

The overall strategy

(1) Consider

$$K = \left\{ x \in [-1,1]^m : \sum_{i=1}^{m} (1+x_i) \cdot |\langle a_i, y \rangle| \leqslant (1+\varepsilon) \sum_{i=1}^{m} |\langle a_i, y \rangle| \, \forall \, y \in \mathbb{R}^n \right\}$$

(2) We will show $\mathrm{Vol}_m(K) \geqslant c^m$.

The overall strategy

(1) Consider

$$K = \left\{ x \in [-1,1]^m : \sum_{i=1}^{m} (1+x_i) \cdot |\langle a_i, y \rangle| \leqslant (1+\varepsilon) \sum_{i=1}^{m} |\langle a_i, y \rangle| \, \forall \, y \in \mathbb{R}^n \right\}$$

(2) We will show $\mathrm{Vol}_m(K) \geqslant c^m$.

(3) Suppose

$$K \cap -K = \left\{ x \in [-1,1]^m : \left| \sum_{i=1}^{m} x_i \cdot |\langle a_i, y \rangle| \right| \leqslant \varepsilon \sum_{i=1}^{m} |\langle a_i, y \rangle| \, \forall \, y \in \mathbb{R}^n \right\}$$

also has $\mathrm{Vol}_m(K \cap -K) \geqslant \tilde{c}^m$.

(4) Then there is $x \in s(\tilde{c}) \cdot (K \cap -K)$ with $\frac{m}{4}$ coordinates $x_i = -1$.

(5) Reduce number of coordinates by $\frac{1}{4}$ at cost of $1 + O(\varepsilon)$ error $\rightarrow$ iterate

The overall strategy

(1) Consider

$$K = \left\{ x \in [-1,1]^m : \sum_{i=1}^{m} (1+x_i) \cdot |\langle a_i, y \rangle| \leqslant (1+\varepsilon) \sum_{i=1}^{m} |\langle a_i, y \rangle| \, \forall \, y \in \mathbb{R}^n \right\}$$

(2) We will show $\mathrm{Vol}_m(K) \geqslant c^m$.

GAP

(3) Suppose

$$K \cap -K = \left\{ x \in [-1,1]^m : \left| \sum_{i=1}^{m} x_i \cdot |\langle a_i, y \rangle| \right| \leqslant \varepsilon \sum_{i=1}^{m} |\langle a_i, y \rangle| \, \forall \, y \in \mathbb{R}^n \right\}$$

also has $\mathrm{Vol}_m(K \cap -K) \geqslant \tilde{c}^m$.

(4) Then there is $x \in s(\tilde{c}) \cdot (K \cap -K)$ with $\frac{m}{4}$ coordinates $x_i = -1$.


(5) Reduce number of coordinates by $\frac{1}{4}$ at cost of $1 + O(\varepsilon)$ error $\rightarrow$ iterate

Size of the symmetrizer (1)

Theorem

Let $K \subseteq [-1,1]^m$ be convex with $\mathrm{Vol}_S(K_S) \geqslant 2^{-\varepsilon^2 m} \cdot 2^{|S|}$ for all $S \subseteq [m]$ and $[-\varepsilon,\varepsilon]^m \subseteq K$. Then

$$\mathrm{Vol}_m(K \cap -K) \geqslant 2^{-5m}.$$




Size of the symmetrizer (1)

Theorem

Let $K \subseteq [-1,1]^m$ be convex with $\mathrm{Vol}_S(K_S) \geqslant 2^{-\varepsilon^2 m} \cdot 2^{|S|}$ for all $S \subseteq [m]$ and $[-\varepsilon,\varepsilon]^m \subseteq K$. Then

$$\mathrm{Vol}_m(K \cap -K) \geqslant 2^{-5m}.$$




Size of the symmetrizer (1)

Theorem

Let $K \subseteq [-1,1]^m$ be convex with $\mathrm{Vol}_S(K_S) \geqslant 2^{-\varepsilon^2 m} \cdot 2^{|S|}$ for all $S \subseteq [m]$ and $[-\varepsilon,\varepsilon]^m \subseteq K$. Then

$$\mathrm{Vol}_m(K \cap -K) \geqslant 2^{-5m}.$$




Size of the symmetrizer (1)

Theorem

Let $K \subseteq [-1,1]^m$ be convex with $\mathrm{Vol}_S(K_S) \geqslant 2^{-\varepsilon^2 m} \cdot 2^{|S|}$ for all $S \subseteq [m]$ and $[-\varepsilon,\varepsilon]^m \subseteq K$. Then


$$\mathrm{Vol}_m(K \cap -K) \geqslant 2^{-5m}.$$



Size of the symmetrizer (2)

Theorem (Milman, Pajor)


For any convex body $K \subseteq \mathbb{R}^m$ with **barycenter** at **0** one has $\mathrm{Vol}_m(K \cap -K) \geqslant 2^{-m}\mathrm{Vol}_m(K)$.



Size of the symmetrizer (2)

Theorem (Milman, Pajor)

For any convex body $K \subseteq \mathbb{R}^m$ with **barycenter** at **0** one has $\mathrm{Vol}_m(K \cap -K) \geqslant 2^{-m}\mathrm{Vol}_m(K)$.




Size of the symmetrizer (2)

Theorem (Milman, Pajor)

For any convex body $K \subseteq \mathbb{R}^m$ with **barycenter** at **0** one has $\mathrm{Vol}_m(K \cap -K) \geqslant 2^{-m}\mathrm{Vol}_m(K)$.

- Consider

$$f_K(z) := \mathrm{Vol}_m((K - z) \cap -(K - z))$$




Size of the symmetrizer (2)

Theorem (Milman, Pajor)

For any convex body $K \subseteq \mathbb{R}^m$ with **barycenter** at **0** one has $\mathrm{Vol}_m(K \cap -K) \geqslant 2^{-m}\mathrm{Vol}_m(K)$.

- Consider
$$f_K(z) := \mathrm{Vol}_m((K - z) \cap -(K - z))$$
- Then $f_K(\mathrm{bary}(K)) \geqslant 2^{-m}\mathrm{Vol}_m(K)$ by [Milman-Pajor]




Size of the symmetrizer (2)

Theorem (Milman, Pajor)

For any convex body $K \subseteq \mathbb{R}^m$ with **barycenter** at **0** one has $\mathrm{Vol}_m(K \cap -K) \geqslant 2^{-m}\mathrm{Vol}_m(K)$.


- Consider
$$f_K(z) := \mathrm{Vol}_m((K - z) \cap -(K - z))$$
- Then $f_K(\mathrm{bary}(K)) \geqslant 2^{-m}\mathrm{Vol}_m(K)$ by [Milman-Pajor]
- For any $z \in K \cap -K$ one has $\mathrm{Vol}_m(K \cap -K) \geqslant 2^{-m} f_K(z)$
$\left[\ \textbf{Proof.}\ \frac{1}{2}((K - z) \cap -(K - z)) \subseteq K \cap -K\ \right]$.



Size of the symmetrizer (3)

Lemma

There is a $z \in [-\varepsilon, \varepsilon]^m$ so that $f_K(z) \geqslant 2^{-4m}$.




Size of the symmetrizer (3)

Lemma

There is a $z \in [-\varepsilon, \varepsilon]^m$ so that $f_K(z) \geqslant 2^{-4m}$.

- Suppose there is no such z.




Size of the symmetrizer (3)

Lemma

There is a $z \in [-\varepsilon, \varepsilon]^m$ so that $f_K(z) \geqslant 2^{-4m}$.

- Suppose there is no such z.
- The set $T := \left\{ z \in K \mid f_K(z) \geqslant 2^{-4m} \right\}$ is convex (Brunn-Minkowski).




Size of the symmetrizer (3)

Lemma

There is a $z \in [-\varepsilon, \varepsilon]^m$ so that $f_K(z) \geqslant 2^{-4m}$.

- Suppose there is no such z.
- The set $T := \left\{ z \in K \mid f_K(z) \geqslant 2^{-4m} \right\}$ is convex (Brunn-Minkowski).
- By hyperplane separation there exists $a \in \mathbb{R}^m$ so that

$$\forall z \in T: \quad \langle a, z \rangle > \max_{x \in \varepsilon B_\infty^m} \langle a, x \rangle = \varepsilon \|a\|_1$$



Size of the symmetrizer (3)


Lemma

There is a $z \in [-\varepsilon, \varepsilon]^m$ so that $f_K(z) \geqslant 2^{-4m}$.

- Suppose there is no such z.
- The set $T := \left\{ z \in K \mid f_K(z) \geqslant 2^{-4m} \right\}$ is convex (Brunn-Minkowski).
- By hyperplane separation there exists $a \in \mathbb{R}^m$ so that

$$\forall z \in T: \quad \langle a, z \rangle > \max_{x \in \varepsilon B_\infty^m} \langle a, x \rangle = \varepsilon \|a\|_1$$


- Let $P := \{x \in K \mid \langle a, x \rangle \leqslant \varepsilon \|a\|_1\}$.



Size of the symmetrizer (4)

Claim


$\mathrm{Vol}_m(P) \geqslant 2^{-3m}$.



Size of the symmetrizer (4)

Claim

$\mathrm{Vol}_m(P) \geqslant 2^{-3m}$.



- With the claim,

$$\begin{aligned} f_K(\mathrm{bary}(P)) &\overset{P\subseteq K}{\geqslant} f_P(\mathrm{bary}(P)) \\ &\geqslant 2^{-m}\mathrm{Vol}_m(P) \\ &\geqslant 2^{-4m} \end{aligned}$$

- Then $\mathrm{bary}(P) \in T \to$ contradiction!


Size of the symmetrizer (5)

Claim

$\mathrm{Vol}_m(P) = \mathrm{Vol}_m(\{x \in K \mid \langle a, x\rangle \leqslant \varepsilon \|a\|_1\}) \geqslant 2^{-3m}$.

- Use concentration:


$$\Pr_{x \sim [-1,1]^m}[\langle a, x\rangle \geqslant \lambda \|a\|_2] \leqslant 2^{-2\lambda^2}$$



Size of the symmetrizer (5)

Claim

$\mathrm{Vol}_m(P) = \mathrm{Vol}_m(\{x \in K \mid \langle a, x\rangle \leqslant \varepsilon \|a\|_1\}) \geqslant 2^{-3m}$.



- Use concentration:

$$\Pr_{x \sim [-1,1]^m}[\langle a, x\rangle \geqslant \lambda \|a\|_2] \leqslant 2^{-2\lambda^2}$$

- W.l.o.g. $\|a\|_1 = 1$; for simplicity assume


$$a_i = \begin{cases} \pm\frac{1}{|S|} & \text{for } i \in S \\ 0 & \text{otherwise} \end{cases}$$

- $\lambda := \varepsilon\sqrt{m}$ gives $\Pr_{x \sim [-1,1]^S}\left[\langle a, x\rangle \geqslant \sqrt{\frac{m}{|S|}} \cdot \varepsilon\right] \leqslant 2^{-2\varepsilon^2 m} \leqslant \frac{1}{2}2^{-\varepsilon^2 m}$

Size of the symmetrizer (5)

Claim

$\mathrm{Vol}_m(P) = \mathrm{Vol}_m(\{x \in K \mid \langle a, x\rangle \leqslant \varepsilon \|a\|_1\}) \geqslant 2^{-3m}$.



- Use concentration:

$$\Pr_{x \sim [-1,1]^m}[\langle a, x\rangle \geqslant \lambda \|a\|_2] \leqslant 2^{-2\lambda^2}$$

- W.l.o.g. $\|a\|_1 = 1$; for simplicity assume


$$a_i = \begin{cases} \pm\frac{1}{|S|} & \text{for } i \in S \\ 0 & \text{otherwise} \end{cases}$$

- $\lambda := \varepsilon\sqrt{m}$ gives $\Pr_{x \sim [-1,1]^S}\left[\langle a, x\rangle \geqslant \sqrt{\frac{m}{|S|}} \cdot \varepsilon\right] \leqslant 2^{-2\varepsilon^2 m} \leqslant \frac{1}{2}2^{-\varepsilon^2 m}$

Size of the symmetrizer (5)

Claim

$\mathrm{Vol}_m(P) = \mathrm{Vol}_m(\{x \in K \mid \langle a, x\rangle \leqslant \varepsilon \|a\|_1\}) \geqslant 2^{-3m}$.



- Use concentration:

$$\Pr_{x\sim[-1,1]^m}[\langle a, x\rangle \geqslant \lambda\|a\|_2] \leqslant 2^{-2\lambda^2}$$

- W.l.o.g. $\|a\|_1 = 1$; for simplicity assume

$$a_i = \begin{cases} \pm\frac{1}{|S|} & \text{for } i \in S \\ 0 & \text{otherwise} \end{cases}$$

- $\lambda := \varepsilon\sqrt{m}$ gives $\Pr_{x\sim[-1,1]^S}\left[\langle a, x\rangle \geqslant \sqrt{\frac{m}{|S|}}\cdot\varepsilon\right] \leqslant 2^{-2\varepsilon^2 m} \leqslant \frac{1}{2}2^{-\varepsilon^2 m}$
- Then

$$Q := \left(\sqrt{\frac{|S|}{m}}\cdot\left\{x \in K_S \mid \langle a, x\rangle \leqslant \sqrt{\frac{m}{|S|}}\cdot\varepsilon\right\}\right) \times K_{[m]\setminus S} \subseteq 2P$$

- $\mathrm{Vol}_m(\frac{1}{2}Q) \geqslant 2^{-m}\left(\sqrt{\frac{|S|}{m}}\right)^{|S|}\cdot\frac{1}{2}2^{-\varepsilon^2 m}\cdot 2^{|S|}\cdot 2^{-\varepsilon^2 m}\cdot 2^{m-|S|} \geqslant 2^{-3m}$. □

The one-sided volume

Theorem

Let $K \subseteq [-1,1]^m$ be convex with $\mathrm{Vol}_S(K_S) \geqslant 2^{-\varepsilon^2 m} \cdot 2^{|S|}$ for all $S \subseteq [m]$ and $[-\varepsilon,\varepsilon]^m \subseteq K$. Then

$$\mathrm{Vol}_m(K \cap -K) \geqslant 2^{-5m}.$$

The one-sided volume

Theorem

Let $K \subseteq [-1,1]^m$ be convex with $\mathrm{Vol}_S(K_S) \geqslant 2^{-\varepsilon^2 m} \cdot 2^{|S|}$ for all $S \subseteq [m]$ and $[-\varepsilon,\varepsilon]^m \subseteq K$. Then

$$\mathrm{Vol}_m(K \cap -K) \geqslant 2^{-5m}.$$

- Recall

$$K = \left\{ x \in [-1,1]^m : \sum_{i=1}^{m} x_i \cdot |\langle a_i, y\rangle| \leqslant \varepsilon \sum_{i=1}^{m} |\langle a_i, y\rangle| \ \forall\, y \in \mathbb{R}^n \right\}$$

The one-sided volume

Theorem

Let $K \subseteq [-1,1]^m$ be convex with $\mathrm{Vol}_S(K_S) \geqslant 2^{-\varepsilon^2 m} \cdot 2^{|S|}$ for all $S \subseteq [m]$ and $[-\varepsilon,\varepsilon]^m \subseteq K$. Then

$$\mathrm{Vol}_m(K \cap -K) \geqslant 2^{-5m}.$$

- Recall

$$K = \left\{ x \in [-1,1]^m : \sum_{i=1}^m x_i \cdot |\langle a_i, y\rangle| \leqslant \varepsilon \sum_{i=1}^m |\langle a_i, y\rangle| \ \forall\, y \in \mathbb{R}^n \right\}$$

- How to lower bound $\mathrm{Vol}_m(K)$?

The one-sided volume

Theorem

Let $K \subseteq [-1,1]^m$ be convex with $\mathrm{Vol}_S(K_S) \geqslant 2^{-\varepsilon^2 m} \cdot 2^{|S|}$ for all $S \subseteq [m]$ and $[-\varepsilon,\varepsilon]^m \subseteq K$. Then

$$\mathrm{Vol}_m(K \cap -K) \geqslant 2^{-5m}.$$

- Recall

$$K = \left\{ x \in [-1,1]^m : \sum_{i=1}^m x_i \cdot |\langle a_i, y\rangle| \leqslant \varepsilon \sum_{i=1}^m |\langle a_i, y\rangle| \; \forall\, y \in \mathbb{R}^n \right\}$$

- How to lower bound $\mathrm{Vol}_m(K)$?
- It suffices to lower bound

$$\Pr_{x \in \{-1,1\}^m} \left[\sum_{i=1}^m x_i \cdot |\langle a_i, y\rangle| \leqslant 0 \; \forall\, y \in \mathbb{R}^n \right]$$

The one-sided volume

Theorem

Let $K \subseteq [-1,1]^m$ be convex with $\mathrm{Vol}_S(K_S) \geqslant 2^{-\varepsilon^2 m} \cdot 2^{|S|}$ for all $S \subseteq [m]$ and $[-\varepsilon,\varepsilon]^m \subseteq K$. Then

$$\mathrm{Vol}_m(K \cap -K) \geqslant 2^{-5m}.$$

- Recall

$$K = \left\{ x \in [-1,1]^m : \sum_{i=1}^m x_i \cdot |\langle a_i, y\rangle| \leqslant \varepsilon \sum_{i=1}^m |\langle a_i, y\rangle| \; \forall\, y \in \mathbb{R}^n \right\}$$

- How to lower bound $\mathrm{Vol}_m(K)$?
- It suffices to lower bound


$$\Pr_{x \in \{-1,1\}^m} \left[\sum_{i=1}^m x_i \cdot |\langle a_i, y\rangle| \leqslant 0 \; \forall\, y \in \mathbb{R}^n \right]$$

- Idea: find a function f so this is "$f(x) > 0$" and use Boolean analysis.

Minkowski subtraction

- For convex sets $K, L \subseteq \mathbb{R}^n$, the **Minkowski sum** is

$$K + L = \{x + y \mid x \in K, y \in L\}.$$




Minkowski subtraction

- For convex sets $K, L \subseteq \mathbb{R}^n$, the **Minkowski sum** is

$$K + L = \{x + y \mid x \in K, y \in L\}.$$

- For convex sets $K, L \subseteq \mathbb{R}^n$, the **Minkowski subtraction** is

$$K \ominus L := \{x \in \mathbb{R}^n \mid x + L \subseteq K\}.$$



Minkowski subtraction


- For convex sets $K, L \subseteq \mathbb{R}^n$, the **Minkowski sum** is

$$K + L = \{x + y \mid x \in K, y \in L\}.$$

- For convex sets $K, L \subseteq \mathbb{R}^n$, the **Minkowski subtraction** is

$$K \ominus L := \{x \in \mathbb{R}^n \mid x + L \subseteq K\}.$$


- Useful because: $K \ominus L \neq \emptyset \iff L \subseteq K$ for symmetric K, L.



Convexity of volume decrease

Lemma


Let $K \subseteq \mathbb{R}^n$ be convex and let $v \in \mathbb{R}^n$. The function $f(s) := \mathrm{Vol}_n(K \ominus s[-v, v])$ is convex for $s \in [0, \infty)$.



Convexity of volume decrease

Lemma

Let $K \subseteq \mathbb{R}^n$ be convex and let $v \in \mathbb{R}^n$. The function $f(s) := \mathrm{Vol}_n(K \ominus s[-v, v])$ is convex for $s \in [0, \infty)$.




- Let $\|v\|_2 = 1$. For $z \in v^\perp$, consider the line $z + \mathbb{R}v$.

Convexity of volume decrease

Lemma

Let $K \subseteq \mathbb{R}^n$ be convex and let $v \in \mathbb{R}^n$. The function $f(s) := \mathrm{Vol}_n(K \ominus s[-v, v])$ is convex for $s \in [0, \infty)$.




- Let $\|v\|_2 = 1$. For $z \in v^\perp$, consider the line $z + \mathbb{R}v$.
- Let $\ell(z)$ be length of intersection with K.

Convexity of volume decrease

Lemma

Let $K \subseteq \mathbb{R}^n$ be convex and let $v \in \mathbb{R}^n$. The function $f(s) := \mathrm{Vol}_n(K \ominus s[-v, v])$ is convex for $s \in [0, \infty)$.





- Let $\|v\|_2 = 1$. For $z \in v^\perp$, consider the line $z + \mathbb{R}v$.
- Let $\ell(z)$ be length of intersection with K.
- Intersection with $K \ominus s[-v, v]$ has length $(\ell(z) - 2s)_+$

Convexity of volume decrease

Lemma

Let $K \subseteq \mathbb{R}^n$ be convex and let $v \in \mathbb{R}^n$. The function $f(s) := \mathrm{Vol}_n(K \ominus s[-v, v])$ is convex for $s \in [0, \infty)$.





- Let $\|v\|_2 = 1$. For $z \in v^\perp$, consider the line $z + \mathbb{R}v$.
- Let $\ell(z)$ be length of intersection with K.
- Intersection with $K \ominus s[-v, v]$ has length $(\ell(z) - 2s)_+$
- Then

$$\mathrm{Vol}_n(K \ominus s[-v, v]) = \int_{v^\perp} \underbrace{(\ell(z) - 2s\|v\|_2)_+}_{\text{convex in } s} \, dz$$

is an average of convex functions.

Inclusion of random zonotopes

- Consider zonotope $Z = \{\sum_{i=1}^{m} y_i A_i : |y_i| \leqslant 1\}$
- For $t \in \mathbb{R}^m_{\geqslant 0}$, let $Z_t := \{\sum_{i=1}^{m} y_i A_i : |y_i| \leqslant t_i \ \forall i \in [m]\}$.

Inclusion of random zonotopes

- Consider zonotope $Z = \{\sum_{i=1}^{m} y_i A_i : |y_i| \leqslant 1\}$
- For $t \in \mathbb{R}^m_{\geqslant 0}$, let $Z_t := \{\sum_{i=1}^{m} y_i A_i : |y_i| \leqslant t_i \ \forall i \in [m]\}$.

Theorem

Draw $x_i \sim \{-1, 1\}$ independently for $i \sim [m]$. Then

$$\Pr[Z_{\mathbf{1}+x} \subseteq Z] \geqslant e^{-2n}.$$

- Define a function $f : [-1, 1]^m \to \mathbb{R}$ with

$$f(x) := \mathrm{Vol}_n(Z \ominus Z_{\mathbf{1}+x})$$

Inclusion of random zonotopes

- Consider zonotope $Z = \{\sum_{i=1}^{m} y_i A_i : |y_i| \leqslant 1\}$
- For $t \in \mathbb{R}^m_{\geqslant 0}$, let $Z_t := \{\sum_{i=1}^{m} y_i A_i : |y_i| \leqslant t_i \ \forall i \in [m]\}$.

Theorem

Draw $x_i \sim \{-1, 1\}$ independently for $i \sim [m]$. Then

$$\Pr[Z_{\mathbf{1}+x} \subseteq Z] \geqslant e^{-2n}.$$

- Define a function $f : [-1, 1]^m \to \mathbb{R}$ with

$$f(x) := \mathrm{Vol}_n(Z \ominus Z_{\mathbf{1}+x})$$

- Then (i) f is separately convex and (ii) $f(\lambda x) = \lambda^n f(x)$ for $\lambda \geqslant 0$.

Inclusion of random zonotopes

- Consider zonotope $Z = \{\sum_{i=1}^{m} y_i A_i : |y_i| \leqslant 1\}$
- For $t \in \mathbb{R}^m_{\geqslant 0}$, let $Z_t := \{\sum_{i=1}^{m} y_i A_i : |y_i| \leqslant t_i \ \forall i \in [m]\}$.

Theorem

Draw $x_i \sim \{-1, 1\}$ independently for $i \sim [m]$. Then

$$\Pr[Z_{\mathbf{1}+x} \subseteq Z] \geqslant e^{-2n}.$$

- Define a function $f : [-1, 1]^m \to \mathbb{R}$ with

$$f(x) := \mathrm{Vol}_n(Z \ominus Z_{\mathbf{1}+x})$$

- Then (i) f is separately convex and (ii) $f(\lambda x) = \lambda^n f(x)$ for $\lambda \geqslant 0$.

Lemma

Any f satisfying (i) and (ii) has $\Pr_{x \sim \{-1,1\}^m}[f(x) > 0] \geqslant e^{-2n}$.

- Uses Bonami's 1970 **hypercontractivity** theorem.

Finishing the argument

- It remains to note that $Z_{1+x} \subseteq Z$ is equivalent to

$$\sum_{i=1}^{m} x_i \cdot |\langle a_i, y \rangle| \leqslant \varepsilon \sum_{i=1}^{m} |\langle a_i, y \rangle| \; \forall \, y \in \mathbb{R}^n.$$

Finishing the argument

- It remains to note that $Z_{1+x} \subseteq Z$ is equivalent to

$$\sum_{i=1}^{m} x_i \cdot |\langle a_i, y \rangle| \leqslant \varepsilon \sum_{i=1}^{m} |\langle a_i, y \rangle| \; \forall \, y \in \mathbb{R}^n.$$

- Thus $\mathrm{Vol}_S(K_S) \geqslant c^n 2^{|S|} \geqslant c^{m\varepsilon^2} 2^{|S|}$ for every $S \subseteq [m]$.

Finishing the argument

- It remains to note that $Z_{1+x} \subseteq Z$ is equivalent to

$$\sum_{i=1}^{m} x_i \cdot |\langle a_i, y \rangle| \leqslant \varepsilon \sum_{i=1}^{m} |\langle a_i, y \rangle| \; \forall \, y \in \mathbb{R}^n.$$

- Thus $\mathrm{Vol}_S(K_S) \geqslant c^n 2^{|S|} \geqslant c^{m\varepsilon^2} 2^{|S|}$ for every $S \subseteq [m]$.
- We get $\mathrm{Vol}_m(K \cap -K) \geqslant 2^{-5m}$.

Finishing the argument

- It remains to note that $Z_{1+x} \subseteq Z$ is equivalent to

$$\sum_{i=1}^{m} x_i \cdot |\langle a_i, y \rangle| \leqslant \varepsilon \sum_{i=1}^{m} |\langle a_i, y \rangle| \; \forall \, y \in \mathbb{R}^n.$$

- Thus $\mathrm{Vol}_S(K_S) \geqslant c^n 2^{|S|} \geqslant c^{m\varepsilon^2} 2^{|S|}$ for every $S \subseteq [m]$.
- We get $\mathrm{Vol}_m(K \cap -K) \geqslant 2^{-5m}$.
- Find $x \in O(1) \cdot (K \cap -K)$ with $\geqslant \frac{m}{4}$ coordinates $x_i = -1$.

Finishing the argument

- It remains to note that $Z_{1+x} \subseteq Z$ is equivalent to

$$\sum_{i=1}^{m} x_i \cdot |\langle a_i, y \rangle| \leqslant \varepsilon \sum_{i=1}^{m} |\langle a_i, y \rangle| \; \forall \, y \in \mathbb{R}^n.$$

- Thus $\mathrm{Vol}_S(K_S) \geqslant c^n 2^{|S|} \geqslant c^{m\varepsilon^2} 2^{|S|}$ for every $S \subseteq [m]$.
- We get $\mathrm{Vol}_m(K \cap -K) \geqslant 2^{-5m}$.
- Find $x \in O(1) \cdot (K \cap -K)$ with $\geqslant \frac{m}{4}$ coordinates $x_i = -1$.
- Iterate on the remaining coordinates until $m \leqslant O(n/\varepsilon^2)$.

Reproving [BSS'08]

Theorem

For PSD matrices $A_1, \ldots, A_m \in \mathbb{R}^{n \times n}$ with $\sum_{i=1}^{m} A_i = B$ and $0 < \varepsilon \leqslant \frac{1}{2}$ there is $w \in \mathbb{R}^m_{\geqslant 0}$ with $|\mathrm{supp}(w)| \leqslant O\left(\frac{n}{\varepsilon^2}\right)$ so that

$$(1-\varepsilon)B \preceq \sum_{i=1}^{m} w_i A_i \preceq (1+\varepsilon)B$$

Reproving [BSS'08]

Theorem

For PSD matrices $A_1, \ldots, A_m \in \mathbb{R}^{n \times n}$ with $\sum_{i=1}^m A_i = B$ and $0 < \varepsilon \leqslant \frac{1}{2}$ there is $w \in \mathbb{R}^m_{\geqslant 0}$ with $|\mathrm{supp}(w)| \leqslant O\left(\frac{n}{\varepsilon^2}\right)$ so that

$$(1-\varepsilon)B \preceq \sum_{i=1}^m w_i A_i \preceq (1+\varepsilon)B$$

- The function

$$\Phi(t) := \begin{cases} \det(B - \sum_{i=1}^m t_i A_i) & \text{if } \sum_{i=1}^m t_i A_i \preceq B \\ 0 & \text{otherwise.} \end{cases}$$

  is separately convex for $t \geqslant \mathbf{0}$.

Reproving [BSS'08]

Theorem

For PSD matrices $A_1, \ldots, A_m \in \mathbb{R}^{n \times n}$ with $\sum_{i=1}^m A_i = B$ and $0 < \varepsilon \leqslant \frac{1}{2}$ there is $w \in \mathbb{R}^m_{\geqslant 0}$ with $|\mathrm{supp}(w)| \leqslant O\left(\frac{n}{\varepsilon^2}\right)$ so that

$$(1-\varepsilon)B \preceq \sum_{i=1}^m w_i A_i \preceq (1+\varepsilon)B$$

- The function

$$\Phi(t) := \begin{cases} \det(B - \sum_{i=1}^m t_i A_i) & \text{if } \sum_{i=1}^m t_i A_i \preceq B \\ 0 & \text{otherwise.} \end{cases}$$

  is separately convex for $t \geqslant \mathbf{0}$.
- Then

$$\Pr_{x \sim \{-1,1\}^m}\left[\sum_{i=1}^m x_i A_i \preceq 0\right] \geqslant c^n$$

Other results + Follow-up

- Can reprove [BSS'08] (+ poly-time algo)
- Optimal Sparsifiers for Abelian Caley graphs
  [Basu, Kothari, Meka, Tudose '26]
- For sum $C := C_1 + \cdots + C_m \subseteq \mathbb{R}^n$ of symmetric convex sets, there is $s \geqslant \mathbf{0}$ with $|\mathrm{supp}(s)| \leqslant O(\frac{n}{\varepsilon^2})$ so that

$$(1-\varepsilon)C \subseteq \sum_{i=1}^{m} s_i C_i \subseteq (1+\varepsilon)C$$

  [Basu, Brakensiek, Chen, Putterman, R., Zhang '26]
- $O(n/\varepsilon^2)$ hypergraph cut sparsifiers: $C_e = \mathrm{conv}\{\mathbf{1}_u - \mathbf{1}_v : u, v \in e\}$
- Can this be made algorithmic? What about ℓ_p for $p \in (1,2)$?

Other results + Follow-up

- Can reprove [BSS'08] (+ poly-time algo)
- Optimal Sparsifiers for Abelian Caley graphs
  [Basu, Kothari, Meka, Tudose '26]
- For sum $C := C_1 + \cdots + C_m \subseteq \mathbb{R}^n$ of symmetric convex sets, there is $s \geqslant \mathbf{0}$ with $|\mathrm{supp}(s)| \leqslant O(\frac{n}{\varepsilon^2})$ so that

$$(1-\varepsilon)C \subseteq \sum_{i=1}^{m} s_i C_i \subseteq (1+\varepsilon)C$$

  [Basu, Brakensiek, Chen, Putterman, R., Zhang '26]
- $O(n/\varepsilon^2)$ hypergraph cut sparsifiers: $C_e = \mathrm{conv}\{\mathbf{1}_u - \mathbf{1}_v : u, v \in e\}$
- Can this be made algorithmic? What about ℓ_p for $p \in (1,2)$?

Thanks for your attention

Fourier analysis

Lemma

Let $f : [-1,1]^m \to \mathbb{R}_{\geqslant 0}$ be (i) separately convex, (ii) $f(\rho x) = \rho^n f(x)$ for all $x \in \{-1,1\}^m$ and $0 \leqslant \rho \leqslant 1$ and (iii) $f \neq 0$. Then $\Pr_{x \sim \{-1,1\}^m}[f(x) > 0] \geqslant c^n$.

- Let $\alpha := \Pr_{x \sim \{-1,1\}^m}[f(x) > 0]$.
- Pick $\rho \in (0,1)$ and $p \in (1,2)$ with $\rho = \sqrt{p-1}$.
- For any $x \in \{-1,1\}^m$,

$$T_\rho f(x) = \mathop{\mathbb{E}}_{z \sim N_\rho(x)}[f(z)] \overset{(i)}{\geqslant} f\Big(\mathop{\mathbb{E}}_{z \sim N_\rho(x)}[z]\Big) = f(\rho x) \overset{(ii)}{=} \rho^n f(x) \quad (*)$$

- Then

$$\rho^n \|f\|_2 \overset{(*)}{\leqslant} \|T_\rho f\|_2 \overset{(**)}{\leqslant} \|f\|_p \overset{(***)}{\leqslant} \|f\|_2 \cdot \alpha^{\frac{1}{p}-\frac{1}{2}}$$

  $(**)$ = hypercontractivity, $(***)$ =Hölder.
- Rearrange for α. □