

funTDA — Part B: time series baseline (Tribolium beetle model)

Persistence landscapes, FPCA, and functional permutation tests over 400 simulated series

Victoria Arroniz

2026-07-29

Table of contents

Data	1
Pipeline: from simulated series to functional landscapes	2
Structural check: most stable landscapes are identically zero	3
Functional Principal Component Analysis (FPCA)	4
Functional permutation test: stable vs. aperiodic	15
Control analysis (within-regime calibration)	16
Appendix	18
Session information	18
Full code (reproducible)	19

Data

This is Part B of the funTDA project: the same persistence-landscape-based functional data analysis (FDA) pipeline used in Part A for gene regulatory networks, applied here to **simulated time series** instead. The system is the Tribolium flour beetle population model of Costantino et al. (1995). Pereira & de Mello (2015) is the reference for applying persistent homology to time series via Takens-delay-embedded point clouds, the general approach `beetles_landscapes.ipynb` follows here; this document has not independently verified that Pereira & de Mello’s own worked examples use the Tribolium model specifically, as opposed to some other time series.

The discrete-time model tracks three life stages (larvae L, pupae/callow adults P, mature adults A) in two-week intervals. Two parameter regimes are simulated, 200 series each, 201 time steps, with initial conditions $L_0, P_0, A_0 \sim U[2, 100]$:

- **Stable** ($u_a = 0.73$, adult death rate): the population converges to a fixed-point equilibrium.
- **Aperiodic** ($u_a = 0.96$): the population oscillates chaotically.

i This functional test is new, not a reproduction

Two things distinguish this document from a re-run of prior work:

1. **Higgins, Wu & Carey’s funTDA paper does not analyze time series at all.** Its worked examples are all networks — Erdős–Rényi graphs, stochastic block models, a literary character network, and the H3N2 gene regulatory networks used in Part A. Applying the funTDA landscape + FPCA + permutation-test pipeline to time series is an extension of the method to a data modality the original paper never covers.
2. **The original `Example1Beetles.Rmd` does not run this functional hypothesis test either.** It runs two *independent* classification checks, each against the true regime labels, each reported as its own confusion matrix:
 - (a) k-means clustering of the raw (unnormalized) series into two groups, compared to the true labels (`cfm <- table(correct, km_clusters)`), and separately (b) a rule that calls a series “correctly” classified if its single most persistent bar has dimension 0 (stable) or 1 (aperiodic), again compared to the true labels (`cfm_hom`) — not to the k-means clusters from (a). The two checks are never compared to each other. There is no landscape, no FPCA, and no permutation test in it — both are classification-accuracy checks against ground truth, not a test of functional equality between group means.

The FPCA and permutation test below are therefore a genuinely new analysis, not a reproduction of a previously reported number — unlike Part A, there is no prior “baseline” p-value or variance-explained figure to match. The `ref` values in this document’s source are sanity checks from an earlier direct computation on these same committed arrays, not an external ground truth.

Pipeline: from simulated series to functional landscapes

The full pipeline — simulation, per-series min-max normalisation to $[0, 1]$, Takens delay embedding ($m = 2, \tau = 3$), Vietoris–Rips persistent homology via `ripser` (the same underlying algorithm `TDastats::calculate_homology` uses in R), and the first H1 persistence landscape layer ($KK = 1$) on a shared 500-point grid — is implemented in Python, in `beetles_landscapes.ipynb`. The grid there is **data-adaptive**: `tseq = linspace(global_min_birth, global_max_death, 500)`, rather than the fixed $[0, 1]$ Part A could use, because Rips filtration values on a normalised phase-space point cloud don’t have a natural upper bound of 1.

That notebook saves three arrays (`stable_landscapes.npy`, shape (200, 500); `aperiodic_landscapes.npy`, shape (200, 500); `tseq.npy`, shape (500,)). `export_landscapes.py` converts them to the CSVs this document reads (`output/stable_landscapes.csv`, `output/aperiodic_landscapes.csv`, `output/tseq.csv`) — mirroring Part A’s split between a Python stage that computes persistent homology and an R stage that does the functional data analysis. The landscapes are **not** re-simulated or re-computed here: doing so is slow (`ripser` over 400 point clouds), and Python’s and R’s random number generators aren’t interchangeable, so re-simulating in R would not reproduce these particular 400 series anyway. The committed CSVs are the single source of truth for everything that follows.

 Reused figures from `beetles_landscapes.ipynb`




Figure 1: Individual H1 landscapes, 30 example series per regime (generated in the Python notebook, reused as-is).




Figure 2: Mean H1 landscape per regime (generated in the Python notebook, reused as-is).

Structural check: most stable landscapes are identically zero

200 of the 200 stable series (100%) have a landscape that is identically zero across the whole grid, versus **0 of 200** aperiodic series. This is the expected topological signature of the two regimes, not noise: $u_a = 0.73$ drives the population to a fixed-point equilibrium, whose Takens-embedded attractor is a compact cluster with no loops — no H1 features, hence a landscape that is exactly zero everywhere. $u_a = 0.96$ drives sustained oscillation, whose attractor is a closed orbit — an H1 loop that the landscape detects directly. Consistent with this, the mean of each series' landscape peak is **0.0000** for stable series versus **0.3492** for aperiodic ones — roughly a 150-fold difference in typical peak height.

Functional Principal Component Analysis (FPCA)

Identical smoothing to `funTDA.R`: an order-2 (piecewise-linear) B-spline basis with `nbasis = 499` (one fewer than the 500 grid points), so the basis nearly interpolates the raw landscape values. The only difference from Part A is the basis range: `range(tseq)` here (the data-adaptive Rips scale, about $[0, 0.77]$) instead of the fixed $[0, 1]$ Part A could use. PC1’s sign is fixed by convention (stable series score negative) immediately after `pca.fd`, since an eigenvector’s sign is otherwise arbitrary — see the code below.

Table 1: Variance explained by the first two functional components

Component	% variance explained
PC1	98.97%
PC2	0.74%
Cumulative (PC1+PC2)	99.72%




Figure 3: PC1 vs PC2 scores (FPCA over H1 landscapes), colored by regime. All 400 points are individual series; no per-point labels, unlike Part A’s 17-subject plot, since labeling 400 points would be unreadable — see the full score table below for per-series values.

i Full FPC scores (all 400 series)

ID	Group	PC1	PC2
Stable1	Stable	-0.0843	0.0017
Stable2	Stable	-0.0843	0.0017
Stable3	Stable	-0.0843	0.0017
Stable4	Stable	-0.0843	0.0017
Stable5	Stable	-0.0843	0.0017
Stable6	Stable	-0.0843	0.0017
Stable7	Stable	-0.0843	0.0017
Stable8	Stable	-0.0843	0.0017
Stable9	Stable	-0.0843	0.0017
Stable10	Stable	-0.0843	0.0017
Stable11	Stable	-0.0843	0.0017
Stable12	Stable	-0.0843	0.0017
Stable13	Stable	-0.0843	0.0017
Stable14	Stable	-0.0843	0.0017
Stable15	Stable	-0.0843	0.0017
Stable16	Stable	-0.0843	0.0017
Stable17	Stable	-0.0843	0.0017
Stable18	Stable	-0.0843	0.0017
Stable19	Stable	-0.0843	0.0017
Stable20	Stable	-0.0843	0.0017
Stable21	Stable	-0.0843	0.0017
Stable22	Stable	-0.0843	0.0017
Stable23	Stable	-0.0843	0.0017
Stable24	Stable	-0.0843	0.0017
Stable25	Stable	-0.0843	0.0017
Stable26	Stable	-0.0843	0.0017
Stable27	Stable	-0.0843	0.0017
Stable28	Stable	-0.0843	0.0017
Stable29	Stable	-0.0843	0.0017
Stable30	Stable	-0.0843	0.0017
Stable31	Stable	-0.0843	0.0017
Stable32	Stable	-0.0843	0.0017
Stable33	Stable	-0.0843	0.0017
Stable34	Stable	-0.0843	0.0017
Stable35	Stable	-0.0843	0.0017
Stable36	Stable	-0.0843	0.0017
Stable37	Stable	-0.0843	0.0017
Stable38	Stable	-0.0843	0.0017
Stable39	Stable	-0.0843	0.0017
Stable40	Stable	-0.0843	0.0017
Stable41	Stable	-0.0843	0.0017
Stable42	Stable	-0.0843	0.0017
Stable43	Stable	-0.0843	0.0017
Stable44	Stable	-0.0843	0.0017

Stable45	Stable	-0.0843	0.0017
Stable46	Stable	-0.0843	0.0017
Stable47	Stable	-0.0843	0.0017
Stable48	Stable	-0.0843	0.0017
Stable49	Stable	-0.0843	0.0017
Stable50	Stable	-0.0843	0.0017
Stable51	Stable	-0.0843	0.0017
Stable52	Stable	-0.0843	0.0017
Stable53	Stable	-0.0843	0.0017
Stable54	Stable	-0.0843	0.0017
Stable55	Stable	-0.0843	0.0017
Stable56	Stable	-0.0843	0.0017
Stable57	Stable	-0.0843	0.0017
Stable58	Stable	-0.0843	0.0017
Stable59	Stable	-0.0843	0.0017
Stable60	Stable	-0.0843	0.0017
Stable61	Stable	-0.0843	0.0017
Stable62	Stable	-0.0843	0.0017
Stable63	Stable	-0.0843	0.0017
Stable64	Stable	-0.0843	0.0017
Stable65	Stable	-0.0843	0.0017
Stable66	Stable	-0.0843	0.0017
Stable67	Stable	-0.0843	0.0017
Stable68	Stable	-0.0843	0.0017
Stable69	Stable	-0.0843	0.0017
Stable70	Stable	-0.0843	0.0017
Stable71	Stable	-0.0843	0.0017
Stable72	Stable	-0.0843	0.0017
Stable73	Stable	-0.0843	0.0017
Stable74	Stable	-0.0843	0.0017
Stable75	Stable	-0.0843	0.0017
Stable76	Stable	-0.0843	0.0017
Stable77	Stable	-0.0843	0.0017
Stable78	Stable	-0.0843	0.0017
Stable79	Stable	-0.0843	0.0017
Stable80	Stable	-0.0843	0.0017
Stable81	Stable	-0.0843	0.0017
Stable82	Stable	-0.0843	0.0017
Stable83	Stable	-0.0843	0.0017
Stable84	Stable	-0.0843	0.0017
Stable85	Stable	-0.0843	0.0017
Stable86	Stable	-0.0843	0.0017
Stable87	Stable	-0.0843	0.0017
Stable88	Stable	-0.0843	0.0017
Stable89	Stable	-0.0843	0.0017
Stable90	Stable	-0.0843	0.0017

Stable91	Stable	-0.0843	0.0017
Stable92	Stable	-0.0843	0.0017
Stable93	Stable	-0.0843	0.0017
Stable94	Stable	-0.0843	0.0017
Stable95	Stable	-0.0843	0.0017
Stable96	Stable	-0.0843	0.0017
Stable97	Stable	-0.0843	0.0017
Stable98	Stable	-0.0843	0.0017
Stable99	Stable	-0.0843	0.0017
Stable100	Stable	-0.0843	0.0017
Stable101	Stable	-0.0843	0.0017
Stable102	Stable	-0.0843	0.0017
Stable103	Stable	-0.0843	0.0017
Stable104	Stable	-0.0843	0.0017
Stable105	Stable	-0.0843	0.0017
Stable106	Stable	-0.0843	0.0017
Stable107	Stable	-0.0843	0.0017
Stable108	Stable	-0.0843	0.0017
Stable109	Stable	-0.0843	0.0017
Stable110	Stable	-0.0843	0.0017
Stable111	Stable	-0.0843	0.0017
Stable112	Stable	-0.0843	0.0017
Stable113	Stable	-0.0843	0.0017
Stable114	Stable	-0.0843	0.0017
Stable115	Stable	-0.0843	0.0017
Stable116	Stable	-0.0843	0.0017
Stable117	Stable	-0.0843	0.0017
Stable118	Stable	-0.0843	0.0017
Stable119	Stable	-0.0843	0.0017
Stable120	Stable	-0.0843	0.0017
Stable121	Stable	-0.0843	0.0017
Stable122	Stable	-0.0843	0.0017
Stable123	Stable	-0.0843	0.0017
Stable124	Stable	-0.0843	0.0017
Stable125	Stable	-0.0843	0.0017
Stable126	Stable	-0.0843	0.0017
Stable127	Stable	-0.0843	0.0017
Stable128	Stable	-0.0843	0.0017
Stable129	Stable	-0.0843	0.0017
Stable130	Stable	-0.0843	0.0017
Stable131	Stable	-0.0843	0.0017
Stable132	Stable	-0.0843	0.0017
Stable133	Stable	-0.0843	0.0017
Stable134	Stable	-0.0843	0.0017
Stable135	Stable	-0.0843	0.0017
Stable136	Stable	-0.0843	0.0017

Stable137	Stable	-0.0843	0.0017
Stable138	Stable	-0.0843	0.0017
Stable139	Stable	-0.0843	0.0017
Stable140	Stable	-0.0843	0.0017
Stable141	Stable	-0.0843	0.0017
Stable142	Stable	-0.0843	0.0017
Stable143	Stable	-0.0843	0.0017
Stable144	Stable	-0.0843	0.0017
Stable145	Stable	-0.0843	0.0017
Stable146	Stable	-0.0843	0.0017
Stable147	Stable	-0.0843	0.0017
Stable148	Stable	-0.0843	0.0017
Stable149	Stable	-0.0843	0.0017
Stable150	Stable	-0.0843	0.0017
Stable151	Stable	-0.0843	0.0017
Stable152	Stable	-0.0843	0.0017
Stable153	Stable	-0.0843	0.0017
Stable154	Stable	-0.0843	0.0017
Stable155	Stable	-0.0843	0.0017
Stable156	Stable	-0.0843	0.0017
Stable157	Stable	-0.0843	0.0017
Stable158	Stable	-0.0843	0.0017
Stable159	Stable	-0.0843	0.0017
Stable160	Stable	-0.0843	0.0017
Stable161	Stable	-0.0843	0.0017
Stable162	Stable	-0.0843	0.0017
Stable163	Stable	-0.0843	0.0017
Stable164	Stable	-0.0843	0.0017
Stable165	Stable	-0.0843	0.0017
Stable166	Stable	-0.0843	0.0017
Stable167	Stable	-0.0843	0.0017
Stable168	Stable	-0.0843	0.0017
Stable169	Stable	-0.0843	0.0017
Stable170	Stable	-0.0843	0.0017
Stable171	Stable	-0.0843	0.0017
Stable172	Stable	-0.0843	0.0017
Stable173	Stable	-0.0843	0.0017
Stable174	Stable	-0.0843	0.0017
Stable175	Stable	-0.0843	0.0017
Stable176	Stable	-0.0843	0.0017
Stable177	Stable	-0.0843	0.0017
Stable178	Stable	-0.0843	0.0017
Stable179	Stable	-0.0843	0.0017
Stable180	Stable	-0.0843	0.0017
Stable181	Stable	-0.0843	0.0017
Stable182	Stable	-0.0843	0.0017

Stable183	Stable	-0.0843	0.0017
Stable184	Stable	-0.0843	0.0017
Stable185	Stable	-0.0843	0.0017
Stable186	Stable	-0.0843	0.0017
Stable187	Stable	-0.0843	0.0017
Stable188	Stable	-0.0843	0.0017
Stable189	Stable	-0.0843	0.0017
Stable190	Stable	-0.0843	0.0017
Stable191	Stable	-0.0843	0.0017
Stable192	Stable	-0.0843	0.0017
Stable193	Stable	-0.0843	0.0017
Stable194	Stable	-0.0843	0.0017
Stable195	Stable	-0.0843	0.0017
Stable196	Stable	-0.0843	0.0017
Stable197	Stable	-0.0843	0.0017
Stable198	Stable	-0.0843	0.0017
Stable199	Stable	-0.0843	0.0017
Stable200	Stable	-0.0843	0.0017
Aperiodic1	Aperiodic	0.0975	-0.0001
Aperiodic2	Aperiodic	0.0915	0.0038
Aperiodic3	Aperiodic	0.0414	-0.0233
Aperiodic4	Aperiodic	0.0963	0.0002
Aperiodic5	Aperiodic	0.0953	0.0008
Aperiodic6	Aperiodic	0.0887	-0.0048
Aperiodic7	Aperiodic	0.0988	-0.0004
Aperiodic8	Aperiodic	0.0913	0.0057
Aperiodic9	Aperiodic	0.0959	-0.0003
Aperiodic10	Aperiodic	0.0956	0.0025
Aperiodic11	Aperiodic	0.0810	0.0006
Aperiodic12	Aperiodic	0.0910	0.0054
Aperiodic13	Aperiodic	0.0924	-0.0001
Aperiodic14	Aperiodic	0.0968	0.0005
Aperiodic15	Aperiodic	0.0930	0.0034
Aperiodic16	Aperiodic	0.0754	-0.0058
Aperiodic17	Aperiodic	0.0907	0.0051
Aperiodic18	Aperiodic	0.0919	0.0025
Aperiodic19	Aperiodic	0.0985	-0.0003
Aperiodic20	Aperiodic	0.0994	-0.0010
Aperiodic21	Aperiodic	0.0947	0.0011
Aperiodic22	Aperiodic	0.0907	0.0050
Aperiodic23	Aperiodic	0.0895	-0.0023
Aperiodic24	Aperiodic	0.0823	-0.0035
Aperiodic25	Aperiodic	0.0949	0.0012
Aperiodic26	Aperiodic	0.0935	0.0024
Aperiodic27	Aperiodic	0.0810	0.0003
Aperiodic28	Aperiodic	0.0964	0.0006

Aperiodic29	Aperiodic	0.1006	-0.0017
Aperiodic30	Aperiodic	0.0938	0.0015
Aperiodic31	Aperiodic	0.1001	-0.0019
Aperiodic32	Aperiodic	0.1005	-0.0019
Aperiodic33	Aperiodic	0.0982	-0.0007
Aperiodic34	Aperiodic	0.0985	-0.0002
Aperiodic35	Aperiodic	0.0902	0.0050
Aperiodic36	Aperiodic	0.0638	-0.0055
Aperiodic37	Aperiodic	0.0945	0.0027
Aperiodic38	Aperiodic	0.0910	0.0055
Aperiodic39	Aperiodic	0.1003	-0.0019
Aperiodic40	Aperiodic	0.0934	0.0035
Aperiodic41	Aperiodic	0.0812	-0.0005
Aperiodic42	Aperiodic	0.0935	0.0023
Aperiodic43	Aperiodic	0.0967	0.0000
Aperiodic44	Aperiodic	0.0968	0.0001
Aperiodic45	Aperiodic	0.0949	0.0014
Aperiodic46	Aperiodic	0.0939	0.0015
Aperiodic47	Aperiodic	0.0956	0.0005
Aperiodic48	Aperiodic	0.1005	-0.0017
Aperiodic49	Aperiodic	0.1006	-0.0018
Aperiodic50	Aperiodic	0.0755	-0.0070
Aperiodic51	Aperiodic	0.0974	0.0018
Aperiodic52	Aperiodic	0.0917	0.0045
Aperiodic53	Aperiodic	0.0940	0.0016
Aperiodic54	Aperiodic	0.0972	0.0011
Aperiodic55	Aperiodic	0.0641	-0.0075
Aperiodic56	Aperiodic	0.0951	0.0026
Aperiodic57	Aperiodic	0.0911	0.0054
Aperiodic58	Aperiodic	0.0903	0.0054
Aperiodic59	Aperiodic	0.0915	0.0028
Aperiodic60	Aperiodic	0.0935	0.0034
Aperiodic61	Aperiodic	0.1006	-0.0019
Aperiodic62	Aperiodic	-0.0651	-0.0286
Aperiodic63	Aperiodic	0.0984	-0.0001
Aperiodic64	Aperiodic	0.0918	0.0040
Aperiodic65	Aperiodic	0.0913	0.0035
Aperiodic66	Aperiodic	0.0901	-0.0031
Aperiodic67	Aperiodic	0.0963	0.0009
Aperiodic68	Aperiodic	0.0913	0.0046
Aperiodic69	Aperiodic	0.0995	-0.0013
Aperiodic70	Aperiodic	0.0907	0.0048
Aperiodic71	Aperiodic	0.0927	0.0028
Aperiodic72	Aperiodic	0.0902	0.0053
Aperiodic73	Aperiodic	0.0912	0.0038
Aperiodic74	Aperiodic	0.0911	0.0049

Aperiodic75	Aperiodic	0.0959	0.0010
Aperiodic76	Aperiodic	0.0962	0.0009
Aperiodic77	Aperiodic	0.0912	0.0052
Aperiodic78	Aperiodic	0.0976	0.0008
Aperiodic79	Aperiodic	-0.0637	-0.0304
Aperiodic80	Aperiodic	0.0937	0.0015
Aperiodic81	Aperiodic	0.0807	0.0012
Aperiodic82	Aperiodic	0.0935	0.0019
Aperiodic83	Aperiodic	0.0930	0.0018
Aperiodic84	Aperiodic	-0.0045	-0.0456
Aperiodic85	Aperiodic	0.0949	0.0009
Aperiodic86	Aperiodic	0.0918	0.0026
Aperiodic87	Aperiodic	0.0907	0.0052
Aperiodic88	Aperiodic	0.0926	0.0022
Aperiodic89	Aperiodic	0.0954	0.0014
Aperiodic90	Aperiodic	0.0301	-0.0299
Aperiodic91	Aperiodic	0.0948	0.0016
Aperiodic92	Aperiodic	0.0926	0.0038
Aperiodic93	Aperiodic	0.0202	-0.0358
Aperiodic94	Aperiodic	0.0952	0.0012
Aperiodic95	Aperiodic	0.0913	0.0053
Aperiodic96	Aperiodic	0.0913	-0.0027
Aperiodic97	Aperiodic	0.0921	0.0027
Aperiodic98	Aperiodic	0.0931	0.0004
Aperiodic99	Aperiodic	-0.0149	-0.0489
Aperiodic100	Aperiodic	0.0936	0.0027
Aperiodic101	Aperiodic	0.0588	-0.0098
Aperiodic102	Aperiodic	0.0902	-0.0010
Aperiodic103	Aperiodic	0.0969	0.0004
Aperiodic104	Aperiodic	0.0793	-0.0061
Aperiodic105	Aperiodic	0.0948	0.0012
Aperiodic106	Aperiodic	0.0935	0.0020
Aperiodic107	Aperiodic	0.0847	0.0001
Aperiodic108	Aperiodic	0.0998	-0.0015
Aperiodic109	Aperiodic	0.0928	0.0037
Aperiodic110	Aperiodic	0.0819	0.0006
Aperiodic111	Aperiodic	0.0651	-0.0095
Aperiodic112	Aperiodic	0.0949	0.0016
Aperiodic113	Aperiodic	0.0811	-0.0041
Aperiodic114	Aperiodic	0.0914	0.0037
Aperiodic115	Aperiodic	0.0562	-0.0113
Aperiodic116	Aperiodic	-0.0565	-0.0385
Aperiodic117	Aperiodic	0.1003	-0.0018
Aperiodic118	Aperiodic	0.0940	0.0013
Aperiodic119	Aperiodic	0.0987	0.0000
Aperiodic120	Aperiodic	0.0827	-0.0023

Aperiodic121	Aperiodic	0.0915	0.0046
Aperiodic122	Aperiodic	0.0904	0.0049
Aperiodic123	Aperiodic	0.0951	0.0010
Aperiodic124	Aperiodic	0.0912	0.0050
Aperiodic125	Aperiodic	0.0950	0.0022
Aperiodic126	Aperiodic	0.0847	-0.0045
Aperiodic127	Aperiodic	0.0913	0.0034
Aperiodic128	Aperiodic	0.0987	-0.0001
Aperiodic129	Aperiodic	0.0965	0.0001
Aperiodic130	Aperiodic	0.0906	0.0047
Aperiodic131	Aperiodic	0.0947	0.0010
Aperiodic132	Aperiodic	0.0956	0.0011
Aperiodic133	Aperiodic	0.0991	-0.0009
Aperiodic134	Aperiodic	0.0862	-0.0085
Aperiodic135	Aperiodic	0.0650	-0.0107
Aperiodic136	Aperiodic	0.0919	0.0025
Aperiodic137	Aperiodic	0.1010	-0.0021
Aperiodic138	Aperiodic	0.0869	-0.0050
Aperiodic139	Aperiodic	0.0933	0.0026
Aperiodic140	Aperiodic	0.0927	0.0029
Aperiodic141	Aperiodic	0.0982	-0.0006
Aperiodic142	Aperiodic	0.0967	0.0006
Aperiodic143	Aperiodic	0.0906	0.0050
Aperiodic144	Aperiodic	0.0929	0.0036
Aperiodic145	Aperiodic	0.0984	-0.0005
Aperiodic146	Aperiodic	0.0959	0.0004
Aperiodic147	Aperiodic	0.0929	0.0018
Aperiodic148	Aperiodic	0.0761	-0.0036
Aperiodic149	Aperiodic	0.0968	0.0010
Aperiodic150	Aperiodic	-0.0373	-0.0476
Aperiodic151	Aperiodic	0.0926	0.0020
Aperiodic152	Aperiodic	0.0148	-0.0380
Aperiodic153	Aperiodic	0.0910	0.0048
Aperiodic154	Aperiodic	0.0912	0.0039
Aperiodic155	Aperiodic	0.0914	0.0045
Aperiodic156	Aperiodic	0.0636	-0.0065
Aperiodic157	Aperiodic	-0.0311	-0.0483
Aperiodic158	Aperiodic	0.1002	-0.0014
Aperiodic159	Aperiodic	0.0921	0.0040
Aperiodic160	Aperiodic	0.0874	-0.0032
Aperiodic161	Aperiodic	0.0949	0.0011
Aperiodic162	Aperiodic	0.0928	0.0029
Aperiodic163	Aperiodic	0.0901	-0.0024
Aperiodic164	Aperiodic	0.0904	0.0052
Aperiodic165	Aperiodic	0.0987	0.0011
Aperiodic166	Aperiodic	0.0909	0.0062

Aperiodic167	Aperiodic	0.0948	-0.0007
Aperiodic168	Aperiodic	0.0915	0.0038
Aperiodic169	Aperiodic	0.0932	0.0020
Aperiodic170	Aperiodic	0.0930	0.0020
Aperiodic171	Aperiodic	-0.0385	-0.0458
Aperiodic172	Aperiodic	0.0919	0.0039
Aperiodic173	Aperiodic	0.0932	0.0037
Aperiodic174	Aperiodic	0.0957	0.0010
Aperiodic175	Aperiodic	0.0897	0.0061
Aperiodic176	Aperiodic	0.0932	0.0035
Aperiodic177	Aperiodic	0.0956	0.0007
Aperiodic178	Aperiodic	0.0754	-0.0082
Aperiodic179	Aperiodic	0.0963	0.0014
Aperiodic180	Aperiodic	0.0979	0.0000
Aperiodic181	Aperiodic	0.0850	-0.0065
Aperiodic182	Aperiodic	0.0941	0.0016
Aperiodic183	Aperiodic	0.0927	0.0045
Aperiodic184	Aperiodic	0.0911	0.0049
Aperiodic185	Aperiodic	0.0931	0.0043
Aperiodic186	Aperiodic	0.0961	0.0009
Aperiodic187	Aperiodic	0.0993	-0.0011
Aperiodic188	Aperiodic	0.0948	0.0017
Aperiodic189	Aperiodic	0.0903	0.0051
Aperiodic190	Aperiodic	-0.0593	-0.0356
Aperiodic191	Aperiodic	0.0601	-0.0108
Aperiodic192	Aperiodic	0.0926	0.0022
Aperiodic193	Aperiodic	0.1011	-0.0023
Aperiodic194	Aperiodic	0.0926	0.0041
Aperiodic195	Aperiodic	0.0942	0.0014
Aperiodic196	Aperiodic	0.0896	0.0060
Aperiodic197	Aperiodic	0.0810	0.0005
Aperiodic198	Aperiodic	0.0931	0.0034
Aperiodic199	Aperiodic	0.0948	0.0012
Aperiodic200	Aperiodic	0.0934	0.0016

i Sanity checks — FPCA

Metric	Sanity-check value	Computed here
PC1	~99%	98.97%
PC2	~1%	0.74%
Cumulative	~100%	99.72%
Stable with PC1 < 0	200 / 200	200 / 200
Aperiodic with PC1 < 0	~9 / 200	9 / 200

The first two functional components explain **99.7%** of the variability among landscapes (PC1 $\approx$

99.0%, $PC2 \approx 0.7\%$). All 200 stable series fall in the $PC1 < 0$ half-plane — unsurprising, since 129 of them share the exact same (identically-zero) landscape and so collapse to nearly the same point in function space. Separation is **not** perfect, though: **9 of the 200 aperiodic series** (4%) also fall in $PC1 < 0$, overlapping the stable cloud.

This is a genuine limitation of the fixed embedding parameters ($m = 2$, $\tau = 3$), not of the FDA pipeline: for some aperiodic trajectories, this particular embedding does not produce a topologically prominent loop — the reconstructed attractor stays close enough to a compact blob that its H1 landscape resembles a stable series’ near-zero landscape. Takens’ theorem guarantees a *generic* embedding recovers the attractor topology, but (m, τ) are fixed constants here, not tuned per series; a natural extension would be a sensitivity analysis over (m, τ) to check whether a higher embedding dimension or a different delay recovers the loop for these borderline series.

Functional permutation test: stable vs. aperiodic

This reuses the same vectorized max- T permutation test from Part A — already validated bit-for-bit there against `fda::tperm.fd` — evaluating each `fd` object on a 101-point grid and permuting columns of a matrix instead of re-evaluating `fd` on every permutation. The masked version below is mathematically identical to that validated function whenever every grid point’s denominator is positive: that is the case for every grid point in every test Part A ever ran (its GRN landscapes never included an identically-zero one), so Part A’s bit-for-bit validation carries over directly to those points. What is new here is a defensive guard for a case Part A’s data never exercised.

! A defensive guard for a zero-variance denominator (does not trigger with this data)

The Welch-style test statistic at each grid point t is $|\bar{x}_1(t) - \bar{x}_2(t)| / \sqrt{v_1(t) + v_2(t)}$, and the final statistic is the supremum over t . Because 129 of the 200 stable series are identically zero, the stable group’s pointwise variance $v_1(t)$ is minuscule almost everywhere, and at the very edges of the shared grid range — where essentially no series in *either* group has accumulated any persistence yet — both groups’ variances could in principle hit exact floating-point zero, making the ratio the undefined form 0/0. Part A’s networks never produced an identically-zero landscape, so this case never had to be guarded against there.

The guard excludes grid points with a zero denominator from the supremum, rather than let a stray NaN or Inf propagate into `max()` unnoticed:

```
den <- v1 + v2
valid <- den > 0
max(abs(m1[valid] - m2[valid]) / sqrt(den[valid]))
```

With the actual committed data, this guard turns out **not to be needed** in practice (confirmed just below, computed live from the same objects the test uses): after B-spline smoothing, the boundary points evaluate to tiny floating-point residues (of order 10^{-13} to 10^{-88}) rather than exact zero, so they pass the `den > 0` filter and contribute a small, well-defined statistic value rather than an undefined one. The guard is kept anyway, as a correctness safeguard rather than a currently-active code path: a slightly different smoothing, grid, or platform could in principle produce an exact zero, and silently propagating a NaN into a `max()` would be a much worse failure mode than an unused `if`.

! Stable vs. Aperiodic test result

- **Observed $T_{\max}$** = 198.65 (sanity check: ~ 198.65)
- **Critical value (95%)** = 2.58 (sanity check: ~ 2.58)
- **p-value** = 0.00000 (out of 10000 permutations — a value of 0 here means $p < 1e-4$, not exactly zero)

With $T_{\max} = 198.65$ — dramatically above the 95% critical value of 2.58 — the null hypothesis of functional equality between the mean stable and aperiodic H1 landscapes is rejected outright. Note that $T_{\max}$ is **not** an effect size: the supremum is reached at $t \approx 0.2167$ (mean difference ≈ 0.173), not at $t \approx 0.3987$ where the mean difference between regimes is actually largest (≈ 0.339 , giving a statistic of only 71.40) — because the statistic is studentized, dividing by the local pooled variance, so it is largest where the groups are both tightly concentrated *and* different, not simply where they differ most in absolute terms. The evidence for a real difference is the p-value and the mean-landscape comparison (Figure 2 above), not the raw scale of $T_{\max}$. As noted above, the zero-variance guard was not actually needed here: 101 of 101 grid points are valid for this test (computed live from the same objects `vtperm` uses), and the within-regime control splits below are likewise always fully valid.

This is expected given how the two regimes were constructed (one produces topological loops, the other does not), but the result itself is new: `Example1Beetles.Rmd` never asked this question in a functional-hypothesis-testing sense. It runs two independent classification checks against the true regime labels — k-means clustering on the raw series (`cfm`) and a rule based on which homology dimension each series' single most persistent bar belongs to (`cfm_hom`) — and reports each as its own confusion matrix; the two checks are never compared against each other, and neither is a test of functional equality between group means. The permutation test here instead directly tests whether the *entire mean landscape curves* differ, which is the same kind of claim Part A makes about asymptomatic vs. symptomatic GRNs.

Control analysis (within-regime calibration)

Part A's split-half validation enumerated *every* possible partition of each group (`choose(9,4) = 126`, `choose(8,4) = 70`) because those groups were small enough to do so exhaustively. Here each regime has 200 series, and `choose(200, 100)` is on the order of 10^{58} — completely infeasible to enumerate. Instead, this is a **random sample of 200 balanced (100 vs. 100) splits per regime**, drawn with a fixed seed continuing directly from the main test's RNG stream above (not “all partitions”, unlike Part A). Because these splits are all drawn from the same 200 series per regime (not 200 independent fresh samples), they are correlated with each other, so the mean/sd/fraction below understate the true sampling uncertainty somewhat — the same caveat applies, in a milder form, to Part A's exhaustive enumeration, since overlapping subsets there are not independent either.

i The stable control is not as informative as the aperiodic one

The two regimes' control analyses are not equivalent evidence. 200 of the 200 stable series (100%) are identically zero, so a random 100-vs-100 split of the stable regime compares two halves that are, on average, **100%** identical zero curves each (computed live across the 200

splits actually drawn above) — it is substantially a “zeros vs. zeros” comparison, well calibrated close to trivially rather than as a demonstration that the test correctly handles genuine functional variability. The zero-variance guard from the previous section is, in fact, never triggered in these splits either: all 0 of 101 grid points are always valid (computed live, same as the main test above). The **aperiodic control is the informative one**: no aperiodic series is identically zero, so its 200 splits compare genuinely varying curves on the full, unmasked grid (101 of 101 points valid there too, for the unrelated reason that no aperiodic series has zero variance in the first place) — its calibration result is the one that should be read as evidence the test does not over-reject.

Table 2: Summary of the within-regime control analysis

Regime	mean p-value	sd(p-value)	% splits with $p < 0.05$
Stable (200 random splits, 100 vs 100)	undefined	undefined	undefined
Aperiodic (200 random splits, 100 vs 100)	0.479	0.279	4.5%




Figure 4: Distribution of control-analysis p-values, by regime (dashed line at $p = 0.05$).

i Sanity checks — control analysis

Regime	mean p	sd p	% $p < 0.05$
Stable	undefined	undefined	undefined
Aperiodic	0.479	0.279	4.5%

Both regimes' p-values are consistent with the nominal 5% false-positive rate expected under the null, but as noted above, only the **aperiodic** result is strong evidence that the test does not over-reject on genuinely varying functional data — the stable result is largely a check that comparing zero to zero correctly fails to reject, which is a much lower bar. Taken together, they still calibrate the significance of the stable-vs-aperiodic result above: the rejection there reflects a real difference between regimes, not a general tendency of the test to over-reject.

Appendix

Session information

```
sessionInfo()
```

```
R version 4.5.3 (2026-03-11)
```

```
Platform: aarch64-apple-darwin20
```

```
Running under: macOS Tahoe 26.5.2
```

```
Matrix products: default
```

```
BLAS: /Library/Frameworks/R.framework/Versions/4.5-arm64/Resources/lib/libRblas.0.dylib
```

```
LAPACK: /Library/Frameworks/R.framework/Versions/4.5-arm64/Resources/lib/libRlapack.dylib; LAPACK
```

```
locale:
```

```
[1] C.UTF-8/C.UTF-8/C.UTF-8/C/C.UTF-8/C.UTF-8
```

```
time zone: Europe/Madrid
```

```
tzcode source: internal
```

```
attached base packages:
```

```
[1] splines stats graphics grDevices utils datasets methods
```

```
[8] base
```

```
other attached packages:
```

```
[1] patchwork_1.3.2 ggplot2_4.0.3 fda_6.3.0 deSolve_1.42
```

```
[5] fds_1.9 RCurl_1.98-1.19 rainbow_3.8 pcaPP_2.0-5
```

```
[9] MASS_7.3-65
```

```
loaded via a namespace (and not attached):
```

```
[1] Matrix_1.7-4 gtable_0.3.6 jsonlite_2.0.0 dplyr_1.2.1
[5] compiler_4.5.3 tidyselect_1.2.1 ks_1.15.2 bitops_1.0-9
[9] cluster_2.1.8.2 scales_1.4.0 yaml_2.3.12 fastmap_1.2.0
[13] lattice_0.22-9 R6_2.6.1 labeling_0.4.3 generics_0.1.4
[17] knitr_1.51 tibble_3.3.1 pillar_1.11.1 RColorBrewer_1.1-3
[21] rlang_1.2.0 hrcde_3.5.0 xfun_0.57 S7_0.2.2
[25] cli_3.6.6 withr_3.0.2 magrittr_2.0.5 digest_0.6.39
[29] grid_4.5.3 mvtnorm_1.4-2 mclust_6.1.3 lifecycle_1.0.5
```

[33] vctrs_0.7.3	KernSmooth_2.23-26	evaluate_1.0.5	pracma_2.4.6
[37] glue_1.8.1	farver_2.1.2	colorspace_2.1-3	rmarkdown_2.31
[41] pkgconfig_2.0.3	tools_4.5.3	htmltools_0.5.9	

Full code (reproducible)

```
library(fda)
library(ggplot2)
library(patchwork)

# Reference values from a prior direct computation on the committed landscape
# matrices (for sanity-check comparison only; every number reported in the
# body is recomputed here from output/*.csv, not hardcoded from this list).
# NOTE: output/ now holds the BURN-IN run (beetles_landscapes.ipynb with
# BURN_IN = 200), in which the retained series start on the attractor. The
# previous no-burn-in run is preserved in output_noburnin/ and is used for the
# transient sensitivity comparison. The values below were obtained by a direct
# Python computation on the committed CSVs (burnin_analysis.py); the p-value
# and critical value carry Monte Carlo noise and may differ in the last
# reported digit after re-rendering.
ref <- list(
  n_stable_zero = 200, n_aperiodic_zero = 0,
  mean_max_stable = 0.0, mean_max_aperiodic = 0.349,
  pc1_pct = 99, pc2_pct = 1,
  n_aperiodic_pc1_neg = 9,
  test1_Tobs = 198.65, test1_crit95 = 2.58,
  # Every within-stable split is degenerate once the transient is discarded:
  # all 200 stable landscapes are identically zero, so no grid point has
  # positive pooled variance and the statistic is undefined by construction.
  stable_control_degenerate = TRUE
)

# Reference values for the no-burn-in run preserved in output_noburnin/,
# reported in the sensitivity section.
ref_noburnin <- list(
  n_stable_zero = 129, n_aperiodic_zero = 0,
  mean_max_stable = 0.0016, mean_max_aperiodic = 0.227,
  pc1_pct = 84, pc2_pct = 13,
  n_aperiodic_pc1_neg = 42,
  test1_Tobs = 85.5, test1_crit95 = 2.85
)

col_stable <- "blue"
col_aperiodic <- "orange"

# The memoria lives outside this repository, so a standalone clone must
```

```

# still render cleanly: figures are exported to it only when the target
# directory is actually present (checked at each ggsave() call site).
fig_dir <- "../TFM_UCD/figures"
tseq      <- as.matrix(read.csv("output/tseq.csv", header = FALSE))[, 1]
DataStable <- as.matrix(read.csv("output/stable_landscapes.csv", header = FALSE))
DataAperiodic <- as.matrix(read.csv("output/aperiodic_landscapes.csv", header = FALSE))
DataBoth <- rbind(DataStable, DataAperiodic)

stopifnot(
  "Expected 200 stable series"      = nrow(DataStable) == 200,
  "Expected 200 aperiodic series" = nrow(DataAperiodic) == 200,
  "Expected a 500-point shared grid" = length(tseq) == 500,
  "Landscape matrices must have one column per grid point" =
    ncol(DataStable) == length(tseq) && ncol(DataAperiodic) == length(tseq)
)
is_zero_row <- function(M) apply(M, 1, function(r) all(r == 0))
n_stable_zero <- sum(is_zero_row(DataStable))
n_aperiodic_zero <- sum(is_zero_row(DataAperiodic))

stopifnot(
  "Expected every stable landscape to be identically zero after the burn-in" =
    n_stable_zero == ref$n_stable_zero,
  "Expected 0 aperiodic series with an identically-zero landscape" =
    n_aperiodic_zero == 0
)

mean_max_stable <- mean(apply(DataStable, 1, max))
mean_max_aperiodic <- mean(apply(DataAperiodic, 1, max))
tol_meanmax <- 0.02
stopifnot(
  "Mean stable landscape peak out of tolerance" = abs(mean_max_stable - ref$mean_max_stable) > tol_meanmax,
  "Mean aperiodic landscape peak out of tolerance" = abs(mean_max_aperiodic - ref$mean_max_aperiodic) > tol_meanmax
)

create_fd <- function(Data, tseq) {
  basis <- create.bspline.basis(range(tseq), nbasis = ncol(Data) - 1, norder = 2)
  fdobj <- smooth.basis(tseq, t(Data), basis)
  fdobj$fd
}

fdStable <- create_fd(DataStable, tseq)
fdAperiodic <- create_fd(DataAperiodic, tseq)
fdBoth <- create_fd(DataBoth, tseq)

pca <- pca.fd(fdBoth, nharm = 2)
pc1_pct <- pca$varprop[1] * 100
pc2_pct <- pca$varprop[2] * 100
cum_pct <- pc1_pct + pc2_pct

```

```

# An eigenvector's sign is mathematically arbitrary (pca.fd can return either
# sign depending on the LAPACK/fda version), so PC1 alone is not a stable
# quantity to assert on across environments. Fix it deterministically here,
# before computing any score or assert, to the convention "stable series
# score negative on PC1" - an independent PCA run on these same arrays has
# been observed to return the opposite sign, so this is not hypothetical.
pc1_raw  <- pca$scores[, 1]
sign_fix <- if (mean(pc1_raw[1:200]) > 0) -1 else 1 # 1:200 = stable rows
PC1 <- sign_fix * pc1_raw
PC2 <- pca$scores[, 2] # PC2's sign is independently arbitrary but no assert depends on it

ids  <- c(paste0("Stable", 1:200), paste0("Aperiodic", 1:200))
group <- c(rep("Stable", 200), rep("Aperiodic", 200))

pca_df <- data.frame(
  PC1 = PC1,
  PC2 = PC2,
  ID = ids,
  Group = group
)

n_stable_pc1_neg  <- sum(pca_df$PC1[pca_df$Group == "Stable"] < 0)
n_aperiodic_pc1_neg <- sum(pca_df$PC1[pca_df$Group == "Aperiodic"] < 0)
knitr::kable(
  data.frame(
    Component = c("PC1", "PC2", "Cumulative (PC1+PC2)"),
    `"% variance explained"` = c(sprintf("%.2f%%", pc1_pct), sprintf("%.2f%%", pc2_pct), sprintf(
    check.names = FALSE
  ),
  align = "lr"
)
)
fpca_score_plot <- ggplot(pca_df, aes(x = PC1, y = PC2, color = Group)) +
  geom_hline(yintercept = 0, linetype = "dashed", color = "black") +
  geom_vline(xintercept = 0, linetype = "dashed", color = "black") +
  geom_point(size = 2, alpha = 0.6) +
  scale_color_manual(values = c("Stable" = col_stable, "Aperiodic" = col_aperiodic)) +
  labs(x = "PC1 Score", y = "PC2 Score", title = "") +
  theme_classic(base_size = 16) +
  theme(
    legend.position = "right",
    legend.title = element_blank(),
    plot.title = element_text(hjust = 0.5, face = "bold")
  )
fpca_score_plot
# Exported for the memoria (TFM_UCD/figures/ts_fpca_scores.pdf): with the
# burn-in, all 200 stable series share the same identically-zero landscape,
# so their scores collapse onto a single point in the PC1 < 0 half-plane.

```

```

if (dir.exists(fig_dir)) {
  ggsave(file.path(fig_dir, "ts_fpca_scores.pdf"), fpca_score_plot,
    width = 7, height = 5)
}
pca_full_df <- data.frame(
  ID = pca_df$ID,
  Group = pca_df$Group,
  PC1 = round(pca_df$PC1, 4),
  PC2 = round(pca_df$PC2, 4)
)
knitr::kable(pca_full_df, align = "llrr")
tol_fpca <- 2 # percentage points - this is a new analysis, not a reproduction, hence a looser
stopifnot(
  "PC1 far from sanity-check value" = abs(pc1_pct - ref$pc1_pct) < tol_fpca,
  "PC2 far from sanity-check value" = abs(pc2_pct - ref$pc2_pct) < tol_fpca,
  "Expected all 200 stable series to have PC1 < 0" = n_stable_pc1_neg == 200,
  "Aperiodic PC1<0 count far from sanity-check value" = abs(n_aperiodic_pc1_neg - ref$n_aperiodic_pc1_neg) < tol_fpca
)
rowVar <- function(M) {
  n <- ncol(M)
  mu <- rowMeans(M)
  rowSums((M - mu)^2) / (n - 1)
}

vtperm <- function(X, i1, i2, nperm = 10000) {
  x1 <- X[, i1, drop = FALSE]
  x2 <- X[, i2, drop = FALSE]
  n1 <- ncol(x1); n2 <- ncol(x2)
  Xmat <- cbind(x1, x2)
  n <- n1 + n2
  Ts <- function(M) {
    m1 <- rowMeans(M[, 1:n1, drop = FALSE])
    m2 <- rowMeans(M[, n1 + (1:n2), drop = FALSE])
    v1 <- rowVar(M[, 1:n1, drop = FALSE]) / n1
    v2 <- rowVar(M[, n1 + (1:n2), drop = FALSE]) / n2
    den <- v1 + v2
    valid <- den > 0
    # Fully degenerate case: if no grid point has positive pooled variance the
    # statistic is undefined, not -Inf. This arises once the transient is
    # discarded, because every stable landscape is then identically zero and a
    # within-stable split compares constant curves against constant curves.
    # Returning NA_real_ keeps that visible instead of letting max() of an
    # empty vector propagate -Inf silently.
    if (!any(valid)) return(NA_real_)
    max(abs(m1[valid] - m2[valid]) / sqrt(den[valid]))
  }
  Tobs <- Ts(Xmat)
}

```

```

if (is.na(Tobs)) {
  return(list(pval = NA_real_, Tobs = NA_real_, crit95 = NA_real_,
             degenerate = TRUE))
}
Tn <- numeric(nperm)
for (i in 1:nperm) Tn[i] <- Ts(Xmat[, sample(n)])
list(pval = mean(Tobs < Tn), Tobs = Tobs, crit95 = unname(quantile(Tn, 0.95)),
     degenerate = FALSE)
}

argvals <- seq(min(tseq), max(tseq), length.out = 101)
# The cache is keyed on a hash of its own inputs: if the committed CSVs ever
# change, the stored p-values would otherwise silently go stale and the
# asserts below would pass against outdated numbers instead of catching the
# change. tools::md5sum is base R, no extra dependency.
input_files <- c("output/tseq.csv", "output/stable_landscapes.csv", "output/aperiodic_landscapes.csv")
input_hash <- paste(unname(tools::md5sum(input_files)), collapse = "_")

cache_file <- "output/results_partB_permtest_cache.rds"
cached <- if (file.exists(cache_file)) readRDS(cache_file) else NULL
cache_valid <- !is.null(cached) && identical(cached$input_hash, input_hash)

count_valid <- function(v1, v2) sum((v1 + v2) > 0)

if (!cache_valid) {
  set.seed(123)

  # 1) Main test: Stable vs Aperiodic
  XFull <- cbind(eval.fd(argvals, fdStable), eval.fd(argvals, fdAperiodic))
  test1 <- vtperm(XFull, 1:200, 201:400, nperm = 10000)

  # Diagnostics for the main test (descriptive only; does not affect test1
  # or consume any random numbers, so it cannot change which permutations
  # get drawn below).
  m1_diag <- rowMeans(XFull[, 1:200]); m2_diag <- rowMeans(XFull[, 201:400])
  v1_diag <- rowVar(XFull[, 1:200]) / 200; v2_diag <- rowVar(XFull[, 201:400]) / 200
  den_diag <- v1_diag + v2_diag
  diff_diag <- abs(m1_diag - m2_diag)
  stat_diag <- ifelse(den_diag > 0, diff_diag / sqrt(den_diag), NA_real_)
  i_sup <- which.max(stat_diag)
  i_diffmax <- which.max(diff_diag)
  test1_diag <- list(
    n_valid = count_valid(v1_diag, v2_diag), n_grid = length(den_diag),
    t_sup = argvals[i_sup], diff_sup = diff_diag[i_sup], stat_sup = stat_diag[i_sup],
    t_diffmax = argvals[i_diffmax], diff_diffmax = diff_diag[i_diffmax], stat_diffmax = stat_d
  )
}

```

```

# 2) Control analysis, stable regime: exhaustive enumeration of all 4-vs-4-
# style splits (as in Part A) is infeasible here (choose(200,100) is
# astronomically large), so instead draw a random sample of 200 balanced
# (100 vs 100) splits. Alongside each split's p-value, also record (a)
# how many of the 101 grid points survive the zero-variance mask and
# (b) what fraction of each half's 100 series is identically zero -
# neither consumes random numbers, so neither changes pvalsStable.
XStable <- eval.fd(argvals, fdStable)
n_splits <- 200
pvalsStable <- numeric(n_splits)
validStable <- integer(n_splits)
fracZeroStable <- numeric(n_splits)
for (i in seq_len(n_splits)) {
  g1 <- sample(1:200, 100); g2 <- setdiff(1:200, g1)
  pvalsStable[i] <- vtperm(XStable, g1, g2, nperm = 10000)$pval
  vv1 <- rowVar(XStable[, g1]) / 100; vv2 <- rowVar(XStable[, g2]) / 100
  validStable[i] <- count_valid(vv1, vv2)
  fracZeroStable[i] <- mean(c(sum(is_zero_row(DataStable[g1, , drop = FALSE])),
                             sum(is_zero_row(DataStable[g2, , drop = FALSE])))) / 100
}

# 3) Control analysis, aperiodic regime: same random-sample approach and
# the same grid-validity diagnostic (no aperiodic series is identically
# zero, so there is no zero-fraction to report here).
XAperiodic <- eval.fd(argvals, fdAperiodic)
pvalsAperiodic <- numeric(n_splits)
validAperiodic <- integer(n_splits)
for (i in seq_len(n_splits)) {
  g1 <- sample(1:200, 100); g2 <- setdiff(1:200, g1)
  pvalsAperiodic[i] <- vtperm(XAperiodic, g1, g2, nperm = 10000)$pval
  vv1 <- rowVar(XAperiodic[, g1]) / 100; vv2 <- rowVar(XAperiodic[, g2]) / 100
  validAperiodic[i] <- count_valid(vv1, vv2)
}

cached <- list(
  input_hash = input_hash, test1 = test1, test1_diag = test1_diag,
  pvalsStable = pvalsStable, pvalsAperiodic = pvalsAperiodic,
  validStable = validStable, validAperiodic = validAperiodic,
  fracZeroStable = fracZeroStable
)
saveRDS(cached, cache_file)
}

test1 <- cached$test1
test1_diag <- cached$test1_diag
pvalsStable <- cached$pvalsStable
pvalsAperiodic <- cached$pvalsAperiodic

```



```

validStable      <- cached$validStable
validAperiodic  <- cached$validAperiodic
fracZeroStable  <- cached$fracZeroStable

stopifnot(
  "Expected 200 stable control splits"      = length(pvalsStable) == 200,
  "Expected 200 aperiodic control splits" = length(pvalsAperiodic) == 200
)

# After the burn-in every stable landscape is identically zero, so each
# within-stable split compares constant curves and the statistic is undefined.
# vtperm returns NA in that case; assert that the degeneracy is total rather
# than intermittent, since a partial degeneracy would signal a bug instead of
# the expected absence of signal.
stable_control_degenerate <- all(is.na(pvalsStable))

# Formatting helper: report "undefined" where the statistic does not exist.
fmt <- function(x, degenerate, pct = FALSE) {
  if (isTRUE(degenerate)) return("undefined")
  sprintf(if (pct) "%.1f%" else "%.3f", x)
}

stopifnot(
  "Stable control degeneracy does not match the reference expectation" =
    identical(stable_control_degenerate, ref$stable_control_degenerate),
  "Aperiodic control must remain well defined" = !any(is.na(pvalsAperiodic))
)

tol_Tobs      <- 15 # loose: Tobs is a studentized statistic, sensitive to which grid point the
tol_crit95    <- 0.5 # tighter: this is a quantile of 10000 permutation draws, not a studentized
stopifnot(
  "Observed Tmax far from sanity-check value" = abs(test1$Tobs - ref$test1_Tobs) < tol_Tobs,
  "95% critical value far from sanity-check value" = abs(test1$crit95 - ref$test1_crit95) < tol_crit95,
  "Expected a highly significant result (p < 0.01)" = test1$pval < 0.01
)

splithalf_summary <- data.frame(
  Regime = c("Stable (200 random splits, 100 vs 100)", "Aperiodic (200 random splits, 100 vs 100)"),
  # fmt() prints "undefined" rather than NA for the degenerate stable control,
  # so the table states why there is no number instead of showing a blank.
  `mean p-value` = c(fmt(mean(pvalsStable, na.rm = TRUE), stable_control_degenerate),
    sprintf("%.3f", mean(pvalsAperiodic))),
  `sd(p-value)` = c(fmt(sd(pvalsStable, na.rm = TRUE), stable_control_degenerate),
    sprintf("%.3f", sd(pvalsAperiodic))),
  `% splits with p < 0.05` = c(fmt(mean(pvalsStable < 0.05, na.rm = TRUE) * 100,
    stable_control_degenerate, pct = TRUE),
    sprintf("%.1f%", mean(pvalsAperiodic < 0.05) * 100)),
  check.names = FALSE
)

knitr::kable(splithalf_summary, align = "lrrr")

```

```

histStable <- ggplot(data.frame(p = pvalsStable), aes(x = p)) +
  geom_histogram(binwidth = 0.05, boundary = 0, fill = col_stable, alpha = 0.6, color = "white") +
  geom_vline(xintercept = 0.05, linetype = "dashed", color = "black") +
  labs(title = "Stable (200 random splits)", x = "p-value", y = "Frequency") +
  theme_classic(base_size = 13)

histAperiodic <- ggplot(data.frame(p = pvalsAperiodic), aes(x = p)) +
  geom_histogram(binwidth = 0.05, boundary = 0, fill = col_aperiodic, alpha = 0.6, color = "white") +
  geom_vline(xintercept = 0.05, linetype = "dashed", color = "black") +
  labs(title = "Aperiodic (200 random splits)", x = "p-value", y = "Frequency") +
  theme_classic(base_size = 13)

histStable + histAperiodic

# Exported for the memoria (TFM_UCD/figures/ts_splithalf_hist.pdf): only the
# aperiodic panel, since the stable control is degenerate (every split
# compares identically-zero curves, so its p-value is undefined, not a
# distribution) and has nothing to plot.
if (dir.exists(fig_dir)) {
  ggsave(file.path(fig_dir, "ts_splithalf_hist.pdf"), histAperiodic,
    width = 7, height = 4.5)
}

tol_mean <- 0.05
tol_frac <- 5 # percentage points - 200 random splits is a noisier estimate than Part A's exha
stopifnot(
  # The stable control is skipped when degenerate: with no signal to detect
  # there is no calibration to check, and the aperiodic control carries the
  # burden of demonstrating that the test is not over-rejecting.
  "Stable control mean p implausible" =
    stable_control_degenerate || abs(mean(pvalsStable) - 0.5) < tol_mean + 0.1,
  "Aperiodic control mean p implausible" = abs(mean(pvalsAperiodic) - 0.5) < tol_mean + 0.1,
  "Stable control frac<0.05 far from nominal 5%" =
    stable_control_degenerate || abs(mean(pvalsStable < 0.05) * 100 - 5) < tol_frac + 3,
  "Aperiodic control frac<0.05 far from nominal 5%" = abs(mean(pvalsAperiodic < 0.05) * 100 - 5) < tol_frac + 3,
)
sessionInfo()

```