

Free-NVT brackets: highest T solid / lowest T melted in 6-ps runs of a perfect crystal (superheating possible: upper bounds). TDI melting H2O from the liquid-fcc G crossing (400, 500 GPa only). fcc-bcc: squares = bcc lower G, dots = fcc lower G; near melting the two are within 1-2 sigma. GRACE dg points: insertion of 1 H (some without the pressure correction). Toy (fitted pair potential, errors 40 meV/atom): trends only; its ice melts lower than GRACE's.



Free-NVT brackets: highest T solid / lowest T melted in 6-ps runs of a perfect crystal (superheating possible: upper bounds). TDI melting H2O from the liquid-fcc G crossing (400, 500 GPa only). fcc-bcc: squares = bcc lower G, dots = fcc lower G; near melting the two are within 1-2 sigma. GRACE dg points: insertion of 1 H (some without the pressure correction). Toy (fitted pair potential, errors 40 meV/atom): trends only; its ice melts lower than GRACE's.



Free-NVT brackets: highest T solid / lowest T melted in 6-ps runs of a perfect crystal (superheating possible: upper bounds). TDI melting H2O from the liquid-fcc G crossing (400, 500 GPa only). fcc-bcc: squares = bcc lower G, dots = fcc lower G; near melting the two are within 1-2 sigma. GRACE dg points: insertion of 1 H (some without the pressure correction). Toy (fitted pair potential, errors 40 meV/atom): trends only; its ice melts lower than GRACE's.