Barcode No	89781009	Lab No	16952412190004
Patient Name	Mr.LAKHVINDER SINGH	Reg Date	19/Dec/2024 04:35PM
Age/Sex	32 YRS/Male	Sample Coll. Date	19/Dec/2024 03:38 PM
Refered By	SELF	Sample Rec.Date	20/Dec/2024 07:04 AM
		Report Date	20/Dec/2024 10:40AM

IMMUNOASSAY

Test Name With Methodology	Result	Unit	Biological Ref.Interval
C-Peptide-F			
C-Peptide Serum, CLIA	0.94	ng/mL	0.92-3.73

Comments:

Comment: C-peptde is useful in distinguishing insulinomas from exogenous insulin administration. When insulin secretion is diminished, as in insulin dependent diabetes, low cpeptide levels are to be expected. Elevated c-peptide levels may result from increased beta cell activity associated with insulinomas. C-Peptide is also useful in monitoring patients who have received islet cell or pancreatic transplants.

C-peptide orginates in pancreatic beta cells as an inert byproduct in the synthesis of insulin from proinsulin. Insulin and c-peptide are released from proinsulin in equimolar concentration into the circulation. C-peptide levels can therefore serve as an index of insulin secretion. Anti-insulin antibodies are commonly found in patients who have underfore insulin therepy. These antibodies may interfere with insulin assay. C-peptide measurments are therefore used as an alternative measurment index in this context.

Prashant.
Dr. Prashant Goyal *(DCP)*
(Director & Chief Pathologist)
Reg. No. DMC-53016