

Barcode No	87139729	Lab No	16952408100011
Patient Name	Mrs.RANI DEVI	Reg Date	10/Aug/2024 07:37PM
Age/Sex	60 YRS/Female	Sample Coll. Date	10/Aug/2024 06:42 PM
Referred By	DR.PALVI	Sample Rec.Date	11/Aug/2024 08:11 AM
		Report Date	11/Aug/2024 10:25AM

.IMMUNO BIOCHEMISTRY-1

Test Name With Methodology	Result	Unit	Biological Ref.Interval
PTH-Intact (Parathormone Intact)			
Parathyroid Hormone (PTH)	126.2	pg/ml	15-65.0
Serum/ EDTA Plasma, ECLIA			

Comment:


Parathyroid hormone (PTH) produced by the parathyroid gland, is the major circulating factor regulating extra cellular calcium concentration. Quantification of circulating intact PTH assists in the difference diagnosis of hypercalcemia in conjunction with the measurement of Ionised calcium, intact PTH valuations can be used to distinguish between patients with hypoparathyroidism, hypoparathyroidism or hypercalcemia of malignancy. The diagnosis of primary hyperparathyroidism, a common cause of hyper calcemias confirmed by elevated ionized calcium concentrations and elevated parathyroid hormone concentrations. Intact PTH levels are also used to assess and manage other metabolic bone disorders including osteoporosis and renal osteodystrophy.

It is recommended that the intact PTH results should always be interpreted with caution & with consideration of the overall manifestations even when used in conjunction with calcium and phosphorus values. Measurement of intact PTH is useful in differentiation between hypercalcemia due to hyperparathyroidism and hypercalcemia of malignancy. However the assay is not intended as and should not be relied upon as a diagnostic indicator of malignancy.

Elevated PTH with normal serum calcium levels may be indicative of Secondary causes of hyperparathyroidism like vitamin D deficiency. It may not always be indicative of Primary hyperparathyroidism. PTH is secreted in a pulsatile manner with an overall circadian rhythm characterized by a nocturnal rise.

Note: Because of the short half-life of PTH, it is recommended that, when serum is needed, the blood should be centrifuged immediately. Preference should be given to EDTA plasma, as it is stable longer than serum.


 Dr. Anuja Adarsh, MBBS, MD
 (Consultant Biochemist)
 Reg. No. DMC-90705


 Dr. Prashant Goyal (DCP)
 (Director & Chief Pathologist)
 Reg. No. DMC-53016

Page 1 of 1

A Unit Of Hexamed Healthcare Services LLP

Corporate Office: Flat No 203, 2nd Floor, Sagacity Heights, Plot No 27, Kavuri Hills Rd, CBI Colony, Madhapur, Telangana 500033
Jammu Address: #25-270/6, Main Bypass Road, NH-1A, Channi Rama, Jammu, J & K 180015

✉ feedback@hexamedhealthcare.com @ enquiry@hexamedhealthcare.com 📧 support@hexamedhealthcare.com

☎ +91 95419 01145

☎ 1800 212 6547

🌐 www.hexamedhealthcare.com



Scan QR for Location



Scan QR for Enquiry Form