

Barcode No	87943592	Lab No	16952411050005
Patient Name	Ms.MISBHA	Reg Date	05/Nov/2024 05:12PM
Age/Sex	17 YRS/Female	Sample Coll. Date	05/Nov/2024 04:14 PM
Referred By	DR. ADNAN RAFIQ	Sample Rec.Date	06/Nov/2024 07:53 AM
Report Date		06/Nov/2024 12:18PM	

IMMUNOASSAY

Test Name With Methodology	Result	Unit	Biological Ref.Interval
TTG- IgA (Tissue Transglutaminase IgA Antibody)			
Tissue Transglutaminase Ab IgA Serum, ELISA	1.15	U/mL	<4.0 Negative 4-10 Weak Positive >10 Positive


Comments:

Useful for Evaluating patients suspected of having celiac disease, as Screening test for dermatitis herpetiformis, in conjunction with endomysial antibody test & Monitoring adherence to gluten-free diet in patients with dermatitis herpetiformis and celiac disease. A definitive diagnosis of celiac disease requires a jejunal biopsy demonstrating villous atrophy. Those individuals with positive laboratory results should be referred for small intestinal biopsy to confirm the diagnosis. If patients strictly adhere to a gluten-free diet, the unit value of IgA-anti-tTG should begin to decrease within 6 to 12 months of onset of dietary therapy.

CAUTIONS

This test should not be solely relied upon to establish a diagnosis of celiac disease. It should be used to identify patients who have an increased probability of having celiac disease and in whom a small intestinal biopsy is recommended. Affected individuals who have been on a gluten-free diet prior to testing may have a negative result. For individuals who test negative, IgA deficiency should be considered. If total IgA is normal and tissue transglutaminase (tTG)-IgA is negative, there is a low probability of the patient having celiac disease and a biopsy may not be necessary.

If serology is negative or there is substantial clinical doubt remaining, then further investigation should be performed with endoscopy and bowel biopsy. This is especially important in patients with frank malabsorptive symptoms since many syndromes can mimic celiac disease. For the patient with frank malabsorptive symptoms, bowel biopsy should be performed regardless of serologic test results.


 Dr. Prashant Goyal (DCP)
 (Director & Chief Pathologist)
 Reg. No. DMC-53016

Page 1 of

A Unit Of Hexamed Healthcare Services LLP

Corporate Office: Flat No 203, 2nd Floor, Sagacity Heights, Plot No 27, Kavuri Hills Rd, CBI Colony, Madhapur, Telangana 500033

Jammu Address: #25-270/6, Main Bypass Road, NH-1A, Channi Rama, Jammu, J & K 180015

✉ feedback@hexamedhealthcare.com @ enquiry@hexamedhealthcare.com 🗨 support@hexamedhealthcare.com

☎ +91 95419 01145

☎ 1800 212 6547

🌐 www.hexamedhealthcare.com



QR for Location



Scan QR for Enquiry Form