




Barcode No	87943635	Lab No	16952411160003
Patient Name	Ms.SAPNA	Reg Date	16/Nov/2024 04:10PM
Age/Sex	17 YRS/Female	Sample Coll. Date	16/Nov/2024 03:14 PM
Referred By	DR. SURINDER KUMAR MD	Sample Rec.Date	17/Nov/2024 08:57 AM
		Report Date	17/Nov/2024 12:08PM

.IMMUNO BIOCHEMISTRY-1

Test Name With Methodology	Result	Unit	Biological Ref.Interval
Insulin Fasting Insulin Fasting Serum, Electro Chemi Luminescent Immuno Assay	15.59	uIU/mL	2.6-24.9

Comment:

This test is used to estimate the levels of Insulin in serum. This may be used to diagnose insulinomas and to help diagnose acute or chronic (fasting) hypoglycemia. Insulin levels may also be used to monitor endogenous insulin, check for insulin resistance, and to help determine when a type 2 diabetic might need to start taking insulin injections to supplement oral medications.


Dr. Prashant Goyal (DCP)
(Director & Chief Pathologist)

Reg. No. DMC-53016

A Unit Of Hexamed Healthcare Services LLP

Corporate Office: Flat No 203, 2nd Floor, Sagacity Heights, Plot No 27, Kavuri Hills Rd, CBI Colony, Madhapur, Telangana 500033

Jammu Address: #25-270/6, Main Bypass Road, NH-1A, Channi Rama, Jammu, J & K 180015

✉ feedback@hexamedhealthcare.com @ enquiry@hexamedhealthcare.com 📞 support@hexamedhealthcare.com

☎ +91 95419 01145

☎ 1800 212 6547

🌐 www.hexamedhealthcare.com



Scan QR for Location



Scan QR for Enquiry Form




Barcode No	87943635	Lab No	16952411160003
Patient Name	Ms.SAPNA	Reg Date	16/Nov/2024 04:10PM
Age/Sex	17 YRS/Female	Sample Coll. Date	16/Nov/2024 03:14 PM
Referred By	DR. SURINDER KUMAR MD	Sample Rec.Date	17/Nov/2024 08:57 AM
		Report Date	17/Nov/2024 11:34AM

Test Name With Methodology	Result	Unit	Biological Ref.Interval
Testosterone- Free			
Free Testosterone Serum, CLIA	2.12	pg/mL	0.1-4.899

SUMMARY AND EXPLANATION OF THE TEST

Testosterone is a male sex hormone, secreted by leydig or interstitial cells of the testes. The amount of testosterone synthesized is regulated and controlled through negative feedback on the hypothalamus and pituitary gland by the pituitary hormone and luteinizing hormone (LH). Testosterone circulating in the blood binds to three proteins: sex hormone binding globulin (SHBG, 60-80%), albumin and cortisol binding globulin'. About 1-2% of the total circulating testosterone remains unbound or free. Only measurement of free testosterone permits the estimating of the biologically active hormone. Free testosterone determinations are recommended to overcome the influences caused by variations in transport proteins on the total testosterone concentration SHBG-bound testosterone remains in the circulation with no binding function in this form. Higher testosterone and lower SHBG levels can increase free testosterone. Free testosterone can be diagnostically useful when testosterone does not correspond with the clinical presentation of hypogonadism, especially in aging men with borderline low levels of testosterone and men in whom levels of SHBG are suspected to be altered?

Measuring free testosterone is an indirect way to measure a far more important health indicator-the level of SHBG in the bloodstream. Free testosterone measurements have been very helpful in monitoring Alzheimer's disease for older men and women. Lower levels of free testosterone may have a higher risk of developing Alzheimer's disease.


Dr. Prashant Goyal (DCP)
(Director & Chief Pathologist)

Reg. No. DMC 53010

A Unit Of Hexamed Healthcare Services LLP

Corporate Office: Flat No 203, 2nd Floor, Sagacity Heights, Plot No 27, Kavuri Hills Rd, CBI Colony, Madhapur, Telangana 500033

Jammu Address: #25-270/6, Main Bypass Road, NH-1A, Channi Rama, Jammu, J & K 180015

✉ feedback@hexamedhealthcare.com @ enquiry@hexamedhealthcare.com 📧 support@hexamedhealthcare.com

☎ +91 95419 01145

☎ 1800 212 6547

🌐 www.hexamedhealthcare.com



Scan QR for Location



Scan QR for Enquiry Form