



|                   |                  |                         |                     |                            |                  |
|-------------------|------------------|-------------------------|---------------------|----------------------------|------------------|
| <b>Name</b>       | Mrs.ANITA CHARAK | <b>Sample Received</b>  | 08-12-2024 10:19 AM | <b>Sample Collected At</b> | Health Sure Labs |
| <b>Age/Gender</b> | 61 Y / Female    | <b>Sample Processed</b> | 08-12-2024 10:19 AM | <b>Referred By</b>         | Dr Rajeev Bhasin |
| <b>UID</b>        | HLD0812202445556 | <b>Approved Date</b>    | 08-12-2024 03:09 PM |                            |                  |

## LABORATORY REPORT

### \*HBA1C, GLYCATED HEMOGLOBIN

**RESULT : 7.7 %**

#### Interpretation

HbA1c result is suggestive of non diabetic adults ( $\geq 18$  years)/ well controlled Diabetes in a known Diabetic.

#### Interpretation as per American Diabetes Association (ADA) Guidelines

| Reference Group   | Non diabetic adults $\geq 18$ years | At risk (Prediabetes) | Diagnosing Diabetes | Therapeutic goals for glycemic control |
|-------------------|-------------------------------------|-----------------------|---------------------|----------------------------------------|
| <b>HbA1c in %</b> | 4.0-5.6                             | 5.7-6.4               | $\geq 6.5$          | $< 7.0$                                |

Note: Presence of Hemoglobin variants and/or conditions that affect red cell turnover must be considered, particularly when the HbA1C result does not correlate with the patient's blood glucose levels

| FACTORS THAT INTERFERE WITH HbA1C MEASUREMENT                                                                                                                                                   | FACTORS THAT AFFECT INTERPRETATION OF HBA1C RESULTS                                                                                                                                                                                                                                                |
|-------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------|----------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------|
| Hemoglobin variants, elevated fetal hemoglobin (HbF) and chemically modified derivatives of e.g., carbamylated Hb in patients with renal failure) can affect the accuracy of HbA1c measurements | Any condition that shortens erythrocyte survival or decreases mean erythrocyte age (e.g., recovery from acute blood loss hemolytic anemia, HbSS, HbCC, and HbSC) will falsely lower HbA1c test results regardless of the assay method used. Iron deficiency anemia is associated with higher HbA1c |

The biological reference range pertains to the age and sex of the Patient.

**Note:** As the reports are computer generated, it must be treated as an original.

**Dr. Anuja Sharma**  
 Consultant Pathologist  
 Reg.No. 455

## A Unit Of Hexamed Healthcare Services LLP

**Corporate Office:** Flat No 203, 2nd Floor, Sagacity Heights, Plot No 27, Kavuri Hills Rd, CBI Colony, Madhapur, Telangana 500033

**Jammu Address:** 25-270/6, Main Bypass Road, NH-1A, Channi Rama, Jammu, J & K 180015

✉ feedback@hexamedhealthcare.com

@ enquiry@hexamedhealthcare.com

📞 support@hexamedhealthcare.com

☎ +91 95419 01145

☎ 1800 212 6547

🌐 www.hexamedhealthcare.com



Scan QR for Location



Scan QR for Enquiry Form